

Second-Line Treatment

| | | | | | | | | | |
|-------------------------------|-------|-------|-------|------|-------|------|------|-------|------|
| Int/Disc n= 3,240 (8.8%) | 11.1% | 7.6% | 8.3% | 4.8% | 4.5% | 4.9% | 2.6% | 56.2% | |
| AD; AP; MS n= 1,180 (3.2%) | 7.6% | 5.9% | 7.2% | 9.3% | 16.5% | 8.9% | | 43.6% | 0.8% |
| AP; MS n= 1,750 (4.7%) | 2.9% | 10.0% | 19.1% | 2.9% | 4.3% | | 7.7% | 50.3% | 2.9% |
| AD; MS n= 1,880 (5.1%) | 15.7% | 1.3% | 15.4% | 3.7% | | 2.7% | 6.4% | 51.6% | 3.2% |
| AD; AP n= 2,565 (6.9%) | 18.5% | 8.0% | 1.4% | | 3.5% | 2.3% | 5.1% | 58.1% | 3.1% |
| MS n= 3,170 (8.6%) | 9.9% | 7.3% | | 1.1% | 8.2% | 5.8% | 3.9% | 59.3% | 4.4% |
| AP n= 3,015 (8.2%) | 11.4% | | 7.8% | 7.1% | 1.3% | 6.5% | 3.8% | 57.4% | 4.6% |
| AD n= 3,540 (9.6%) | | 8.1% | 7.9% | 7.8% | 7.2% | 1.3% | 4.1% | 59.2% | 4.5% |

AD

AP

MS

AD; AP

AD; MS

AP; MS

AD; AP; MS

No Switch

Int/Disc

Third-Line Treatment

Percentage (%)

