

ON GENERATOR AND PARITY-CHECK POLYNOMIAL MATRICES OF GENERALIZED QUASI-CONSTACYCLIC CODES

Sümeysra Bedir

Yıldız Technical University
Department of Mathematics

Thesis Report-4
03.11.2017

Base Study:

This study is based on the following article;

- ▶ Matsui, H. "On Generator and Parity-check Polynomial Matrices of Generalized Quasi-cyclic Codes", Finite Fields and Their Applications 34 (2015):280-304.

In the cited work, a complete theory of generator polynomial matrices of GQC codes, including a relation formula between generator polynomial matrices and parity-check polynomial matrices through their equations, is provided. As the author noted; "Background knowledge of this paper is required only on linear codes, cyclic codes and basic polynomial arithmetic over finite fields."

We extended this work to the constacyclic case, namely; we showed that the facts and the theory for the quasi-cyclic codes obtained from cyclic components, also hold for quasi-codes obtained from constacyclic components. We are trying to prove a similar fact for quasi-cyclic codes obtained from pseudo-cyclic components.

Matsui shows that each GQC code obtained from l cyclic components, can be described by an upper triangular generator matrix $G = (g_{i,j} \in F_q[x])$ of the form

$$G = \begin{bmatrix} g_{1,1} & g_{1,2} & \cdots & g_{1,l} \\ 0 & g_{2,2} & \cdots & g_{2,l} \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \cdots & 0 & g_{l,l} \end{bmatrix}_{l \times l}$$

which satisfies the identical equation of G ;

$$AG = \text{diag}[x^{n_1} - 1, \dots, x^{n_l} - 1]$$

where $A = (a_{i,j})$ is another upper triangular $l \times l$ polynomial matrix. This identical equation generalizes a cyclic code's $ag = x^n - 1$ for its generator polynomial g , to the quasi-cyclic case.

Introduction

Overview

Definitions and
Algorithms

The Identical
Equation of G
The Duality Theorem

Our Studies

Application of the
Theory to the
Constacyclic Case

Future Studies

References

Further, he generalizes the well known fact $h = x^{\deg h}a(x^{-1})$ for the dual of a cyclic code to the dual of the quasi-cyclic code obtained from cyclic components (GQC). He shows that the generator polynomial matrix for the dual GQC code (which is the parity-check polynomial matrix for the GQC code) can be calculated from the matrix A .

Introduction

Overview

Definitions and
Algorithms

The Identical

Equation of G

The Duality Theorem

Our Studies

Application of the
Theory to the
Constacyclic Case

Future Studies

References

Generator Polynomial Matrix of GQC Codes

Definition

Let C be a GQC code, and let $G = (g_{i,j})$ be an $l \times l$ matrix whose entries are in $F_q[x]$ and whose rows are codewords of C . If $g_{i,j} = 0$ for all $1 \leq i, j \leq l$ with $i > j$, namely, G is of the form

$$G = \begin{bmatrix} g_{1,1} & g_{1,2} & \cdots & g_{1,l} \\ 0 & g_{2,2} & \cdots & g_{2,l} \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \cdots & 0 & g_{l,l} \end{bmatrix}_{l \times l}$$

and moreover, for all $1 \leq i \leq l$, $g_{i,i}$ has the minimum degree among all codewords of the form $(0, \dots, 0, c_i, \dots, c_l) \in C$ with $c_i \neq 0$, then we call G a **generator polynomial matrix** of C . If $g_{i,i}$ is monic for all $1 \leq i \leq l$ and G satisfies $\deg g_{i,j} < \deg g_{j,j}$ for all $1 \leq i \neq j \leq l$, then we say that G is **reduced**.

Introduction

Overview

Definitions and
Algorithms

The Identical
Equation of G
The Duality Theorem

Our Studies

Application of the
Theory to the
Constacyclic Case

Future Studies

References

Parity-check Polynomial Matrix of GQC Codes

Definition

Let C be a GQC code, and let $H = (h_{i,j})$ be an $l \times l$ matrix whose entries are in $F_q[x]$ and whose rows are codewords of C^\perp . If $h_{i,j} = 0$ for all $1 \leq i, j \leq l$ with $i < j$, namely, H is of the form

$$H = \begin{bmatrix} h_{1,1} & 0 & \cdots & 0 \\ h_{2,1} & h_{2,2} & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ h_{l,1} & h_{l,2} & \cdots & h_{l,l} \end{bmatrix}_{l \times l}$$

and moreover, for all $1 \leq i \leq l$, $h_{i,i}$ has the minimum degree among all codewords of the form $(c_1, \dots, c_i, 0, \dots, 0) \in C^\perp$ with $c_i \neq 0$, then we call H a **parity-check polynomial matrix** of C . If $h_{i,i}$ is monic for all $1 \leq i \leq l$ and H satisfies $\deg h_{i,j} < \deg h_{j,j}$ for all $1 \leq i \neq j \leq l$, then we say that H is **reduced**.

[Introduction](#)

[Overview](#)

[Definitions and Algorithms](#)

[The Identical Equation of \$G\$](#)
[The Duality Theorem](#)

[Our Studies](#)

[Application of the Theory to the Constacyclic Case](#)

[Future Studies](#)

[References](#)

Fact

For each GQC code, the reduced generator polynomial matrix is uniquely determined, and moreover, the reduced parity-check polynomial matrix is also uniquely determined. From any generator polynomial matrix and parity-check polynomial matrix, we can obtain the reduced ones by elementary row operations of polynomial matrices.

[Introduction](#)[Overview](#)[Definitions and Algorithms](#)[The Identical](#)[Equation of \$G\$](#) [The Duality Theorem](#)[Our Studies](#)[Application of the Theory to the Constacyclic Case](#)[Future Studies](#)[References](#)

Buchberger's Algorithm for GQC Codes

The algorithm for obtaining the reduced generator polynomial matrix from a generator matrix G of a GQC code is described as follows;

- ▶ We start with the polynomial representation

$$G' = \begin{bmatrix} c_{1,1} & c_{1,2} & \cdots & c_{1,l} \\ c_{2,1} & c_{2,2} & \cdots & c_{2,l} \\ \vdots & \ddots & \ddots & \vdots \\ c_{k,1} & \cdots & 0 & c_{k,l} \end{bmatrix}_{k \times l}$$

where $c_{i,j} \in F_q[x]$ for $1 \leq i \leq k$ and $1 \leq j \leq l$. Let c_i denote the i^{th} row of G' for $1 \leq i \leq k$. In this algorithm, the following manipulations of the polynomial matrix are carried out inductively.

Introduction

Overview

Definitions and Algorithms

The Identical

Equation of G

The Duality Theorem

Our Studies

Application of the Theory to the Constacyclic Case

Future Studies

References

Buchberger's Algorithm for GQC Codes

1. If $c_{1,1} = \dots = c_{k,1} = 0$, then set $c_1 = (x^{n_1} - 1, 0, \dots, 0)$ and stop. If $c_{1,1} \neq 0$ and $c_{2,1} = \dots = c_{k,1} = 0$, then stop.

After the above manipulations, $c_1 = (c_{1,1}, \dots, c_{1,l})$ is denoted by $g_1 = (g_{1,1}, \dots, g_{1,l})$ and then we have $g_{1,1} = \gcd(c_{1,1}, \dots, c_{k,1})$ from the initial matrix G' .

Introduction

Overview

Definitions and
Algorithms

The Identical
Equation of G

The Duality Theorem

Our Studies

Application of the
Theory to the
Constacyclic Case

Future Studies

References

Buchberger's Algorithm for GQC Codes

1. If $c_{1,1} = \dots = c_{k,1} = 0$, then set $c_1 = (x^{n_1} - 1, 0, \dots, 0)$ and stop. If $c_{1,1} \neq 0$ and $c_{2,1} = \dots = c_{k,1} = 0$, then stop.
2. By exchanging c_1 for another row of c_2, \dots, c_k if it is required, we can assume that $c_{1,1}$ has the minimum degree among nonzero $c_{1,1}, \dots, c_{k,1}$.

After the above manipulations, $c_1 = (c_{1,1}, \dots, c_{1,l})$ is denoted by $g_1 = (g_{1,1}, \dots, g_{1,l})$ and then we have $g_{1,1} = \gcd(c_{1,1}, \dots, c_{k,1})$ from the initial matrix G' .

Introduction

Overview

Definitions and
Algorithms

The Identical

Equation of G

The Duality Theorem

Our Studies

Application of the
Theory to the
Constacyclic Case

Future Studies

References

Buchberger's Algorithm for GQC Codes

1. If $c_{1,1} = \dots = c_{k,1} = 0$, then set $c_1 = (x^{n_1} - 1, 0, \dots, 0)$ and stop. If $c_{1,1} \neq 0$ and $c_{2,1} = \dots = c_{k,1} = 0$, then stop.
2. By exchanging c_1 for another row of c_2, \dots, c_k if it is required, we can assume that $c_{1,1}$ has the minimum degree among nonzero $c_{1,1}, \dots, c_{k,1}$.
3. Compute $p_i, r_i \in F_q[x]$ such that $c_{i,1} = p_i c_{1,1} + r_i$ with $\deg r_i < \deg c_{1,1}$ for all $2 \leq i \leq k$ and replace c_i with $c_i - p_i c_1$ for all $2 \leq i \leq k$, and go to step 1.

After the above manipulations, $c_1 = (c_{1,1}, \dots, c_{1,l})$ is denoted by $g_1 = (g_{1,1}, \dots, g_{1,l})$ and then we have $g_{1,1} = \gcd(c_{1,1}, \dots, c_{k,1})$ from the initial matrix G' .

Introduction

Overview

Definitions and
Algorithms

The Identical
Equation of G
The Duality Theorem

Our Studies

Application of the
Theory to the
Constacyclic Case

Future Studies

References

Buchberger's Algorithm for GQC Codes

Now, G' is converted to;

$$G'' = \begin{bmatrix} g_{1,1} & g_{1,2} & \cdots & g_{1,l} \\ 0 & c_{2,2} & \cdots & c_{2,l} \\ \vdots & \ddots & \ddots & \vdots \\ 0 & c_{k,2} & \cdots & c_{k,l} \end{bmatrix}_{k \times l}$$

where $c_{i,j}$ in G'' is generally unequal to $c_{i,j}$ in G' .

Next, we apply the above manipulation to the submatrix;

$$\begin{bmatrix} c_{2,2} & \cdots & c_{2,l} \\ \vdots & \ddots & \vdots \\ c_{k,2} & \cdots & c_{k,l} \end{bmatrix}$$

and continuing recursively we obtain the reduced form G .

The Identical Equation of G

Fact

As a consequence of the fact that upper triangular matrices over the quotient field of $F_q[x]$ form a group, the matrix A satisfying the equation

$$AG = \text{diag}[x^{n_1} - 1, \dots, x^{n_l} - 1]$$

is also an upper triangular matrix.

Introduction

Overview

Definitions and
Algorithms

The Identical
Equation of G
The Duality Theorem

Our Studies

Application of the
Theory to the
Constacyclic Case

Future Studies

References

The Duality Theorem

Theorem

Let $G = (g_{i,j})$ be the reduced generator polynomial matrix of a GQC code C , and let A be the polynomial matrix which satisfies $AG = \text{diag}[x^{n_1} - 1, \dots, x^{n_l} - 1]$. Then

$$H = \begin{bmatrix} x^{\deg a_{1,1}} a_{1,1}^{<n_1>} & 0 & \dots & 0 \\ x^{\deg a_{2,2}} a_{1,2}^{<n_1>} & x^{\deg a_{2,2}} a_{2,2}^{<n_2>} & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ x^{\deg a_{l,l}} a_{1,l}^{<n_1>} & x^{\deg a_{l,l}} a_{2,l}^{<n_2>} & \dots & x^{\deg a_{l,l}} a_{l,l}^{<n_l>} \end{bmatrix}_{l \times l}$$

where each $a_{i,j}^{<\omega>}$ is the polynomial with coefficient vector as the first row of transpose of the circulant matrix obtained from $a_{i,j}$, and each column i of H is considered modulo $x^{n_i} - 1$.

Introduction

Overview

Definitions and
Algorithms

The Identical
Equation of G
The Duality Theorem

Our Studies

Application of the
Theory to the
Constacyclic Case

Future Studies

References

Application of the Theory to the Constacyclic Case

The proof of the main theorem above was relying mainly on the well-known fact below;

Fact

$x^{n_i} - 1 \mid x^N - 1$ if and only if $n_i \mid N$.

In order to make use of this fact in the concept of constacyclic codes we prove the following corollary;

Corollary

$x^{n_i} - \alpha_i \mid x^N - 1$ if and only if
 $N = \text{lcm}(n_1, \dots, n_l) \cdot \text{lcm}(\text{ord}(\alpha_1), \dots, \text{ord}(\alpha_l))$, where $\alpha_i \in F_q$.

[Introduction](#)

[Overview](#)

[Definitions and Algorithms](#)

[The Identical](#)

[Equation of G](#)

[The Duality Theorem](#)

[Our Studies](#)

[Application of the Theory to the Constacyclic Case](#)

[Future Studies](#)

[References](#)

Proof.

We have

$$\begin{aligned} & (x^{n_i} - \alpha_i)(\alpha_i^{-1} + \alpha_i^{-2}x^{n_i} + \dots + \alpha_i^{-\text{ord}(\alpha_i)}x^{n_i\text{ord}(\alpha_i)}) \\ &= x^{n_i\text{ord}(\alpha_i)} - 1 \\ &\iff (x^{n_i} - \alpha_i) | (x^{n_i\text{ord}(\alpha_i)} - 1) \dots (*) \end{aligned}$$

We also have

$$\begin{aligned} & n_i\text{ord}(\alpha_i) | \text{lcm}(n_1, \dots, n_l) \cdot \text{lcm}(\text{ord}(\alpha_1), \dots, \text{ord}(\alpha_l)) \\ &\iff (x^{n_i\text{ord}(\alpha_i)} - 1) | (x^{\text{lcm}(n_1, \dots, n_l) \cdot \text{lcm}(\text{ord}(\alpha_1), \dots, \text{ord}(\alpha_l))} - 1) \dots (**) \end{aligned}$$

By (*) and (**),

$$\iff x^{n_i} - \alpha_i | x^N - 1$$



Application of the Theory to the Constacyclic Case

Another base concept to implement is the definition of $a_{i,j}^{<\omega>}$ and the modulo $x^\omega - 1$ from the duality theorem.

The implementations should consider the fact that we use constacyclic shift instead of cyclic shift.

So we define $a_{i,j}^{<\omega>}$ as follows, for simplicity we denote $a_{i,j}$ simply by a .

Definition

Let $a \in F_q[x]$ with $\deg a < \omega$ have the extended coefficient vector $(a_0, a_1, \dots, a_{\omega-1})$. The coefficient vector of $a^{<\omega>}$ is the first row of transpose of the α^{-1} - *twistulant* matrix of a .

Introduction

Overview

Definitions and

Algorithms

The Identical

Equation of G

The Duality Theorem

Our Studies

Application of the

Theory to the

Constacyclic Case

Future Studies

References

In this case; α^{-1} – *twistulant* matrix of a is

$$\begin{bmatrix} a_0 & a_1 & \cdots & a_{\omega-1} \\ \alpha^{-1}a_{\omega-1} & a_0 & \cdots & a_{\omega-2} \\ \alpha^{-1}a_{\omega-2} & \alpha^{-1}a_{\omega-1} & \ddots & \vdots \\ \vdots & \ddots & \ddots & \vdots \\ \alpha^{-1}a_1 & \cdots & \alpha^{-1}a_{\omega-1} & a_0 \end{bmatrix}$$

so $a^{<\omega>} = a_0 + \alpha^{-1}a_{\omega-1}x + \alpha^{-1}a_{\omega-2}x^2 + \cdots + \alpha^{-1}a_1x^{\omega-1}$.

Introduction

Overview

Definitions and
Algorithms

The Identical

Equation of G

The Duality Theorem

Our Studies

Application of the
Theory to the
Constacyclic Case

Future Studies

References

Application of the Theory to the Constacyclic Case

We also implemented the following fact to the constacyclic case;

$$AG = \text{diag}[x^{n_1} - 1, \dots, x^{n_l} - 1]$$

This time we should have

$$AG = \text{diag}[x^{n_1} - \alpha_1, \dots, x^{n_l} - \alpha_l]$$

where each constacyclic component i , is α_i - *constacyclic*.

The Modulo

When we look back at the parity-check matrix

$$H = \begin{bmatrix} x^{\deg a_{1,1}} a_{1,1}^{<n_1>} & 0 & \dots & 0 \\ x^{\deg a_{2,2}} a_{1,2}^{<n_1>} & x^{\deg a_{2,2}} a_{2,2}^{<n_2>} & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ x^{\deg a_{l,l}} a_{1,l}^{<n_1>} & x^{\deg a_{l,l}} a_{2,l}^{<n_2>} & \dots & x^{\deg a_{l,l}} a_{l,l}^{<n_l>} \end{bmatrix}_{l \times l}$$

the i^{th} column is considered modulo $x^{n_i} - 1$.

To implement this fact to the constacyclic case, we should be careful that we are talking about the dual code, so we consider each column i modulo $x^{n_i} - \alpha_i^{-1}$.

Introduction

Overview

Definitions and
Algorithms

The Identical
Equation of G
The Duality Theorem

Our Studies

Application of the
Theory to the
Constacyclic Case

Future Studies

References

The Duality Theorem for Constacyclic Case

Theorem

Let $G = (g_{i,j})$ be the reduced generator polynomial matrix of a generalized quazi constacyclic code C , and let A be the polynomial matrix which satisfies $AG = \text{diag}[x^{n_1} - \alpha_1, \dots, x^{n_l} - \alpha_l]$. Then

$$H = \begin{bmatrix} x^{\deg a_{1,1}} a_{1,1}^{<n_1>} & 0 & \dots & 0 \\ x^{\deg a_{2,2}} a_{1,2}^{<n_1>} & x^{\deg a_{2,2}} a_{2,2}^{<n_2>} & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ x^{\deg a_{l,l}} a_{1,l}^{<n_1>} & x^{\deg a_{l,l}} a_{2,1}^{<n_2>} & \dots & x^{\deg a_{l,l}} a_{l,l}^{<n_l>} \end{bmatrix}_{l \times l}$$

where each $a_{i,j}^{<\omega>}$ is the polynomial with coefficient vector as the first row of transpose of the α_i^{-1} - twistulant matrix obtained from $a_{i,j}$, and each column i of H is considered modulo $x^{n_i} - \alpha_i$.

Introduction

Overview

Definitions and
Algorithms

The Identical
Equation of G
The Duality Theorem

Our Studies


Application of the
Theory to the
Constacyclic Case

Future Studies


References


Future Studies


- ▶ We aim to find the necessary and sufficient implementations for the quasi-polycyclic case.


 A. Alahamdi, S. Dougherty, A. Leroy, P. Sole, (2016). On the duality and the direction of polycyclic codes, Adv. in Math. of Com., 10(4):921-929.


 A. Asamov, N. Aydın, (2007). The database for \mathbb{Z}_4 codes, <http://www.asamov.com/Z4Codes/>

 S. Jitman, (2013). Vector-circulant matrices over finite fields and related codes, arXiv:1408.2059 [math.RA]

 S.R. Lopez-Permouth, B.R. Parra-Avila, S. Szabo, (2009). Dual generalizations of the concept of cyclicity of codes, Adv. in Math. of Com. (2009) 227–234.

 William Wesley Peterson, E. J. Jr Weldon, Error Correcting codes: second edition, MIT Press (1972).

 D. Radkova, A.J. Van Zanten, Constacyclic codes as invariant subspaces, Linear Algebra and its Applications, 430(2009), 855-864.

 M. Wu, Free cyclic codes as invariant submodules over finite chain rings, International Mathematical Forum, 8-37(2013), 1835 - 1838.

 V. Gritsenko, A. Maevskiy, (2014). On a construction of optimal codes in term rank metric via $p(x)$ -circulants. Fourteenth