

ELEC 442 - Introduction to Robotics

Assignment 1

Sondre Myrberg
sondrmyr@alumni.ubc.ca
81113433



THE UNIVERSITY
OF BRITISH COLUMBIA

1

Given the homogenous transformation

$$\begin{bmatrix} \mathbf{y} \\ 1 \end{bmatrix} = \underbrace{\begin{bmatrix} Q & \mathbf{d} \\ \mathbf{0}^\top & 1 \end{bmatrix}}_T \begin{bmatrix} \mathbf{x} \\ 1 \end{bmatrix}$$

where Q and \mathbf{d} accounts for rotation and translation, respectively. We have that the inverse is on the form

$$T^{-1} = \begin{bmatrix} \tilde{Q} & \tilde{\mathbf{d}} \\ \mathbf{0}^\top & 1 \end{bmatrix}$$

where we know that $T^{-1}T$ is equal to the 4×4 identity matrix. This yields

$$\begin{aligned} T^{-1}T &= \begin{bmatrix} \tilde{Q} & \tilde{\mathbf{d}} \\ \mathbf{0}^\top & 1 \end{bmatrix} \begin{bmatrix} Q & \mathbf{d} \\ \mathbf{0}^\top & 1 \end{bmatrix} \\ &= \begin{bmatrix} \tilde{Q}Q & \tilde{Q}\mathbf{d} + \tilde{\mathbf{d}} \\ \mathbf{0}^\top & 1 \end{bmatrix} = \mathbf{I}_{4 \times 4} \\ &\Rightarrow \begin{cases} \tilde{Q}Q &= \mathbf{I}_{3 \times 3} \\ \tilde{Q}\mathbf{d} + \tilde{\mathbf{d}} &= \mathbf{0} \end{cases} \\ &\Rightarrow \begin{cases} \tilde{Q} &= Q^{-1} = Q^\top \\ \tilde{\mathbf{d}} &= -\tilde{Q}\mathbf{d} = -Q^\top \mathbf{d} \end{cases} \\ &\Rightarrow T^{-1} = \underline{\begin{bmatrix} Q^\top & -Q^\top \mathbf{d} \\ \mathbf{0} & 1 \end{bmatrix}} \end{aligned}$$

2

Considering the homogenous transformation matrix

$${}^0T_1 = \begin{bmatrix} Q & \mathbf{d} \\ \mathbf{0}^\top & 1 \end{bmatrix}$$

with

$$Q = \underbrace{\begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ 0 & 0 & 1 \end{bmatrix}}_{Q_1} \underbrace{\begin{bmatrix} 1 & 0 & 0 \\ 0 & -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ 0 & \frac{\sqrt{3}}{2} & -\frac{1}{2} \end{bmatrix}}_{Q_2}$$

and

$$\mathbf{d} = \begin{bmatrix} -\frac{5}{\sqrt{2}} \\ \frac{5}{\sqrt{2}} \\ 4 \end{bmatrix} \text{ cm}$$