

CARDIFF UNIVERSITY

EXAMINATION PAPER

Academic Year: 2012/2013
Examination Period AUTUMN
Examination Paper Number: BST169
Examination Paper Title: ECONOMETRICS
Duration: 3 HOURS

Do not turn this page over until instructed to do so by the Senior Invigilator

Structure of Examination Paper:

There are 3 pages.

There are 6 questions in total.

There are no appendices

The maximum mark for the examination paper is 100%. The percentage for each question is given in parentheses.

Students to be provided with:

Answer book
Statistical Tables

Instructions to Students:

All questions must be answered.

The use of non-electronic translation dictionaries between English or Welsh and a foreign language bearing an appropriate departmental stamp is permitted in this examination.

Please turn over

- 1) For the linear regression model:
 $y = \alpha \mathbf{1} + X\beta + \varepsilon$
 where $y = (y_1, y_2, \dots, y_N)'$, $\mathbf{1}$ is an $N \times 1$ vector of ones, $\beta = (\beta_1, \beta_2, \dots, \beta_K)'$ and X is an $N \times K$ matrix of explanatory variables.
 - a) Find separately the OLS estimators of $\hat{\alpha}$ and $\hat{\beta}$. (6)
 - b) Calculate the statistic to test $H_0: \beta = 0$ under homoskedasticity (6)
 - c) Suppose $\varepsilon|X \sim N(0, \sigma^2 I_N)$, where I_N is the identity matrix of dimension N . Show the distribution of the test statistic for the previous question. (6)

- 2) Explain the principle of maximum likelihood estimation (MLE). (4) Are the estimators obtained from MLE unbiased? (4) How can hypothesis tests be implemented? (8)

- 3) For the linear regression model,

$$y_i = \alpha + \beta x_i + \varepsilon_i, \quad i = 1, 2, \dots, N,$$

suppose x_i is a scalar and $\{(x_i, \varepsilon_i)\}$ is an i.i.d. sequence with $E(\varepsilon_i | x_i) = 0$. y_i is in fact generated by the above model.

 - a) How do you calculate the coefficient of determination for the model? Why is it also called R-squared? What does it measure? (5)
 - b) If N is fixed, what is the effect of including irrelevant regressors on R-squared? What is the maximum value of R-squared one can obtain by including more regressors and why? (5)
 - c) Suppose the variances of x_i and ε_i respectively are σ_x^2 and σ^2 , which are finite. What is the probability limit of R-squared when N tends to infinity? (6)

- 4) Describe the situations under which we cannot include some of the important explanatory variables into our linear regression model. (3) What would be the potential problems? (3) How can we remedy such problems? (Show the exact estimation procedure.) (10)

- 5) For a sample of 506 communities in the Boston area of the United States, we estimate a model relating median housing price (price) in the community to various community characteristics: nox is the amount of nitrous oxide in the air, in parts per million; dist is a weighted distance of the community from five employment centers, in miles; rooms is the average number of rooms in houses in the community; and stratio is the average student-teacher ratio of schools in the community. The population model is:

$$\log(\text{price}) = 11.084 - 0.9535 \log(\text{nox}) - 0.1343 \log(\text{dist}) + 0.2545 \text{rooms} - 0.05245 \text{stratio} + \varepsilon$$

The moments for the error term ε are: $E(\varepsilon) = 0$ and $\text{Var}(\varepsilon) = \sigma^2$. Suppose Gauss-Markov assumptions are satisfied for the model.

- a) How will you interpret the estimated slope coefficients for $\log(\text{dist})$ and rooms? (4)
 - b) If a house has one more room while other house characteristics remain unchanged, what is the exact percentage change of the house price? (5)
 - c) For the exact house price percentage change estimator in the previous question, is it unbiased? Is it consistent? Why? (4)
 - d) Applying the Jarque-Bera test to the estimated residuals gives a test statistic of 480.1425. Given that $\sum_{i=1}^{506} \exp(\varepsilon_i) = 524.53$ and $\widehat{\sigma^2} = 0.1$, how would you predict the price of the house with the following characteristics: nox = 5.54, dist = 3.21, rooms = 6 and stratio = 19? (5)
- 6) Suppose $\{Z_i\}$ is a sequence of uncorrelated random variables from the same distribution,
- a) What are the conditions for $\frac{1}{N} \sum_{i=1}^N Z_i$ to converge in probability to μ , where $\mu = E(Z_i)$? (6)
 - b) Can you use the results from the previous question to state the conditions under which the OLS estimators in Question 1 of this paper are consistent? Prove it. (10)

Table 1



[illegible]

Table 3

T Distribution



Example:

$$P(t_{30} \leq 1.697) = .95$$


$$P(t_{30} > 1.697) = .05$$

Percentiles of the *t*-distribution

<i>df</i> \ <i>p</i>	.60	.70	.80	.90	.95	.975	.99	.995
1	.325	.727	1.367	3.078	6.314	12.706	31.821	63.657
2	.289	.617	1.061	1.886	2.920	4.303	6.965	9.925
3	.277	.584	.978	1.638	2.353	3.182	4.541	5.841
4	.271	.569	.941	1.533	2.132	2.776	3.747	4.604
5	.267	.559	.920	1.476	2.015	2.571	3.365	4.032
6	.265	.553	.906	1.440	1.943	2.447	3.143	3.707
7	.263	.549	.896	1.415	1.895	2.365	2.998	3.499
8	.262	.546	.889	1.397	1.860	2.306	2.896	3.355
9	.261	.543	.883	1.383	1.833	2.262	2.821	3.250
10	.260	.542	.879	1.372	1.812	2.228	2.764	3.169
11	.260	.540	.876	1.363	1.796	2.201	2.718	3.106
12	.259	.539	.873	1.356	1.782	2.179	2.681	3.055
13	.259	.538	.870	1.350	1.771	2.160	2.650	3.012
14	.258	.537	.868	1.345	1.761	2.145	2.624	2.977
15	.258	.536	.866	1.341	1.753	2.131	2.602	2.947
16	.258	.535	.865	1.337	1.746	2.120	2.583	2.921
17	.257	.534	.863	1.333	1.740	2.110	2.567	2.898
18	.257	.534	.862	1.330	1.734	2.101	2.552	2.878
19	.257	.533	.861	1.328	1.729	2.093	2.539	2.861
20	.257	.533	.860	1.325	1.725	2.086	2.528	2.845
21	.257	.532	.859	1.323	1.721	2.080	2.518	2.831
22	.256	.532	.858	1.321	1.717	2.074	2.508	2.819
23	.256	.532	.858	1.319	1.714	2.069	2.500	2.807
24	.256	.531	.857	1.316	1.708	2.060	2.485	2.787
25	.256	.531	.856	1.316	1.708	2.060	2.485	2.787
26	.256	.531	.856	1.315	1.706	2.056	2.479	2.779
27	.256	.531	.855	1.314	1.703	2.052	2.473	2.771
28	.256	.530	.855	1.313	1.701	2.048	2.467	2.763
29	.256	.530	.854	1.310	1.697	2.042	2.457	2.750
30	.256	.530	.854	1.310	1.697	2.042	2.457	2.750
infinity	.253	.524	.842	1.282	1.645	1.960	2.326	2.576

Chi-square Distribution

Table 4



Example:

$$P(\chi^2_4 < 9.488) = .95$$


$$P(\chi^2_4 > 9.488) = .05$$

Percentiles of the Chi-square distribution

df/p	.005	.01	.025	.05	.10	.90	.95	.975	.99	.995
1	.00004	.00016	.00098	.0039	.0158	2.71	3.84	5.02	6.63	7.88
2	.0100	.0201	.0506	.1026	.2107	4.61	5.99	7.38	9.21	10.60
3	.0717	.115	.216	.352	.584	6.25	7.81	9.35	11.34	12.84
4	.207	.297	.484	.711	1.064	7.78	9.49	11.14	13.28	14.86
5	.412	.554	.831	1.15	1.61	9.24	11.07	12.83	15.09	16.75
6	.676	.872	1.24	1.64	2.20	10.64	12.59	14.45	16.81	18.55
7	.989	1.24	1.69	2.17	2.83	12.02	14.07	16.01	18.48	20.28
8	1.34	1.65	2.18	2.73	3.49	13.36	15.51	17.53	20.09	21.96
9	1.73	2.09	2.70	3.33	4.17	14.68	16.92	19.02	21.67	23.59
10	2.16	2.56	3.25	3.94	4.87	15.99	18.31	20.48	23.21	25.19
11	2.60	3.05	3.82	4.57	5.58	17.28	19.68	21.92	24.73	26.76
12	3.07	3.57	4.40	5.23	6.30	18.55	21.03	23.34	26.22	28.30
13	3.57	4.11	5.01	5.89	7.04	19.81	22.36	24.74	27.69	29.82
14	4.07	4.66	5.63	6.57	7.79	21.06	23.68	26.12	29.14	31.32
15	4.6	5.23	6.26	7.26	8.55	22.31	25	27.49	30.58	32.80
16	5.14	5.81	6.91	7.96	9.31	23.54	26.30	28.85	32.00	34.27
18	6.26	7.01	8.23	9.39	10.86	25.99	28.87	31.53	34.81	37.16
20	7.43	8.26	9.59	10.85	12.44	28.41	31.41	34.17	37.57	40.00
24	9.89	10.86	12.40	13.85	15.66	33.20	36.42	39.36	42.98	45.56
30	13.79	14.95	16.79	18.49	20.60	40.26	43.77	46.98	50.89	53.67
40	20.71	22.16	24.43	26.51	29.05	51.81	55.76	59.34	63.69	66.77
60	35.53	37.48	40.48	43.19	46.46	74.40	79.08	83.30	88.38	91.95
120	83.85	86.92	91.58	95.70	100.62	140.23	146.57	152.21	158.95	163.64

Table 5

F-Distribution



Example:

$$P(F_{10,20} < 2.47) = .95$$

$$P(F_{10,20} > 2.47) = .05$$

95th Percentile for the F-distribution

(df1 < 11)

	df ₁ =1	2	3	4	5	6	7	8	9	10
df ₂ =1	161.4476	199.5000	215.7073	224.5832	230.1619	233.9860	236.7684	238.8827	240.5433	241.8817
2	18.5128	19.0000	19.1643	19.2468	19.2964	19.3295	19.3532	19.3710	19.3848	19.3959
3	10.1280	9.5521	9.2766	9.1172	9.0135	8.9406	8.8867	8.8452	8.8123	8.7855
4	7.7086	6.9443	6.5914	6.3882	6.2561	6.1631	6.0942	6.0410	5.9988	5.9644
5	6.6079	5.7861	5.4095	5.1922	5.0503	4.9503	4.8759	4.8183	4.7725	4.7351
6	5.9874	5.1433	4.7571	4.5337	4.3874	4.2839	4.2067	4.1468	4.0990	4.0600
7	5.5914	4.7374	4.3468	4.1203	3.9715	3.8660	3.7870	3.7257	3.6767	3.6365
8	5.3177	4.4590	4.0662	3.8379	3.6875	3.5806	3.5005	3.4381	3.3881	3.3472
9	5.1174	4.2565	3.8625	3.6331	3.4817	3.3738	3.2927	3.2296	3.1789	3.1373
10	4.9646	4.1028	3.7083	3.4780	3.3258	3.2172	3.1355	3.0717	3.0204	2.9782
11	4.8443	3.9823	3.5874	3.3567	3.2039	3.0946	3.0123	2.9480	2.8962	2.8536
12	4.7472	3.8853	3.4903	3.2592	3.1059	2.9961	2.9134	2.8486	2.7964	2.7534
13	4.6672	3.8056	3.4105	3.1791	3.0254	2.9153	2.8321	2.7669	2.7144	2.6710
14	4.6001	3.7389	3.3439	3.1122	2.9582	2.8477	2.7642	2.6987	2.6458	2.6022
15	4.5431	3.6823	3.2874	3.0556	2.9013	2.7905	2.7066	2.6408	2.5876	2.5437
16	4.4940	3.6337	3.2389	3.0069	2.8524	2.7413	2.6572	2.5911	2.5377	2.4935
17	4.4513	3.5915	3.1968	2.9647	2.8100	2.6987	2.6143	2.5480	2.4943	2.4499
18	4.4139	3.5546	3.1599	2.9277	2.7729	2.6613	2.5767	2.5102	2.4563	2.4117
19	4.3807	3.5219	3.1274	2.8951	2.7401	2.6283	2.5435	2.4768	2.4227	2.3779
20	4.3512	3.4928	3.0984	2.8661	2.7109	2.5990	2.5140	2.4471	2.3928	2.3479
21	4.3248	3.4668	3.0725	2.8401	2.6848	2.5727	2.4876	2.4205	2.3660	2.3210
22	4.3009	3.4434	3.0491	2.8167	2.6613	2.5491	2.4638	2.3965	2.3419	2.2967
23	4.2793	3.4221	3.0280	2.7955	2.6400	2.5277	2.4422	2.3748	2.3201	2.2747
24	4.2597	3.4028	3.0088	2.7763	2.6207	2.5082	2.4226	2.3551	2.3002	2.2547
25	4.2417	3.3852	2.9912	2.7587	2.6030	2.4904	2.4047	2.3371	2.2821	2.2365
26	4.2252	3.3690	2.9752	2.7426	2.5868	2.4741	2.3883	2.3205	2.2655	2.2197
27	4.2100	3.3541	2.9604	2.7278	2.5719	2.4591	2.3732	2.3053	2.2501	2.2043
28	4.1960	3.3404	2.9467	2.7141	2.5581	2.4453	2.3593	2.2913	2.2360	2.1900
29	4.1830	3.3277	2.9340	2.7014	2.5454	2.4324	2.3463	2.2783	2.2229	2.1768
30	4.1709	3.3158	2.9223	2.6896	2.5336	2.4205	2.3343	2.2662	2.2107	2.1646
40	4.0847	3.2317	2.8387	2.6060	2.4495	2.3359	2.2490	2.1802	2.1240	2.0772
60	4.0012	3.1504	2.7581	2.5252	2.3683	2.2541	2.1665	2.0970	2.0401	1.9926
120	3.9201	3.0718	2.6802	2.4472	2.2899	2.1750	2.0868	2.0164	1.9588	1.9105
∞	3.8415	2.9957	2.6049	2.3719	2.2141	2.0986	2.0096	1.9384	1.8799	1.8307

(df1 > 11)

	Df ₁ =12	15	20	24	30	40	60	120	∞
df ₂ =1	243.9060	245.9499	248.0131	249.0518	250.0951	251.1432	252.1957	253.2529	254.3144
2	19.4125	19.4291	19.4458	19.4541	19.4624	19.4707	19.4791	19.4874	19.4957
3	8.7446	8.7029	8.6602	8.6385	8.6166	8.5944	8.5720	8.5494	8.5264
4	5.9117	5.8578	5.8025	5.7744	5.7459	5.7170	5.6877	5.6581	5.6281
5	4.6777	4.6188	4.5581	4.5272	4.4957	4.4638	4.4314	4.3985	4.3650
6	3.9999	3.9381	3.8742	3.8415	3.8082	3.7743	3.7398	3.7047	3.6689
7	3.5747	3.5107	3.4445	3.4105	3.3758	3.3404	3.3043	3.2674	3.2298
8	3.2839	3.2184	3.1503	3.1152	3.0794	3.0428	3.0053	2.9669	2.9276
9	3.0729	3.0061	2.9365	2.9005	2.8637	2.8259	2.7872	2.7475	2.7067
10	2.9130	2.8450	2.7740	2.7372	2.6996	2.6609	2.6211	2.5801	2.5379
11	2.7876	2.7186	2.6464	2.6090	2.5705	2.5309	2.4901	2.4480	2.4045
12	2.6866	2.6169	2.5436	2.5055	2.4663	2.4259	2.3842	2.3410	2.2962
13	2.6037	2.5331	2.4589	2.4202	2.3803	2.3392	2.2966	2.2524	2.2064
14	2.5342	2.4630	2.3879	2.3487	2.3082	2.2664	2.2229	2.1778	2.1307
15	2.4753	2.4034	2.3275	2.2878	2.2468	2.2043	2.1601	2.1141	2.0658
16	2.4247	2.3522	2.2756	2.2354	2.1938	2.1507	2.1058	2.0589	2.0096
17	2.3807	2.3077	2.2304	2.1898	2.1477	2.1040	2.0584	2.0107	1.9604
18	2.3421	2.2686	2.1906	2.1497	2.1071	2.0629	2.0166	1.9681	1.9168
19	2.3080	2.2341	2.1555	2.1141	2.0712	2.0264	1.9795	1.9302	1.8780
20	2.2776	2.2033	2.1242	2.0825	2.0391	1.9938	1.9464	1.8963	1.8432
21	2.2504	2.1757	2.0960	2.0540	2.0102	1.9645	1.9165	1.8657	1.8117
22	2.2258	2.1508	2.0707	2.0283	1.9842	1.9380	1.8894	1.8380	1.7831
23	2.2036	2.1282	2.0476	2.0050	1.9605	1.9139	1.8648	1.8128	1.7570
24	2.1834	2.1077	2.0267	1.9838	1.9390	1.8920	1.8424	1.7896	1.7330
25	2.1649	2.0889	2.0075	1.9643	1.9192	1.8718	1.8217	1.7684	1.7110
30	2.0921	2.0148	1.9317	1.8874	1.8409	1.7918	1.7396	1.6835	1.6223
40	2.0035	1.9245	1.8389	1.7929	1.7444	1.6928	1.6373	1.5766	1.5089
60	1.9174	1.8364	1.7480	1.7001	1.6491	1.5943	1.5343	1.4673	1.3893
120	1.8337	1.7505	1.6587	1.6084	1.5543	1.4952	1.4290	1.3519	1.2539
∞	1.7522	1.6664	1.5705	1.5173	1.4591	1.3940	1.3180	1.2214	1.0000

Table 6

Durbin - Watson Statistic

FIVE PERCENT SIGNIFICANCE POINTS OF d_i AND d_u FOR DURBIN-WATSON TEST†

N	k = 1		k = 2		k = 3		k = 4		k = 5	
	d_i	d_u	d_i	d_u	d_i	d_u	d_i	d_u	d_i	d_u
15	1.08	1.36	.95	1.54	.82	1.75	.69	1.97	.56	2.21
16	1.10	1.37	.98	1.54	.86	1.73	.74	1.93	.62	2.15
17	1.13	1.38	1.02	1.54	.90	1.71	.78	1.90	.67	2.10
18	1.16	1.39	1.05	1.53	.93	1.69	.82	1.87	.71	2.06
19	1.18	1.40	1.08	1.53	.97	1.68	.86	1.85	.75	2.02
20	1.20	1.41	1.10	1.54	1.00	1.66	.90	1.83	.79	1.99
21	1.22	1.42	1.13	1.54	1.03	1.67	.93	1.81	.83	1.98
22	1.24	1.43	1.15	1.54	1.05	1.66	.96	1.80	.86	1.94
23	1.26	1.44	1.17	1.54	1.08	1.66	.99	1.79	.90	1.92
24	1.27	1.45	1.19	1.55	1.10	1.66	1.01	1.78	.93	1.90
25	1.29	1.45	1.21	1.55	1.12	1.66	1.04	1.77	.95	1.89
26	1.30	1.46	1.22	1.55	1.14	1.66	1.06	1.76	.98	1.88
27	1.32	1.47	1.24	1.56	1.16	1.65	1.08	1.75	1.01	1.86
28	1.33	1.48	1.26	1.56	1.18	1.65	1.10	1.75	1.03	1.85
29	1.34	1.48	1.27	1.56	1.20	1.65	1.12	1.74	1.05	1.84
30	1.35	1.49	1.28	1.57	1.21	1.65	1.14	1.74	1.07	1.83
31	1.36	1.50	1.30	1.57	1.23	1.65	1.16	1.74	1.09	1.83
32	1.37	1.50	1.31	1.57	1.24	1.65	1.18	1.73	1.11	1.82
33	1.38	1.51	1.32	1.58	1.26	1.65	1.19	1.73	1.13	1.81
34	1.39	1.51	1.33	1.58	1.27	1.65	1.21	1.73	1.15	1.81
35	1.40	1.52	1.34	1.53	1.28	1.65	1.22	1.73	1.16	1.80
36	1.41	1.52	1.35	1.59	1.29	1.65	1.24	1.73	1.18	1.80
37	1.42	1.53	1.36	1.59	1.31	1.66	1.25	1.72	1.19	1.80
38	1.43	1.54	1.37	1.59	1.32	1.66	1.26	1.72	1.21	1.79
39	1.43	1.54	1.38	1.60	1.33	1.66	1.27	1.72	1.22	1.79
40	1.44	1.54	1.39	1.60	1.34	1.66	1.29	1.72	1.23	1.79
45	1.48	1.57	1.43	1.62	1.38	1.67	1.34	1.72	1.29	1.78
50	1.50	1.59	1.46	1.63	1.42	1.67	1.38	1.72	1.34	1.77
55	1.53	1.60	1.49	1.64	1.45	1.68	1.41	1.72	1.38	1.77
60	1.55	1.62	1.51	1.65	1.48	1.69	1.44	1.73	1.41	1.77
65	1.57	1.63	1.54	1.66	1.50	1.70	1.47	1.73	1.44	1.77
70	1.58	1.64	1.55	1.67	1.52	1.70	1.49	1.74	1.46	1.77
75	1.60	1.65	1.57	1.68	1.54	1.71	1.51	1.74	1.49	1.77
80	1.61	1.66	1.59	1.69	1.56	1.72	1.53	1.74	1.51	1.77
85	1.62	1.67	1.60	1.70	1.57	1.72	1.55	1.75	1.52	1.77
90	1.63	1.68	1.61	1.70	1.59	1.73	1.57	1.75	1.54	1.78
95	1.64	1.69	1.62	1.71	1.60	1.73	1.58	1.75	1.56	1.78
100	1.65	1.69	1.63	1.72	1.61	1.74	1.59	1.76	1.57	1.78

† N = number of observations; k = number of explanatory variables (excluding the constant term)