

Machine Learning in Genomics: Containerised
tutorials demonstrating best practises, pitfalls,
and reproducibility

Sach Nehal

2024-08-07

Contents

About	5
I Introduction	7
1 Epigenetic Data	9
1.1 What is epigenetic data?	9
1.2 What does epigenetic data look like?	12
1.3 Sources of epigenetic data	15
1.4 UCSC'S Genome Browser	16
1.5 Handling bigWig files	16
1.6 Data loaders and pre-processing	17
1.7 Dealing with missing data (oversampling, undersampling, weight- ing)	17
II Training models with DNA input	19
2 Loss functions, and peak metrics	21
3 Base pair averaging	23
4 Training tricks	25
5 Choosing which genomic regions to train on	27
6 Effect of differences in sequencing depths	29
7 Reproducibility of machine learning models	31
7.1 Seeding	31
7.2 Dashboarding	31
8 Testing	33

III Software libraries for model building	35
9 gReLU	37
10 Kipoi	39
11 Weights and Biases	41
IV ML pitfalls in genomics	43
12 Pitfalls overview	45
12.1 Distributional differences	45
12.2 Dependent examples	45
12.3 Confounding	45
12.4 Leaky pre-processing	45
12.5 Unbalanced classes	45
12.6 Balancing the proportion of peaks / no-peaks in validation sets .	45
V Model interpretability	47
13 Creating and visualising a simple model	49
14 TF mo-Disco	51
VI Using existing models	53
15 Using the gReLU model zoo	55
16 Fine tuning of Enformer	57
VII Predicting in novel cell types	59
17 Incorporating ATAC-seq info	61
18 Use of cell type averages	63
VIII More complex models	65
19 Training multi-headed models	67
20 Training siamese twin models	69

About

Applied machine learning utilising vast amounts of data has aided in pattern identification, predictive analytics, and solving complex problems across a multitude of fields. Solving these complex problems within these fields, researchers would find differing answers to the following questions; **what machine learning techniques can we apply to the problem, how do we apply the techniques in the context of this field, and why do we need to apply them in this way?** In any case, applied machine learning requires an interdisciplinary understanding of computing techniques and the field in question.

The aim of this project is to provide you with **a set of reproducible, containerized tutorials that include all necessary data, code, and descriptions to replicate key results, along with demonstrations of common pitfalls, in the field of genomics.** It is designed for users with knowledge of machine learning but little or no background in biology as a process to learn about applying machine learning techniques in genomics.

Part I

Introduction

Chapter 1

Epigenetic Data

1.1 What is epigenetic data?

As you may already know, typically all of the cells in your body contain the same DNA. How, then, do we have different cell types in our body? Your DNA contains a script that is able to produce the proteins required for each specific cell in your body. Which proteins, and subsequently which cells are made, depends on gene expression and regulation, i.e. “the way each cell deploys its genome.”¹

Epigenetic data arises from “the study of heritable and stable changes in gene expression that occur through alterations in the chromosome rather than in the DNA sequence.”²

¹Ralston and Shaw [2008]

²Al-Aboud et al. [2023]



commonfund.nih.gov

The image above shows quite simply the basics of genetic structures. Several more complex processes are involved during cell replication such as DNA transcription and translation in order to make proteins. A key takeaway in coming closer to understanding gene expression is that **Chromatin** is a complex structure made up of DNA wound around histone proteins, with some segments of DNA being accessible/inaccessible to further processes. **Euchromatin** refers to the accessible state, while **Heterochromatin** refers to a chromatin state in which DNA cannot be transcribed (inaccessible).³ There are many different epigenetic modifications that affect chromatin accessibility.

Some common epigenetic modifications include:

1. **DNA Methylation:** Addition of methyl groups to DNA, affecting gene expression regulation⁴.
2. **Histone Modifications:** Chemical changes to histone proteins that DNA wraps around, including acetylation, methylation, or phosphorylation. These changes influence chromatin structure and gene accessibility.⁵
3. **Chromatin Accessibility:** Regions of open chromatin that are accessible to transcription factors (special types of proteins that bind to DNA sequences and regulate gene expression) further dictate which regions of DNA can be expressed⁶.

³Shahid et al. [2023]

⁴Al-Aboud et al. [2023]

⁵T. [2007]

⁶Kappelmann-Fenzl [2021]

In studying gene expression and epigenetic modifications, we aim to more closely understand biological mechanisms that regulate development, disease, and how cells respond to epigenetic factors.

1.1.1 What Does DNA Look Like?

As illustrated in the image above, DNA is structured as a double helix, with two complementary strands intertwined to form the characteristic helical shape. DNA consists of an extremely long sequence composed of four types of nucleotides: Adenine (A), Cytosine (C), Thymine (T), and Guanine (G).

According to the National Cancer Institute (USA), nucleotides within the DNA double helix form complementary pairs—Adenine pairs with Thymine, and Guanine pairs with Cytosine⁷. These pairs are commonly referred to as base pairs. For example, if one strand of the double helix has the sequence “ATCGG”, the complementary strand will have the sequence “TAGCC”.

Genes are sequences of DNA located at specific positions on chromosomes and can vary in length. Each gene encodes information necessary for producing proteins or RNA molecules, which are essential for the structure, function, and regulation of an organism⁸. The complete set of genetic material in an organism is known as its genome.




Image highlighting part of a dna sequence and base pairs.⁹

1.1.2 Common Epigenetic Sequencing Techniques:

1. **ATAC-Seq** (Assay for Transposase-Accessible Chromatin with Sequencing): oMeasures chromatin accessibility to identify open regions of the genome where transcription factors can bind.
oOutput: Peaks indicating accessible chromatin regions.
2. **ChIP-Seq** (Chromatin Immunoprecipitation Sequencing): oUsed to identify DNA regions bound by specific proteins (e.g., transcription factors, histones with specific modifications).

⁷Board

⁸Board

⁹Kanani and Padole [2020]

oOutput: Peaks indicating binding sites or modification locations.

1.2 What does epigenetic data look like?

Epigenetic data can be represented in various forms, depending on the type of modification being studied and the methods used to gather the data. **ATAC-Seq** and **ChIP-Seq** are the common methods I will focus on, but there are others that may produce different forms of data, such as WGS (whole-genome sequencing) which produces nucleotide sequencing data, or Bisulfite conversion of DNA producing data on methylation levels across the genome.

1.2.1 Representing epigenetic data




Image showing the sequencing pipeline from high-throughput sequencing methods¹⁰.

1. **Raw Sequence Reads:** oThese are the basic output of sequencing experiments, such as those from ChIP-Seq or ATAC-Seq. oReads are processed and aligned to a reference genome before undergoing peak calling.

Lets look at what a few lines of raw sequence read data consists of: The data is taken from Encode Experiment ENCSR817LUF (chIP-Seq). The accession ID

¹⁰ Akalin [2020]

of the raw sequence read data is ENCFF397NRK. Genomic data comes in many file formats. This specific raw sequence read data is a compressed FASTQ file.

```
import gzip
from Bio import SeqIO
import requests
import io

# Define the URL of the FASTQ file
url = "https://www.encodeproject.org/files/ENCFF397NRK/@download/ENCFF397NRK.fastq.gz"

# Stream the file from the URL
response = requests.get(url, stream=True)
if response.status_code == 200:
    # Use BytesIO to handle the file content in memory
    with io.BytesIO(response.content) as file_content:
        with gzip.open(file_content, "rt") as handle:
            for i, record in enumerate(SeqIO.parse(handle, "fastq")):
                if i >= 3:
                    break
                print(f"ID: {record.id}")
                print(f"Sequence: {record.seq}")
                print(f"Quality Scores: {record.letter_annotations['phred_quality']}")
                print() # Newline for better readability
else:
    print(f"Failed to retrieve the file: Status code {response.status_code}")

## ID: B091JABXX110402:1:2204:12975:184709
## Sequence: GTTAGGGTTAGGGTTAGGGTTAGGGTTAGGGTTAGG
## Quality Scores: [31, 30, 30, 32, 36, 37, 36, 32, 33, 32, 35, 36, 37, 33, 33, 34, 37, 37, 36, 3
##
## ID: B091JABXX110402:1:2205:18641:8399
## Sequence: GGTTAGGGTTAGGGTTAGGGTTAGGGTTAGGGTTAG
## Quality Scores: [22, 27, 31, 30, 30, 33, 31, 32, 31, 31, 24, 36, 37, 36, 33, 34, 33, 37, 37, 3
##
## ID: B091JABXX110402:1:1207:12202:100922
## Sequence: AGGGTTAGGGTTAGGGTTAGGGTTAGGGTTAGGGTT
## Quality Scores: [33, 33, 36, 37, 31, 34, 34, 36, 37, 36, 29, 33, 35, 36, 39, 37, 32, 33, 35, 3
```


As you can see, each data entry is a DNA sequence (read). While I'm only showing the first three entries, for each experiment there are millions of sequencing reads. The quality scores indicate the confidence of each base call in the sequence. Higher scores suggest higher confidence. The scores are in Phred format, where a score of 20 corresponds to a 99% base call accuracy, 30 corresponds to 99.9%, and so on.¹¹

2. **Peak Call-**

¹¹Green and Ewing

ing: oA method used to identify regions in the genome where there is significant enrichment of sequencing reads. This indicates the presence of DNA-protein interactions (e.g., transcription factor binding sites or accessible chromatin regions). oPeaks represent areas where epigenetic marks or chromatin accessibility are concentrated.

Representing Peaks: o **P-value or Fold-change:** P-value: Indicates the statistical significance of the peak, helping to distinguish true peaks from background noise. Fold-change: Represents the difference in read density between treated and control samples, indicating the strength of the signal.



o **Types of Peaks:** Categorical Peaks: Simple yes/no indication of a peak's presence. Continuous Peaks: More nuanced representation that includes the intensity or enrichment level of the peak, often visualized as a signal track. Thresholded/Pseudoreplicated Peaks: Usually categorical, these peaks are of high confidence regions from multiple replicates (experiments) or pseudoreplicates (artificial data splits), to ensure reliability and reproducibility.

Example Data Pipeline

encodeproject.org

This example data pipeline originates from a ChIP-seq experiment targeting the H3K36me3 histone modification in brain tissue. The aim of the experiment is to map the locations where the H3K36me3 histone modification is present along the genome. The yellow bubbles represent downloadable data sets of different types, while the blue boxes represent step types (e.g. peak calling). In the left column are multiple data sets of raw sequence reads, which then undergo data quality steps before being aligned (first blue box) to the reference human genome GRCh38 (denoted by ENCFF110MCL below the reads). The next steps include Peak calling (categorical peaks) and signal generation (continuous peaks) to produce the data we normally use in our machine learning models. This data pipeline process aids in normalisation, noise reduction, and dimensionality reduction of the data.


1.2.2 Example Data Representations:

SIMPLE PICTURES

1.2.3 Transformations to stop extreme p-values

When utilising genomic data which incorporates p-values, it is important to consider and deal with extreme p-values. One way this is done is through using an Arcsinh-transformation (inverse hyperbolic sine). $\text{arsinh}(x) = \ln(x + \sqrt{x^2 + 1})$

The arcsinh-transformation as a logarithmic function helps in reducing the significance of outliers and sequencing depth while maintaining variance by compressing the range of the data. This transformation can be used in the data preprocessing stage. The graph below visualises how the transformation works. While extreme values are transformed logarithmically, the smaller values are barely transformed as the function for smaller values is more linear in nature.



Plot of Arcsinh Transformation compared to a log function, made with Desmos available under CC BY-SA 4.0. Text, arrows, and box shape added to image.

1.3 Sources of epigenetic data

There are numerous public data banks which contain genomic datasets ready to be downloaded. Blueprint's genomic datasets are focused on gene expression in healthy and diseased cells mostly relating to haematopoietic cells (cells which develop into different types of blood cells).

Roadmap The National Institute of Health's Roadmap Epigenomics Project contains sample datasets from multiple experiments as well as reference and mapping datasets.

Encode The Encode Project contains a large amount of publicly available genomic data easily filtered and downloaded. The genomic data used in this markdown book is sourced from Encode.

The largest genomic data bank is the UK Biobank, however they require that you apply for access to their datasets.

1.4 UCSC'S Genome Browser

The UCSC Genome Browser is a powerful and versatile tool that allows the visualisation and exploration of many sets of genomic data, especially bigWig files. It offers an extensive collection of genome assemblies, annotation tracks, and functional data, enabling users to examine gene structures, regulatory elements, and genetic variations. With its user-friendly interface and customisable display options, the UCSC Genome Browser facilitates detailed genomic analyses and supports a wide range of applications in genomics and bioinformatics. Whether you're investigating gene functions, exploring genetic variants, or studying comparative genomics, the UCSC Genome Browser serves as an essential resource for understanding complex genomic information. It is also possible to load and visualise genomic data from other sources such as Encode. While the visualisations are extensive, as you can explore below, the browser can be quite overwhelming for first time users.

The following is an example of what the same chIP-Seq data targeting the H3K36me3 histone modification in brain tissue looks like using UCSC's Genome Browser. The pseudoreplicated peaks represent categorically, the significant locations along the genome where the H3K36me3 histone modification is present.

UCSC Genome Browser

The following is an example of ATAC-Seq data from an experiment on T-helper 17 cells (a type of immune system cell). Recall that the ATAC-Seq method aims to find chromatin regions that are accessible for transcription factor binding. The p-value and fold change graphs show continuous peaks, while the IDR thresholded peaks and pseudoreplicated peaks represent the significant locations of accessible chromatin along the genome.

UCSC Genome Browser

1.5 Handling bigWig files

BigWig files can be quite tricky to deal with. However, libraries such as pyBigWig enable easier access of data.

Tutorial 1: Dealing with bigWigs (interactive)

Tutorial 1: Dealing with bigWigs (nbviewer)

1.6 Data loaders and pre-processing

1.7 Dealing with missing data (oversampling, undersampling, weighting)

Part II

Training models with DNA input

Chapter 2

Loss functions, and peak metrics

Chapter 3

Base pair averaging

Chapter 4

Training tricks

Chapter 5

Choosing which genomic regions to train on

Chapter 6

Effect of differences in sequencing depths

Chapter 7

Reproducibility of machine learning models

7.1 Seeding

7.2 Dashboarding

Chapter 8

Testing

Part III

Software libraries for model building

Chapter 9

gReLU

Chapter 10

Kipoi

Chapter 11

Weights and Biases

Part IV

ML pitfalls in genomics

Chapter 12

Pitfalls overview

12.1 Distributional differences

12.2 Dependent examples

12.3 Confounding

12.4 Leaky pre-processing

12.5 Unbalanced classes

12.6 Balancing the proportion of peaks / no-peaks in validation sets

Part V

Model interpretability

Chapter 13

Creating and visualising a simple model

Chapter 14

TF mo-Disco

Part VI

Using existing models

Chapter 15

Using the gReLU model zoo

Chapter 16

Fine tuning of Enformer

Part VII

Predicting in novel cell types

Chapter 17

Incorporating ATAC-seq info

Chapter 18

Use of cell type averages

Part VIII

More complex models

Chapter 19

Training multi-headed models

Chapter 20

Training siamese twin models

Bibliography

- Altuna Akalin. *Computational Genomics with R*. Github Pages, 2020. URL <https://compgenomr.github.io/book/>.
- Nora M. Al-Aboud, Connor Tupper, and Ishwarlal Jialal. *Genetics, Epigenetic Mechanism*. National Library of Medicine, 2023. URL <https://www.ncbi.nlm.nih.gov/books/NBK532999/#article-22137.r1>.
- PDQ® Cancer Genetics Editorial Board. *genetics-dictionary*. National Cancer Institute USA. URL <https://www.cancer.gov/publications/dictionaries/genetics-dictionary>.
- Phil Green and Brent Ewing. *Phred - Quality Base Calling*. CodonCode Corporation. URL [https://www.phrap.com/phred/#:~:text=Phred%20is%20a%20base%2Dcalling,%22\)%20to%20each%20base%20call](https://www.phrap.com/phred/#:~:text=Phred%20is%20a%20base%2Dcalling,%22)%20to%20each%20base%20call).
- Pratik Kanani and Mamta Padole. *Improving Pattern Matching performance in Genome sequences using Run Length Encoding in Distributed Raspberry Pi Clustering Environment*. Procedia Computer Science, 2020. URL <https://www.sciencedirect.com/science/article/pii/S1877050920311601?via%3Dihub>. <https://doi.org/10.1016/j.procs.2020.04.179>.
- Melanie Kappelman-Fenzl. *Design and Analysis of Epigenetics and ChIP-Sequencing Data*. Springer, 2021. URL https://doi.org/10.1007/978-3-030-62490-3_12. ISBN 978-3-030-62490-3.
- Amy Ralston and Kenna Shaw. *Gene Expression Regulates Cell Differentiation*. Nature Education, 2008. URL <https://www.nature.com/scitable/topicpage/gene-expression-regulates-cell-differentiation-931/#:~:text=All%20of%20the%20cells%20within,each%20cell%20deploys%20its%20genome>. Nature Education 1(1):127.
- Zainab Shahid, Brittany Simpson, Kathleen H. Miao, and Gurdeep Singh. *Genetics, Histone Code*. StatPearls Publishing LLC, 2023. URL <https://www.ncbi.nlm.nih.gov/books/NBK538477/>. PMID: 30860712.
- Kouzarides T. *Chromatin modifications and their function*. National Library of Medicine, 2007. URL <https://doi.org/10.1016/j.cell.2007.02.005>. PMID: 17320507.