



CS F402: Computational Geometry


Lecture 14: Upward Planar Drawings



Siddharth Gupta


April 28, 2025

Upward Planar Drawings – Motivation





Upward Planar Drawings – Motivation

- What may the direction of edges in a digraph represent?



Upward Planar Drawings – Motivation



- What may the direction of edges in a digraph represent?
 - Time




PERT diagram

Upward Planar Drawings – Motivation

- What may the direction of edges in a digraph represent?
 - Time
 - Flow





PERT diagram




Petri net

Upward Planar Drawings – Motivation


- What may the direction of edges in a digraph represent?
 - Time
 - Flow
 - Hierarchy



PERT diagram





Petri net




Phylogenetic network

Upward Planar Drawings – Motivation


- What may the direction of edges in a digraph represent?
 - Time
 - Flow
 - Hierarchy
 - ...



PERT diagram





Petri net




Phylogenetic network

Upward Planar Drawings – Motivation


- What may the direction of edges in a digraph represent?
 - Time
 - Flow
 - Hierarchy
 - ...
- Would be nice to have general direction preserved in drawing.



PERT diagram




Petri net



Phylogenetic network

Upward Planar Drawings – Definition


A directed graph $G = (V, E)$ is **upward planar** when it admits a drawing Γ that is



Upward Planar Drawings – Definition

A directed graph $G = (V, E)$ is **upward planar** when it admits a drawing Γ that is


- planar and



Upward Planar Drawings – Definition

A directed graph $G = (V, E)$ is **upward planar** when it admits a drawing Γ that is


- planar and
- where each edge is drawn as an upward, y-monotone curve.



Upward Planar Drawings – Definition


A directed graph $G = (V, E)$ is **upward planar** when it admits a drawing Γ that is

- planar and
- where each edge is drawn as an upward, y-monotone curve.




Upward Planarity – Necessary Conditions

- For a digraph G to be upward planar, it has to be:





Upward Planarity – Necessary Conditions

- For a digraph G to be upward planar, it has to be:
 - planar






Upward Planarity – Necessary Conditions

- For a digraph G to be upward planar, it has to be:
 - planar
 - acyclic







Upward Planarity – Necessary Conditions

- For a digraph G to be upward planar, it has to be:
 - planar
 - acyclic







Upward Planarity – Necessary Conditions

- For a digraph G to be upward planar, it has to be:
 - planar
 - acyclic







Upward Planarity – Necessary Conditions

- For a digraph G to be upward planar, it has to be:
 - planar
 - acyclic







Upward Planarity – Necessary Conditions

- For a digraph G to be upward planar, it has to be:
 - planar
 - acyclic







Upward Planarity – Necessary Conditions

- For a digraph G to be upward planar, it has to be:
 - planar
 - acyclic







Upward Planarity – Necessary Conditions

- For a digraph G to be upward planar, it has to be:
 - planar
 - acyclic





Upward Planarity – Necessary Conditions

- For a digraph G to be upward planar, it has to be:
 - planar
 - acyclic




Upward Planarity – Necessary Conditions



- For a digraph G to be upward planar, it has to be:
 - planar
 - acyclic



bimodal vertex





not bimodal




Upward Planarity – Necessary Conditions



- For a digraph G to be upward planar, it has to be:
 - planar
 - acyclic
 - bimodal



bimodal vertex





not bimodal




Upward Planarity – Necessary Conditions



- For a digraph G to be upward planar, it has to be:
 - planar
 - acyclic
 - bimodal
- ...but these conditions are *not sufficient*.



bimodal vertex



not bimodal



Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]
For a digraph G the following statements are equivalent:

Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.

Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.

Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.



no crossings

Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.



Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]


For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

no crossings

↓

acyclic digraph with
a single **source** s and single **sink** t



Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]


For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

no crossings

↓

acyclic digraph with
a single **source** s and single **sink** t



Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]


For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

no crossings

↓

acyclic digraph with
a single **source** s and single **sink** t



Upward Planarity – Characterization


Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

no crossings

acyclic digraph with
a single **source** s and single **sink** t



Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]


For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

no crossings

↓

acyclic digraph with
a single **source** s and single **sink** t



Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]


For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

no crossings

↓

acyclic digraph with
a single **source** s and single **sink** t



Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]


For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

no crossings

↓

acyclic digraph with
a single **source** s and single **sink** t



Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]


For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

no crossings

↓

acyclic digraph with
a single **source** s and single **sink** t




Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.



Additionally:

Embedded such that
 s and t are on the
 outerface f_0 .

no crossings


acyclic digraph with
 a single **source** s and single **sink** t

Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.



Additionally:

Embedded such that
 s and t are on the
 outerface f_0 .

or:

Edge (s, t) exists.

no crossings

acyclic digraph with
 a single **source** s and single **sink** t


Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

Proof.



Upward Planarity – Characterization


Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

Proof.

(2) \Rightarrow (1) By definition.



Upward Planarity – Characterization


Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

Proof.

(2) \Rightarrow (1) By definition. **(1) \Leftrightarrow (3)** Example:



Upward Planarity – Characterization


Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

Proof.

(2) \Rightarrow (1) By definition. (1) \Leftrightarrow (3) Example:



Upward Planarity – Characterization


Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

Proof.

(2) \Rightarrow (1) By definition. (1) \Leftrightarrow (3) Example:



Upward Planarity – Characterization


Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

Proof.

(2) \Rightarrow (1) By definition. (1) \Leftrightarrow (3) Example:



Upward Planarity – Characterization


Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

Proof.

(2) \Rightarrow (1) By definition. (1) \Leftrightarrow (3) Example:



Upward Planarity – Characterization


Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

Proof.

(2) \Rightarrow (1) By definition. (1) \Leftrightarrow (3) Example:



Upward Planarity – Characterization


Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

Proof.

(2) \Rightarrow (1) By definition. (1) \Leftrightarrow (3) Example:



Upward Planarity – Characterization


Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

Proof.

(2) \Rightarrow (1) By definition. (1) \Leftrightarrow (3) Example:



Upward Planarity – Characterization


Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

Proof.

(2) \Rightarrow (1) By definition. (1) \Leftrightarrow (3) Example:



Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]


For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

Proof.

(2) \Rightarrow (1) By definition. (1) \Leftrightarrow (3) Example:

(3) \Rightarrow (2) Triangulate & construct drawing:



Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]


For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

Proof.

(2) \Rightarrow (1) By definition. (1) \Leftrightarrow (3) Example:

(3) \Rightarrow (2) Triangulate & construct drawing:



Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.


Proof.

(2) \Rightarrow (1) By definition. (1) \Leftrightarrow (3) Example:

(3) \Rightarrow (2) Triangulate & construct drawing:

Claim.

Can draw in
prespecified
triangle.



Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.



Proof.

(2) \Rightarrow (1) By definition. (1) \Leftrightarrow (3) Example:

(3) \Rightarrow (2) Triangulate & construct drawing:

Claim.

Can draw in
prespecified
triangle.



Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

Proof.



(2) \Rightarrow (1) By definition. (1) \Leftrightarrow (3) Example:

(3) \Rightarrow (2) Triangulate & construct drawing:

Claim.

Can draw in
prespecified
triangle.

Induction on n .



Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

Proof.



(2) \Rightarrow (1) By definition. (1) \Leftrightarrow (3) Example:

(3) \Rightarrow (2) Triangulate & construct drawing:

Claim.

Can draw in
prespecified
triangle.

Induction on n .



Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

Proof.



(2) \Rightarrow (1) By definition. (1) \Leftrightarrow (3) Example:

(3) \Rightarrow (2) Triangulate & construct drawing:

Claim.

Can draw in
prespecified
triangle.

Induction on n .



Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

Proof.

(2) \Rightarrow (1) By definition. (1) \Leftrightarrow (3) Example:



(3) \Rightarrow (2) Triangulate & construct drawing:

Claim.

Can draw in
prespecified
triangle.

Induction on n .

Case 1:
chord



Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

Proof.

(2) \Rightarrow (1) By definition. (1) \Leftrightarrow (3) Example:



(3) \Rightarrow (2) Triangulate & construct drawing:

Claim.

Can draw in
prespecified
triangle.

Induction on n .

Case 1:
chord



Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

Proof.

(2) \Rightarrow (1) By definition. (1) \Leftrightarrow (3) Example:



(3) \Rightarrow (2) Triangulate & construct drawing:

Claim.

Can draw in
prespecified
triangle.

Induction on n .

Case 1:
chord



Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

Proof.

(2) \Rightarrow (1) By definition. (1) \Leftrightarrow (3) Example:


(3) \Rightarrow (2) Triangulate & construct drawing:

Claim.



Can draw in
prespecified
triangle.

Induction on n .

Case 1:
chord



Case 2:
no chord



Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

Proof.

(2) \Rightarrow (1) By definition. (1) \Leftrightarrow (3) Example:


(3) \Rightarrow (2) Triangulate & construct drawing:

Claim.



Can draw in
prespecified
triangle.

Induction on n .

Case 1:
chord



Case 2:
no chord



Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

Proof.

(2) \Rightarrow (1) By definition. (1) \Leftrightarrow (3) Example:


(3) \Rightarrow (2) Triangulate & construct drawing:

Claim.



Can draw in
prespecified
triangle.

Induction on n .

Case 1:
chord



Case 2:
no chord



Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

Proof.

(2) \Rightarrow (1) By definition. (1) \Leftrightarrow (3) Example:


(3) \Rightarrow (2) Triangulate & construct drawing:

Claim.



Can draw in
prespecified
triangle.

Induction on n .

Case 1:
chord



Case 2:
no chord



Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

Proof.

(2) \Rightarrow (1) By definition. (1) \Leftrightarrow (3) Example:


(3) \Rightarrow (2) Triangulate & construct drawing:

Claim.



Can draw in
prespecified
triangle.

Induction on n .

Case 1:
chord



Case 2:
no chord



Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

Proof.

(2) \Rightarrow (1) By definition. (1) \Leftrightarrow (3) Example:


(3) \Rightarrow (2) Triangulate & construct drawing:

Claim.



Can draw in
prespecified
triangle.

Induction on n .

Case 1:
chord



Case 2:
no chord



Upward Planarity – Characterization

Theorem 1. [Kelly 1987, Di Battista & Tamassia 1988]

For a digraph G the following statements are equivalent:

1. G is upward planar.
2. G admits an upward planar straight-line drawing.
3. G is the spanning subgraph of a planar st -digraph.

Proof.

(2) \Rightarrow (1) By definition. (1) \Leftrightarrow (3) Example:


(3) \Rightarrow (2) Triangulate & construct drawing:

Claim.



Can draw in
prespecified
triangle.

Induction on n .

Case 1:
chord



Case 2:
no chord



Upward Planarity – Complexity

Theorem.

[Garg & Tamassia, 1995]

For a *planar acyclic* digraph it is in general NP-hard to decide whether it is upward planar.

Upward Planarity – Complexity

Theorem.

[Garg & Tamassia, 1995]

For a *planar acyclic* digraph it is in general NP-hard to decide whether it is upward planar.

Theorem.

[Bertolazzi et al., 1994]

For a *combinatorially embedded* planar digraph it can be tested in $\mathcal{O}(n^2)$ time whether it is upward planar.

Upward Planarity – Complexity

Theorem.

[Garg & Tamassia, 1995]

For a *planar acyclic* digraph it is in general NP-hard to decide whether it is upward planar.

Theorem.

[Bertolazzi et al., 1994]

For a *combinatorially embedded* planar digraph it can be tested in $\mathcal{O}(n^2)$ time whether it is upward planar.

Corollary.

For a *triconnected* planar digraph it can be tested in $\mathcal{O}(n^2)$ time whether it is upward planar.

Upward Planarity – Complexity

Theorem.

[Garg & Tamassia, 1995]

For a *planar acyclic* digraph it is in general NP-hard to decide whether it is upward planar.

Theorem.

[Bertolazzi et al., 1994]

For a *combinatorially embedded* planar digraph it can be tested in $\mathcal{O}(n^2)$ time whether it is upward planar.

Corollary.

For a *triconnected* planar digraph it can be tested in $\mathcal{O}(n^2)$ time whether it is upward planar.

Theorem.

[Hutton & Lubiw, 1996]

For a *single-source* acyclic digraph it can be tested in $\mathcal{O}(n)$ time whether it is upward planar.

Upward Planarity – Complexity

Theorem.

[Garg & Tamassia, 1995]

For a *planar acyclic* digraph it is in general NP-hard to decide whether it is upward planar.

Theorem.

[Bertolazzi et al., 1994]

For a *combinatorially embedded* planar digraph it can be tested in $\mathcal{O}(n^2)$ time whether it is upward planar.

Corollary.

For a *triconnected* planar digraph it can be tested in $\mathcal{O}(n^2)$ time whether it is upward planar.

Theorem.

[Hutton & Lubiw, 1996]

For a *single-source* acyclic digraph it can be tested in $\mathcal{O}(n)$ time whether it is upward planar.

The Problem

Fixed Embedding Upward Planarity Testing.

Let $G = (V, E)$ be a plane digraph with set of faces F and outer face f_0 .

Test whether G is upward planar (wrt to F, f_0).

The Problem

Fixed Embedding Upward Planarity Testing.

Let $G = (V, E)$ be a plane digraph with set of faces F and outer face f_0 .

Test whether G is upward planar (wrt to F, f_0).

Idea.

- Find property that any upward planar drawing of G satisfies.

The Problem

Fixed Embedding Upward Planarity Testing.

Let $G = (V, E)$ be a plane digraph with set of faces F and outer face f_0 .

Test whether G is upward planar (wrt to F, f_0).

Idea.

- Find property that any upward planar drawing of G satisfies.
- Formalize property.

The Problem

Fixed Embedding Upward Planarity Testing.

Let $G = (V, E)$ be a plane digraph with set of faces F and outer face f_0 .


Test whether G is upward planar (wrt to F, f_0).

Idea.

- Find property that any upward planar drawing of G satisfies.
- Formalize property.
- Find algorithm to test property.

Angles, Local Sources & Sinks


Definitions.



Angles, Local Sources & Sinks

Definitions.


- A vertex v is a **local source** wrt a face f if v has two outgoing edges on ∂f .



Angles, Local Sources & Sinks

Definitions.


- A vertex v is a **local source** wrt a face f if v has two outgoing edges on ∂f .



Angles, Local Sources & Sinks

Definitions.


- A vertex v is a **local source** wrt a face f if v has two outgoing edges on ∂f .



Angles, Local Sources & Sinks

Definitions.


- A vertex v is a **local source** wrt a face f if v has two outgoing edges on ∂f .



Angles, Local Sources & Sinks

Definitions.


- A vertex v is a **local source** wrt a face f if v has two outgoing edges on ∂f .



Angles, Local Sources & Sinks

Definitions.


- A vertex v is a **local source** wrt a face f if v has two outgoing edges on ∂f .



Angles, Local Sources & Sinks

Definitions.


- A vertex v is a **local source** wrt a face f if v has two outgoing edges on ∂f .



Angles, Local Sources & Sinks

Definitions.


- A vertex v is a **local source** wrt a face f if v has two outgoing edges on ∂f .
- A vertex v is a **local sink** wrt a face f if v has two incoming edges on ∂f .



Angles, Local Sources & Sinks

Definitions.


- A vertex v is a **local source** wrt a face f if v has two outgoing edges on ∂f .
- A vertex v is a **local sink** wrt a face f if v has two incoming edges on ∂f .



Angles, Local Sources & Sinks

Definitions.


- A vertex v is a **local source** wrt a face f if v has two outgoing edges on ∂f .
- A vertex v is a **local sink** wrt a face f if v has two incoming edges on ∂f .



Angles, Local Sources & Sinks

Definitions.


- A vertex v is a **local source** wrt a face f if v has two outgoing edges on ∂f .
- A vertex v is a **local sink** wrt a face f if v has two incoming edges on ∂f .



Angles, Local Sources & Sinks

Definitions.


- A vertex v is a **local source** wrt a face f if v has two outgoing edges on ∂f .
- A vertex v is a **local sink** wrt a face f if v has two incoming edges on ∂f .



Angles, Local Sources & Sinks

Definitions.


- A vertex v is a **local source** wrt a face f if v has two outgoing edges on ∂f .
- A vertex v is a **local sink** wrt a face f if v has two incoming edges on ∂f .



Angles, Local Sources & Sinks

Definitions.


- A vertex v is a **local source** wrt a face f if v has two outgoing edges on ∂f .
- A vertex v is a **local sink** wrt a face f if v has two incoming edges on ∂f .
- An angle α at a local **source** / **sink** is **large** when $\alpha > \pi$ and **small** otherwise.



Angles, Local Sources & Sinks

Definitions.


- A vertex v is a **local source** wrt a face f if v has two outgoing edges on ∂f .
- A vertex v is a **local sink** wrt a face f if v has two incoming edges on ∂f .
- An angle α at a local **source** / **sink** is **large** when $\alpha > \pi$ and **small** otherwise.



Angles, Local Sources & Sinks

Definitions.


- A vertex v is a **local source** wrt a face f if v has two outgoing edges on ∂f .
- A vertex v is a **local sink** wrt a face f if v has two incoming edges on ∂f .
- An angle α at a local **source** / **sink** is **large** when $\alpha > \pi$ and **small** otherwise.



Angles, Local Sources & Sinks

Definitions.


- A vertex v is a **local source** wrt a face f if v has two outgoing edges on ∂f .
- A vertex v is a **local sink** wrt a face f if v has two incoming edges on ∂f .
- An angle α at a local **source** / **sink** is **large** when $\alpha > \pi$ and **small** otherwise.



Angles, Local Sources & Sinks

Definitions.


- A vertex v is a **local source** wrt a face f if v has two outgoing edges on ∂f .
- A vertex v is a **local sink** wrt a face f if v has two incoming edges on ∂f .
- An angle α at a local **source** / **sink** is **large** when $\alpha > \pi$ and **small** otherwise.



Angles, Local Sources & Sinks

Definitions.


- A vertex v is a **local source** wrt a face f if v has two outgoing edges on ∂f .
- A vertex v is a **local sink** wrt a face f if v has two incoming edges on ∂f .
- An angle α at a local **source** / **sink** is **large** when $\alpha > \pi$ and **small** otherwise.



Angles, Local Sources & Sinks

Definitions.


- A vertex v is a **local source** wrt a face f if v has two outgoing edges on ∂f .
- A vertex v is a **local sink** wrt a face f if v has two incoming edges on ∂f .
- An angle α at a local **source** / **sink** is **large** when $\alpha > \pi$ and **small** otherwise.
- $L(v) = \#$ large angles at v



Angles, Local Sources & Sinks

Definitions.


- A vertex v is a **local source** wrt a face f if v has two outgoing edges on ∂f .
- A vertex v is a **local sink** wrt a face f if v has two incoming edges on ∂f .
- An angle α at a local **source** / **sink** is **large** when $\alpha > \pi$ and **small** otherwise.
- $L(v) = \#$ large angles at v
- $L(f) = \#$ large angles in f



Angles, Local Sources & Sinks

Definitions.


- A vertex v is a **local source** wrt a face f if v has two outgoing edges on ∂f .
- A vertex v is a **local sink** wrt a face f if v has two incoming edges on ∂f .
- An angle α at a local **source** / **sink** is **large** when $\alpha > \pi$ and **small** otherwise.
- $L(v)$ = # large angles at v
- $L(f)$ = # large angles in f
- $S(v)$ & $S(f)$ for # small angles



Angles, Local Sources & Sinks

Definitions.


- A vertex v is a **local source** wrt a face f if v has two outgoing edges on ∂f .
- A vertex v is a **local sink** wrt a face f if v has two incoming edges on ∂f .
- An angle α at a local **source** / **sink** is **large** when $\alpha > \pi$ and **small** otherwise.
- $L(v)$ = # large angles at v
- $L(f)$ = # large angles in f
- $S(v)$ & $S(f)$ for # small angles
- $A(f)$ = # **local sources** wrt f



Angles, Local Sources & Sinks

Definitions.


- A vertex v is a **local source** wrt a face f if v has two outgoing edges on ∂f .
- A vertex v is a **local sink** wrt a face f if v has two incoming edges on ∂f .
- An angle α at a local **source** / **sink** is **large** when $\alpha > \pi$ and **small** otherwise.
- $L(v)$ = # large angles at v
- $L(f)$ = # large angles in f
- $S(v)$ & $S(f)$ for # small angles
- $A(f)$ = # **local sources** wrt f
= # **local sinks** wrt f



Angles, Local Sources & Sinks

Definitions.

- A vertex v is a **local source** wrt a face f if v has two outgoing edges on ∂f .
- A vertex v is a **local sink** wrt a face f if v has two incoming edges on ∂f .
- An angle α at a local **source** / **sink** is **large** when $\alpha > \pi$ and **small** otherwise.
- $L(v)$ = # large angles at v
- $L(f)$ = # large angles in f
- $S(v)$ & $S(f)$ for # small angles
- $A(f)$ = # **local sources** wrt f
= # **local sinks** wrt f




Lemma 1.

$$L(f) + S(f) = 2A(f)$$


Assignment Problem

- Vertex v is a **global source**.



Assignment Problem

- Vertex v is a **global source**.
- At which face does v have a **large** angle?



Angle Relations

Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

Angle Relations

Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

Proof by induction.

Angle Relations

Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

Proof by induction.

■ $L(f) = 0$


Angle Relations

Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

Proof by induction.

■ $L(f) = 0$




Angle Relations

Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

Proof by induction.

■ $L(f) = 0$



$$\Rightarrow S(f) = 2$$

Angle Relations


Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

■ $L(f) \geq 1$

Proof by induction.

■ $L(f) = 0$



$\Rightarrow S(f) = 2$

Angle Relations

Lemma 2.


$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

■ $L(f) \geq 1$

Split f with **edge** from a large angle at a “low” **sink** u to

Proof by induction.

■ $L(f) = 0$



$\Rightarrow S(f) = 2$


Angle Relations

Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

Proof by induction.

■ $L(f) = 0$




$\Rightarrow S(f) = 2$

■ $L(f) \geq 1$

Split f with **edge** from a large angle at a “low” **sink** u to

■ **sink** v with small angle:




Angle Relations

Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

Proof by induction.

■ $L(f) = 0$




$\Rightarrow S(f) = 2$

■ $L(f) \geq 1$

Split f with **edge** from a large angle at a “low” **sink** u to

■ **sink** v with small angle:




Angle Relations

Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

Proof by induction.

■ $L(f) = 0$




$\Rightarrow S(f) = 2$

■ $L(f) \geq 1$

Split f with **edge** from a large angle at a “low” **sink** u to

■ **sink** v with small angle:



$L(f) - S(f)$


Angle Relations

Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

Proof by induction.

■ $L(f) = 0$




$\Rightarrow S(f) = 2$

■ $L(f) \geq 1$

Split f with **edge** from a large angle at a “low” **sink** u to

■ **sink** v with small angle:



$$L(f) - S(f) = L(f_1) + L(f_2) + 1$$

Angle Relations


Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

■ $L(f) \geq 1$


Split f with **edge** from a large angle at a “low” **sink** u to

■ **sink** v with small angle:



Proof by induction.

■ $L(f) = 0$



$\Rightarrow S(f) = 2$

$$L(f) - S(f) = L(f_1) + L(f_2) + 1 - (S(f_1) + S(f_2) - 1)$$

Angle Relations


Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

■ $L(f) \geq 1$


Split f with **edge** from a large angle at a “low” **sink** u to

■ **sink** v with small angle:



Proof by induction.

■ $L(f) = 0$



$\Rightarrow S(f) = 2$

$$L(f) - S(f) = \overset{-2}{L(f_1)} + \overset{-2}{L(f_2)} + 1 - (S(f_1) + S(f_2) - 1)$$

Angle Relations


Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

■ $L(f) \geq 1$


Split f with **edge** from a large angle at a “low” **sink** u to

■ **sink** v with small angle:



Proof by induction.

■ $L(f) = 0$



$\Rightarrow S(f) = 2$

$$\begin{aligned} L(f) - S(f) &= \overset{-2}{L(f_1)} + \overset{-2}{L(f_2)} + 1 \\ &\quad - (S(f_1) + S(f_2) - 1) \\ &= -2 \end{aligned}$$

Angle Relations


Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

■ $L(f) \geq 1$


Split f with **edge** from a large angle at a “low” **sink** u to

■ **sink** v with small/large angle:



Proof by induction.

■ $L(f) = 0$



$\Rightarrow S(f) = 2$

$$\begin{aligned} L(f) - S(f) &= \overset{-2}{L(f_1)} + \overset{-2}{L(f_2)} + 1 \\ &\quad - (S(f_1) + S(f_2) - 1) \\ &= -2 \end{aligned}$$

Angle Relations


Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

■ $L(f) \geq 1$


Split f with **edge** from a large angle at a “low” **sink** u to

■ **sink** v with small/large angle:



Proof by induction.

■ $L(f) = 0$



$\Rightarrow S(f) = 2$

$$\begin{aligned} L(f) - S(f) &= \overset{-2}{L(f_1)} + \overset{-2}{L(f_2)} + 1 \\ &\quad - (S(f_1) + S(f_2) - 1) \\ &= -2 \end{aligned}$$

Angle Relations


Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

■ $L(f) \geq 1$


Split f with **edge** from a large angle at a “low” **sink** u to

■ **source** v with small angle:



Proof by induction.

■ $L(f) = 0$



$\Rightarrow S(f) = 2$

Angle Relations


Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

■ $L(f) \geq 1$


Split f with **edge** from a large angle at a “low” **sink** u to

■ **source** v with small angle:



Proof by induction.

■ $L(f) = 0$



$\Rightarrow S(f) = 2$

Angle Relations


Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

■ $L(f) \geq 1$


Split f with **edge** from a large angle at a “low” **sink** u to

■ **source** v with ~~small~~ angle:



Proof by induction.

■ $L(f) = 0$



$\Rightarrow S(f) = 2$


Angle Relations

Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

Proof by induction.

■ $L(f) = 0$




$\Rightarrow S(f) = 2$

■ $L(f) \geq 1$

Split f with **edge** from a large angle at a “low” **sink** u to

■ **source** v with ~~small~~/large angle:




Angle Relations

Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

Proof by induction.

■ $L(f) = 0$




$\Rightarrow S(f) = 2$

■ $L(f) \geq 1$

Split f with **edge** from a large angle at a “low” **sink** u to

■ **source** v with ~~small~~/large angle:




Angle Relations

Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

Proof by induction.

■ $L(f) = 0$





$\Rightarrow S(f) = 2$

■ $L(f) \geq 1$

Split f with **edge** from a large angle at a “low” **sink** u to

■ **source** v with ~~small~~/large angle:



$L(f) - S(f)$


Angle Relations

Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

Proof by induction.

■ $L(f) = 0$




$\Rightarrow S(f) = 2$

■ $L(f) \geq 1$

Split f with **edge** from a large angle at a “low” **sink** u to

■ **source** v with ~~small~~/large angle:



$$L(f) - S(f) = L(f_1) + L(f_2) + 2$$


Angle Relations

Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

Proof by induction.

■ $L(f) = 0$




$\Rightarrow S(f) = 2$

■ $L(f) \geq 1$

Split f with **edge** from a large angle at a “low” **sink** u to

■ **source** v with ~~small~~/large angle:



$$L(f) - S(f) = L(f_1) + L(f_2) + 2 - (S(f_1) + S(f_2))$$


Angle Relations

Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

Proof by induction.

■ $L(f) = 0$




$\Rightarrow S(f) = 2$

■ $L(f) \geq 1$

Split f with **edge** from a large angle at a “low” **sink** u to

■ **source** v with ~~small~~/large angle:



$$L(f) - S(f) = \overset{-2}{L(f_1)} + \overset{-2}{L(f_2)} + 2 - (S(f_1) + S(f_2))$$


Angle Relations

Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

Proof by induction.

■ $L(f) = 0$




$\Rightarrow S(f) = 2$

■ $L(f) \geq 1$

Split f with **edge** from a large angle at a “low” **sink** u to

■ **source** v with ~~small~~/large angle:



$$\begin{aligned} L(f) - S(f) &= \overset{-2}{L(f_1)} + \overset{-2}{L(f_2)} + 2 \\ &\quad - (S(f_1) + S(f_2)) \\ &= -2 \end{aligned}$$


Angle Relations

Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

Proof by induction.

■ $L(f) = 0$




$\Rightarrow S(f) = 2$

■ $L(f) \geq 1$

Split f with **edge** from a large angle at a “low” **sink** u to

■ vertex v that is neither source nor sink:




Angle Relations

Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

Proof by induction.

■ $L(f) = 0$




$\Rightarrow S(f) = 2$

■ $L(f) \geq 1$

Split f with **edge** from a large angle at a “low” **sink** u to

■ vertex v that is neither source nor sink:




Angle Relations

Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

Proof by induction.

■ $L(f) = 0$




$\Rightarrow S(f) = 2$

■ $L(f) \geq 1$

Split f with **edge** from a large angle at a “low” **sink** u to

■ vertex v that is neither source nor sink:



$L(f) - S(f)$


Angle Relations

Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

Proof by induction.

■ $L(f) = 0$




$\Rightarrow S(f) = 2$

■ $L(f) \geq 1$

Split f with **edge** from a large angle at a “low” **sink** u to

■ vertex v that is neither source nor sink:



$$L(f) - S(f) = L(f_1) + L(f_2) + 1$$


Angle Relations

Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

Proof by induction.

■ $L(f) = 0$




$\Rightarrow S(f) = 2$

■ $L(f) \geq 1$

Split f with **edge** from a large angle at a “low” **sink** u to

■ vertex v that is neither source nor sink:



$$L(f) - S(f) = L(f_1) + L(f_2) + 1 \\ - (S(f_1) + S(f_2) - 1)$$


Angle Relations

Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

Proof by induction.

■ $L(f) = 0$




$$\Rightarrow S(f) = 2$$

■ $L(f) \geq 1$

Split f with **edge** from a large angle at a “low” **sink** u to

■ vertex v that is neither source nor sink:



$$L(f) - S(f) = \overset{-2}{L(f_1)} + \overset{-2}{L(f_2)} + 1 - (S(f_1) + S(f_2) - 1)$$


Angle Relations

Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

Proof by induction.

■ $L(f) = 0$




$\Rightarrow S(f) = 2$

■ $L(f) \geq 1$

Split f with **edge** from a large angle at a “low” **sink** u to

■ vertex v that is neither source nor sink:



$$\begin{aligned} L(f) - S(f) &= \overset{-2}{L(f_1)} + \overset{-2}{L(f_2)} + 1 \\ &\quad - (S(f_1) + S(f_2) - 1) \\ &= -2 \end{aligned}$$

Angle Relations


Lemma 2.

$$L(f) - S(f) = \begin{cases} -2, & f \neq f_0 \\ +2, & f = f_0 \end{cases}$$

■ $L(f) \geq 1$


Split f with **edge** from a large angle at a “low” **sink** u to

■ vertex v that is neither source nor sink:



Proof by induction.

■ $L(f) = 0$



$\Rightarrow S(f) = 2$

$$\begin{aligned} L(f) - S(f) &= \overset{-2}{L(f_1)} + \overset{-2}{L(f_2)} + 1 \\ &\quad - (S(f_1) + S(f_2) - 1) \\ &= -2 \end{aligned}$$

■ Otherwise “high” **source** u exists.

Number of Large Angles

Lemma 3.

In every upward planar drawing of G holds that

Number of Large Angles

Lemma 3.

In every upward planar drawing of G holds that

- for each vertex $v \in V$: $L(v) =$

Number of Large Angles

Lemma 3.

In every upward planar drawing of G holds that

- for each vertex $v \in V$: $L(v) = \begin{cases} 0 & v \text{ inner vertex,} \end{cases}$

Number of Large Angles

Lemma 3.

In every upward planar drawing of G holds that

- for each vertex $v \in V$: $L(v) = \begin{cases} 0 & v \text{ inner vertex,} \\ 1 & v \text{ source / sink;} \end{cases}$

Number of Large Angles

Lemma 3.

In every upward planar drawing of G holds that

- for each vertex $v \in V$: $L(v) = \begin{cases} 0 & v \text{ inner vertex,} \\ 1 & v \text{ source / sink;} \end{cases}$
- for each face f : $L(f) =$

Number of Large Angles

Lemma 3.

In every upward planar drawing of G holds that

- for each vertex $v \in V$: $L(v) = \begin{cases} 0 & v \text{ inner vertex,} \\ 1 & v \text{ source / sink;} \end{cases}$
- for each face f : $L(f) = \begin{cases} A(f) - 1 & f \neq f_0, \end{cases}$

Number of Large Angles

Lemma 3.

In every upward planar drawing of G holds that


- for each vertex $v \in V$: $L(v) = \begin{cases} 0 & v \text{ inner vertex,} \\ 1 & v \text{ source / sink;} \end{cases}$
- for each face f : $L(f) = \begin{cases} A(f) - 1 & f \neq f_0, \\ A(f) + 1 & f = f_0. \end{cases}$

Number of Large Angles

Lemma 3.

In every upward planar drawing of G holds that

- for each vertex $v \in V$: $L(v) = \begin{cases} 0 & v \text{ inner vertex,} \\ 1 & v \text{ source / sink;} \end{cases}$
- for each face f : $L(f) = \begin{cases} A(f) - 1 & f \neq f_0, \\ A(f) + 1 & f = f_0. \end{cases}$




Number of Large Angles

Lemma 3.

In every upward planar drawing of G holds that

- for each vertex $v \in V$: $L(v) = \begin{cases} 0 & v \text{ inner vertex,} \\ 1 & v \text{ source / sink;} \end{cases}$
- for each face f : $L(f) = \begin{cases} A(f) - 1 & f \neq f_0, \\ A(f) + 1 & f = f_0. \end{cases}$




Number of Large Angles

Lemma 3.

In every upward planar drawing of G holds that

- for each vertex $v \in V$: $L(v) = \begin{cases} 0 & v \text{ inner vertex,} \\ 1 & v \text{ source / sink;} \end{cases}$
- for each face f : $L(f) = \begin{cases} A(f) - 1 & f \neq f_0, \\ A(f) + 1 & f = f_0. \end{cases}$





Number of Large Angles

Lemma 3.

In every upward planar drawing of G holds that

- for each vertex $v \in V$: $L(v) = \begin{cases} 0 & v \text{ inner vertex,} \\ 1 & v \text{ source / sink;} \end{cases}$
- for each face f : $L(f) = \begin{cases} A(f) - 1 & f \neq f_0, \\ A(f) + 1 & f = f_0. \end{cases}$




Number of Large Angles

Lemma 3.

In every upward planar drawing of G holds that

- for each vertex $v \in V$: $L(v) = \begin{cases} 0 & v \text{ inner vertex,} \\ 1 & v \text{ source / sink;} \end{cases}$
- for each face f : $L(f) = \begin{cases} A(f) - 1 & f \neq f_0, \\ A(f) + 1 & f = f_0. \end{cases}$




Number of Large Angles

Lemma 3.

In every upward planar drawing of G holds that

- for each vertex $v \in V$: $L(v) = \begin{cases} 0 & v \text{ inner vertex,} \\ 1 & v \text{ source / sink;} \end{cases}$
- for each face f : $L(f) = \begin{cases} A(f) - 1 & f \neq f_0, \\ A(f) + 1 & f = f_0. \end{cases}$

Proof.




Number of Large Angles

Lemma 3.

In every upward planar drawing of G holds that

- for each vertex $v \in V$: $L(v) = \begin{cases} 0 & v \text{ inner vertex,} \\ 1 & v \text{ source / sink;} \end{cases}$
- for each face f : $L(f) = \begin{cases} A(f) - 1 & f \neq f_0, \\ A(f) + 1 & f = f_0. \end{cases}$

Proof. Lemma 1: $L(f) + S(f) = 2A(f)$



Number of Large Angles


Lemma 3.

In every upward planar drawing of G holds that

- for each vertex $v \in V$: $L(v) = \begin{cases} 0 & v \text{ inner vertex,} \\ 1 & v \text{ source / sink;} \end{cases}$
- for each face f : $L(f) = \begin{cases} A(f) - 1 & f \neq f_0, \\ A(f) + 1 & f = f_0. \end{cases}$

Proof. Lemma 1: $L(f) + S(f) = 2A(f)$

Lemma 2: $L(f) - S(f) = \pm 2$.



Number of Large Angles

Lemma 3.


In every upward planar drawing of G holds that

- for each vertex $v \in V$: $L(v) = \begin{cases} 0 & v \text{ inner vertex,} \\ 1 & v \text{ source / sink;} \end{cases}$
- for each face f : $L(f) = \begin{cases} A(f) - 1 & f \neq f_0, \\ A(f) + 1 & f = f_0. \end{cases}$

Proof. Lemma 1: $L(f) + S(f) = 2A(f)$

Lemma 2: $L(f) - S(f) = \pm 2$.

$\Rightarrow 2L(f) = 2A(f) \pm 2$.



Assignment of Large Angles to Faces

Let S and T be the sets of **sources** and **sinks**, respectively.

Assignment of Large Angles to Faces

Let S and T be the sets of **sources** and **sinks**, respectively.

Definition.

A **consistent assignment** $\Phi: S \cup T \rightarrow F$ is a mapping where

Assignment of Large Angles to Faces

Let S and T be the sets of **sources** and **sinks**, respectively.

Definition.

A **consistent assignment** $\Phi: S \cup T \rightarrow F$ is a mapping where

$\Phi: v \mapsto$ incident face, where v forms **large angle**

such that

Assignment of Large Angles to Faces

Let S and T be the sets of **sources** and **sinks**, respectively.

Definition.

A **consistent assignment** $\Phi: S \cup T \rightarrow F$ is a mapping where

$\Phi: v \mapsto$ incident face, where v forms **large angle**

such that

$$|\Phi^{-1}(f)| =$$

Assignment of Large Angles to Faces

Let S and T be the sets of **sources** and **sinks**, respectively.

Definition.

A **consistent assignment** $\Phi: S \cup T \rightarrow F$ is a mapping where

$\Phi: v \mapsto$ incident face, where v forms **large angle**

such that

$$|\Phi^{-1}(f)| = L(f) =$$

Assignment of Large Angles to Faces

Let S and T be the sets of **sources** and **sinks**, respectively.

Definition.


A **consistent assignment** $\Phi: S \cup T \rightarrow F$ is a mapping where

$\Phi: v \mapsto$ incident face, where v forms **large angle**


such that

$$|\Phi^{-1}(f)| = L(f) = \begin{cases} A(f) - 1 & \text{if } f \neq f_0, \\ A(f) + 1 & \text{if } f = f_0. \end{cases}$$

Example of Angle to Face Assignment




Example of Angle to Face Assignment




global **sources**

Example of Angle to Face Assignment




■ global **sources** & **sinks**

Example of Angle to Face Assignment



Example of Angle to Face Assignment




■ global **sources** & **sinks**

$A(f)$ # **sources** / **sinks** of f

$L(f)$ # **large angles** of f

Example of Angle to Face Assignment



■ global **sources** & **sinks**

$A(f)$ # **sources** / **sinks** of f

$L(f)$ # **large angles** of f

assignment

$\Phi : S \cup T \rightarrow F$

Finding a Consistent Assignment

Idea.

Flow $(v, f) = 1$ from global **source** / **sink** v to the incident face f its **large angle** gets assigned to.

Finding a Consistent Assignment

Idea.

Flow $(v, f) = 1$ from global **source** / **sink** v to the incident face f its **large angle** gets assigned to.

Flow network.

$$N_{F,f_0}(G) = ((W, E'); b; \ell; u)$$

■ $W =$

■ $E' =$

■ $\ell(e) =$

■ $u(e) =$

■ $b(w) =$

Finding a Consistent Assignment

Idea.

Flow $(v, f) = 1$ from global **source** / **sink** v to the incident face f its **large angle** gets assigned to.


Flow network.

$$N_{F,f_0}(G) = ((W, E'); b; \ell; u)$$

- $W =$
- $E' =$
- $\ell(e) =$
- $u(e) =$

- $b(w) =$

Example.



Finding a Consistent Assignment

Idea.

Flow $(v, f) = 1$ from global **source** / **sink** v to the incident face f its **large angle** gets assigned to.

Flow network.

$$N_{F,f_0}(G) = ((W, E'); b; \ell; u)$$

$$\blacksquare W = \{v \in V \mid v \text{ **source** or **sink**}\} \cup F$$


$$\blacksquare E' =$$

$$\blacksquare \ell(e) =$$

$$\blacksquare u(e) =$$

$$\blacksquare b(w) =$$

Example.



Finding a Consistent Assignment

Idea.

Flow $(v, f) = 1$ from global **source** / **sink** v to the incident face f its **large angle** gets assigned to.

Flow network.

$$N_{F,f_0}(G) = ((W, E'); b; \ell; u)$$

$$\blacksquare W = \{v \in V \mid v \text{ **source** or **sink**}\} \cup F$$


$$\blacksquare E' =$$

$$\blacksquare \ell(e) =$$

$$\blacksquare u(e) =$$

$$\blacksquare b(w) =$$

Example.



Finding a Consistent Assignment

Idea.


Flow $(v, f) = 1$ from global **source** / **sink** v to the incident face f its **large angle** gets assigned to.

Flow network.

$$N_{F,f_0}(G) = ((W, E'); b; \ell; u)$$

- $W = \{v \in V \mid v \text{ **source** or **sink**}\} \cup F$
- $E' = \{(v, f) \mid v \text{ incident to } f\}$
- $\ell(e) =$
- $u(e) =$
- $b(w) =$

Example.



Finding a Consistent Assignment

Idea.


Flow $(v, f) = 1$ from global **source** / **sink** v to the incident face f its **large angle** gets assigned to.

Flow network.

$$N_{F,f_0}(G) = ((W, E'); b; \ell; u)$$

- $W = \{v \in V \mid v \text{ **source** or **sink**}\} \cup F$
- $E' = \{(v, f) \mid v \text{ incident to } f\}$
- $\ell(e) = 0 \ \forall e \in E'$
- $u(e) = 1 \ \forall e \in E'$
- $b(w) =$

Example.



Finding a Consistent Assignment

Idea.

Flow $(v, f) = 1$ from global **source** / **sink** v to the incident face f its **large angle** gets assigned to.

Flow network.

$$N_{F,f_0}(G) = ((W, E'); b; \ell; u)$$

$$\blacksquare W = \{v \in V \mid v \text{ source or sink}\} \cup F$$


$$\blacksquare E' = \{(v, f) \mid v \text{ incident to } f\}$$

$$\blacksquare \ell(e) = 0 \quad \forall e \in E'$$

$$\blacksquare u(e) = 1 \quad \forall e \in E'$$

$$\blacksquare b(w) = \begin{cases} 1 \\ 0 \end{cases} \quad \forall w \in W \cap V$$

Example.



Finding a Consistent Assignment

Idea.

Flow $(v, f) = 1$ from global **source** / **sink** v to the incident face f its **large angle** gets assigned to.

Flow network.

$$N_{F,f_0}(G) = ((W, E'); b; \ell; u)$$

$$\blacksquare W = \{v \in V \mid v \text{ source or sink}\} \cup F$$


$$\blacksquare E' = \{(v, f) \mid v \text{ incident to } f\}$$

$$\blacksquare \ell(e) = 0 \quad \forall e \in E'$$

$$\blacksquare u(e) = 1 \quad \forall e \in E'$$

$$\blacksquare b(w) = \begin{cases} 1 & \forall w \in W \cap V \\ -(A(w) - 1) & \forall w \in F \setminus \{f_0\} \end{cases}$$

Example.



Finding a Consistent Assignment

Idea.

Flow $(v, f) = 1$ from global **source** / **sink** v to the incident face f its **large angle** gets assigned to.

Flow network.

$$N_{F, f_0}(G) = ((W, E'); b; \ell; u)$$

$$\blacksquare W = \{v \in V \mid v \text{ source or sink}\} \cup F$$


$$\blacksquare E' = \{(v, f) \mid v \text{ incident to } f\}$$

$$\blacksquare \ell(e) = 0 \quad \forall e \in E'$$

$$\blacksquare u(e) = 1 \quad \forall e \in E'$$

$$\blacksquare b(w) = \begin{cases} 1 & \forall w \in W \cap V \\ -(A(w) - 1) & \forall w \in F \setminus \{f_0\} \end{cases}$$

Example.



Flow $(v, f) = 1$ from global **source** / **sink** v to the incident face f its **large angle** gets assigned to.

$$N_{F,f_0}(G) = ((W, E'); b; \ell; u)$$

- $W = \{v \in V \mid v \text{ source or sink}\} \cup F$
- $E' = \{(v, f) \mid v \text{ incident to } f\}$
- $\ell(e) = 0 \ \forall e \in E'$
- $u(e) = 1 \ \forall e \in E'$
- $b(w) = \begin{cases} 1 & \forall w \in W \cap V \\ -(A(w) - 1) & \forall w \in F \setminus \{f_0\} \end{cases}$

The graph consists of 10 nodes and several edges. The nodes are categorized by color and shape: purple squares (labeled 1), orange squares (labeled 1), light blue diamonds (labeled 0 and -2), and white circles. Blue arrows highlight specific paths and weights: a path from the top-left purple square to the bottom-left light blue diamond (labeled 1), a path from the top-left purple square to the top-right purple square (labeled 1), a path from the top-right purple square to the top-right orange square (labeled 1), a path from the top-right orange square to the bottom-right orange square (labeled 1), a path from the bottom-right orange square to the bottom-right light blue diamond (labeled -2), a path from the bottom-right light blue diamond to the bottom-left light blue diamond (labeled -2), and a path from the bottom-left light blue diamond to the top-left purple square (labeled 1). There are also several other edges connecting the nodes, some of which are highlighted with blue arrows.

Finding a Consistent Assignment

Idea.


Flow $(v, f) = 1$ from global **source** / **sink** v to the incident face f its **large angle** gets assigned to.

Flow network.

$$N_{F,f_0}(G) = ((W, E'); b; \ell; u)$$

- $W = \{v \in V \mid v \text{ source or sink}\} \cup F$
- $E' = \{(v, f) \mid v \text{ incident to } f\}$
- $\ell(e) = 0 \ \forall e \in E'$
- $u(e) = 1 \ \forall e \in E'$
- $b(w) = \begin{cases} 1 & \forall w \in W \cap V \\ -(A(w) - 1) & \forall w \in F \setminus \{f_0\} \end{cases}$

Example.



Finding a Consistent Assignment

Idea.

Flow $(v, f) = 1$ from global **source** / **sink** v to the incident face f its **large angle** gets assigned to.

Flow network.

$$N_{F, f_0}(G) = ((W, E'); b; \ell; u)$$

$$\blacksquare W = \{v \in V \mid v \text{ source or sink}\} \cup F$$


$$\blacksquare E' = \{(v, f) \mid v \text{ incident to } f\}$$

$$\blacksquare \ell(e) = 0 \quad \forall e \in E'$$

$$\blacksquare u(e) = 1 \quad \forall e \in E'$$

$$\blacksquare b(w) = \begin{cases} 1 & \forall w \in W \cap V \\ -(A(w) - 1) & \forall w \in F \setminus \{f_0\} \\ -(A(w) + 1) & w = f_0 \end{cases}$$

Example.



Finding a Consistent Assignment

Idea.


Flow $(v, f) = 1$ from global **source** / **sink** v to the incident face f its **large angle** gets assigned to.

Flow network.

$$N_{F, f_0}(G) = ((W, E'); b; \ell; u)$$

- $W = \{v \in V \mid v \text{ source or sink}\} \cup F$
- $E' = \{(v, f) \mid v \text{ incident to } f\}$
- $\ell(e) = 0 \ \forall e \in E'$
- $u(e) = 1 \ \forall e \in E'$
- $b(w) = \begin{cases} 1 & \forall w \in W \cap V \\ -(A(w) - 1) & \forall w \in F \setminus \{f_0\} \\ -(A(w) + 1) & w = f_0 \end{cases}$

Example.



Finding a Consistent Assignment

Idea.

Flow $(v, f) = 1$ from global **source** / **sink** v to the incident face f its **large angle** gets assigned to.

Flow network.

$$N_{F,f_0}(G) = ((W, E'); b; \ell; u)$$

$$\blacksquare W = \{v \in V \mid v \text{ source or sink}\} \cup F$$


$$\blacksquare E' = \{(v, f) \mid v \text{ incident to } f\}$$

$$\blacksquare \ell(e) = 0 \quad \forall e \in E'$$

$$\blacksquare u(e) = 1 \quad \forall e \in E'$$

$$\blacksquare b(w) = \begin{cases} 1 & \forall w \in W \cap V \\ -(A(w) - 1) & \forall w \in F \setminus \{f_0\} \\ -(A(w) + 1) & w = f_0 \end{cases}$$

Example.



Finding a Consistent Assignment

Idea.

Flow $(v, f) = 1$ from global **source** / **sink** v to the incident face f its **large angle** gets assigned to.

Flow network.

$$N_{F, f_0}(G) = ((W, E'); b; \ell; u)$$

$$\blacksquare W = \{v \in V \mid v \text{ source or sink}\} \cup F$$



$$\blacksquare E' = \{(v, f) \mid v \text{ incident to } f\}$$

$$\blacksquare \ell(e) = 0 \quad \forall e \in E'$$

$$\blacksquare u(e) = 1 \quad \forall e \in E'$$

$$\blacksquare b(w) = \begin{cases} 1 & \forall w \in W \cap V \\ -(A(w) - 1) & \forall w \in F \setminus \{f_0\} \\ -(A(w) + 1) & w = f_0 \end{cases}$$

Example.



Finding a Consistent Assignment

Idea.

Flow $(v, f) = 1$ from global **source** / **sink** v to the incident face f its **large angle** gets assigned to.

Flow network.

$$N_{F, f_0}(G) = ((W, E'); b; \ell; u)$$

$$\blacksquare W = \{v \in V \mid v \text{ source or sink}\} \cup F$$



$$\blacksquare E' = \{(v, f) \mid v \text{ incident to } f\}$$

$$\blacksquare \ell(e) = 0 \quad \forall e \in E'$$

$$\blacksquare u(e) = 1 \quad \forall e \in E'$$

$$\blacksquare b(w) = \begin{cases} 1 & \forall w \in W \cap V \\ -(A(w) - 1) & \forall w \in F \setminus \{f_0\} \\ -(A(w) + 1) & w = f_0 \end{cases}$$

Example.



Finding a Consistent Assignment

Idea.

Flow $(v, f) = 1$ from global **source** / **sink** v to the incident face f its **large angle** gets assigned to.

Flow network.

$$N_{F, f_0}(G) = ((W, E'); b; \ell; u)$$

$$\blacksquare W = \{v \in V \mid v \text{ source or sink}\} \cup F$$



$$\blacksquare E' = \{(v, f) \mid v \text{ incident to } f\}$$

$$\blacksquare \ell(e) = 0 \quad \forall e \in E'$$

$$\blacksquare u(e) = 1 \quad \forall e \in E'$$

$$\blacksquare b(w) = \begin{cases} 1 & \forall w \in W \cap V \\ -(A(w) - 1) & \forall w \in F \setminus \{f_0\} \\ -(A(w) + 1) & w = f_0 \end{cases}$$

Example.



Result Characterization

Theorem 3.

Let $G = (V, E)$ be an acyclic plane digraph with embedding given by F, f_0 .

Result Characterization

Theorem 3.

Let $G = (V, E)$ be an acyclic plane digraph with embedding given by F, f_0 .

Then G is upward planar (respecting F, f_0) if and only if G is bimodal and there exists consistent assignment Φ .

Result Characterization

Theorem 3.

Let $G = (V, E)$ be an acyclic plane digraph with embedding given by F, f_0 .

Then G is upward planar (respecting F, f_0) if and only if G is bimodal and there exists consistent assignment Φ .

Proof.

\Rightarrow : As constructed before.

Result Characterization

Theorem 3.

Let $G = (V, E)$ be an acyclic plane digraph with embedding given by F, f_0 .

Then G is upward planar (respecting F, f_0) if and only if G is bimodal and there exists consistent assignment Φ .

Proof.

\Rightarrow : As constructed before.

\Leftarrow : Idea:

- Construct planar st-digraph that is supergraph of G .

Result Characterization

Theorem 3.

Let $G = (V, E)$ be an acyclic plane digraph with embedding given by F, f_0 .

Then G is upward planar (respecting F, f_0) if and only if G is bimodal and there exists consistent assignment Φ .

Proof.

\Rightarrow : As constructed before.

\Leftarrow : Idea:

- Construct planar st-digraph that is supergraph of G .
- Apply equivalence from Theorem 1.

Result Characterization

Theorem 3.

Let $G = (V, E)$ be an acyclic plane digraph with embedding given by F, f_0 .

Then G is upward planar (respecting F, f_0) if and only if G is bimodal and there exists consistent assignment Φ .

Proof.

\Rightarrow : As constructed before.

\Leftarrow : Idea:

- Construct planar st -digraph that is supergraph of G .
- Apply equivalence from Theorem 1.

Theorem 1.

[Kelly 1987, Di Battista & Tamassia 1988]


[...] G is upward planar

$\Leftrightarrow G$ is the spanning subgraph of a planar st -digraph.

Refinement Algorithm – $\Phi, F, f_0 \rightarrow \text{st-digraph}$


Refinement Algorithm – $\Phi, F, f_0 \rightarrow \text{st-digraph}$

Let f be a face. Consider the clockwise angle sequence σ_f of L/S on local **sources** and **sinks** of f .



Refinement Algorithm – $\Phi, F, f_0 \rightarrow \text{st-digraph}$


Let f be a face. Consider the clockwise angle sequence σ_f of L/S on local **sources** and **sinks** of f .



Refinement Algorithm – $\Phi, F, f_0 \rightarrow \text{st-digraph}$

Let f be a face. Consider the clockwise angle sequence σ_f of L/S on local **sources** and **sinks** of f .


- Goal: Add edges to break **large angles** (**sources** and **sinks**).



Refinement Algorithm – $\Phi, F, f_0 \rightarrow \text{st-digraph}$

Let f be a face. Consider the clockwise angle sequence σ_f of L/S on local **sources** and **sinks** of f .


- Goal: Add edges to break **large angles** (**sources** and **sinks**).
- For $f \neq f_0$ with $|\sigma_f| \geq 2$ containing $\langle L, S, S \rangle$ at vertices x, y, z :



Refinement Algorithm – $\Phi, F, f_0 \rightarrow \text{st-digraph}$

Let f be a face. Consider the clockwise angle sequence σ_f of L/S on local **sources** and **sinks** of f .


- Goal: Add edges to break **large angles** (**sources** and **sinks**).
- For $f \neq f_0$ with $|\sigma_f| \geq 2$ containing $\langle L, S, S \rangle$ at vertices x, y, z :



Refinement Algorithm – $\Phi, F, f_0 \rightarrow \text{st-digraph}$

Let f be a face. Consider the clockwise angle sequence σ_f of L/S on local **sources** and **sinks** of f .


- Goal: Add edges to break **large angles** (**sources** and **sinks**).
- For $f \neq f_0$ with $|\sigma_f| \geq 2$ containing $\langle L, S, S \rangle$ at vertices x, y, z :
- x **source** \Rightarrow insert **edge** (z, x)



Refinement Algorithm – $\Phi, F, f_0 \rightarrow \text{st-digraph}$

Let f be a face. Consider the clockwise angle sequence σ_f of L/S on local **sources** and **sinks** of f .


- Goal: Add edges to break **large angles** (**sources** and **sinks**).
- For $f \neq f_0$ with $|\sigma_f| \geq 2$ containing $\langle L, S, S \rangle$ at vertices x, y, z :
- x **source** \Rightarrow insert **edge** (z, x)



Refinement Algorithm – $\Phi, F, f_0 \rightarrow \text{st-digraph}$

Let f be a face. Consider the clockwise angle sequence σ_f of L/S on local **sources** and **sinks** of f .


- Goal: Add edges to break **large angles** (**sources** and **sinks**).
- For $f \neq f_0$ with $|\sigma_f| \geq 2$ containing $\langle L, S, S \rangle$ at vertices x, y, z :
- x **source** \Rightarrow insert **edge** (z, x)



Refinement Algorithm – $\Phi, F, f_0 \rightarrow \text{st-digraph}$

Let f be a face. Consider the clockwise angle sequence σ_f of L/S on local **sources** and **sinks** of f .


- Goal: Add edges to break **large angles** (**sources** and **sinks**).
- For $f \neq f_0$ with $|\sigma_f| \geq 2$ containing $\langle L, S, S \rangle$ at vertices x, y, z :
- x **source** \Rightarrow insert **edge** (z, x)



Refinement Algorithm – $\Phi, F, f_0 \rightarrow \text{st-digraph}$

Let f be a face. Consider the clockwise angle sequence σ_f of L/S on local **sources** and **sinks** of f .


- Goal: Add edges to break **large angles** (**sources** and **sinks**).
- For $f \neq f_0$ with $|\sigma_f| \geq 2$ containing $\langle L, S, S \rangle$ at vertices x, y, z :
- x **source** \Rightarrow insert **edge** (z, x)



Refinement Algorithm – $\Phi, F, f_0 \rightarrow \text{st-digraph}$

Let f be a face. Consider the clockwise angle sequence σ_f of L/S on local **sources** and **sinks** of f .


- Goal: Add edges to break **large angles** (**sources** and **sinks**).
- For $f \neq f_0$ with $|\sigma_f| \geq 2$ containing $\langle L, S, S \rangle$ at vertices x, y, z :
- x **source** \Rightarrow insert **edge** (z, x)
- x **sink** \Rightarrow insert **edge** (x, z) .



Refinement Algorithm – $\Phi, F, f_0 \rightarrow \text{st-digraph}$

Let f be a face. Consider the clockwise angle sequence σ_f of L/S on local **sources** and **sinks** of f .


- Goal: Add edges to break **large angles** (**sources** and **sinks**).
- For $f \neq f_0$ with $|\sigma_f| \geq 2$ containing $\langle L, S, S \rangle$ at vertices x, y, z :
- x **source** \Rightarrow insert **edge** (z, x)
- x **sink** \Rightarrow insert **edge** (x, z) .



Refinement Algorithm – $\Phi, F, f_0 \rightarrow \text{st-digraph}$

Let f be a face. Consider the clockwise angle sequence σ_f of L/S on local **sources** and **sinks** of f .


- Goal: Add edges to break **large angles** (**sources** and **sinks**).
- For $f \neq f_0$ with $|\sigma_f| \geq 2$ containing $\langle L, S, S \rangle$ at vertices x, y, z :
- x **source** \Rightarrow insert **edge** (z, x)
- x **sink** \Rightarrow insert **edge** (x, z) .



Refinement Algorithm – $\Phi, F, f_0 \rightarrow \text{st-digraph}$

Let f be a face. Consider the clockwise angle sequence σ_f of L/S on local **sources** and **sinks** of f .


- Goal: Add edges to break **large angles** (**sources** and **sinks**).
- For $f \neq f_0$ with $|\sigma_f| \geq 2$ containing $\langle L, S, S \rangle$ at vertices x, y, z :
- x **source** \Rightarrow insert **edge** (z, x)
- x **sink** \Rightarrow insert **edge** (x, z) .



Refinement Algorithm – $\Phi, F, f_0 \rightarrow \text{st-digraph}$

Let f be a face. Consider the clockwise angle sequence σ_f of L/S on local **sources** and **sinks** of f .


- Goal: Add edges to break **large angles** (**sources** and **sinks**).
- For $f \neq f_0$ with $|\sigma_f| \geq 2$ containing $\langle L, S, S \rangle$ at vertices x, y, z :
- x **source** \Rightarrow insert **edge** (z, x)
- x **sink** \Rightarrow insert **edge** (x, z) .
- Refine outer face f_0 .



Refinement Algorithm – $\Phi, F, f_0 \rightarrow \text{st-digraph}$

Let f be a face. Consider the clockwise angle sequence σ_f of L/S on local **sources** and **sinks** of f .

- Goal: Add edges to break **large angles** (**sources** and **sinks**).
- For $f \neq f_0$ with $|\sigma_f| \geq 2$ containing $\langle L, S, S \rangle$ at vertices x, y, z :
- x **source** \Rightarrow insert **edge** (z, x)
- x **sink** \Rightarrow insert **edge** (x, z) .
- Refine outer face f_0 .




- Refine all faces. \Rightarrow G is contained in a planar st-digraph.

Refinement Algorithm – $\Phi, F, f_0 \rightarrow \text{st-digraph}$


Let f be a face. Consider the clockwise angle sequence σ_f of L/S on local **sources** and **sinks** of f .

- Goal: Add edges to break **large angles** (**sources** and **sinks**).
- For $f \neq f_0$ with $|\sigma_f| \geq 2$ containing $\langle L, S, S \rangle$ at vertices x, y, z :
- x **source** \Rightarrow insert **edge** (z, x)
- x **sink** \Rightarrow insert **edge** (x, z) .
- Refine outer face f_0 .




- Refine all faces. \Rightarrow G is contained in a planar st-digraph.
- Planarity, acyclicity, bimodality are invariants under construction.


Refinement Example




Refinement Example




Refinement Example




Refinement Example




Refinement Example




Refinement Example




Refinement Example




Refinement Example




Refinement Example




Refinement Example




Refinement Example




Refinement Example




Refinement Example



Refinement Example



Refinement Example



Result Upward Planarity Test

Theorem 2.

[Bertolazzi et al., 1994]

For a *combinatorially embedded* planar digraph G it can be tested in $\mathcal{O}(n^2)$ time whether it is upward planar.

Result Upward Planarity Test

Theorem 2.

[Bertolazzi et al., 1994]

For a *combinatorially embedded* planar digraph G it can be tested in $\mathcal{O}(n^2)$ time whether it is upward planar.

Proof.

- Test for bimodality.

Result Upward Planarity Test

Theorem 2.

[Bertolazzi et al., 1994]

For a *combinatorially embedded* planar digraph G it can be tested in $\mathcal{O}(n^2)$ time whether it is upward planar.

Proof.

- Test for bimodality.
- Test for a consistent assignment Φ (via flow network).

Result Upward Planarity Test

Theorem 2.

[Bertolazzi et al., 1994]

For a *combinatorially embedded* planar digraph G it can be tested in $\mathcal{O}(n^2)$ time whether it is upward planar.

Proof.

- Test for bimodality.
- Test for a consistent assignment Φ (via flow network).
- If G bimodal and Φ exists, refine G to plane st-digraph H .

Result Upward Planarity Test

Theorem 2.

[Bertolazzi et al., 1994]

For a *combinatorially embedded* planar digraph G it can be tested in $\mathcal{O}(n^2)$ time whether it is upward planar.

Proof.

- Test for bimodality.
- Test for a consistent assignment Φ (via flow network).
- If G bimodal and Φ exists, refine G to plane st-digraph H .
- Draw H upward planar.

Result Upward Planarity Test

Theorem 2.

[Bertolazzi et al., 1994]

For a *combinatorially embedded* planar digraph G it can be tested in $\mathcal{O}(n^2)$ time whether it is upward planar.

Proof.

- Test for bimodality.
- Test for a consistent assignment Φ (via flow network).
- If G bimodal and Φ exists, refine G to plane st-digraph H .
- Draw H upward planar.
- Deleted edges added in refinement step.

Discussion

- There exist fixed-parameter tractable algorithms to test upward planarity of general digraphs with the parameter being the number of triconnected components.

[Healy, Lynch 2005, Didimo et al. 2009]

Discussion

- There exist fixed-parameter tractable algorithms to test upward planarity of general digraphs with the parameter being the number of triconnected components.
[Healy, Lynch 2005, Didimo et al. 2009]
- Finding assignment in Theorem 2 can be sped up to $\mathcal{O}(n + r^{1.5})$ where $r = \# \text{sources} / \text{sinks}$.
[Abbasi, Healy, Rextin 2010]

Discussion

- There exist fixed-parameter tractable algorithms to test upward planarity of general digraphs with the parameter being the number of triconnected components.
[Healy, Lynch 2005, Didimo et al. 2009]
- Finding assignment in Theorem 2 can be sped up to $\mathcal{O}(n + r^{1.5})$ where $r = \# \text{sources} / \text{sinks}$.
[Abbasi, Healy, Rextin 2010]
- Many related concepts have been studied: quasi-planarity, upward drawings of mixed graphs, upward planarity on cylinder/torus, ...