



CS F402: Computational Geometry

Lecture 10: GD - Force-Directed Drawing Algorithms-II



Siddharth Gupta

February 14+17+19, 2025



Variant by Fruchterman & Reingold

■ Repulsive forces

repulsion constant (e.g. 2.0)

$$f_{\text{rep}}(u, v) = \frac{c_{\text{rep}}}{||p_v - p_u||^2} \cdot \overrightarrow{p_v p_u}$$

■ Attractive forces

spring constant (e.g. 1.0)

$$f_{\text{spring}}(u, v) = c_{\text{spring}} \cdot \log \frac{||p_v - p_u||}{\ell} \cdot \overrightarrow{p_u p_v}$$

$$f_{\text{attr}}(u, v) = f_{\text{spring}}(u, v) - f_{\text{rep}}(u, v)$$

■ Resulting displacement vector

$$F_u = \sum_{v \in V} f_{\text{rep}}(u, v) + \sum_{uv \in E} f_{\text{attr}}(u, v)$$

```
ForceDirected( $G = (V, E)$ ,  $p = (p_v)_{v \in V}$ ,  $\varepsilon > 0$ ,  $K \in \mathbb{N}$ )
   $t \leftarrow 1$ 
  while  $t < K$  and  $\max_{v \in V} \|F_v(t)\| > \varepsilon$  do
    foreach  $u \in V$  do
       $F_u(t) \leftarrow \sum_{v \in V} f_{\text{rep}}(u, v) + \sum_{uv \in E} f_{\text{attr}}(u, v)$ 
    foreach  $u \in V$  do
       $p_u \leftarrow p_u + \delta(t) \cdot F_u(t)$ 
     $t \leftarrow t + 1$ 
  return  $p$ 
```

Notation.

- $||p_u - p_v||$ = Euclidean distance between u and v
- $\overrightarrow{p_u p_v}$ = unit vector pointing from u to v
- ℓ = ideal spring length for edges

Variant by Fruchterman & Reingold

■ Repulsive forces

$$f_{\text{rep}}(u, v) = \frac{\ell^2}{||p_v - p_u||} \cdot \overrightarrow{p_v p_u}$$

■ Attractive forces

spring constant (e.g. 1.0)

$$f_{\text{spring}}(u, v) = c_{\text{spring}} \cdot \log \frac{||p_v - p_u||}{\ell} \cdot \overrightarrow{p_u p_v}$$

$$f_{\text{attr}}(u, v) = f_{\text{spring}}(u, v) - f_{\text{rep}}(u, v)$$

■ Resulting displacement vector

$$F_u = \sum_{v \in V} f_{\text{rep}}(u, v) + \sum_{uv \in E} f_{\text{attr}}(u, v)$$

```
ForceDirected( $G = (V, E)$ ,  $p = (p_v)_{v \in V}$ ,  $\varepsilon > 0$ ,  $K \in \mathbb{N}$ )
   $t \leftarrow 1$ 
  while  $t < K$  and  $\max_{v \in V} \|F_v(t)\| > \varepsilon$  do
    foreach  $u \in V$  do
       $F_u(t) \leftarrow \sum_{v \in V} f_{\text{rep}}(u, v) + \sum_{uv \in E} f_{\text{attr}}(u, v)$ 
    foreach  $u \in V$  do
       $p_u \leftarrow p_u + \delta(t) \cdot F_u(t)$ 
     $t \leftarrow t + 1$ 
  return  $p$ 
```

Notation.

- $||p_u - p_v||$ = Euclidean distance between u and v
- $\overrightarrow{p_u p_v}$ = unit vector pointing from u to v
- ℓ = ideal spring length for edges

Variant by Fruchterman & Reingold

■ Repulsive forces

$$f_{\text{rep}}(u, v) = \frac{\ell^2}{||p_v - p_u||} \cdot \overrightarrow{p_v p_u}$$

■ Attractive forces

$$f_{\text{attr}}(u, v) = \frac{||p_v - p_u||^2}{\ell} \cdot \overrightarrow{p_u p_v}$$

■ Resulting displacement vector


$$F_u = \sum_{v \in V} f_{\text{rep}}(u, v) + \sum_{uv \in E} f_{\text{attr}}(u, v)$$

```
ForceDirected( $G = (V, E)$ ,  $p = (p_v)_{v \in V}$ ,  $\varepsilon > 0$ ,  $K \in \mathbb{N}$ )
   $t \leftarrow 1$ 
  while  $t < K$  and  $\max_{v \in V} \|F_v(t)\| > \varepsilon$  do
    foreach  $u \in V$  do
       $F_u(t) \leftarrow \sum_{v \in V} f_{\text{rep}}(u, v) + \sum_{uv \in E} f_{\text{attr}}(u, v)$ 
    foreach  $u \in V$  do
       $p_u \leftarrow p_u + \delta(t) \cdot F_u(t)$ 
     $t \leftarrow t + 1$ 
  return  $p$ 
```


Notation.

- $||p_u - p_v||$ = Euclidean distance between u and v
- $\overrightarrow{p_u p_v}$ = unit vector pointing from u to v
- ℓ = ideal spring length for edges

Fruchterman & Reingold – Force Diagram




Fruchterman & Reingold – Force Diagram




$$f_{\text{rep}}(u, v) = \frac{\ell^2}{\|p_v - p_u\|} \cdot \overrightarrow{p_v p_u}$$


Fruchterman & Reingold – Force Diagram



Fruchterman & Reingold – Force Diagram




Tutte Algorithm: Idea



William T. Tutte
1917 – 2002

Tutte Algorithm: Idea



William T. Tutte
1917 – 2002

HOW TO DRAW A GRAPH

By W. T. TUTTE

[Received 22 May 1962]

1. Introduction



We use the definitions of (11). However, in deference to some recent attempts to unify the terminology of graph theory we replace the term ‘circuit’ by ‘polygon’, and ‘degree’ by ‘valency’.

A graph G is 3-connected (*nodally* 3-connected) if it is simple and non-separable and satisfies the following condition; if G is the union of two proper subgraphs H and K such that $H \cap K$ consists solely of two vertices u and v , then one of H and K is a link-graph (arc-graph) with ends u and v .

It should be noted that the union of two proper subgraphs H and K of G can be the whole of G only if each of H and K includes at least one edge or vertex not belonging to the other. In this paper we are concerned mainly with nodally 3-connected graphs, but a specialization to 3-connected graphs is made in § 12.

Tutte Algorithm: Idea

Consider a fixed triangle (a, b, c)



William T. Tutte
1917 – 2002

HOW TO DRAW A GRAPH

By W. T. TUTTE

[Received 22 May 1962]

1. Introduction



We use the definitions of (11). However, in deference to some recent attempts to unify the terminology of graph theory we replace the term ‘circuit’ by ‘polygon’, and ‘degree’ by ‘valency’.

A graph G is 3-connected (*nodally 3-connected*) if it is simple and non-separable and satisfies the following condition; if G is the union of two proper subgraphs H and K such that $H \cap K$ consists solely of two vertices u and v , then one of H and K is a link-graph (arc-graph) with ends u and v .

It should be noted that the union of two proper subgraphs H and K of G can be the whole of G only if each of H and K includes at least one edge or vertex not belonging to the other. In this paper we are concerned mainly with nodally 3-connected graphs, but a specialization to 3-connected graphs is made in § 12.

Tutte Algorithm: Idea

Consider a fixed triangle (a, b, c) with one common neighbor v



William T. Tutte
1917 – 2002

HOW TO DRAW A GRAPH

By W. T. TUTTE

[Received 22 May 1962]

1. Introduction

We use the definitions of (11). However, in deference to some recent attempts to unify the terminology of graph theory we replace the term ‘circuit’ by ‘polygon’, and ‘degree’ by ‘valency’.



A graph G is 3-connected (*nodally 3-connected*) if it is simple and non-separable and satisfies the following condition; if G is the union of two proper subgraphs H and K such that $H \cap K$ consists solely of two vertices u and v , then one of H and K is a link-graph (arc-graph) with ends u and v .

It should be noted that the union of two proper subgraphs H and K of G can be the whole of G only if each of H and K includes at least one edge or vertex not belonging to the other. In this paper we are concerned mainly with nodally 3-connected graphs, but a specialization to 3-connected graphs is made in § 12.

Tutte Algorithm: Idea

Consider a fixed triangle (a, b, c) with one common neighbor v

Where would you place v ?



William T. Tutte
1917 – 2002

HOW TO DRAW A GRAPH

By W. T. TUTTE

[Received 22 May 1962]

1. Introduction

We use the definitions of (11). However, in deference to some recent attempts to unify the terminology of graph theory we replace the term 'circuit' by 'polygon', and 'degree' by 'valency'.



A graph G is 3-connected (*nodally 3-connected*) if it is simple and non-separable and satisfies the following condition; if G is the union of two proper subgraphs H and K such that $H \cap K$ consists solely of two vertices u and v , then one of H and K is a link-graph (arc-graph) with ends u and v .

It should be noted that the union of two proper subgraphs H and K of G can be the whole of G only if each of H and K includes at least one edge or vertex not belonging to the other. In this paper we are concerned mainly with nodally 3-connected graphs, but a specialization to 3-connected graphs is made in § 12.

Tutte Algorithm: Idea

Consider a fixed triangle (a, b, c) with one common neighbor v

Where would you place v ?



William T. Tutte
1917 – 2002

HOW TO DRAW A GRAPH

By W. T. TUTTE

[Received 22 May 1962]

1. Introduction

We use the definitions of (11). However, in deference to some recent attempts to unify the terminology of graph theory we replace the term ‘circuit’ by ‘polygon’, and ‘degree’ by ‘valency’.



A graph G is 3-connected (*nodally 3-connected*) if it is simple and non-separable and satisfies the following condition; if G is the union of two proper subgraphs H and K such that $H \cap K$ consists solely of two vertices u and v , then one of H and K is a link-graph (arc-graph) with ends u and v .

It should be noted that the union of two proper subgraphs H and K of G can be the whole of G only if each of H and K includes at least one edge or vertex not belonging to the other. In this paper we are concerned mainly with nodally 3-connected graphs, but a specialization to 3-connected graphs is made in § 12.

Tutte Algorithm: Idea

Consider a fixed triangle (a, b, c) with one common neighbor v

Where would you place v ?



William T. Tutte
1917 – 2002

HOW TO DRAW A GRAPH

By W. T. TUTTE

[Received 22 May 1962]

1. Introduction

We use the definitions of (11). However, in deference to some recent attempts to unify the terminology of graph theory we replace the term 'circuit' by 'polygon', and 'degree' by 'valency'.



A graph G is 3-connected (*nodally 3-connected*) if it is simple and non-separable and satisfies the following condition; if G is the union of two proper subgraphs H and K such that $H \cap K$ consists solely of two vertices u and v , then one of H and K is a link-graph (arc-graph) with ends u and v .

It should be noted that the union of two proper subgraphs H and K of G can be the whole of G only if each of H and K includes at least one edge or vertex not belonging to the other. In this paper we are concerned mainly with nodally 3-connected graphs, but a specialization to 3-connected graphs is made in § 12.

Tutte Algorithm: Idea

Consider a fixed triangle (a, b, c) with one common neighbor v

Where would you place v ?



William T. Tutte
1917 – 2002

HOW TO DRAW A GRAPH

By W. T. TUTTE

[Received 22 May 1962]

1. Introduction

We use the definitions of (11). However, in deference to some recent attempts to unify the terminology of graph theory we replace the term ‘circuit’ by ‘polygon’, and ‘degree’ by ‘valency’.



A graph G is 3-connected (*nodally 3-connected*) if it is simple and non-separable and satisfies the following condition; if G is the union of two proper subgraphs H and K such that $H \cap K$ consists solely of two vertices u and v , then one of H and K is a link-graph (arc-graph) with ends u and v .

It should be noted that the union of two proper subgraphs H and K of G can be the whole of G only if each of H and K includes at least one edge or vertex not belonging to the other. In this paper we are concerned mainly with nodally 3-connected graphs, but a specialization to 3-connected graphs is made in § 12.

Tutte Algorithm: Idea

Consider a fixed triangle (a, b, c) with one common neighbor v

Where would you place v ?



William T. Tutte
1917 – 2002

HOW TO DRAW A GRAPH

By W. T. TUTTE

[Received 22 May 1962]

1. Introduction

We use the definitions of (11). However, in deference to some recent attempts to unify the terminology of graph theory we replace the term 'circuit' by 'polygon', and 'degree' by 'valency'.


A graph G is 3-connected (*nodally 3-connected*) if it is simple and non-separable and satisfies the following condition; if G is the union of two proper subgraphs H and K such that $H \cap K$ consists solely of two vertices u and v , then one of H and K is a link-graph (arc-graph) with ends u and v .

It should be noted that the union of two proper subgraphs H and K of G can be the whole of G only if each of H and K includes at least one edge or vertex not belonging to the other. In this paper we are concerned mainly with nodally 3-connected graphs, but a specialization to 3-connected graphs is made in § 12.


Tutte Algorithm: Idea

Consider a fixed triangle (a, b, c) with one common neighbor v

Where would you place v ?



barycenter(x_1, \dots, x_k) = ?



William T. Tutte
1917 – 2002

HOW TO DRAW A GRAPH

By W. T. TUTTE

[Received 22 May 1962]

1. Introduction

We use the definitions of (11). However, in deference to some recent attempts to unify the terminology of graph theory we replace the term 'circuit' by 'polygon', and 'degree' by 'valency'.


A graph G is 3-connected (nodally 3-connected) if it is simple and non-separable and satisfies the following condition; if G is the union of two proper subgraphs H and K such that $H \cap K$ consists solely of two vertices u and v , then one of H and K is a link-graph (arc-graph) with ends u and v .

It should be noted that the union of two proper subgraphs H and K of G can be the whole of G only if each of H and K includes at least one edge or vertex not belonging to the other. In this paper we are concerned mainly with nodally 3-connected graphs, but a specialization to 3-connected graphs is made in § 12.


Tutte Algorithm: Idea

Consider a fixed triangle (a, b, c) with one common neighbor v

Where would you place v ?



$$\text{barycenter}(x_1, \dots, x_k) = \sum_{i=1}^k x_i / k$$



William T. Tutte
1917 – 2002

HOW TO DRAW A GRAPH

By W. T. TUTTE

[Received 22 May 1962]

1. Introduction

We use the definitions of (11). However, in deference to some recent attempts to unify the terminology of graph theory we replace the term 'circuit' by 'polygon', and 'degree' by 'valency'.


A graph G is 3-connected (nodally 3-connected) if it is simple and non-separable and satisfies the following condition; if G is the union of two proper subgraphs H and K such that $H \cap K$ consists solely of two vertices u and v , then one of H and K is a link-graph (arc-graph) with ends u and v .

It should be noted that the union of two proper subgraphs H and K of G can be the whole of G only if each of H and K includes at least one edge or vertex not belonging to the other. In this paper we are concerned mainly with nodally 3-connected graphs, but a specialization to 3-connected graphs is made in § 12.

Tutte Algorithm: Idea

Consider a fixed triangle (a, b, c) with one common neighbor v


Where would you place v ?



$$\text{barycenter}(x_1, \dots, x_k) = \sum_{i=1}^k x_i / k$$

Idea.

Repeatedly place every vertex at barycenter of neighbors.



William T. Tutte
1917 – 2002

HOW TO DRAW A GRAPH

By W. T. TUTTE

[Received 22 May 1962]

1. Introduction

We use the definitions of (11). However, in deference to some recent attempts to unify the terminology of graph theory we replace the term 'circuit' by 'polygon', and 'degree' by 'valency'.

A graph G is 3-connected (nodally 3-connected) if it is simple and non-separable and satisfies the following condition; if G is the union of two proper subgraphs H and K such that $H \cap K$ consists solely of two vertices u and v , then one of H and K is a link-graph (arc-graph) with ends u and v .

It should be noted that the union of two proper subgraphs H and K of G can be the whole of G only if each of H and K includes at least one edge or vertex not belonging to the other. In this paper we are concerned mainly with nodally 3-connected graphs, but a specialization to 3-connected graphs is made in § 12.

Tutte's Forces

```

ForceDirected( $G = (V, E)$ ,  $p = (p_v)_{v \in V}$ ,  $\varepsilon > 0$ ,  $K \in \mathbb{N}$ )
   $t \leftarrow 1$ 
  while  $t < K$  and  $\max_{v \in V} \|F_v(t)\| > \varepsilon$  do
    foreach  $u \in V$  do
       $F_u(t) \leftarrow \sum_{v \in V} f_{\text{rep}}(u, v) + \sum_{uv \in E} f_{\text{attr}}(u, v)$ 
    foreach  $u \in V$  do
       $p_u \leftarrow p_u + \delta(t) \cdot F_u(t)$ 
     $t \leftarrow t + 1$ 
  return  $p$ 

```

Tutte's Forces

ForceDirected($G = (V, E)$, $p = (p_v)_{v \in V}$, $\varepsilon > 0$, $K \in \mathbb{N}$)

```

 $t \leftarrow 1$ 
while  $t < K$  and  $\max_{v \in V} \|F_v(t)\| > \varepsilon$  do
    foreach  $u \in V$  do
         $F_u(t) \leftarrow \sum_{v \in V} f_{\text{rep}}(u, v) + \sum_{uv \in E} f_{\text{attr}}(u, v)$ 
    foreach  $u \in V$  do
         $p_u \leftarrow p_u + \cancel{\delta(t)} \cdot 1 \cdot F_u(t)$ 
     $t \leftarrow t + 1$ 
return  $p$ 

```

Tutte's Forces

Goal.

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v)$$

ForceDirected($G = (V, E)$, $p = (p_v)_{v \in V}$, $\varepsilon > 0$, $K \in \mathbb{N}$)

```

 $t \leftarrow 1$ 
while  $t < K$  and  $\max_{v \in V} \|F_v(t)\| > \varepsilon$  do
    foreach  $u \in V$  do
         $F_u(t) \leftarrow \sum_{v \in V} f_{\text{rep}}(u, v) + \sum_{uv \in E} f_{\text{attr}}(u, v)$ 
    foreach  $u \in V$  do
         $p_u \leftarrow p_u + \cancel{\delta(t)} \cdot F_u(t)$ 
     $t \leftarrow t + 1$ 
return  $p$ 

```

barycenter(x_1, \dots, x_k) = $\sum_{i=1}^k x_i / k$

Tutte's Forces

Goal.

$$\begin{aligned} p_u &= \text{barycenter}(\bigcup_{uv \in E} v) \\ &= \sum_{uv \in E} p_v / \end{aligned}$$

ForceDirected($G = (V, E)$, $p = (p_v)_{v \in V}$, $\varepsilon > 0$, $K \in \mathbb{N}$)

```

 $t \leftarrow 1$ 
while  $t < K$  and  $\max_{v \in V} \|F_v(t)\| > \varepsilon$  do
    foreach  $u \in V$  do
         $F_u(t) \leftarrow \sum_{v \in V} f_{\text{rep}}(u, v) + \sum_{uv \in E} f_{\text{attr}}(u, v)$ 
    foreach  $u \in V$  do
         $p_u \leftarrow p_u + \cancel{\delta(t)} \cdot 1 \cdot F_u(t)$ 
     $t \leftarrow t + 1$ 
return  $p$ 

```

barycenter(x_1, \dots, x_k) = $\sum_{i=1}^k x_i / k$

Tutte's Forces

Goal.

$$\begin{aligned} p_u &= \text{barycenter}(\bigcup_{uv \in E} v) \\ &= \sum_{uv \in E} p_v / \deg(u) \end{aligned}$$

ForceDirected($G = (V, E)$, $p = (p_v)_{v \in V}$, $\varepsilon > 0$, $K \in \mathbb{N}$)

```

 $t \leftarrow 1$ 
while  $t < K$  and  $\max_{v \in V} \|F_v(t)\| > \varepsilon$  do
    foreach  $u \in V$  do
         $F_u(t) \leftarrow \sum_{v \in V} f_{\text{rep}}(u, v) + \sum_{uv \in E} f_{\text{attr}}(u, v)$ 
    foreach  $u \in V$  do
         $p_u \leftarrow p_u + \cancel{\delta(t)} \cdot 1 \cdot F_u(t)$ 
     $t \leftarrow t + 1$ 
return  $p$ 

```

barycenter(x_1, \dots, x_k) = $\sum_{i=1}^k x_i / k$

Tutte's Forces

Goal.

$$\begin{aligned} p_u &= \text{barycenter}(\bigcup_{uv \in E} v) \\ &= \sum_{uv \in E} p_v / \deg(u) \end{aligned}$$

$$F_u(t) = \sum_{uv \in E} p_v / \deg(u) - p_u$$

ForceDirected($G = (V, E)$, $p = (p_v)_{v \in V}$, $\varepsilon > 0$, $K \in \mathbb{N}$)

```

 $t \leftarrow 1$ 
while  $t < K$  and  $\max_{v \in V} \|F_v(t)\| > \varepsilon$  do
    foreach  $u \in V$  do
         $F_u(t) \leftarrow \sum_{v \in V} f_{\text{rep}}(u, v) + \sum_{uv \in E} f_{\text{attr}}(u, v)$ 
    foreach  $u \in V$  do
         $p_u \leftarrow p_u + \cancel{\delta(t)} \cdot 1 \cdot F_u(t)$ 
     $t \leftarrow t + 1$ 
return  $p$ 

```

barycenter(x_1, \dots, x_k) = $\sum_{i=1}^k x_i / k$

Tutte's Forces

Goal.

$$\begin{aligned} p_u &= \text{barycenter}(\bigcup_{uv \in E} v) \\ &= \sum_{uv \in E} p_v / \deg(u) \end{aligned}$$

$$\begin{aligned} F_u(t) &= \sum_{uv \in E} p_v / \deg(u) - p_u \\ &= \sum_{uv \in E} (p_v - p_u) / \deg(u) \end{aligned}$$

ForceDirected($G = (V, E)$, $p = (p_v)_{v \in V}$, $\varepsilon > 0$, $K \in \mathbb{N}$)

```

 $t \leftarrow 1$ 
while  $t < K$  and  $\max_{v \in V} \|F_v(t)\| > \varepsilon$  do
    foreach  $u \in V$  do
         $F_u(t) \leftarrow \sum_{v \in V} f_{\text{rep}}(u, v) + \sum_{uv \in E} f_{\text{attr}}(u, v)$ 
    foreach  $u \in V$  do
         $p_u \leftarrow p_u + \cancel{\delta(t)} \cdot 1 \cdot F_u(t)$ 
     $t \leftarrow t + 1$ 
return  $p$ 

```

barycenter(x_1, \dots, x_k) = $\sum_{i=1}^k x_i / k$

Tutte's Forces

Goal.

$$\begin{aligned} p_u &= \text{barycenter}(\bigcup_{uv \in E} v) \\ &= \sum_{uv \in E} p_v / \deg(u) \end{aligned}$$

$$\begin{aligned} F_u(t) &= \sum_{uv \in E} p_v / \deg(u) - p_u \\ &= \sum_{uv \in E} (p_v - p_u) / \deg(u) \\ &= \sum_{uv \in E} \|p_u - p_v\| / \deg(u) \end{aligned}$$

ForceDirected($G = (V, E)$, $p = (p_v)_{v \in V}$, $\varepsilon > 0$, $K \in \mathbb{N}$)

```

 $t \leftarrow 1$ 
while  $t < K$  and  $\max_{v \in V} \|F_v(t)\| > \varepsilon$  do
    foreach  $u \in V$  do
         $F_u(t) \leftarrow \sum_{v \in V} f_{\text{rep}}(u, v) + \sum_{uv \in E} f_{\text{attr}}(u, v)$ 
    foreach  $u \in V$  do
         $p_u \leftarrow p_u + \cancel{\delta(t)} \cdot 1 \cdot F_u(t)$ 
     $t \leftarrow t + 1$ 
return  $p$ 

```

barycenter(x_1, \dots, x_k) = $\sum_{i=1}^k x_i / k$

Tutte's Forces

Goal.

$$\begin{aligned} p_u &= \text{barycenter}(\bigcup_{uv \in E} v) \\ &= \sum_{uv \in E} p_v / \deg(u) \end{aligned}$$

$$\begin{aligned} F_u(t) &= \sum_{uv \in E} p_v / \deg(u) - p_u \\ &= \sum_{uv \in E} (p_v - p_u) / \deg(u) \\ &= \sum_{uv \in E} \|p_u - p_v\| / \deg(u) \end{aligned}$$

ForceDirected($G = (V, E)$, $p = (p_v)_{v \in V}$, $\varepsilon > 0$, $K \in \mathbb{N}$)

```

 $t \leftarrow 1$ 
while  $t < K$  and  $\max_{v \in V} \|F_v(t)\| > \varepsilon$  do
    foreach  $u \in V$  do
         $F_u(t) \leftarrow \sum_{v \in V} f_{\text{rep}}(u, v) + \sum_{uv \in E} f_{\text{attr}}(u, v)$ 
    foreach  $u \in V$  do
         $p_u \leftarrow p_u + \cancel{\delta(t)} \cdot 1 \cdot F_u(t)$ 
     $t \leftarrow t + 1$ 
return  $p$ 

```

barycenter(x_1, \dots, x_k) = $\sum_{i=1}^k x_i / k$

Tutte's Forces

Goal.

$$\begin{aligned} p_u &= \text{barycenter}(\bigcup_{uv \in E} v) \\ &= \sum_{uv \in E} p_v / \deg(u) \end{aligned}$$

$$\begin{aligned} F_u(t) &= \sum_{uv \in E} p_v / \deg(u) - p_u \\ &= \sum_{uv \in E} (p_v - p_u) / \deg(u) \\ &= \sum_{uv \in E} \|p_u - p_v\| / \deg(u) \end{aligned}$$

■ Repulsive forces

$$f_{\text{rep}}(u, v) = 0$$

```
ForceDirected( $G = (V, E)$ ,  $p = (p_v)_{v \in V}$ ,  $\varepsilon > 0$ ,  $K \in \mathbb{N}$ )
   $t \leftarrow 1$ 
  while  $t < K$  and  $\max_{v \in V} \|F_v(t)\| > \varepsilon$  do
    foreach  $u \in V$  do
       $F_u(t) \leftarrow \sum_{v \in V} f_{\text{rep}}(u, v) + \sum_{uv \in E} f_{\text{attr}}(u, v)$ 
    foreach  $u \in V$  do
       $p_u \leftarrow p_u + \cancel{\delta(t)} \cdot 1 \cdot F_u(t)$ 
     $t \leftarrow t + 1$ 
  return  $p$ 
barycenter( $x_1, \dots, x_k$ ) =  $\sum_{i=1}^k x_i / k$ 
```

Tutte's Forces

Goal.

$$\begin{aligned} p_u &= \text{barycenter}(\bigcup_{uv \in E} v) \\ &= \sum_{uv \in E} p_v / \deg(u) \end{aligned}$$

$$\begin{aligned} F_u(t) &= \sum_{uv \in E} p_v / \deg(u) - p_u \\ &= \sum_{uv \in E} (p_v - p_u) / \deg(u) \\ &= \sum_{uv \in E} \|p_u - p_v\| / \deg(u) \end{aligned}$$

```

ForceDirected( $G = (V, E)$ ,  $p = (p_v)_{v \in V}$ ,  $\varepsilon > 0$ ,  $K \in \mathbb{N}$ )
   $t \leftarrow 1$ 
  while  $t < K$  and  $\max_{v \in V} \|F_v(t)\| > \varepsilon$  do
    foreach  $u \in V$  do
       $F_u(t) \leftarrow \sum_{v \in V} f_{\text{rep}}(u, v) + \sum_{uv \in E} f_{\text{attr}}(u, v)$ 
    foreach  $u \in V$  do
       $p_u \leftarrow p_u + \cancel{\delta(t)} \cdot 1 \cdot F_u(t)$ 
     $t \leftarrow t + 1$ 
  return  $p$ 

```

$\text{barycenter}(x_1, \dots, x_k) = \sum_{i=1}^k x_i / k$

■ Repulsive forces

$$f_{\text{rep}}(u, v) = 0$$

■ Attractive forces

$$f_{\text{attr}}(u, v) = \frac{1}{\deg(u)} \cdot \|p_u - p_v\|$$

Tutte's Forces

Goal.

$$\begin{aligned} p_u &= \text{barycenter}(\bigcup_{uv \in E} v) \\ &= \sum_{uv \in E} p_v / \deg(u) \end{aligned}$$

$$\begin{aligned} F_u(t) &= \sum_{uv \in E} p_v / \deg(u) - p_u \\ &= \sum_{uv \in E} (p_v - p_u) / \deg(u) \\ &= \sum_{uv \in E} \|p_u - p_v\| / \deg(u) \end{aligned}$$

ForceDirected($G = (V, E)$, $p = (p_v)_{v \in V}$, $\varepsilon > 0$, $K \in \mathbb{N}$)

```

 $t \leftarrow 1$ 
while  $t < K$  and  $\max_{v \in V} \|F_v(t)\| > \varepsilon$  do
    foreach  $u \in V$  do
         $F_u(t) \leftarrow \sum_{v \in V} f_{\text{rep}}(u, v) + \sum_{uv \in E} f_{\text{attr}}(u, v)$ 
    foreach  $u \in V$  do
         $p_u \leftarrow p_u + \cancel{\delta(t)} \cdot 1 \cdot F_u(t)$ 
     $t \leftarrow t + 1$ 
return  $p$ 

```

barycenter(x_1, \dots, x_k) = $\sum_{i=1}^k x_i / k$

■ Repulsive forces

Solution: $p_u = (0, 0) \ \forall u \in V$

$$f_{\text{rep}}(u, v) = 0$$

■ Attractive forces

$$f_{\text{attr}}(u, v) = \frac{1}{\deg(u)} \cdot \|p_u - p_v\|$$

Tutte's Forces

Goal.

$$\begin{aligned} p_u &= \text{barycenter}(\bigcup_{uv \in E} v) \\ &= \sum_{uv \in E} p_v / \deg(u) \end{aligned}$$

$$\begin{aligned} F_u(t) &= \sum_{uv \in E} p_v / \deg(u) - p_u \\ &= \sum_{uv \in E} (p_v - p_u) / \deg(u) \\ &= \sum_{uv \in E} \|p_u - p_v\| / \deg(u) \end{aligned}$$

```

ForceDirected( $G = (V, E)$ ,  $p = (p_v)_{v \in V}$ ,  $\varepsilon > 0$ ,  $K \in \mathbb{N}$ )
   $t \leftarrow 1$ 
  while  $t < K$  and  $\max_{v \in V} \|F_v(t)\| > \varepsilon$  do
    foreach  $u \in V$  do
       $F_u(t) \leftarrow \sum_{v \in V} f_{\text{rep}}(u, v) + \sum_{uv \in E} f_{\text{attr}}(u, v)$ 
    foreach  $u \in V$  do
       $p_u \leftarrow p_u + \cancel{\delta(t)} \cdot 1 \cdot F_u(t)$ 
     $t \leftarrow t + 1$ 
  return  $p$ 

```

barycenter(x_1, \dots, x_k) = $\sum_{i=1}^k x_i / k$


■ Repulsive forces

$$f_{\text{rep}}(u, v) = 0$$

■ Attractive forces

$$f_{\text{attr}}(u, v) = \frac{1}{\deg(u)} \cdot \|p_u - p_v\|$$

Solution: $p_u = (0, 0) \quad \forall u \in V$



Tutte's Forces

Goal.

$$\begin{aligned} p_u &= \text{barycenter}(\bigcup_{uv \in E} v) \\ &= \sum_{uv \in E} p_v / \deg(u) \end{aligned}$$

$$\begin{aligned} F_u(t) &= \sum_{uv \in E} p_v / \deg(u) - p_u \\ &= \sum_{uv \in E} (p_v - p_u) / \deg(u) \\ &= \sum_{uv \in E} \|p_u - p_v\| / \deg(u) \end{aligned}$$

ForceDirected($G = (V, E)$, $p = (p_v)_{v \in V}$, $\varepsilon > 0$, $K \in \mathbb{N}$)

```

 $t \leftarrow 1$ 
while  $t < K$  and  $\max_{v \in V} \|F_v(t)\| > \varepsilon$  do
    foreach  $u \in V$  do
         $F_u(t) \leftarrow \sum_{v \in V} f_{\text{rep}}(u, v) + \sum_{uv \in E} f_{\text{attr}}(u, v)$ 
    foreach  $u \in V$  do
         $p_u \leftarrow p_u + \cancel{\delta(t)} \cdot 1 \cdot F_u(t)$ 
     $t \leftarrow t + 1$ 
return  $p$ 

```

barycenter(x_1, \dots, x_k) = $\sum_{i=1}^k x_i / k$


■ Repulsive forces

$$f_{\text{rep}}(u, v) = 0$$

■ Attractive forces

$$f_{\text{attr}}(u, v) = \frac{1}{\deg(u)} \cdot \|p_u - p_v\|$$

Solution: $p_u = (0, 0) \forall u \in V$



Fix coordinates
of outer face!

Tutte's Forces

Goal.

$$\begin{aligned} p_u &= \text{barycenter}(\bigcup_{uv \in E} v) \\ &= \sum_{uv \in E} p_v / \deg(u) \end{aligned}$$

$$\begin{aligned} F_u(t) &= \sum_{uv \in E} p_v / \deg(u) - p_u \\ &= \sum_{uv \in E} (p_v - p_u) / \deg(u) \\ &= \sum_{uv \in E} \|p_u - p_v\| / \deg(u) \end{aligned}$$

```
ForceDirected( $G = (V, E)$ ,  $p = (p_v)_{v \in V}$ ,  $\varepsilon > 0$ ,  $K \in \mathbb{N}$ )
   $t \leftarrow 1$ 
  while  $t < K$  and  $\max_{v \in V} \|F_v(t)\| > \varepsilon$  do
    foreach  $u \in V$  do
       $F_u(t) \leftarrow \sum_{v \in V} f_{\text{rep}}(u, v) + \sum_{uv \in E} f_{\text{attr}}(u, v)$ 
    foreach  $u \in V$  do
       $p_u \leftarrow p_u + \cancel{\delta(t)} \cdot 1 \cdot F_u(t)$ 
     $t \leftarrow t + 1$ 
  return  $p$ 
```

barycenter(x_1, \dots, x_k) = $\sum_{i=1}^k x_i / k$


■ **Repulsive forces**

$$f_{\text{rep}}(u, v) = 0$$

■ **Attractive forces**

$$f_{\text{attr}}(u, v) = \begin{cases} 0 & u \text{ fixed} \\ \frac{1}{\deg(u)} \cdot \|p_u - p_v\| & \text{else} \end{cases}$$

Solution: $p_u = (0, 0) \forall u \in V$



Fix coordinates
of outer face!

Tutte's Forces

Goal.

$$\begin{aligned} p_u &= \text{barycenter}(\bigcup_{uv \in E} v) \\ &= \sum_{uv \in E} p_v / \deg(u) \end{aligned}$$

$$\begin{aligned} F_u(t) &= \sum_{uv \in E} p_v / \deg(u) - p_u \\ &= \sum_{uv \in E} (p_v - p_u) / \deg(u) \\ &= \sum_{uv \in E} \|p_u - p_v\| / \deg(u) \end{aligned}$$

```
ForceDirected( $G = (V, E)$ ,  $p = (p_v)_{v \in V}$ ,  $\varepsilon > 0$ ,  $K \in \mathbb{N}$ )
   $t \leftarrow 1$ 
  while  $t < K$  and  $\max_{v \in V} \|F_v(t)\| > \varepsilon$  do
    foreach  $u \in V$  do
       $F_u(t) \leftarrow \sum_{v \in V} f_{\text{rep}}(u, v) + \sum_{uv \in E} f_{\text{attr}}(u, v)$ 
    foreach  $u \in V$  do
       $p_u \leftarrow p_u + \cancel{\delta(t)} \cdot 1 \cdot F_u(t)$ 
     $t \leftarrow t + 1$ 
  return  $p$ 
```

$\text{barycenter}(x_1, \dots, x_k) = \sum_{i=1}^k x_i / k$


■ **Repulsive forces**

$$f_{\text{rep}}(u, v) = 0$$

■ **Attractive forces**

$$f_{\text{attr}}(u, v) = \begin{cases} 0 & u \text{ fixed} \\ \frac{1}{\deg(u)} \cdot \|p_u - p_v\| & \text{else} \end{cases}$$

Solution: $p_u = (0, 0) \forall u \in V$



Fix coordinates
of outer face!

Linear System of Equations

Goal.

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u)$$

$$y_u = \sum_{uv \in E} y_v / \deg(u)$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$

$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$

$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$

$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$

$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$

$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$

$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



A

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



A

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



	u_1	u_2	u_3	u_4	u_5	u_6	A
u_1	1	0	0	0	1	1	
u_2	0	1	0	1	0	0	
u_3	1	0	1	0	0	0	
u_4	0	1	0	1	0	0	
u_5	0	0	0	0	1	0	
u_6	1	0	0	1	0	1	

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



	u_1	u_2	u_3	u_4	u_5	u_6	A
u_1	3						
u_2							
u_3							
u_4							
u_5							
u_6							

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



$$A = \begin{pmatrix} u_1 & u_2 & u_3 & u_4 & u_5 & u_6 \\ u_1 & 3 & -1 & 0 & 0 & 0 \\ u_2 & -1 & 0 & 1 & 0 & 0 \\ u_3 & 0 & 1 & 0 & 1 & 0 \\ u_4 & 0 & 0 & 1 & 0 & 1 \\ u_5 & 0 & 0 & 0 & 1 & 0 \\ u_6 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



$$A = \begin{pmatrix} u_1 & u_2 & u_3 & u_4 & u_5 & u_6 \\ u_1 & 3 & -1 & -1 & & \\ u_2 & -1 & 3 & -1 & & \\ u_3 & -1 & -1 & 3 & & \\ u_4 & & & & 3 & -1 \\ u_5 & & & & -1 & 3 \\ u_6 & & & & & -1 \end{pmatrix}$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



$$A = \begin{pmatrix} u_1 & u_2 & u_3 & u_4 & u_5 & u_6 \\ u_1 & 3 & -1 & -1 & 0 & \\ u_2 & & & & & \\ u_3 & & & & & \\ u_4 & & & & & \\ u_5 & & & & & \\ u_6 & & & & & \end{pmatrix}$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



$$A = \begin{pmatrix} u_1 & u_2 & u_3 & u_4 & u_5 & u_6 \\ u_1 & 3 & -1 & -1 & 0 & -1 \\ u_2 & -1 & 3 & -1 & 0 & -1 \\ u_3 & -1 & -1 & 3 & 0 & -1 \\ u_4 & 0 & 0 & 0 & 3 & -1 \\ u_5 & -1 & -1 & -1 & -1 & 3 \\ u_6 & -1 & -1 & -1 & -1 & -1 & 3 \end{pmatrix}$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



$$A = \begin{pmatrix} u_1 & u_2 & u_3 & u_4 & u_5 & u_6 \\ u_1 & 3 & -1 & -1 & 0 & -1 & 0 \\ u_2 & & & & & & \\ u_3 & & & & & & \\ u_4 & & & & & & \\ u_5 & & & & & & \\ u_6 & & & & & & \end{pmatrix}$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



$$A = \begin{pmatrix} u_1 & u_2 & u_3 & u_4 & u_5 & u_6 \\ u_1 & 3 & -1 & -1 & 0 & -1 & 0 \\ u_2 & & 3 & & & & \\ u_3 & & & & & & \\ u_4 & & & & & & \\ u_5 & & & & & & \\ u_6 & & & & & & \end{pmatrix}$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



$$A = \begin{pmatrix} u_1 & u_2 & u_3 & u_4 & u_5 & u_6 \\ u_1 & 3 & -1 & -1 & 0 & -1 & 0 \\ u_2 & -1 & 3 & -1 & -1 & 0 & 0 \\ u_3 & & & & & & \\ u_4 & & & & & & \\ u_5 & & & & & & \\ u_6 & & & & & & \end{pmatrix}$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



$$A = \begin{pmatrix} u_1 & u_2 & u_3 & u_4 & u_5 & u_6 \\ u_1 & 3 & -1 & -1 & 0 & -1 & 0 \\ u_2 & -1 & 3 & -1 & -1 & 0 & 0 \\ u_3 & -1 & -1 & 3 & 0 & 0 & -1 \\ u_4 & & & & & & \\ u_5 & & & & & & \\ u_6 & & & & & & \end{pmatrix}$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



$$A = \begin{pmatrix} u_1 & u_2 & u_3 & u_4 & u_5 & u_6 \\ u_1 & 3 & -1 & -1 & 0 & -1 & 0 \\ u_2 & -1 & 3 & -1 & -1 & 0 & 0 \\ u_3 & -1 & -1 & 3 & 0 & 0 & -1 \\ u_4 & 0 & -1 & 0 & 3 & -1 & -1 \\ u_5 & & & & & & \\ u_6 & & & & & & \end{pmatrix}$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



$$A = \begin{pmatrix} u_1 & u_2 & u_3 & u_4 & u_5 & u_6 \\ u_1 & 3 & -1 & -1 & 0 & -1 & 0 \\ u_2 & -1 & 3 & -1 & -1 & 0 & 0 \\ u_3 & -1 & -1 & 3 & 0 & 0 & -1 \\ u_4 & 0 & -1 & 0 & 3 & -1 & -1 \\ u_5 & -1 & 0 & 0 & -1 & 2 & 0 \\ u_6 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



$$\begin{array}{cccccc}
 & u_1 & u_2 & u_3 & u_4 & u_5 & u_6 \\
 \begin{matrix} u_1 \\ u_2 \\ u_3 \\ u_4 \\ u_5 \\ u_6 \end{matrix} & \left(\begin{array}{cccccc}
 3 & -1 & -1 & 0 & -1 & 0 \\
 -1 & 3 & -1 & -1 & 0 & 0 \\
 -1 & -1 & 3 & 0 & 0 & -1 \\
 0 & -1 & 0 & 3 & -1 & -1 \\
 -1 & 0 & 0 & -1 & 2 & 0 \\
 0 & 0 & -1 & -1 & 0 & 2
 \end{array} \right)
 \end{array}$$

A

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



$$\begin{array}{cccccc}
 & u_1 & u_2 & u_3 & u_4 & u_5 & u_6 & A \\
 \begin{matrix} u_1 \\ u_2 \\ u_3 \\ u_4 \\ u_5 \\ u_6 \end{matrix} & \left(\begin{array}{cccccc}
 3 & -1 & -1 & 0 & -1 & 0 \\
 -1 & 3 & -1 & -1 & 0 & 0 \\
 -1 & -1 & 3 & 0 & 0 & -1 \\
 0 & -1 & 0 & 3 & -1 & -1 \\
 -1 & 0 & 0 & -1 & 2 & 0 \\
 0 & 0 & -1 & -1 & 0 & 2
 \end{array} \right)
 \end{array}$$

$A_{ii} = \deg(u_i)$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



$$A = \begin{pmatrix} u_1 & u_2 & u_3 & u_4 & u_5 & u_6 \\ u_1 & 3 & -1 & -1 & 0 & -1 & 0 \\ u_2 & -1 & 3 & -1 & -1 & 0 & 0 \\ u_3 & -1 & -1 & 3 & 0 & 0 & -1 \\ u_4 & 0 & -1 & 0 & 3 & -1 & -1 \\ u_5 & -1 & 0 & 0 & -1 & 2 & 0 \\ u_6 & 0 & 0 & -1 & -1 & 0 & 2 \end{pmatrix}$$

$$A_{ii} = \deg(u_i)$$

$$A_{ij, i \neq j} = \begin{cases} -1 & u_i u_j \in E \\ 0 & u_i u_j \notin E \end{cases}$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



$$A = \begin{pmatrix} u_1 & u_2 & u_3 & u_4 & u_5 & u_6 \\ u_1 & 3 & -1 & -1 & 0 & -1 & 0 \\ u_2 & -1 & 3 & -1 & -1 & 0 & 0 \\ u_3 & -1 & -1 & 3 & 0 & 0 & -1 \\ u_4 & 0 & -1 & 0 & 3 & -1 & -1 \\ u_5 & -1 & 0 & 0 & -1 & 2 & 0 \\ u_6 & 0 & 0 & -1 & -1 & 0 & 2 \end{pmatrix}$$

Laplacian matrix of G

$$A_{ii} = \deg(u_i)$$

$$A_{ij, i \neq j} = \begin{cases} -1 & u_i u_j \in E \\ 0 & u_i u_j \notin E \end{cases}$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



$$A = \begin{pmatrix} u_1 & u_2 & u_3 & u_4 & u_5 & u_6 \\ u_1 & 3 & -1 & -1 & 0 & -1 & 0 \\ u_2 & -1 & 3 & -1 & -1 & 0 & 0 \\ u_3 & -1 & -1 & 3 & 0 & 0 & -1 \\ u_4 & 0 & -1 & 0 & 3 & -1 & -1 \\ u_5 & -1 & 0 & 0 & -1 & 2 & 0 \\ u_6 & 0 & 0 & -1 & -1 & 0 & 2 \end{pmatrix}$$

Laplacian matrix of G

$$A_{ii} = \deg(u_i)$$

$$A_{ij, i \neq j} = \begin{cases} -1 & u_i u_j \in E \\ 0 & u_i u_j \notin E \end{cases}$$

unique solution

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



$$A = \begin{pmatrix} u_1 & u_2 & u_3 & u_4 & u_5 & u_6 \\ u_1 & 3 & -1 & -1 & 0 & -1 & 0 \\ u_2 & -1 & 3 & -1 & -1 & 0 & 0 \\ u_3 & -1 & -1 & 3 & 0 & 0 & -1 \\ u_4 & 0 & -1 & 0 & 3 & -1 & -1 \\ u_5 & -1 & 0 & 0 & -1 & 2 & 0 \\ u_6 & 0 & 0 & -1 & -1 & 0 & 2 \end{pmatrix}$$

Laplacian matrix of G

variables, constraints, $\det(A) =$
unique solution

$$A_{ii} = \deg(u_i)$$

$$A_{ij, i \neq j} = \begin{cases} -1 & u_i u_j \in E \\ 0 & u_i u_j \notin E \end{cases}$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



$$A = \begin{pmatrix} u_1 & u_2 & u_3 & u_4 & u_5 & u_6 \\ u_1 & 3 & -1 & -1 & 0 & -1 & 0 \\ u_2 & -1 & 3 & -1 & -1 & 0 & 0 \\ u_3 & -1 & -1 & 3 & 0 & 0 & -1 \\ u_4 & 0 & -1 & 0 & 3 & -1 & -1 \\ u_5 & -1 & 0 & 0 & -1 & 2 & 0 \\ u_6 & 0 & 0 & -1 & -1 & 0 & 2 \end{pmatrix}$$

Laplacian matrix of G

n variables, constraints, $\det(A) =$
unique solution

$$A_{ii} = \deg(u_i)$$

$$A_{ij, i \neq j} = \begin{cases} -1 & u_i u_j \in E \\ 0 & u_i u_j \notin E \end{cases}$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



$$A = \begin{pmatrix} u_1 & u_2 & u_3 & u_4 & u_5 & u_6 \\ u_1 & 3 & -1 & -1 & 0 & -1 & 0 \\ u_2 & -1 & 3 & -1 & -1 & 0 & 0 \\ u_3 & -1 & -1 & 3 & 0 & 0 & -1 \\ u_4 & 0 & -1 & 0 & 3 & -1 & -1 \\ u_5 & -1 & 0 & 0 & -1 & 2 & 0 \\ u_6 & 0 & 0 & -1 & -1 & 0 & 2 \end{pmatrix}$$

Laplacian matrix of G

n variables, n constraints, $\det(A) =$
unique solution

$$A_{ii} = \deg(u_i)$$

$$A_{ij, i \neq j} = \begin{cases} -1 & u_i u_j \in E \\ 0 & u_i u_j \notin E \end{cases}$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



$$A = \begin{pmatrix} u_1 & u_2 & u_3 & u_4 & u_5 & u_6 \\ u_1 & 3 & -1 & -1 & 0 & -1 & 0 \\ u_2 & -1 & 3 & -1 & -1 & 0 & 0 \\ u_3 & -1 & -1 & 3 & 0 & 0 & -1 \\ u_4 & 0 & -1 & 0 & 3 & -1 & -1 \\ u_5 & -1 & 0 & 0 & -1 & 2 & 0 \\ u_6 & 0 & 0 & -1 & -1 & 0 & 2 \end{pmatrix}$$

Laplacian matrix of G

n variables, n constraints, $\det(A) = 0$
unique solution

$$A_{ii} = \deg(u_i)$$

$$A_{ij, i \neq j} = \begin{cases} -1 & u_i u_j \in E \\ 0 & u_i u_j \notin E \end{cases}$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



$$A = \begin{pmatrix} u_1 & u_2 & u_3 & u_4 & u_5 & u_6 \\ u_1 & 3 & -1 & -1 & 0 & -1 & 0 \\ u_2 & -1 & 3 & -1 & -1 & 0 & 0 \\ u_3 & -1 & -1 & 3 & 0 & 0 & -1 \\ u_4 & 0 & -1 & 0 & 3 & -1 & -1 \\ u_5 & -1 & 0 & 0 & -1 & 2 & 0 \\ u_6 & 0 & 0 & -1 & -1 & 0 & 2 \end{pmatrix}$$

Laplacian matrix of G

n variables, n constraints, $\det(A) = 0$

\Rightarrow no unique solution



$$A_{ii} = \deg(u_i)$$

$$A_{ij, i \neq j} = \begin{cases} -1 & u_i u_j \in E \\ 0 & u_i u_j \notin E \end{cases}$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



$$A = \begin{pmatrix} u_1 & u_2 & u_3 & u_4 & u_5 & u_6 \\ u_1 & 3 & -1 & -1 & 0 & -1 & 0 \\ u_2 & -1 & 3 & -1 & -1 & 0 & 0 \\ u_3 & -1 & -1 & 3 & 0 & 0 & -1 \\ u_4 & 0 & -1 & 0 & 3 & -1 & -1 \\ u_5 & -1 & 0 & 0 & -1 & 2 & 0 \\ u_6 & 0 & 0 & -1 & -1 & 0 & 2 \end{pmatrix}$$

Laplacian matrix of G

n variables, n constraints, $\det(A) = 0$

\Rightarrow no unique solution



$$A_{ii} = \deg(u_i)$$

$$A_{ij, i \neq j} = \begin{cases} -1 & u_i u_j \in E \\ 0 & u_i u_j \notin E \end{cases}$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



$$A$$

$$\begin{pmatrix} u_1 & u_2 & u_3 & u_4 & u_5 & u_6 \\ u_1 & 3 & -1 & -1 & 0 & -1 & 0 \\ u_2 & -1 & 3 & -1 & -1 & 0 & 0 \\ u_3 & -1 & -1 & 3 & 0 & 0 & -1 \\ u_4 & 0 & -1 & 0 & 3 & -1 & -1 \\ u_5 & -1 & 0 & 0 & -1 & 2 & 0 \\ u_6 & 0 & 0 & -1 & -1 & 0 & 2 \end{pmatrix}$$

Laplacian matrix of G

n variables, k constraints, $\det(A) = 0$

$k = \#\text{free vertices}$

\Rightarrow no unique solution



$$A_{ii} = \deg(u_i)$$

$$A_{ij, i \neq j} = \begin{cases} -1 & u_i u_j \in E \\ 0 & u_i u_j \notin E \end{cases}$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



	u_1	u_2	u_3	u_4	u_5	u_6
u_1	3	-1	-1	0	-1	0
u_2	-1	3	-1	-1	0	0
u_3	-1	-1	3	0	0	-1
u_4	0	-1	0	3	-1	-1
u_5	-1	0	0	-1	2	0
u_6	0	0	-1	-1	0	2

Laplacian matrix of G

A

$$A_{ii} = \deg(u_i)$$

$$A_{ij, i \neq j} = \begin{cases} -1 & u_i u_j \in E \\ 0 & u_i u_j \notin E \end{cases}$$

n variables, k constraints, $\det(A) = 0$

$k = \#\text{free vertices}$

\Rightarrow no unique solution

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



	u_1	u_2	u_3	u_4	u_5	u_6
u_1	3	-1	-1	0	-1	0
u_2	-1	3	-1	-1	0	0
u_3	-1	-1	3	0	0	-1
u_4	0	-1	0	3	-1	-1
u_5	-1	0	0	-1	2	0
u_6	0	0	-1	-1	0	2

Laplacian matrix of G

k variables, k constraints, $\det(A) = 0$

$k = \#\text{free vertices}$

\Rightarrow no unique solution

A

$$A_{ii} = \deg(u_i)$$

$$A_{ij, i \neq j} = \begin{cases} -1 & u_i u_j \in E \\ 0 & u_i u_j \notin E \end{cases}$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



	u_1	u_2	u_3	u_4	u_5	u_6
u_1	3	-1	-1	0	-1	0
u_2	-1	3	-1	-1	0	0
u_3	-1	-1	3	0	0	-1
u_4	0	-1	0	3	-1	-1
u_5	-1	0	0	-1	2	0
u_6	0	0	-1	-1	0	2

Laplacian matrix of G

k variables, k constraints, $\det(A) > 0$

$k = \#\text{free vertices}$

\Rightarrow no unique solution

A

$$A_{ii} = \deg(u_i)$$

$$A_{ij, i \neq j} = \begin{cases} -1 & u_i u_j \in E \\ 0 & u_i u_j \notin E \end{cases}$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$

$$p_u = \text{barycenter}(\bigcup_{uv \in E} v) = \sum_{uv \in E} p_v / \deg(u)$$

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v$$


$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v$$

$$Ax = b \quad Ay = b \quad b = (0)_n$$

2 Systems of linear equations

$$\Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$\Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



	u_1	u_2	u_3	u_4	u_5	u_6
u_1	3	-1	-1	0	-1	0
u_2	-1	3	-1	-1	0	0
u_3	-1	-1	3	0	0	-1
u_4	0	-1	0	3	-1	-1
u_5	-1	0	0	-1	2	0
u_6	0	0	-1	-1	0	2

Laplacian matrix of G

k variables, k constraints, $\det(A) > 0$

$k = \#\text{free vertices}$

\Rightarrow unique solution 😊

A

$$A_{ii} = \deg(u_i)$$

$$A_{ij, i \neq j} = \begin{cases} -1 & u_i u_j \in E \\ 0 & u_i u_j \notin E \end{cases}$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$


$p_u = \text{barycenter}(\bigcup_{uv \in E} v)$

Theorem.

Tutte's barycentric algorithm admits a unique solution.
It can be computed in polynomial time.

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v \Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v \Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



	u_1	u_2	u_3	u_4	u_5	u_6	A
u_1	3	-1	-1	0	-1	0	
u_2	-1	3	-1	-1	0	0	
u_3	-1	-1	3	0	0	-1	
u_4	0	-1	0	3	-1	-1	
u_5	-1	0	0	-1	2	0	
u_6	0	0	-1	-1	0	2	

Laplacian matrix of G

k variables, k constraints, $\det(A) > 0$

$k = \#\text{free vertices}$

\Rightarrow unique solution 😊

$$A_{ii} = \deg(u_i)$$

$$A_{ij, i \neq j} = \begin{cases} -1 & u_i u_j \in E \\ 0 & u_i u_j \notin E \end{cases}$$

Linear System of Equations

Goal. $p_u = (x_u, y_u)$


$p_u = \text{barycenter}(\bigcup_{uv \in E} v)$

Theorem.

Tutte's barycentric algorithm admits a **unique solution**.
It can be computed in polynomial time.

$$x_u = \sum_{uv \in E} x_v / \deg(u) \Leftrightarrow \deg(u) \cdot x_u = \sum_{uv \in E} x_v \Leftrightarrow \deg(u) \cdot x_u - \sum_{uv \in E} x_v = 0$$

$$y_u = \sum_{uv \in E} y_v / \deg(u) \Leftrightarrow \deg(u) \cdot y_u = \sum_{uv \in E} y_v \Leftrightarrow \deg(u) \cdot y_u - \sum_{uv \in E} y_v = 0$$



	u_1	u_2	u_3	u_4	u_5	u_6	A
u_1	3	-1	-1	0	-1	0	
u_2	-1	3	-1	-1	0	0	
u_3	-1	-1	3	0	0	-1	
u_4	0	-1	0	3	-1	-1	
u_5	-1	0	0	-1	2	0	
u_6	0	0	-1	-1	0	2	

Laplacian matrix of G

k variables, k constraints, $\det(A) > 0$

$k = \#\text{free vertices}$


\Rightarrow unique solution 😊

Tutte drawing

3-Connected Planar Graphs

planar: G can be drawn in such a way
that no edges cross each other

connected: There is a u - v -path for every $u, v \in V$




3-Connected Planar Graphs

planar: G can be drawn in such a way
that no edges cross each other

connected: There is a u - v -path for every $u, v \in V$

k -connected:




3-Connected Planar Graphs

planar: G can be drawn in such a way that no edges cross each other

connected: There is a u - v -path for every $u, v \in V$

k -connected: $G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$




3-Connected Planar Graphs

planar: G can be drawn in such a way that no edges cross each other

connected: There is a u - v -path for every $u, v \in V$

k -connected: $G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$




3-Connected Planar Graphs

planar: G can be drawn in such a way that no edges cross each other

connected: There is a u - v -path for every $u, v \in V$

k -connected: $G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$




3-Connected Planar Graphs

planar: G can be drawn in such a way that no edges cross each other

connected: There is a u - v -path for every $u, v \in V$

k -connected: $G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$




3-Connected Planar Graphs

planar: G can be drawn in such a way that no edges cross each other

connected: There is a u - v -path for every $u, v \in V$

k -connected: $G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$




3-Connected Planar Graphs

planar: G can be drawn in such a way
that no edges cross each other

connected: There is a u - v -path for every $u, v \in V$

k -connected: $G - \{v_1, \dots, v_{k-1}\}$ is connected
for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)




3-Connected Planar Graphs

planar: G can be drawn in such a way that no edges cross each other

connected: There is a u - v -path for every $u, v \in V$

k -connected: $G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)

There are at least k vertex-disjoint u - v -paths for every $u, v \in V$




3-Connected Planar Graphs

planar: G can be drawn in such a way that no edges cross each other

connected: There is a u - v -path for every $u, v \in V$

k -connected: $G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)

There are at least k vertex-disjoint u - v -paths for every $u, v \in V$




3-Connected Planar Graphs

planar: G can be drawn in such a way that no edges cross each other

connected: There is a u - v -path for every $u, v \in V$

k -connected: $G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)

There are at least k vertex-disjoint u - v -paths for every $u, v \in V$




3-Connected Planar Graphs

planar: G can be drawn in such a way that no edges cross each other

connected: There is a u - v -path for every $u, v \in V$

k -connected: $G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)

There are at least k vertex-disjoint u - v -paths for every $u, v \in V$




3-Connected Planar Graphs

planar: G can be drawn in such a way that no edges cross each other

connected: There is a u - v -path for every $u, v \in V$

k -connected: $G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)

There are at least k vertex-disjoint u - v -paths for every $u, v \in V$




3-Connected Planar Graphs

planar: G can be drawn in such a way that no edges cross each other

connected: There is a u - v -path for every $u, v \in V$

k -connected: $G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)

There are at least k vertex-disjoint u - v -paths for every $u, v \in V$




3-Connected Planar Graphs

planar: G can be drawn in such a way that no edges cross each other

connected: There is a u - v -path for every $u, v \in V$

k -connected: $G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)

There are at least k vertex-disjoint u - v -paths for every $u, v \in V$




3-Connected Planar Graphs

planar: G can be drawn in such a way that no edges cross each other

connected: There is a u - v -path for every $u, v \in V$

k -connected: $G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)

There are at least k vertex-disjoint u - v -paths for every $u, v \in V$




3-Connected Planar Graphs

planar: G can be drawn in such a way that no edges cross each other

connected: There is a u - v -path for every $u, v \in V$

k -connected: $G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)

There are at least k vertex-disjoint u - v -paths for every $u, v \in V$



3-Connected Planar Graphs

planar:

G can be drawn in such a way that no edges cross each other

connected:

There is a u - v -path for every $u, v \in V$

k -connected:


$G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)

There are at least k vertex-disjoint u - v -paths for every $u, v \in V$

Theorem.

[Whitney 1933]

Every 3-connected planar graph has a unique planar embedding.



3-Connected Planar Graphs

planar:

G can be drawn in such a way that no edges cross each other


connected:

There is a u - v -path for every $u, v \in V$

k -connected:

$G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)

There are at least k vertex-disjoint u - v -paths for every $u, v \in V$



Theorem.

[Whitney 1933]

Every 3-connected planar graph has a unique planar embedding.

Proof sketch.

3-Connected Planar Graphs

planar:

G can be drawn in such a way that no edges cross each other


connected:

There is a u - v -path for every $u, v \in V$

k -connected:

$G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)

There are at least k vertex-disjoint u - v -paths for every $u, v \in V$



Theorem.

[Whitney 1933]

Every 3-connected planar graph has a unique planar embedding.

Proof sketch.

Γ_1, Γ_2 embeddings of G

3-Connected Planar Graphs

planar:

G can be drawn in such a way that no edges cross each other


connected:

There is a u - v -path for every $u, v \in V$

k -connected:

$G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)

There are at least k vertex-disjoint u - v -paths for every $u, v \in V$



Theorem.

[Whitney 1933]

Every 3-connected planar graph has a unique planar embedding.

Proof sketch.

Γ_1, Γ_2 embeddings of G

C face of Γ_2 , but not Γ_1

3-Connected Planar Graphs

planar:

G can be drawn in such a way that no edges cross each other


connected:

There is a u - v -path for every $u, v \in V$

k -connected:

$G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)

There are at least k vertex-disjoint u - v -paths for every $u, v \in V$



Γ_1

Theorem.

[Whitney 1933]

Every 3-connected planar graph has a unique planar embedding.

Proof sketch.

Γ_1, Γ_2 embeddings of G

C face of Γ_2 , but not Γ_1

3-Connected Planar Graphs

planar:

G can be drawn in such a way that no edges cross each other


connected:

There is a u - v -path for every $u, v \in V$


k -connected:

$G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)

There are at least k vertex-disjoint u - v -paths for every $u, v \in V$



Γ_1



Theorem.

[Whitney 1933]

Every 3-connected planar graph has a unique planar embedding.

Proof sketch.

Γ_1, Γ_2 embeddings of G

C face of Γ_2 , but not Γ_1

3-Connected Planar Graphs

planar:

G can be drawn in such a way that no edges cross each other

connected:


There is a u - v -path for every $u, v \in V$

k -connected:


$G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$

or (equivalently)

There are at least k vertex-disjoint u - v -paths for every $u, v \in V$



Γ_1



Theorem.

[Whitney 1933]

Every 3-connected planar graph has a unique planar embedding.

Proof sketch.

Γ_1, Γ_2 embeddings of G

C face of Γ_2 , but not Γ_1

u inside C in Γ_1

3-Connected Planar Graphs

planar:

G can be drawn in such a way that no edges cross each other

connected:



There is a u - v -path for every $u, v \in V$

k -connected:

$G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$

or (equivalently)

There are at least k vertex-disjoint u - v -paths for every $u, v \in V$



Theorem.

[Whitney 1933]

Every 3-connected planar graph has a unique planar embedding.

Proof sketch.

Γ_1, Γ_2 embeddings of G

C face of Γ_2 , but not Γ_1

u inside C in Γ_1 , v outside C in Γ_1

3-Connected Planar Graphs

planar:

G can be drawn in such a way that no edges cross each other

connected:



There is a u - v -path for every $u, v \in V$

k -connected:

$G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$

or (equivalently)

There are at least k vertex-disjoint u - v -paths for every $u, v \in V$



Theorem.

[Whitney 1933]

Every 3-connected planar graph has a unique planar embedding.

Proof sketch.

Γ_1, Γ_2 embeddings of G

C face of Γ_2 , but not Γ_1

u inside C in Γ_1 , v outside C in Γ_1

3-Connected Planar Graphs

planar:

G can be drawn in such a way that no edges cross each other



connected:

There is a u - v -path for every $u, v \in V$

k -connected:

$G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)

There are at least k vertex-disjoint u - v -paths for every $u, v \in V$



Theorem.

[Whitney 1933]

Every 3-connected planar graph has a unique planar embedding.

Proof sketch.

Γ_1, Γ_2 embeddings of G

C face of Γ_2 , but not Γ_1

u inside C in Γ_1 , v outside C in Γ_1

3-Connected Planar Graphs

planar:

G can be drawn in such a way that no edges cross each other



connected:

There is a u - v -path for every $u, v \in V$

k -connected:

$G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)

There are at least k vertex-disjoint u - v -paths for every $u, v \in V$



Theorem.

[Whitney 1933]

Every 3-connected planar graph has a unique planar embedding.

Proof sketch.

Γ_1, Γ_2 embeddings of G

C face of Γ_2 , but not Γ_1

u inside C in Γ_1 , v outside C in Γ_1

3-Connected Planar Graphs

planar:

G can be drawn in such a way that no edges cross each other


connected:

There is a u - v -path for every $u, v \in V$


k -connected:

$G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)


There are at least k vertex-disjoint u - v -paths for every $u, v \in V$



Γ_1



Γ_2



Theorem.

[Whitney 1933]

Every 3-connected planar graph has a unique planar embedding.

Proof sketch.

Γ_1, Γ_2 embeddings of G

C face of Γ_2 , but not Γ_1

u inside C in Γ_1 , v outside C in Γ_1

3-Connected Planar Graphs

planar:

G can be drawn in such a way that no edges cross each other


connected:

There is a u - v -path for every $u, v \in V$


k -connected:

$G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)


There are at least k vertex-disjoint u - v -paths for every $u, v \in V$



Γ_1



Γ_2



Theorem.

[Whitney 1933]

Every 3-connected planar graph has a unique planar embedding.

Proof sketch.

Γ_1, Γ_2 embeddings of G

C face of Γ_2 , but not Γ_1

u inside C in Γ_1 , v outside C in Γ_1

3-Connected Planar Graphs

planar:

G can be drawn in such a way that no edges cross each other


connected:

There is a u - v -path for every $u, v \in V$


k -connected:

$G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)


There are at least k vertex-disjoint u - v -paths for every $u, v \in V$



Γ_1



Γ_2



Theorem.

[Whitney 1933]

Every 3-connected planar graph has a unique planar embedding.

Proof sketch.

Γ_1, Γ_2 embeddings of G

C face of Γ_2 , but not Γ_1

u inside C in Γ_1 , v outside C in Γ_1
both on same side in Γ_2

3-Connected Planar Graphs

planar:

G can be drawn in such a way that no edges cross each other


connected:

There is a u - v -path for every $u, v \in V$


k -connected:

$G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)


There are at least k vertex-disjoint u - v -paths for every $u, v \in V$



Γ_1



Γ_2



Theorem.

[Whitney 1933]

Every 3-connected planar graph has a unique planar embedding.

Proof sketch.

Γ_1, Γ_2 embeddings of G

C face of Γ_2 , but not Γ_1

u inside C in Γ_1 , v outside C in Γ_1
both on same side in Γ_2

3-Connected Planar Graphs

planar:

G can be drawn in such a way that no edges cross each other

connected:


There is a u - v -path for every $u, v \in V$

k -connected:


$G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$

or (equivalently)


There are at least k vertex-disjoint u - v -paths for every $u, v \in V$



Γ_1



Γ_2



Theorem.

[Whitney 1933]

Every 3-connected planar graph has a unique planar embedding.

Proof sketch.

Γ_1, Γ_2 embeddings of G

C face of Γ_2 , but not Γ_1

u inside C in Γ_1 , v outside C in Γ_1
both on same side in Γ_2

3-Connected Planar Graphs

planar:

G can be drawn in such a way that no edges cross each other


connected:

There is a u - v -path for every $u, v \in V$


k -connected:

$G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)


There are at least k vertex-disjoint u - v -paths for every $u, v \in V$



Γ_1



Γ_2



Theorem.

[Whitney 1933]

Every 3-connected planar graph has a unique planar embedding.

Proof sketch.

Γ_1, Γ_2 embeddings of G

C face of Γ_2 , but not Γ_1

u inside C in Γ_1 , v outside C in Γ_1
both on same side in Γ_2

3-Connected Planar Graphs

planar:

G can be drawn in such a way that no edges cross each other


connected:

There is a u - v -path for every $u, v \in V$


k -connected:

$G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)


There are at least k vertex-disjoint u - v -paths for every $u, v \in V$



Γ_1



Γ_2



Theorem.

[Whitney 1933]

Every 3-connected planar graph has a unique planar embedding.

Proof sketch.

Γ_1, Γ_2 embeddings of G

C face of Γ_2 , but not Γ_1

u inside C in Γ_1 , v outside C in Γ_1
both on same side in Γ_2

3-Connected Planar Graphs

planar:

G can be drawn in such a way that no edges cross each other


connected:

There is a u - v -path for every $u, v \in V$


k -connected:

$G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)


There are at least k vertex-disjoint u - v -paths for every $u, v \in V$



Γ_1



Γ_2



Theorem.

[Whitney 1933]

Every 3-connected planar graph has a unique planar embedding.

Proof sketch.

Γ_1, Γ_2 embeddings of G

C face of Γ_2 , but not Γ_1

u inside C in Γ_1 , v outside C in Γ_1
both on same side in Γ_2

3-Connected Planar Graphs

planar:

G can be drawn in such a way that no edges cross each other


connected:

There is a u - v -path for every $u, v \in V$


k -connected:

$G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)


There are at least k vertex-disjoint u - v -paths for every $u, v \in V$



Γ_1



Γ_2



Theorem.

[Whitney 1933]

Every 3-connected planar graph has a unique planar embedding.

Proof sketch.

Γ_1, Γ_2 embeddings of G

C face of Γ_2 , but not Γ_1

u inside C in Γ_1 , v outside C in Γ_1
both on same side in Γ_2

3-Connected Planar Graphs

planar:

G can be drawn in such a way that no edges cross each other


connected:

There is a u - v -path for every $u, v \in V$


k -connected:

$G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)


There are at least k vertex-disjoint u - v -paths for every $u, v \in V$



Γ_1



Γ_2



Theorem.

[Whitney 1933]

Every 3-connected planar graph has a unique planar embedding.

Proof sketch.

Γ_1, Γ_2 embeddings of G

C face of Γ_2 , but not Γ_1

u inside C in Γ_1 , v outside C in Γ_1
both on same side in Γ_2

3-Connected Planar Graphs

planar:

G can be drawn in such a way that no edges cross each other


connected:

There is a u - v -path for every $u, v \in V$


k -connected:

$G - \{v_1, \dots, v_{k-1}\}$ is connected for **any** $v_1, \dots, v_{k-1} \in V$
or (equivalently)


There are at least k vertex-disjoint u - v -paths for every $u, v \in V$



Γ_1



Γ_2



Theorem.

[Whitney 1933]

Every 3-connected planar graph has a unique planar embedding.

Proof sketch.

Γ_1, Γ_2 embeddings of G

C face of Γ_2 , but not Γ_1


u inside C in Γ_1 , v outside C in Γ_1
both on same side in Γ_2

Tutte's Theorem

Theorem.

Let G be a 3-connected planar graph

[Tutte 1963]




Tutte's Theorem

Theorem.

Let G be a 3-connected planar graph, and let C be a face of its unique embedding.

[Tutte 1963]




Tutte's Theorem

Theorem.

Let G be a 3-connected planar graph, and let C be a face of its unique embedding. If we fix C on a strictly convex polygon,

[Tutte 1963]




Tutte's Theorem

Theorem.

[Tutte 1963]

Let G be a 3-connected planar graph, and let C be a face of its unique embedding.

If we fix C on a strictly convex polygon, then the Tutte drawing of G is planar




Tutte's Theorem

Theorem.

[Tutte 1963]


Let G be a 3-connected planar graph, and let C be a face of its unique embedding.

If we fix C on a strictly convex polygon, then the Tutte drawing of G is planar and all its faces are strictly convex.




Properties of Tutte Drawings

Property 1. Let $v \in V$ free




Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .



Properties of Tutte Drawings


Property 1. Let $v \in V$ free, ℓ line through v .
 $\exists u, v \in E$ on one side of ℓ



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side




Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists u, v \in E$ on one side of $\ell \Rightarrow \exists v, w \in E$ on other side

Otherwise, all forces to same side ...




Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists u, v \in E$ on one side of $\ell \Rightarrow \exists v, w \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .




Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$$\exists uv \in E \text{ on one side of } \ell \Rightarrow \exists vw \in E \text{ on other side}$$

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .




Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .




Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .




Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .




Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$$\exists uv \in E \text{ on one side of } \ell \Rightarrow \exists vw \in E \text{ on other side}$$

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .




Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$$\exists uv \in E \text{ on one side of } \ell \Rightarrow \exists vw \in E \text{ on other side}$$

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .




Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$$\exists uv \in E \text{ on one side of } \ell \Rightarrow \exists vw \in E \text{ on other side}$$

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .



Properties of Tutte Drawings


Property 1. Let $v \in V$ free, ℓ line through v .

$$\exists uv \in E \text{ on one side of } \ell \Rightarrow \exists vw \in E \text{ on other side}$$

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .

Property 3. Let ℓ be any line.



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .


$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .


$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .


$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .


$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side


Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

v furthest away from ℓ



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...


Property 2. All free vertices lie inside C .

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

v furthest away from ℓ

Pick any vertex u



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...


Property 2. All free vertices lie inside C .

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

v furthest away from ℓ

Pick any vertex u , ℓ' parallel to ℓ through u



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...


Property 2. All free vertices lie inside C .

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

v furthest away from ℓ

Pick any vertex u , ℓ' parallel to ℓ through u
 G connected, v not on ℓ'



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .


Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

v furthest away from ℓ

Pick any vertex u , ℓ' parallel to ℓ through u

G connected, v not on $\ell' \Rightarrow \exists w$ on ℓ' with neighbor further away from ℓ



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .


Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

v furthest away from ℓ

Pick any vertex u , ℓ' parallel to ℓ through u

G connected, v not on $\ell' \Rightarrow \exists w$ on ℓ' with neighbor further away from ℓ
 $\Rightarrow \exists$ path from u to v



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .


Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

v furthest away from ℓ

Pick any vertex u , ℓ' parallel to ℓ through u

G connected, v not on $\ell' \Rightarrow \exists w$ on ℓ' with neighbor further away from ℓ
 $\Rightarrow \exists$ path from u to v



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .


Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

v furthest away from ℓ

Pick any vertex u , ℓ' parallel to ℓ through u

G connected, v not on $\ell' \Rightarrow \exists w$ on ℓ' with neighbor further away from ℓ
 $\Rightarrow \exists$ path from u to v



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .


Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

v furthest away from ℓ

Pick any vertex u , ℓ' parallel to ℓ through u

G connected, v not on $\ell' \Rightarrow \exists w$ on ℓ' with neighbor further away from ℓ
 $\Rightarrow \exists$ path from u to v



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .


Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

v furthest away from ℓ

Pick any vertex u , ℓ' parallel to ℓ through u

G connected, v not on $\ell' \Rightarrow \exists w$ on ℓ' with neighbor further away from ℓ
 $\Rightarrow \exists$ path from u to v



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .


Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

v furthest away from ℓ

Pick any vertex u , ℓ' parallel to ℓ through u

G connected, v not on $\ell' \Rightarrow \exists w$ on ℓ' with neighbor further away from ℓ
 $\Rightarrow \exists$ path from u to v



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .

Property 3. Let ℓ be any line.


Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

v furthest away from ℓ

Pick any vertex u , ℓ' parallel to ℓ through u

G connected, v not on $\ell' \Rightarrow \exists w$ on ℓ' with neighbor further away from ℓ
 $\Rightarrow \exists$ path from u to v

Property 4. No vertex is collinear with all of its neighbors.



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists u, v \in E$ on one side of $\ell \Rightarrow \exists v, w \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .

Property 3. Let ℓ be any line.



Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

v furthest away from ℓ

Pick any vertex u , ℓ' parallel to ℓ through u

G connected, v not on $\ell' \Rightarrow \exists w$ on ℓ' with neighbor further away from ℓ
 $\Rightarrow \exists$ path from u to v

Property 4. No vertex is collinear with all of its neighbors.



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists u, v \in E$ on one side of $\ell \Rightarrow \exists v, w \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .

Property 3. Let ℓ be any line.



Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

v furthest away from ℓ

Pick any vertex u , ℓ' parallel to ℓ through u

G connected, v not on $\ell' \Rightarrow \exists w$ on ℓ' with neighbor further away from ℓ
 $\Rightarrow \exists$ path from u to v

Property 4. No vertex is collinear with all of its neighbors.



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.



v furthest away from ℓ

Pick any vertex u , ℓ' parallel to ℓ through u

G connected, v not on $\ell' \Rightarrow \exists w$ on ℓ' with neighbor further away from ℓ
 $\Rightarrow \exists$ path from u to v

Property 4. No vertex is collinear with all of its neighbors.

Not all vertices collinear



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.



v furthest away from ℓ

Pick any vertex u , ℓ' parallel to ℓ through u

G connected, v not on $\ell' \Rightarrow \exists w$ on ℓ' with neighbor further away from ℓ
 $\Rightarrow \exists$ path from u to v

Property 4. No vertex is collinear with all of its neighbors.

Not all vertices collinear



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.



v furthest away from ℓ

Pick any vertex u , ℓ' parallel to ℓ through u

G connected, v not on $\ell' \Rightarrow \exists w$ on ℓ' with neighbor further away from ℓ
 $\Rightarrow \exists$ path from u to v

Property 4. No vertex is collinear with all of its neighbors.

Not all vertices collinear



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.



v furthest away from ℓ

Pick any vertex u , ℓ' parallel to ℓ through u

G connected, v not on $\ell' \Rightarrow \exists w$ on ℓ' with neighbor further away from ℓ
 $\Rightarrow \exists$ path from u to v

Property 4. No vertex is collinear with all of its neighbors.

Not all vertices collinear



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.



v furthest away from ℓ

Pick any vertex u , ℓ' parallel to ℓ through u

G connected, v not on $\ell' \Rightarrow \exists w$ on ℓ' with neighbor further away from ℓ
 $\Rightarrow \exists$ path from u to v

Property 4. No vertex is collinear with all of its neighbors.

Not all vertices collinear
 G 3-connected



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.



v furthest away from ℓ

Pick any vertex u , ℓ' parallel to ℓ through u

G connected, v not on $\ell' \Rightarrow \exists w$ on ℓ' with neighbor further away from ℓ
 $\Rightarrow \exists$ path from u to v

Property 4. No vertex is collinear with all of its neighbors.

Not all vertices collinear
 G 3-connected



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.



v furthest away from ℓ

Pick any vertex u , ℓ' parallel to ℓ through u

G connected, v not on $\ell' \Rightarrow \exists w$ on ℓ' with neighbor further away from ℓ
 $\Rightarrow \exists$ path from u to v

Property 4. No vertex is collinear with all of its neighbors.

Not all vertices collinear
 G 3-connected



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.



v furthest away from ℓ

Pick any vertex u , ℓ' parallel to ℓ through u

G connected, v not on $\ell' \Rightarrow \exists w$ on ℓ' with neighbor further away from ℓ
 $\Rightarrow \exists$ path from u to v

Property 4. No vertex is collinear with all of its neighbors.

Not all vertices collinear
 G 3-connected
 $\Rightarrow K_{3,3}$ minor



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.



v furthest away from ℓ

Pick any vertex u , ℓ' parallel to ℓ through u

G connected, v not on $\ell' \Rightarrow \exists w$ on ℓ' with neighbor further away from ℓ
 $\Rightarrow \exists$ path from u to v

Property 4. No vertex is collinear with all of its neighbors.

Not all vertices collinear
 G 3-connected
 $\Rightarrow K_{3,3}$ minor



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.




v furthest away from ℓ

Pick any vertex u , ℓ' parallel to ℓ through u

G connected, v not on $\ell' \Rightarrow \exists w$ on ℓ' with neighbor further away from ℓ
 $\Rightarrow \exists$ path from u to v

Property 4. No vertex is collinear with all of its neighbors.

Not all vertices collinear
 G 3-connected
 $\Rightarrow K_{3,3}$ minor



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.




v furthest away from ℓ

Pick any vertex u , ℓ' parallel to ℓ through u

G connected, v not on $\ell' \Rightarrow \exists w$ on ℓ' with neighbor further away from ℓ
 $\Rightarrow \exists$ path from u to v

Property 4. No vertex is collinear with all of its neighbors.

Not all vertices collinear
 G 3-connected
 $\Rightarrow K_{3,3}$ minor



Properties of Tutte Drawings

Property 1. Let $v \in V$ free, ℓ line through v .

$\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Otherwise, all forces to same side ...

Property 2. All free vertices lie inside C .

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.




v furthest away from ℓ

Pick any vertex u , ℓ' parallel to ℓ through u

G connected, v not on $\ell' \Rightarrow \exists w$ on ℓ' with neighbor further away from ℓ
 $\Rightarrow \exists$ path from u to v


Property 4. No vertex is collinear with all of its neighbors.

Not all vertices collinear
 G 3-connected
 $\Rightarrow K_{3,3}$ minor




Proof of Tutte's Theorem

Lemma. Let $uv \in E$ be a non-boundary edge




Proof of Tutte's Theorem

Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv .




Proof of Tutte's Theorem

Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .





Proof of Tutte's Theorem

Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .



Proof of Tutte's Theorem

Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .





Proof of Tutte's Theorem

Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.





Proof of Tutte's Theorem

Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.





Proof of Tutte's Theorem

Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.



Proof of Tutte's Theorem

Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .

Property 1. Let $v \in V$ free, ℓ line through v .



$$\exists uv \in E \text{ on one side of } \ell \Rightarrow \exists vw \in E \text{ on other side}$$

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .

Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.



Proof of Tutte's Theorem

Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .

Property 1. Let $v \in V$ free, ℓ line through v .



$$\exists uv \in E \text{ on one side of } \ell \Rightarrow \exists vw \in E \text{ on other side}$$

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .

Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.



Proof of Tutte's Theorem

Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .

Property 1. Let $v \in V$ free, ℓ line through v .



$$\exists uv \in E \text{ on one side of } \ell \Rightarrow \exists vw \in E \text{ on other side}$$

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .

Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.



Proof of Tutte's Theorem

Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .

Property 1. Let $v \in V$ free, ℓ line through v .



$$\exists uv \in E \text{ on one side of } \ell \Rightarrow \exists vw \in E \text{ on other side}$$

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .

Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.



Proof of Tutte's Theorem

Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .

Property 1. Let $v \in V$ free, ℓ line through v .



$$\exists uv \in E \text{ on one side of } \ell \Rightarrow \exists vw \in E \text{ on other side}$$

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .

Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.



Proof of Tutte's Theorem

Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .

Property 1. Let $v \in V$ free, ℓ line through v .



$$\exists uv \in E \text{ on one side of } \ell \Rightarrow \exists vw \in E \text{ on other side}$$

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .

Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.



Proof of Tutte's Theorem

Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .

Property 1. Let $v \in V$ free, ℓ line through v .



$$\exists uv \in E \text{ on one side of } \ell \Rightarrow \exists vw \in E \text{ on other side}$$

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .

Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.



Proof of Tutte's Theorem



Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .

Property 1. Let $v \in V$ free, ℓ line through v .
 $\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Property 3. Let ℓ be any line.
Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.

Lemma. All faces are strictly convex.



Proof of Tutte's Theorem



Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .

Property 1. Let $v \in V$ free, ℓ line through v .
 $\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Property 3. Let ℓ be any line.
Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.

Lemma. All faces are strictly convex.



Proof of Tutte's Theorem



Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .

Property 1. Let $v \in V$ free, ℓ line through v .
 $\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Property 3. Let ℓ be any line.
Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.

Lemma. All faces are strictly convex.



Proof of Tutte's Theorem




Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .

Property 1. Let $v \in V$ free, ℓ line through v .
 $\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Property 3. Let ℓ be any line.
Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.

Lemma. All faces are strictly convex.



Proof of Tutte's Theorem




Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .

Property 1. Let $v \in V$ free, ℓ line through v .
 $\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Property 3. Let ℓ be any line.
Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.

Lemma. All faces are strictly convex.



Proof of Tutte's Theorem




Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .

Property 1. Let $v \in V$ free, ℓ line through v .
 $\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Property 3. Let ℓ be any line.
Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.

Lemma. All faces are strictly convex.



Proof of Tutte's Theorem




Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .

Property 1. Let $v \in V$ free, ℓ line through v .
 $\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Property 3. Let ℓ be any line.
Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.

Lemma. All faces are strictly convex.



Proof of Tutte's Theorem




Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .

Property 1. Let $v \in V$ free, ℓ line through v .
 $\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Property 3. Let ℓ be any line.
Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.

Lemma. All faces are strictly convex.



Proof of Tutte's Theorem




Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .

Property 1. Let $v \in V$ free, ℓ line through v .
 $\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Property 3. Let ℓ be any line.
Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.

Lemma. All faces are strictly convex.



Proof of Tutte's Theorem




Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .

Property 1. Let $v \in V$ free, ℓ line through v .
 $\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Property 3. Let ℓ be any line.
Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.

Lemma. All faces are strictly convex.



Proof of Tutte's Theorem




Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .

Property 1. Let $v \in V$ free, ℓ line through v .
 $\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Property 3. Let ℓ be any line.
Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.

Lemma. All faces are strictly convex.



Lemma. The drawing is planar.

Proof of Tutte's Theorem




Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .

Property 1. Let $v \in V$ free, ℓ line through v .
 $\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side


Property 3. Let ℓ be any line.
Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.

Lemma. All faces are strictly convex.




Lemma. The drawing is planar.



Proof of Tutte's Theorem


Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .




Property 1. Let $v \in V$ free, ℓ line through v .
 $\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Property 3. Let ℓ be any line.
Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.


Property 4. No vertex is collinear with all of its neighbors.



Lemma. All faces are strictly convex.




Lemma. The drawing is planar.



Proof of Tutte's Theorem


Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .




Property 1. Let $v \in V$ free, ℓ line through v .
 $\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Property 3. Let ℓ be any line.
Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.




Lemma. All faces are strictly convex.




p inside two faces

Lemma. The drawing is planar.



Proof of Tutte's Theorem


Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .




Property 1. Let $v \in V$ free, ℓ line through v .
 $\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Property 3. Let ℓ be any line.
Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.




Lemma. All faces are strictly convex.




p inside two faces

Lemma. The drawing is planar.



Proof of Tutte's Theorem


Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .




Property 1. Let $v \in V$ free, ℓ line through v .
 $\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Property 3. Let ℓ be any line.
Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.




Lemma. All faces are strictly convex.




p inside two faces

Lemma. The drawing is planar.



Proof of Tutte's Theorem


Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .




Property 1. Let $v \in V$ free, ℓ line through v .
 $\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Property 3. Let ℓ be any line.
Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.




Lemma. All faces are strictly convex.



Property 2. All free vertices lie inside C .


p inside two faces

Lemma. The drawing is planar.



Proof of Tutte's Theorem

Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .



Property 1. Let $v \in V$ free, ℓ line through v .


$$\exists uv \in E \text{ on one side of } \ell \Rightarrow \exists vw \in E \text{ on other side}$$

Property 3. Let ℓ be any line.


Let V_ℓ be all vertices on one side of ℓ .

Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.




Lemma. All faces are strictly convex.



p inside two faces


Property 2. All free vertices lie inside C .
 $\Rightarrow q$ in one face

Lemma. The drawing is planar.



Proof of Tutte's Theorem


Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .




Property 1. Let $v \in V$ free, ℓ line through v .
 $\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Property 3. Let ℓ be any line.
Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.




Lemma. All faces are strictly convex.




Property 2. All free vertices lie inside C .
 p inside two faces
 $\Rightarrow q$ in one face

Lemma. The drawing is planar.



Proof of Tutte's Theorem


Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .




Property 1. Let $v \in V$ free, ℓ line through v .
 $\exists uv \in E$ on one side of $\ell \Rightarrow \exists vw \in E$ on other side

Property 3. Let ℓ be any line.
Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.




Lemma. All faces are strictly convex.




Property 2. All free vertices lie inside C .
 $\Rightarrow p$ in one face
 $\Rightarrow q$ in one face

Lemma. The drawing is planar.



Proof of Tutte's Theorem

Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .



Property 1. Let $v \in V$ free, ℓ line through v .


$$\exists uv \in E \text{ on one side of } \ell \Rightarrow \exists vw \in E \text{ on other side}$$

Property 3. Let ℓ be any line.


Let V_ℓ be all vertices on one side of ℓ .

Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.



Lemma. All faces are strictly convex.




p inside two faces

Property 2. All free vertices lie inside C .


$\Rightarrow q$ in one face

Lemma. The drawing is planar.



Proof of Tutte's Theorem

Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .



Property 1. Let $v \in V$ free, ℓ line through v .

$$\exists uv \in E \text{ on one side of } \ell \Rightarrow \exists vw \in E \text{ on other side}$$


Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .

Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.

Lemma. All faces are strictly convex.




Property 2. All free vertices lie inside C .

p inside two faces


$\Rightarrow q$ in one face

jumping over edge

$\rightarrow \# \text{faces the same}$




Lemma. The drawing is planar.



Proof of Tutte's Theorem

Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .



Property 1. Let $v \in V$ free, ℓ line through v .

$$\exists uv \in E \text{ on one side of } \ell \Rightarrow \exists vw \in E \text{ on other side}$$


Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .

Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.

Lemma. All faces are strictly convex.




Property 2. All free vertices lie inside C .

p inside two faces


$\Rightarrow q$ in one face

jumping over edge

$\rightarrow \# \text{faces the same}$




Lemma. The drawing is planar.



Proof of Tutte's Theorem

Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .



Property 1. Let $v \in V$ free, ℓ line through v .

$$\exists uv \in E \text{ on one side of } \ell \Rightarrow \exists vw \in E \text{ on other side}$$


Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .

Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.

Lemma. All faces are strictly convex.




Property 2. All free vertices lie inside C .

p inside two faces


$\Rightarrow q$ in one face

jumping over edge

$\rightarrow \# \text{faces the same}$




Lemma. The drawing is planar.



Proof of Tutte's Theorem

Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .



Property 1. Let $v \in V$ free, ℓ line through v .

$$\exists uv \in E \text{ on one side of } \ell \Rightarrow \exists vw \in E \text{ on other side}$$


Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .

Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.

Lemma. All faces are strictly convex.



Property 2. All free vertices lie inside C .


p inside two faces

$\Rightarrow q$ in one face

jumping over edge


$\rightarrow \# \text{faces the same}$

Lemma. The drawing is planar.



Proof of Tutte's Theorem

Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .



Property 1. Let $v \in V$ free, ℓ line through v .


$$\exists uv \in E \text{ on one side of } \ell \Rightarrow \exists vw \in E \text{ on other side}$$

Property 3. Let ℓ be any line.

Let V_ℓ be all vertices on one side of ℓ .
Then $G[V_\ell]$ is connected.


Property 4. No vertex is collinear with all of its neighbors.

Lemma. All faces are strictly convex.




Property 2. All free vertices lie inside C .

p inside two faces
 $\Rightarrow q$ in one face
jumping over edge
 \rightarrow #faces the same
 $\Rightarrow p$ inside one face




Lemma. The drawing is planar.



Proof of Tutte's Theorem

Lemma. Let $uv \in E$ be a non-boundary edge, ℓ line through uv . Then the two faces f_1, f_2 incident to uv lie completely on opposite sides of ℓ .



Property 1. Let $v \in V$ free, ℓ line through v .


$$\exists uv \in E \text{ on one side of } \ell \Rightarrow \exists vw \in E \text{ on other side}$$

Property 3. Let ℓ be any line.


Let V_ℓ be all vertices on one side of ℓ .

Then $G[V_\ell]$ is connected.

Property 4. No vertex is collinear with all of its neighbors.



Lemma. All faces are strictly convex.



Property 2. All free vertices lie inside C .

p inside two faces

$\Rightarrow q$ in one face

jumping over edge

\rightarrow #faces the same

$\Rightarrow p$ inside one face

Lemma. The drawing is planar.

