


CS F402: Computational Geometry




Lecture 12:

Straight-Line Drawings of Planar Graphs II: Schnyder Realizer

Siddharth Gupta

March 21+26 + April 2+4, 2025



Planar Straight-Line Drawings

Theorem.

[De Fraysseix, Pach, Pollack '90]

Every n -vertex planar graph has a planar straight-line drawing of size $(2n - 4) \times (n - 2)$.

Theorem.

[Schnyder '90]

Every n -vertex planar graph has a planar straight-line drawing of size $(n - 2) \times (n - 2)$.

Planar Straight-Line Drawings

Theorem.

[De Fraysseix, Pach, Pollack '90]

Every n -vertex planar graph has a planar straight-line drawing of size $(2n - 4) \times (n - 2)$.

Theorem.

[Schnyder '90]

Every n -vertex planar graph has a planar straight-line drawing of size $(n - 2) \times (n - 2)$.

Planar Straight-Line Drawings

Theorem.

[De Fraysseix, Pach, Pollack '90]

Every n -vertex planar graph has a planar straight-line drawing of size $(2n - 4) \times (n - 2)$.


Theorem.

[Schnyder '90]

Every n -vertex planar graph has a planar straight-line drawing of size $(n - 2) \times (n - 2)$.

Idea.

- Fix outer triangle.



Planar Straight-Line Drawings

Theorem.

[De Fraysseix, Pach, Pollack '90]

Every n -vertex planar graph has a planar straight-line drawing of size $(2n - 4) \times (n - 2)$.


Theorem.

[Schneider '90]

Every n -vertex planar graph has a planar straight-line drawing of size $(n - 2) \times (n - 2)$.

Idea.

- Fix outer triangle.
- Compute coordinates of inner vertices



Planar Straight-Line Drawings

Theorem.

[De Fraysseix, Pach, Pollack '90]

Every n -vertex planar graph has a planar straight-line drawing of size $(2n - 4) \times (n - 2)$.


Theorem.

[Schneider '90]

Every n -vertex planar graph has a planar straight-line drawing of size $(n - 2) \times (n - 2)$.

Idea.

- Fix outer triangle.
- Compute coordinates of inner vertices
 - based on outer triangle



Planar Straight-Line Drawings

Theorem. [De Fraysseix, Pach, Pollack '90]


Every n -vertex planar graph has a planar straight-line drawing of size $(2n - 4) \times (n - 2)$.

Theorem. [Schneider '90]

Every n -vertex planar graph has a planar straight-line drawing of size $(n - 2) \times (n - 2)$.

Idea.

- Fix outer triangle.
- Compute coordinates of inner vertices
 - based on outer triangle and
 - how much space there should be for other vertices



Planar Straight-Line Drawings

Theorem.

[De Fraysseix, Pach, Pollack '90]

Every n -vertex planar graph has a planar straight-line drawing of size $(2n - 4) \times (n - 2)$.


Theorem.

[Schneider '90]

Every n -vertex planar graph has a planar straight-line drawing of size $(n - 2) \times (n - 2)$.

Idea.

- Fix outer triangle.
- Compute coordinates of inner vertices
 - based on outer triangle and
 - how much space there should be for other vertices
 - using weighted barycentric coordinates.



Planar Straight-Line Drawings

Theorem. [De Fraysseix, Pach, Pollack '90]


Every n -vertex planar graph has a planar straight-line drawing of size $(2n - 4) \times (n - 2)$.

Theorem. [Schneider '89]

Every n -vertex planar graph has a planar straight-line drawing of size $(2n - 5) \times (2n - 5)$.


Idea.

- Fix outer triangle.
- Compute coordinates of inner vertices
 - based on outer triangle and
 - how much space there should be for other vertices
 - using weighted barycentric coordinates.



Barycentric Coordinates



Recall: $\text{barycenter}(x_1, \dots, x_k) = \sum_{i=1}^k x_i / k$



Barycentric Coordinates

Recall: $\text{barycenter}(x_1, \dots, x_k) = \sum_{i=1}^k x_i / k$



Let A, B, C form a triangle



Barycentric Coordinates

Recall: $\text{barycenter}(x_1, \dots, x_k) = \sum_{i=1}^k x_i / k$



Let A, B, C form a triangle, let x lie inside $\triangle ABC$.



Barycentric Coordinates

Recall: $\text{barycenter}(x_1, \dots, x_k) = \sum_{i=1}^k x_i / k$



Let A, B, C form a triangle, let x lie inside $\triangle ABC$.



Barycentric Coordinates

Recall: $\text{barycenter}(x_1, \dots, x_k) = \sum_{i=1}^k x_i / k$

Let A, B, C form a triangle, let x lie inside $\triangle ABC$. The **barycentric coordinates** of x with respect to $\triangle ABC$ are a triple $(\alpha, \beta, \gamma) \in \mathbb{R}_{\geq 0}^3$ such that





Barycentric Coordinates

Recall: $\text{barycenter}(x_1, \dots, x_k) = \sum_{i=1}^k x_i / k$

Let A, B, C form a triangle, let x lie inside $\triangle ABC$. The **barycentric coordinates** of x with respect to $\triangle ABC$ are a triple $(\alpha, \beta, \gamma) \in \mathbb{R}_{\geq 0}^3$ such that

■ $x = \alpha A + \beta B + \gamma C$ and





Barycentric Coordinates

Recall: $\text{barycenter}(x_1, \dots, x_k) = \sum_{i=1}^k x_i / k$

Let A, B, C form a triangle, let x lie inside $\triangle ABC$. The **barycentric coordinates** of x with respect to $\triangle ABC$ are a triple $(\alpha, \beta, \gamma) \in \mathbb{R}_{\geq 0}^3$ such that

■ $x = \alpha A + \beta B + \gamma C$ and

■ $\alpha + \beta + \gamma = 1$





Barycentric Coordinates

Recall: $\text{barycenter}(x_1, \dots, x_k) = \sum_{i=1}^k x_i / k$

Let A, B, C form a triangle, let x lie inside $\triangle ABC$. The **barycentric coordinates** of x with respect to $\triangle ABC$ are a triple $(\alpha, \beta, \gamma) \in \mathbb{R}_{\geq 0}^3$ such that

■ $x = \alpha A + \beta B + \gamma C$ and

■ $\alpha + \beta + \gamma = 1$





Barycentric Coordinates

Recall: $\text{barycenter}(x_1, \dots, x_k) = \sum_{i=1}^k x_i / k$

Let A, B, C form a triangle, let x lie inside $\triangle ABC$. The **barycentric coordinates** of x with respect to $\triangle ABC$ are a triple $(\alpha, \beta, \gamma) \in \mathbb{R}_{\geq 0}^3$ such that

■ $x = \alpha A + \beta B + \gamma C$ and

■ $\alpha + \beta + \gamma = 1$





Barycentric Coordinates

Recall: $\text{barycenter}(x_1, \dots, x_k) = \sum_{i=1}^k x_i / k$

Let A, B, C form a triangle, let x lie inside $\triangle ABC$. The **barycentric coordinates** of x with respect to $\triangle ABC$ are a triple $(\alpha, \beta, \gamma) \in \mathbb{R}_{\geq 0}^3$ such that

■ $x = \alpha A + \beta B + \gamma C$ and

■ $\alpha + \beta + \gamma = 1$





Barycentric Coordinates

Recall: $\text{barycenter}(x_1, \dots, x_k) = \sum_{i=1}^k x_i / k$

Let A, B, C form a triangle, let x lie inside $\triangle ABC$. The **barycentric coordinates** of x with respect to $\triangle ABC$ are a triple $(\alpha, \beta, \gamma) \in \mathbb{R}_{\geq 0}^3$ such that

■ $x = \alpha A + \beta B + \gamma C$ and

■ $\alpha + \beta + \gamma = 1$





Barycentric Coordinates

Recall: $\text{barycenter}(x_1, \dots, x_k) = \sum_{i=1}^k x_i / k$

Let A, B, C form a triangle, let x lie inside $\triangle ABC$. The **barycentric coordinates** of x with respect to $\triangle ABC$ are a triple $(\alpha, \beta, \gamma) \in \mathbb{R}_{\geq 0}^3$ such that

■ $x = \alpha A + \beta B + \gamma C$ and

■ $\alpha + \beta + \gamma = 1$





Barycentric Coordinates

Recall: $\text{barycenter}(x_1, \dots, x_k) = \sum_{i=1}^k x_i / k$

Let A, B, C form a triangle, let x lie inside $\triangle ABC$. The **barycentric coordinates** of x with respect to $\triangle ABC$ are a triple $(\alpha, \beta, \gamma) \in \mathbb{R}_{\geq 0}^3$ such that

■ $x = \alpha A + \beta B + \gamma C$ and

■ $\alpha + \beta + \gamma = 1$





Barycentric Coordinates

Recall: $\text{barycenter}(x_1, \dots, x_k) = \sum_{i=1}^k x_i / k$

Let A, B, C form a triangle, let x lie inside $\triangle ABC$. The **barycentric coordinates** of x with respect to $\triangle ABC$ are a triple $(\alpha, \beta, \gamma) \in \mathbb{R}_{\geq 0}^3$ such that

■ $x = \alpha A + \beta B + \gamma C$ and

■ $\alpha + \beta + \gamma = 1$





Barycentric Coordinates

Recall: $\text{barycenter}(x_1, \dots, x_k) = \sum_{i=1}^k x_i / k$

Let A, B, C form a triangle, let x lie inside $\triangle ABC$. The **barycentric coordinates** of x with respect to $\triangle ABC$ are a triple $(\alpha, \beta, \gamma) \in \mathbb{R}_{\geq 0}^3$ such that

■ $x = \alpha A + \beta B + \gamma C$ and

■ $\alpha + \beta + \gamma = 1$





Barycentric Coordinates

Recall: $\text{barycenter}(x_1, \dots, x_k) = \sum_{i=1}^k x_i / k$

Let A, B, C form a triangle, let x lie inside $\triangle ABC$. The **barycentric coordinates** of x with respect to $\triangle ABC$ are a triple $(\alpha, \beta, \gamma) \in \mathbb{R}_{\geq 0}^3$ such that

■ $x = \alpha A + \beta B + \gamma C$ and

■ $\alpha + \beta + \gamma = 1$




Barycentric Representation

A **barycentric representation** of a graph $G = (V, E)$ is an assignment of barycentric coordinates to the vertices of G :

$$f: V \rightarrow \mathbb{R}_{\geq 0}^3, v \mapsto (v_1, v_2, v_3)$$

with the following properties:




Barycentric Representation

A **barycentric representation** of a graph $G = (V, E)$ is an assignment of barycentric coordinates to the vertices of G :

$$f: V \rightarrow \mathbb{R}_{\geq 0}^3, v \mapsto (v_1, v_2, v_3)$$

with the following properties:

(B1) $v_1 + v_2 + v_3 = 1$ for all $v \in V$,



Barycentric Representation


A **barycentric representation** of a graph $G = (V, E)$ is an assignment of barycentric coordinates to the vertices of G :

$$f: V \rightarrow \mathbb{R}_{\geq 0}^3, v \mapsto (\textcolor{red}{v}_1, \textcolor{blue}{v}_2, \textcolor{green}{v}_3)$$

with the following properties:

(B1) $\textcolor{red}{v}_1 + \textcolor{blue}{v}_2 + \textcolor{green}{v}_3 = 1$ for all $v \in V$,

(B2) for each $xy \in E$



Barycentric Representation


A **barycentric representation** of a graph $G = (V, E)$ is an assignment of barycentric coordinates to the vertices of G :

$$f: V \rightarrow \mathbb{R}_{\geq 0}^3, v \mapsto (\textcolor{red}{v}_1, \textcolor{blue}{v}_2, \textcolor{green}{v}_3)$$

with the following properties:

(B1) $\textcolor{red}{v}_1 + \textcolor{blue}{v}_2 + \textcolor{green}{v}_3 = 1$ for all $v \in V$,

(B2) for each $xy \in E$ and each $z \in V \setminus \{x, y\}$




Barycentric Representation

A **barycentric representation** of a graph $G = (V, E)$ is an assignment of barycentric coordinates to the vertices of G :

$$f: V \rightarrow \mathbb{R}_{\geq 0}^3, v \mapsto (\textcolor{red}{v}_1, \textcolor{blue}{v}_2, \textcolor{green}{v}_3)$$

with the following properties:

- (B1) $\textcolor{red}{v}_1 + \textcolor{blue}{v}_2 + \textcolor{green}{v}_3 = 1$ for all $v \in V$,
- (B2) for each $xy \in E$ and each $z \in V \setminus \{x, y\}$
there exists $k \in \{\textcolor{red}{1}, \textcolor{blue}{2}, \textcolor{green}{3}\}$ with $x_k < z_k$ and $y_k < z_k$.




Barycentric Representation

A **barycentric representation** of a graph $G = (V, E)$ is an assignment of barycentric coordinates to the vertices of G :

$$f: V \rightarrow \mathbb{R}_{\geq 0}^3, v \mapsto (\textcolor{red}{v}_1, \textcolor{blue}{v}_2, \textcolor{green}{v}_3)$$

with the following properties:

- (B1) $\textcolor{red}{v}_1 + \textcolor{blue}{v}_2 + \textcolor{green}{v}_3 = 1$ for all $v \in V$,
- (B2) for each $xy \in E$ and each $z \in V \setminus \{x, y\}$
there exists $k \in \{\textcolor{red}{1}, \textcolor{blue}{2}, \textcolor{green}{3}\}$ with $x_k < z_k$ and $y_k < z_k$.




Barycentric Representation

A **barycentric representation** of a graph $G = (V, E)$ is an assignment of barycentric coordinates to the vertices of G :

$$f: V \rightarrow \mathbb{R}_{\geq 0}^3, v \mapsto (\textcolor{red}{v}_1, \textcolor{blue}{v}_2, \textcolor{green}{v}_3)$$

with the following properties:

- (B1) $\textcolor{red}{v}_1 + \textcolor{blue}{v}_2 + \textcolor{green}{v}_3 = 1$ for all $v \in V$,
- (B2) for each $xy \in E$ and each $z \in V \setminus \{x, y\}$
there exists $k \in \{\textcolor{red}{1}, \textcolor{blue}{2}, \textcolor{green}{3}\}$ with $x_k < z_k$ and $y_k < z_k$.




Barycentric Representation

A **barycentric representation** of a graph $G = (V, E)$ is an assignment of barycentric coordinates to the vertices of G :

$$f: V \rightarrow \mathbb{R}_{\geq 0}^3, v \mapsto (\textcolor{red}{v}_1, \textcolor{blue}{v}_2, \textcolor{green}{v}_3)$$

with the following properties:

- (B1) $\textcolor{red}{v}_1 + \textcolor{blue}{v}_2 + \textcolor{green}{v}_3 = 1$ for all $v \in V$,
- (B2) for each $xy \in E$ and each $z \in V \setminus \{x, y\}$
there exists $k \in \{\textcolor{red}{1}, \textcolor{blue}{2}, \textcolor{green}{3}\}$ with $x_k < z_k$ and $y_k < z_k$.




Barycentric Representation

A **barycentric representation** of a graph $G = (V, E)$ is an assignment of barycentric coordinates to the vertices of G :

$$f: V \rightarrow \mathbb{R}_{\geq 0}^3, v \mapsto (\textcolor{red}{v}_1, \textcolor{blue}{v}_2, \textcolor{green}{v}_3)$$

with the following properties:

- (B1) $\textcolor{red}{v}_1 + \textcolor{blue}{v}_2 + \textcolor{green}{v}_3 = 1$ for all $v \in V$,
- (B2) for each $xy \in E$ and each $z \in V \setminus \{x, y\}$
there exists $k \in \{\textcolor{red}{1}, \textcolor{blue}{2}, \textcolor{green}{3}\}$ with $x_k < z_k$ and $y_k < z_k$.



Barycentric Representations of Planar Graphs


Lemma.

Let $f : v \mapsto (v_1, v_2, v_3)$ be a barycentric representation of a planar graph G

Barycentric Representations of Planar Graphs

Lemma.

Let $f : v \mapsto (\color{red}{v_1}, \color{blue}{v_2}, \color{green}{v_3})$ be a barycentric representation of a planar graph G and let $\color{red}{A}, \color{blue}{B}, \color{green}{C} \in \mathbb{R}^2$ be in general position.




Barycentric Representations of Planar Graphs

Lemma.

Let $f : v \mapsto (v_1, v_2, v_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position. Then the mapping

$$\phi : v \in V \mapsto v_1 A + v_2 B + v_3 C$$

gives a **planar** drawing of G inside $\triangle ABC$.



Barycentric Representations of Planar Graphs


Lemma.

Let $f : v \mapsto (v_1, v_2, v_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position. Then the mapping

$$\phi : v \in V \mapsto v_1 A + v_2 B + v_3 C$$

gives a **planar** drawing of G inside $\triangle ABC$.

- No vertex x can lie on an edge $\{u, v\}$.



Barycentric Representations of Planar Graphs


Lemma.

Let $f : v \mapsto (v_1, v_2, v_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position. Then the mapping

$$\phi : v \in V \mapsto v_1 A + v_2 B + v_3 C$$

gives a **planar** drawing of G inside $\triangle ABC$.

- No vertex x can lie on an edge $\{u, v\}$.



Barycentric Representations of Planar Graphs


Lemma.

Let $f : v \mapsto (v_1, v_2, v_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position. Then the mapping

$$\phi : v \in V \mapsto v_1 A + v_2 B + v_3 C$$

gives a **planar** drawing of G inside $\triangle ABC$.

- No vertex x can lie on an edge $\{u, v\}$.



Barycentric Representations of Planar Graphs


Lemma.

Let $f : v \mapsto (\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position. Then the mapping

$$\phi : v \in V \mapsto \mathbf{v}_1 A + \mathbf{v}_2 B + \mathbf{v}_3 C$$

gives a **planar** drawing of G inside $\triangle ABC$.

- No vertex x can lie on an edge $\{u, v\}$.
- No pair of edges $\{u, v\}$ and $\{u', v'\}$ cross:



Barycentric Representations of Planar Graphs


Lemma.

Let $f : v \mapsto (v_1, v_2, v_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position. Then the mapping

$$\phi : v \in V \mapsto v_1 A + v_2 B + v_3 C$$

gives a **planar** drawing of G inside $\triangle ABC$.

- No vertex x can lie on an edge $\{u, v\}$.
- No pair of edges $\{u, v\}$ and $\{u', v'\}$ cross:



Barycentric Representations of Planar Graphs


Lemma.

Let $f : v \mapsto (\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position. Then the mapping

$$\phi : v \in V \mapsto \mathbf{v}_1 A + \mathbf{v}_2 B + \mathbf{v}_3 C$$

gives a **planar** drawing of G inside $\triangle ABC$.

- No vertex x can lie on an edge $\{u, v\}$.
- No pair of edges $\{u, v\}$ and $\{u', v'\}$ cross:



Barycentric Representations of Planar Graphs

Lemma.


Let $f : v \mapsto (\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position. Then the mapping

$$\phi : v \in V \mapsto \mathbf{v}_1 A + \mathbf{v}_2 B + \mathbf{v}_3 C$$

gives a **planar** drawing of G inside $\triangle ABC$.

- No vertex x can lie on an edge $\{u, v\}$.
- No pair of edges $\{u, v\}$ and $\{u', v'\}$ cross:

$$u'_i > u_i, v_i$$



Barycentric Representations of Planar Graphs

Lemma.


Let $f : v \mapsto (\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position. Then the mapping

$$\phi : v \in V \mapsto \mathbf{v}_1 A + \mathbf{v}_2 B + \mathbf{v}_3 C$$

gives a **planar** drawing of G inside $\triangle ABC$.

- No vertex x can lie on an edge $\{u, v\}$.
- No pair of edges $\{u, v\}$ and $\{u', v'\}$ cross:

$$u'_i > u_i, v_i \quad v'_j > u_j, v_j$$



Barycentric Representations of Planar Graphs

Lemma.


Let $f : v \mapsto (\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position. Then the mapping

$$\phi : v \in V \mapsto \mathbf{v}_1 A + \mathbf{v}_2 B + \mathbf{v}_3 C$$

gives a **planar** drawing of G inside $\triangle ABC$.

- No vertex x can lie on an edge $\{u, v\}$.
- No pair of edges $\{u, v\}$ and $\{u', v'\}$ cross:

$$u'_i > u_i, v_i \quad v'_j > u_j, v_j \quad u_k > u'_k, v'_k$$



Barycentric Representations of Planar Graphs

Lemma.


Let $f : v \mapsto (\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position. Then the mapping

$$\phi : v \in V \mapsto \mathbf{v}_1 A + \mathbf{v}_2 B + \mathbf{v}_3 C$$

gives a **planar** drawing of G inside $\triangle ABC$.

- No vertex x can lie on an edge $\{u, v\}$.
- No pair of edges $\{u, v\}$ and $\{u', v'\}$ cross:

$$u'_i > u_i, v_i \quad v'_j > u_j, v_j \quad u_k > u'_k, v'_k \quad v_l > u'_l, v'_l$$



Barycentric Representations of Planar Graphs

Lemma.

Let $f : v \mapsto (\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position. Then the mapping


$$\phi : v \in V \mapsto \mathbf{v}_1 A + \mathbf{v}_2 B + \mathbf{v}_3 C$$

gives a **planar** drawing of G inside $\triangle ABC$.

- No vertex x can lie on an edge $\{u, v\}$.
- No pair of edges $\{u, v\}$ and $\{u', v'\}$ cross:

$$u'_i > u_i, v_i \quad v'_j > u_j, v_j \quad u_k > u'_k, v'_k \quad v_l > u'_l, v'_l$$

$$\Rightarrow \{i, j\} \cap \{k, l\} = \emptyset$$



Barycentric Representations of Planar Graphs

Lemma.

Let $f : v \mapsto (\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position. Then the mapping

$$\phi : v \in V \mapsto \mathbf{v}_1 A + \mathbf{v}_2 B + \mathbf{v}_3 C$$


gives a **planar** drawing of G inside $\triangle ABC$.

- No vertex x can lie on an edge $\{u, v\}$.
- No pair of edges $\{u, v\}$ and $\{u', v'\}$ cross:

$$u'_i > u_i, v_i \quad v'_j > u_j, v_j \quad u_k > u'_k, v'_k \quad v_l > u'_l, v'_l$$

$$\Rightarrow \{i, j\} \cap \{k, l\} = \emptyset$$

$$\text{wlog } i = j = 2$$



Barycentric Representations of Planar Graphs

Lemma.

Let $f : v \mapsto (\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position. Then the mapping

$$\phi : v \in V \mapsto \mathbf{v}_1 A + \mathbf{v}_2 B + \mathbf{v}_3 C$$


gives a **planar** drawing of G inside $\triangle ABC$.

- No vertex x can lie on an edge $\{u, v\}$.
- No pair of edges $\{u, v\}$ and $\{u', v'\}$ cross:

$$u'_i > u_i, v_i \quad v'_j > u_j, v_j \quad u_k > u'_k, v'_k \quad v_l > u'_l, v'_l$$

$$\Rightarrow \{i, j\} \cap \{k, l\} = \emptyset$$

$$\text{wlog } i = j = 2 \Rightarrow u'_2, v'_2 > u_2, v_2$$



Barycentric Representations of Planar Graphs

Lemma.

Let $f : v \mapsto (\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position. Then the mapping

$$\phi : v \in V \mapsto \mathbf{v}_1 A + \mathbf{v}_2 B + \mathbf{v}_3 C$$


gives a **planar** drawing of G inside $\triangle ABC$.

- No vertex x can lie on an edge $\{u, v\}$.
- No pair of edges $\{u, v\}$ and $\{u', v'\}$ cross:

$$u'_i > u_i, v_i \quad v'_j > u_j, v_j \quad u_k > u'_k, v'_k \quad v_l > u'_l, v'_l$$

$$\Rightarrow \{i, j\} \cap \{k, l\} = \emptyset$$

$$\text{wlog } i = j = 2 \Rightarrow u'_2, v'_2 > u_2, v_2 \Rightarrow \text{separated by straight line}$$



Barycentric Representations of Planar Graphs

How to find barycentric representation?

Lemma.

Let $f : v \mapsto (v_1, v_2, v_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position. Then the mapping

$$\phi : v \in V \mapsto v_1 A + v_2 B + v_3 C$$


gives a **planar** drawing of G inside $\triangle ABC$.

- No vertex x can lie on an edge $\{u, v\}$.
- No pair of edges $\{u, v\}$ and $\{u', v'\}$ cross:

$$u'_i > u_i, v_i \quad v'_j > u_j, v_j \quad u_k > u'_k, v'_k \quad v_l > u'_l, v'_l$$


$$\Rightarrow \{i, j\} \cap \{k, l\} = \emptyset$$

$$\text{wlog } i = j = 2 \Rightarrow u'_2, v'_2 > u_2, v_2 \Rightarrow \text{separated by straight line}$$




Schnyder Labeling

Let $\phi : v \mapsto (v_1, v_2, v_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position.




Schnyder Labeling

Let $\phi : v \mapsto (v_1, v_2, v_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position.




Schnyder Labeling

Let $\phi : v \mapsto (v_1, v_2, v_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position.




Schnyder Labeling

Let $\phi : v \mapsto (\color{red}{v_1}, \color{blue}{v_2}, \color{green}{v_3})$ be a barycentric representation of a planar graph G and let $\color{red}{A}, \color{blue}{B}, \color{green}{C} \in \mathbb{R}^2$ be in general position.




Schnyder Labeling

Let $\phi : v \mapsto (v_1, v_2, v_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position.



Schnyder Labeling


Let $\phi : v \mapsto (v_1, v_2, v_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position.



Schnyder Labeling

Let $\phi : v \mapsto (\color{red}{v_1}, \color{blue}{v_2}, \color{green}{v_3})$ be a barycentric representation of a planar graph G and let $\color{red}{A}, \color{blue}{B}, \color{green}{C} \in \mathbb{R}^2$ be in general position.

We can label each angle in $\triangle xyz$ **uniquely** with $k \in \{\color{red}{1}, \color{blue}{2}, \color{green}{3}\}$.




Schnyder Labeling

Let $\phi : v \mapsto (v_1, v_2, v_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position.

We can label each angle in $\triangle xyz$ **uniquely** with $k \in \{1, 2, 3\}$.

A **Schnyder Labeling** of a plane triangulation G is a labeling of all internal angles with labels **1**, **2** and **3** such that:




Schnyder Labeling

Let $\phi : v \mapsto (v_1, v_2, v_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position.

We can label each angle in $\triangle xyz$ **uniquely** with $k \in \{1, 2, 3\}$.

A **Schnyder Labeling** of a plane triangulation G is a labeling of all internal angles with labels **1**, **2** and **3** such that:

Faces: The three angles of an internal face are labeled **1**, **2** and **3** in counterclockwise order.





Schnyder Labeling

Let $\phi : v \mapsto (v_1, v_2, v_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position.

We can label each angle in $\triangle xyz$ **uniquely** with $k \in \{1, 2, 3\}$.

A **Schnyder Labeling** of a plane triangulation G is a labeling of all internal angles with labels **1**, **2** and **3** such that:

Faces: The three angles of an internal face are labeled **1**, **2** and **3** in counterclockwise order.



Schnyder Labeling



Let $\phi : v \mapsto (v_1, v_2, v_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position.

We can label each angle in $\triangle xyz$ **uniquely** with $k \in \{1, 2, 3\}$.

A **Schnyder Labeling** of a plane triangulation G is a labeling of all internal angles with labels **1**, **2** and **3** such that:

Faces: The three angles of an internal face are labeled **1**, **2** and **3** in counterclockwise order.

Vertices: The ccw order of labels around each vertex consists of



Schnyder Labeling

Let $\phi : v \mapsto (v_1, v_2, v_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position.



We can label each angle in $\triangle xyz$ **uniquely** with $k \in \{1, 2, 3\}$.

A **Schnyder Labeling** of a plane triangulation G is a labeling of all internal angles with labels **1**, **2** and **3** such that:

Faces: The three angles of an internal face are labeled **1**, **2** and **3** in counterclockwise order.

Vertices: The ccw order of labels around each vertex consists of

- a nonempty interval of **1**'s



Schnyder Labeling

Let $\phi : v \mapsto (v_1, v_2, v_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position.



We can label each angle in $\triangle xyz$ **uniquely** with $k \in \{1, 2, 3\}$.

A **Schnyder Labeling** of a plane triangulation G is a labeling of all internal angles with labels **1**, **2** and **3** such that:

Faces: The three angles of an internal face are labeled **1**, **2** and **3** in counterclockwise order.

Vertices: The ccw order of labels around each vertex consists of

- a nonempty interval of **1**'s
- followed by a nonempty interval of **2**'s



Schnyder Labeling

Let $\phi : v \mapsto (v_1, v_2, v_3)$ be a barycentric representation of a planar graph G and let $A, B, C \in \mathbb{R}^2$ be in general position.



We can label each angle in $\triangle xyz$ **uniquely** with $k \in \{1, 2, 3\}$.

A **Schnyder Labeling** of a plane triangulation G is a labeling of all internal angles with labels **1**, **2** and **3** such that:

Faces: The three angles of an internal face are labeled **1**, **2** and **3** in counterclockwise order.



Vertices: The ccw order of labels around each vertex consists of

- a nonempty interval of **1**'s
- followed by a nonempty interval of **2**'s
- followed by a nonempty interval of **3**'s.




Schnyder Realizer

A Schnyder labeling induces an edge labeling.




Schnyder Realizer

A Schnyder labeling induces an edge labeling.




Schnyder Realizer

A Schnyder labeling induces an edge labeling.




Schnyder Realizer

A Schnyder labeling induces an edge labeling.




Schnyder Realizer

A Schnyder labeling induces an edge labeling.



Schnyder Realizer


A Schnyder labeling induces an edge labeling.



Schnyder Realizer

A Schnyder labeling induces an edge labeling.


A **Schnyder Realizer** (or **Wood**) of a plane triangulation $G = (V, E)$ is a partition of the inner edges of E into three sets of oriented edges T_1 , T_2 , T_3



Schnyder Realizer

A Schnyder labeling induces an edge labeling.

A **Schnyder Realizer** (or **Wood**) of a plane triangulation $G = (V, E)$ is a partition of the inner edges of E into three sets of oriented edges T_1 , T_2 , T_3 such that for each inner vertex $v \in V$ holds:




Schnyder Realizer

A Schnyder labeling induces an edge labeling.

A **Schnyder Realizer** (or **Wood**) of a plane triangulation $G = (V, E)$ is a partition of the inner edges of E into three sets of oriented edges T_1 , T_2 , T_3 such that for each inner vertex $v \in V$ holds:

- v has one outgoing edge in each of T_1 , T_2 , and T_3 .




Schnyder Realizer

A Schnyder labeling induces an edge labeling.

A **Schnyder Realizer** (or **Wood**) of a plane triangulation $G = (V, E)$ is a partition of the inner edges of E into three sets of oriented edges T_1 , T_2 , T_3 such that for each inner vertex $v \in V$ holds:

- v has one outgoing edge in each of T_1 , T_2 , and T_3 .
- The ccw order of edges around v is:




Schnyder Realizer

A Schnyder labeling induces an edge labeling.

A **Schnyder Realizer** (or **Wood**) of a plane triangulation $G = (V, E)$ is a partition of the inner edges of E into three sets of oriented edges T_1 , T_2 , T_3 such that for each inner vertex $v \in V$ holds:

- v has one outgoing edge in each of T_1 , T_2 , and T_3 .
- The ccw order of edges around v is:
leaving in T_1




Schnyder Realizer

A Schnyder labeling induces an edge labeling.

A **Schnyder Realizer** (or **Wood**) of a plane triangulation $G = (V, E)$ is a partition of the inner edges of E into three sets of oriented edges T_1 , T_2 , T_3 such that for each inner vertex $v \in V$ holds:

- v has one outgoing edge in each of T_1 , T_2 , and T_3 .
- The ccw order of edges around v is:
leaving in T_1 , entering in T_3




Schnyder Realizer

A Schnyder labeling induces an edge labeling.

A **Schnyder Realizer** (or **Wood**) of a plane triangulation $G = (V, E)$ is a partition of the inner edges of E into three sets of oriented edges T_1 , T_2 , T_3 such that for each inner vertex $v \in V$ holds:

- v has one outgoing edge in each of T_1 , T_2 , and T_3 .
- The ccw order of edges around v is:
leaving in T_1 , entering in T_3 , leaving in T_2 .




Schnyder Realizer

A Schnyder labeling induces an edge labeling.

A **Schnyder Realizer** (or **Wood**) of a plane triangulation $G = (V, E)$ is a partition of the inner edges of E into three sets of oriented edges T_1 , T_2 , T_3 such that for each inner vertex $v \in V$ holds:

- v has one outgoing edge in each of T_1 , T_2 , and T_3 .
- The ccw order of edges around v is:
 leaving in T_1 , entering in T_3 , leaving in T_2 ,
 entering in T_1




Schnyder Realizer

A Schnyder labeling induces an edge labeling.

A **Schnyder Realizer** (or **Wood**) of a plane triangulation $G = (V, E)$ is a partition of the inner edges of E into three sets of oriented edges T_1 , T_2 , T_3 such that for each inner vertex $v \in V$ holds:

- v has one outgoing edge in each of T_1 , T_2 , and T_3 .
- The ccw order of edges around v is:
 leaving in T_1 , entering in T_3 , leaving in T_2 ,
 entering in T_1 , leaving in T_3




Schnyder Realizer



A Schnyder labeling induces an edge labeling.

A **Schnyder Realizer** (or **Wood**) of a plane triangulation $G = (V, E)$ is a partition of the inner edges of E into three sets of oriented edges T_1 , T_2 , T_3 such that for each inner vertex $v \in V$ holds:



- v has one outgoing edge in each of T_1 , T_2 , and T_3 .
- The ccw order of edges around v is:
 leaving in T_1 , entering in T_3 , leaving in T_2 ,
 entering in T_1 , leaving in T_3 , entering in T_2 .




Schnyder Realizer – Example and Properties




Schnyder Realizer – Example and Properties




Schnyder Realizer – Example and Properties




Schnyder Realizer – Example and Properties





Schnyder Realizer – Example and Properties



Schnyder Realizer – Example and Properties





Schnyder Realizer – Example and Properties





- All inner edges incident to a , b , and c are incoming in the same color.

Schnyder Realizer – Example and Properties





- All inner edges incident to a , b , and c are incoming in the same color.
- T_1 , T_2 , and T_3 are trees on all inner vertices and one outer vertex each (as its root).

Schnyder Realizer – Example and Properties





- All inner edges incident to a , b , and c are incoming in the same color.
- T_1 , T_2 , and T_3 are trees on all inner vertices and one outer vertex each (as its root).

Schnyder Realizer – Example and Properties




- All inner edges incident to a , b , and c are incoming in the same color.
- T_1 , T_2 , and T_3 are trees on all inner vertices and one outer vertex each (as its root).

Schnyder Realizer – Example and Properties




- All inner edges incident to a , b , and c are incoming in the same color.
- T_1 , T_2 , and T_3 are trees on all inner vertices and one outer vertex each (as its root).


Schnyder Realizer – Existence




Schnyder Realizer – Existence




Schnyder Realizer – Existence



Schnyder Realizer – Existence




Schnyder Realizer – Existence




...requires that a and x have exactly 2 common neighbors.

Schnyder Realizer – Existence



...requires that a and x have exactly 2 common neighbors.

Schnyder Realizer – Existence




...requires that a and x have exactly 2 common neighbors.

Schnyder Realizer – Existence

Lemma.

[Kampen 1976]

Let G be a plane triangulation with vertices a, b, c on the outer face. There exists a **contractible edge** $\{a, x\}$ in G , $x \neq b, c$.



...requires that a and x have exactly 2 common neighbors.

Schnyder Realizer – Existence


Lemma.

[Kampen 1976]

Let G be a plane triangulation with vertices a, b, c on the outer face. There exists a **contractible edge** $\{a, x\}$ in G , $x \neq b, c$.

Theorem.

Every plane triangulation has a Schnyder Labeling and Realizer.



...requires that a and x have exactly 2 common neighbors.

Schnyder Realizer – Existence

Lemma.


[Kampen 1976]

Let G be a plane triangulation with vertices a, b, c on the outer face. There exists a **contractible edge** $\{a, x\}$ in G , $x \neq b, c$.

Theorem.

Every plane triangulation has a Schnyder Labeling and Realizer.

Proof by induction on # vertices via edge contractions.



...requires that a and x have exactly 2 common neighbors.

Schnyder Realizer – Existence

Lemma.


[Kampen 1976]

Let G be a plane triangulation with vertices a, b, c on the outer face. There exists a **contractible edge** $\{a, x\}$ in G , $x \neq b, c$.

Theorem.

Every plane triangulation has a Schnyder Labeling and Realizer.

Proof by induction on # vertices via edge contractions.



...requires that a and x have exactly 2 common neighbors.

Schnyder Realizer – Existence

Lemma.


[Kampen 1976]

Let G be a plane triangulation with vertices a, b, c on the outer face. There exists a **contractible edge** $\{a, x\}$ in G , $x \neq b, c$.

Theorem.

Every plane triangulation has a Schnyder Labeling and Realizer.

Proof by induction on # vertices via edge contractions.



...requires that a and x have exactly 2 common neighbors.

Schnyder Realizer – Existence

Lemma.


[Kampen 1976]

Let G be a plane triangulation with vertices a, b, c on the outer face. There exists a **contractible edge** $\{a, x\}$ in G , $x \neq b, c$.

Theorem.

Every plane triangulation has a Schnyder Labeling and Realizer.

Proof by induction on # vertices via edge contractions.



...requires that a and x have exactly 2 common neighbors.

Schnyder Realizer – Existence

Lemma.


[Kampen 1976]

Let G be a plane triangulation with vertices a, b, c on the outer face. There exists a **contractible edge** $\{a, x\}$ in G , $x \neq b, c$.

Theorem.

Every plane triangulation has a Schnyder Labeling and Realizer.

Proof by induction on # vertices via edge contractions.



...requires that a and x have exactly 2 common neighbors.

Schnyder Realizer – Existence

Lemma.


[Kampen 1976]

Let G be a plane triangulation with vertices a, b, c on the outer face. There exists a **contractible edge** $\{a, x\}$ in G , $x \neq b, c$.

Theorem.

Every plane triangulation has a Schnyder Labeling and Realizer.

Proof by induction on # vertices via edge contractions.



...requires that a and x have exactly 2 common neighbors.

Schnyder Realizer – Existence

Lemma.


[Kampen 1976]

Let G be a plane triangulation with vertices a, b, c on the outer face. There exists a **contractible edge** $\{a, x\}$ in G , $x \neq b, c$.

Theorem.

Every plane triangulation has a Schnyder Labeling and Realizer.

Proof by induction on # vertices via edge contractions.



...requires that a and x have exactly 2 common neighbors.

Schnyder Realizer – Existence

Lemma.


[Kampen 1976]

Let G be a plane triangulation with vertices a, b, c on the outer face. There exists a **contractible edge** $\{a, x\}$ in G , $x \neq b, c$.

Theorem.

Every plane triangulation has a Schnyder Labeling and Realizer.

Proof by induction on # vertices via edge contractions.



...requires that a and x have exactly 2 common neighbors.

Schnyder Realizer – Existence

Lemma.


[Kampen 1976]

Let G be a plane triangulation with vertices a, b, c on the outer face. There exists a **contractible edge** $\{a, x\}$ in G , $x \neq b, c$.

Theorem.

Every plane triangulation has a Schnyder Labeling and Realizer.


Proof by induction on # vertices via edge contractions.



Constructive proof
can be used as
algorithm to compute
a Schnyder labeling.
It can be
implemented in $\mathcal{O}(n)$
time.


...requires that a and x have exactly 2 common neighbors.

Schnyder Realizer – More Properties




Schnyder Realizer – More Properties

■ From each vertex v there exists




Schnyder Realizer – More Properties

- From each vertex v there exists a directed **red** path $P_1(v)$ to a ,




Schnyder Realizer – More Properties

- From each vertex v there exists a directed **red** path $P_1(v)$ to a , a directed **blue** path $P_2(v)$ to b , and




Schnyder Realizer – More Properties

- From each vertex v there exists a directed **red** path $P_1(v)$ to a , a directed **blue** path $P_2(v)$ to b , and a directed **green** path $P_3(v)$ to c .




Schnyder Realizer – More Properties



- From each vertex v there exists a directed **red** path $P_1(v)$ to a , a directed **blue** path $P_2(v)$ to b , and a directed **green** path $P_3(v)$ to c .

$P_i(v)$: path from v to root of T_i .

Schnyder Realizer – More Properties




- From each vertex v there exists a directed **red** path $P_1(v)$ to a , a directed **blue** path $P_2(v)$ to b , and a directed **green** path $P_3(v)$ to c .

$P_i(v)$: path from v to root of T_i .

Lemma.

- $P_1(v)$, $P_2(v)$, $P_3(v)$ cross only at v .

Schnyder Realizer – More Properties




- From each vertex v there exists a directed **red** path $P_1(v)$ to a , a directed **blue** path $P_2(v)$ to b , and a directed **green** path $P_3(v)$ to c .

$P_i(v)$: path from v to root of T_i .

Lemma.

- $P_1(v)$, $P_2(v)$, $P_3(v)$ cross only at v .

Schnyder Realizer – More Properties




- From each vertex v there exists a directed **red** path $P_1(v)$ to a , a directed **blue** path $P_2(v)$ to b , and a directed **green** path $P_3(v)$ to c .

$P_i(v)$: path from v to root of T_i .

Lemma.

- $P_1(v)$, $P_2(v)$, $P_3(v)$ cross only at v .

Schnyder Realizer – More Properties




- From each vertex v there exists a directed **red** path $P_1(v)$ to a , a directed **blue** path $P_2(v)$ to b , and a directed **green** path $P_3(v)$ to c .

$P_i(v)$: path from v to root of T_i .

Lemma.

- $P_1(v)$, $P_2(v)$, $P_3(v)$ cross only at v .

Schnyder Realizer – More Properties




- From each vertex v there exists a directed **red** path $P_1(v)$ to a , a directed **blue** path $P_2(v)$ to b , and a directed **green** path $P_3(v)$ to c .

$P_i(v)$: path from v to root of T_i .

Lemma.

- $P_1(v)$, $P_2(v)$, $P_3(v)$ cross only at v .

Schnyder Realizer – More Properties




- From each vertex v there exists a directed **red** path $P_1(v)$ to a , a directed **blue** path $P_2(v)$ to b , and a directed **green** path $P_3(v)$ to c .

$P_i(v)$: path from v to root of T_i .

Lemma.

- $P_1(v)$, $P_2(v)$, $P_3(v)$ cross only at v .

Schnyder Realizer – More Properties




- From each vertex v there exists a directed **red** path $P_1(v)$ to a , a directed **blue** path $P_2(v)$ to b , and a directed **green** path $P_3(v)$ to c .

$P_i(v)$: path from v to root of T_i .

Lemma.

- $P_1(v)$, $P_2(v)$, $P_3(v)$ cross only at v .

Schnyder Realizer – More Properties



- From each vertex v there exists a directed **red** path $P_1(v)$ to a , a directed **blue** path $P_2(v)$ to b , and a directed **green** path $P_3(v)$ to c .


$P_i(v)$: path from v to root of T_i .

$R_1(v)$: set of faces contained in P_2, bc, P_3 .

Lemma.

- $P_1(v), P_2(v), P_3(v)$ cross only at v .

Schnyder Realizer – More Properties



- From each vertex v there exists a directed **red** path $P_1(v)$ to a , a directed **blue** path $P_2(v)$ to b , and a directed **green** path $P_3(v)$ to c .

$P_i(v)$: path from v to root of T_i .


$R_1(v)$: set of faces contained in P_2, bc, P_3 .

$R_2(v)$: set of faces contained in P_3, ca, P_1 .

Lemma.

- $P_1(v), P_2(v), P_3(v)$ cross only at v .

Schnyder Realizer – More Properties



- From each vertex v there exists a directed **red** path $P_1(v)$ to a , a directed **blue** path $P_2(v)$ to b , and a directed **green** path $P_3(v)$ to c .

$P_i(v)$: path from v to root of T_i .

$R_1(v)$: set of faces contained in P_2, bc, P_3 .


$R_2(v)$: set of faces contained in P_3, ca, P_1 .

$R_3(v)$: set of faces contained in P_1, ab, P_2 .

Lemma.

- $P_1(v), P_2(v), P_3(v)$ cross only at v .

Schnyder Realizer – More Properties



- From each vertex v there exists a directed **red** path $P_1(v)$ to a , a directed **blue** path $P_2(v)$ to b , and a directed **green** path $P_3(v)$ to c .

$P_i(v)$: path from v to root of T_i .

$R_1(v)$: set of faces contained in P_2, bc, P_3 .


$R_2(v)$: set of faces contained in P_3, ca, P_1 .

$R_3(v)$: set of faces contained in P_1, ab, P_2 .

Lemma.

- $P_1(v), P_2(v), P_3(v)$ cross only at v .
- For inner vertices $u \neq v$ it holds that $u \in R_i(v) \Rightarrow R_i(u) \subsetneq R_i(v)$.

Schnyder Realizer – More Properties



- From each vertex v there exists a directed **red** path $P_1(v)$ to a , a directed **blue** path $P_2(v)$ to b , and a directed **green** path $P_3(v)$ to c .

$P_i(v)$: path from v to root of T_i .

$R_1(v)$: set of faces contained in P_2, bc, P_3 .


$R_2(v)$: set of faces contained in P_3, ca, P_1 .

$R_3(v)$: set of faces contained in P_1, ab, P_2 .

Lemma.

- $P_1(v), P_2(v), P_3(v)$ cross only at v .
- For inner vertices $u \neq v$ it holds that $u \in R_i(v) \Rightarrow R_i(u) \subsetneq R_i(v)$.

Schnyder Realizer – More Properties



- From each vertex v there exists a directed **red** path $P_1(v)$ to a , a directed **blue** path $P_2(v)$ to b , and a directed **green** path $P_3(v)$ to c .

$P_i(v)$: path from v to root of T_i .

$R_1(v)$: set of faces contained in P_2, bc, P_3 .


$R_2(v)$: set of faces contained in P_3, ca, P_1 .

$R_3(v)$: set of faces contained in P_1, ab, P_2 .

Lemma.

- $P_1(v), P_2(v), P_3(v)$ cross only at v .
- For inner vertices $u \neq v$ it holds that $u \in R_i(v) \Rightarrow R_i(u) \subsetneq R_i(v)$.

Schnyder Realizer – More Properties



- From each vertex v there exists a directed **red** path $P_1(v)$ to a , a directed **blue** path $P_2(v)$ to b , and a directed **green** path $P_3(v)$ to c .

$P_i(v)$: path from v to root of T_i .

$R_1(v)$: set of faces contained in P_2, bc, P_3 .


$R_2(v)$: set of faces contained in P_3, ca, P_1 .

$R_3(v)$: set of faces contained in P_1, ab, P_2 .

Lemma.

- $P_1(v), P_2(v), P_3(v)$ cross only at v .
- For inner vertices $u \neq v$ it holds that $u \in R_i(v) \Rightarrow R_i(u) \subsetneq R_i(v)$.

Schnyder Realizer – More Properties



- From each vertex v there exists a directed **red** path $P_1(v)$ to a , a directed **blue** path $P_2(v)$ to b , and a directed **green** path $P_3(v)$ to c .

$P_i(v)$: path from v to root of T_i .

$R_1(v)$: set of faces contained in P_2, bc, P_3 .


$R_2(v)$: set of faces contained in P_3, ca, P_1 .

$R_3(v)$: set of faces contained in P_1, ab, P_2 .

Lemma.

- $P_1(v), P_2(v), P_3(v)$ cross only at v .
- For inner vertices $u \neq v$ it holds that $u \in R_i(v) \Rightarrow R_i(u) \subsetneq R_i(v)$.
- $|R_1(v)| + |R_2(v)| + |R_3(v)| =$

Schnyder Realizer – More Properties



- From each vertex v there exists a directed **red** path $P_1(v)$ to a , a directed **blue** path $P_2(v)$ to b , and a directed **green** path $P_3(v)$ to c .

$P_i(v)$: path from v to root of T_i .

$R_1(v)$: set of faces contained in P_2, bc, P_3 .


$R_2(v)$: set of faces contained in P_3, ca, P_1 .

$R_3(v)$: set of faces contained in P_1, ab, P_2 .

Lemma.

- $P_1(v), P_2(v), P_3(v)$ cross only at v .
- For inner vertices $u \neq v$ it holds that $u \in R_i(v) \Rightarrow R_i(u) \subsetneq R_i(v)$.
- $|R_1(v)| + |R_2(v)| + |R_3(v)| = 2n - 5$

Schnyder Realizer – More Properties



- From each vertex v there exists a directed **red** path $P_1(v)$ to a , a directed **blue** path $P_2(v)$ to b , and a directed **green** path $P_3(v)$ to c .

$P_i(v)$: path from v to root of T_i .

$R_1(v)$: set of faces contained in P_2, bc, P_3 .


$R_2(v)$: set of faces contained in P_3, ca, P_1 .

$R_3(v)$: set of faces contained in P_1, ab, P_2 .

Lemma.

- $P_1(v), P_2(v), P_3(v)$ cross only at v .
- For inner vertices $u \neq v$ it holds that $u \in R_i(v) \Rightarrow R_i(u) \subsetneq R_i(v)$.
- $|R_1(v)| + |R_2(v)| + |R_3(v)| = 2n - 5$

Schnyder Realizer – More Properties



- From each vertex v there exists a directed **red** path $P_1(v)$ to a , a directed **blue** path $P_2(v)$ to b , and a directed **green** path $P_3(v)$ to c .

$P_i(v)$: path from v to root of T_i .

$R_1(v)$: set of faces contained in P_2, bc, P_3 .

$R_2(v)$: set of faces contained in P_3, ca, P_1 .

$R_3(v)$: set of faces contained in P_1, ab, P_2 .

Lemma.

- $P_1(v), P_2(v), P_3(v)$ cross only at v .
- For inner vertices $u \neq v$ it holds that $u \in R_i(v) \Rightarrow R_i(u) \subsetneq R_i(v)$.
- $|R_1(v)| + |R_2(v)| + |R_3(v)| = 2n - 5$

Schnyder Drawing

Theorem.

[Schnyder '89]

For a plane triangulation G , the mapping

$$f: v \mapsto (v_1, v_2, v_3) = \frac{1}{2n-5} (|R_1(v)|, |R_2(v)|, |R_3(v)|)$$

is a barycentric representation of G , which thus gives a planar straight-line drawing of G

Schnyder Drawing

Theorem.

[Schnyder '89]

For a plane triangulation G , the mapping

$$f: v \mapsto (v_1, v_2, v_3) = \frac{1}{2n-5} (|R_1(v)|, |R_2(v)|, |R_3(v)|)$$

is a barycentric representation of G , which thus gives a planar straight-line drawing of G

(B1) $v_1 + v_2 + v_3 = 1$ for all $v \in V$

Schnyder Drawing

Theorem.

[Schnyder '89]

For a plane triangulation G , the mapping

$$f: v \mapsto (v_1, v_2, v_3) = \frac{1}{2n-5} (|R_1(v)|, |R_2(v)|, |R_3(v)|)$$

is a barycentric representation of G , which thus gives a planar straight-line drawing of G

(B1) $v_1 + v_2 + v_3 = 1$ for all $v \in V$ ✓

Schnyder Drawing

Theorem.

[Schnyder '89]


For a plane triangulation G , the mapping

$$f: v \mapsto (v_1, v_2, v_3) = \frac{1}{2n-5} (|R_1(v)|, |R_2(v)|, |R_3(v)|)$$

is a barycentric representation of G , which thus gives a planar straight-line drawing of G

(B1) $v_1 + v_2 + v_3 = 1$ for all $v \in V$ ✓

(B2) for each $\{x, y\} \in E$ and each $z \in V \setminus \{x, y\}$
there exists $k \in \{1, 2, 3\}$ with $x_k < z_k$ and $y_k < z_k$



Schnyder Drawing

Theorem.

[Schnyder '89]

For a plane triangulation G , the mapping


$$f: v \mapsto (\color{red}{v_1}, \color{blue}{v_2}, \color{green}{v_3}) = \frac{1}{2n-5} (|\color{red}{R_1}(v)|, |\color{blue}{R_2}(v)|, |\color{green}{R_3}(v)|)$$

is a barycentric representation of G , which thus gives a planar straight-line drawing of G

(B1) $\color{red}{v_1} + \color{blue}{v_2} + \color{green}{v_3} = 1$ for all $v \in V$ ✓

(B2) for each $\{x, y\} \in E$ and each $z \in V \setminus \{x, y\}$
there exists $k \in \{\color{red}{1}, \color{blue}{2}, \color{green}{3}\}$ with $x_k < z_k$ and $y_k < z_k$

■ $\{x, y\}$ must lie in some $R_i(z)$ for $i \in \{\color{red}{1}, \color{blue}{2}, \color{green}{3}\}$



Schnyder Drawing

Theorem.

[Schnyder '89]

For a plane triangulation G , the mapping


$$f: v \mapsto (\color{red}{v_1}, \color{blue}{v_2}, \color{green}{v_3}) = \frac{1}{2n-5} (|\color{red}{R_1(v)}|, |\color{blue}{R_2(v)}|, |\color{green}{R_3(v)}|)$$

is a barycentric representation of G , which thus gives a planar straight-line drawing of G

(B1) $\color{red}{v_1} + \color{blue}{v_2} + \color{green}{v_3} = 1$ for all $v \in V$ ✓

(B2) for each $\{x, y\} \in E$ and each $z \in V \setminus \{x, y\}$
there exists $k \in \{\color{red}{1}, \color{blue}{2}, \color{green}{3}\}$ with $x_k < z_k$ and $y_k < z_k$

■ $\{x, y\}$ must lie in some $R_i(z)$ for $i \in \{\color{red}{1}, \color{blue}{2}, \color{green}{3}\}$



Schnyder Drawing

Theorem.


[Schnyder '89]

For a plane triangulation G , the mapping

$$f: v \mapsto (\color{red}{v_1}, \color{blue}{v_2}, \color{green}{v_3}) = \frac{1}{2n-5} (|\color{red}{R_1(v)}|, |\color{blue}{R_2(v)}|, |\color{green}{R_3(v)}|)$$

is a barycentric representation of G , which thus gives a planar straight-line drawing of G

- (B1) $\color{red}{v_1} + \color{blue}{v_2} + \color{green}{v_3} = 1$ for all $v \in V$ ✓
- (B2) for each $\{x, y\} \in E$ and each $z \in V \setminus \{x, y\}$
 there exists $k \in \{\color{red}{1}, \color{blue}{2}, \color{green}{3}\}$ with $x_k < z_k$ and $y_k < z_k$
 - $\{x, y\}$ must lie in some $R_i(z)$ for $i \in \{\color{red}{1}, \color{blue}{2}, \color{green}{3}\}$



Schnyder Drawing

Theorem.


[Schnyder '89]

For a plane triangulation G , the mapping

$$f: v \mapsto (\color{red}{v_1}, \color{blue}{v_2}, \color{green}{v_3}) = \frac{1}{2n-5} (|\color{red}{R_1(v)}|, |\color{blue}{R_2(v)}|, |\color{green}{R_3(v)}|)$$

is a barycentric representation of G , which thus gives a planar straight-line drawing of G

- (B1) $\color{red}{v_1} + \color{blue}{v_2} + \color{green}{v_3} = 1$ for all $v \in V$ ✓
- (B2) for each $\{x, y\} \in E$ and each $z \in V \setminus \{x, y\}$
there exists $k \in \{\color{red}{1}, \color{blue}{2}, \color{green}{3}\}$ with $x_k < z_k$ and $y_k < z_k$
 - $\{x, y\}$ must lie in some $R_i(z)$ for $i \in \{\color{red}{1}, \color{blue}{2}, \color{green}{3}\}$
 - For inner vertices $u \neq v$ it holds
that $u \in R_i(v) \Rightarrow R_i(u) \subsetneq R_i(v)$.



Schnyder Drawing

Theorem.


[Schnyder '89]

For a plane triangulation G , the mapping

$$f: v \mapsto (\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3) = \frac{1}{2n-5} (|R_1(v)|, |R_2(v)|, |R_3(v)|)$$

is a barycentric representation of G , which thus gives a planar straight-line drawing of G

- (B1) $\mathbf{v}_1 + \mathbf{v}_2 + \mathbf{v}_3 = 1$ for all $v \in V$ ✓
- (B2) for each $\{x, y\} \in E$ and each $z \in V \setminus \{x, y\}$
there exists $k \in \{1, 2, 3\}$ with $x_k < z_k$ and $y_k < z_k$ ✓
 - $\{x, y\}$ must lie in some $R_i(z)$ for $i \in \{1, 2, 3\}$
 - For inner vertices $u \neq v$ it holds
that $u \in R_i(v) \Rightarrow R_i(u) \subsetneq R_i(v)$.



Schnyder Drawing

Set $A = (0, 0)$, $B = (2n - 5, 0)$, and $C = (0, 2n - 5)$.

Theorem.

[Schnyder '89]

For a plane triangulation G , the mapping

$$f: v \mapsto (\color{red}{v_1}, \color{blue}{v_2}, \color{green}{v_3}) = \frac{1}{2n-5} (|\color{red}{R_1(v)}|, |\color{blue}{R_2(v)}|, |\color{green}{R_3(v)}|)$$


is a barycentric representation of G , which thus gives a planar straight-line drawing of G

(B1) $\color{red}{v_1} + \color{blue}{v_2} + \color{green}{v_3} = 1$ for all $v \in V$ ✓

(B2) for each $\{x, y\} \in E$ and each $z \in V \setminus \{x, y\}$
there exists $k \in \{\color{red}{1}, \color{blue}{2}, \color{green}{3}\}$ with $x_k < z_k$ and $y_k < z_k$ ✓

■ $\{x, y\}$ must lie in some $R_i(z)$ for $i \in \{\color{red}{1}, \color{blue}{2}, \color{green}{3}\}$

■ For inner vertices $u \neq v$ it holds
that $u \in R_i(v) \Rightarrow R_i(u) \subsetneq R_i(v)$.



Schnyder Drawing

Set $A = (0,0)$, $B = (2n - 5, 0)$, and $C = (0, 2n - 5)$.

Theorem.

[Schnyder '89]

For a plane triangulation G , the mapping

$$f: v \mapsto (\color{red}{v_1}, \color{blue}{v_2}, \color{green}{v_3}) = \frac{1}{2n-5} (|R_1(v)|, |R_2(v)|, |R_3(v)|)$$


is a barycentric representation of G , which thus gives a planar straight-line drawing of G on the $(2n - 5) \times (2n - 5)$ grid.

(B1) $\color{red}{v_1} + \color{blue}{v_2} + \color{green}{v_3} = 1$ for all $v \in V$ ✓


(B2) for each $\{x, y\} \in E$ and each $z \in V \setminus \{x, y\}$
there exists $k \in \{\color{red}{1}, \color{blue}{2}, \color{green}{3}\}$ with $x_k < z_k$ and $y_k < z_k$ ✓

■ $\{x, y\}$ must lie in some $R_i(z)$ for $i \in \{\color{red}{1}, \color{blue}{2}, \color{green}{3}\}$


■ For inner vertices $u \neq v$ it holds
that $u \in R_i(v) \Rightarrow R_i(u) \subsetneq R_i(v)$.



Schnyder Drawing – Example




Schnyder Drawing – Example



$$n = 7, 2n - 5 = 9$$


Schnyder Drawing – Example



$$n = 7, 2n - 5 = 9$$

$$f(a) = (\textcolor{red}{9}, \textcolor{blue}{0}, \textcolor{green}{0})$$

Schnyder Drawing – Example




$$n = 7, 2n - 5 = 9$$

$$f(a) = (\textcolor{red}{9}, \textcolor{blue}{0}, \textcolor{green}{0})$$

$$f(b) = (\textcolor{red}{0}, \textcolor{blue}{9}, \textcolor{green}{0})$$

Schnyder Drawing – Example




$$n = 7, 2n - 5 = 9$$

$$f(a) = (\textcolor{red}{9}, \textcolor{blue}{0}, \textcolor{green}{0})$$

$$f(b) = (\textcolor{red}{0}, \textcolor{blue}{9}, \textcolor{green}{0})$$

$$f(c) = (\textcolor{red}{0}, \textcolor{blue}{0}, \textcolor{green}{9})$$

Schnyder Drawing – Example




$$n = 7, 2n - 5 = 9 \quad f(d) = (\textcolor{red}{2}, \textcolor{blue}{6}, \textcolor{green}{1})$$

$$f(\textcolor{red}{a}) = (\textcolor{red}{9}, \textcolor{blue}{0}, \textcolor{green}{0})$$

$$f(\textcolor{blue}{b}) = (\textcolor{red}{0}, \textcolor{blue}{9}, \textcolor{green}{0})$$

$$f(\textcolor{green}{c}) = (\textcolor{red}{0}, \textcolor{blue}{0}, \textcolor{green}{9})$$

Schnyder Drawing – Example




$$n = 7, 2n - 5 = 9 \quad f(d) = (\textcolor{red}{2}, \textcolor{blue}{6}, \textcolor{green}{1})$$

$$f(\textcolor{red}{a}) = (\textcolor{red}{9}, \textcolor{blue}{0}, \textcolor{green}{0}) \quad f(e) = (\textcolor{red}{5}, \textcolor{blue}{2}, \textcolor{green}{2})$$

$$f(\textcolor{blue}{b}) = (\textcolor{red}{0}, \textcolor{blue}{9}, \textcolor{green}{0})$$

$$f(\textcolor{green}{c}) = (\textcolor{red}{0}, \textcolor{blue}{0}, \textcolor{green}{9})$$

Schnyder Drawing – Example




$$n = 7, 2n - 5 = 9 \quad f(d) = (\textcolor{red}{2}, \textcolor{blue}{6}, \textcolor{green}{1})$$

$$f(\textcolor{red}{a}) = (\textcolor{red}{9}, \textcolor{blue}{0}, \textcolor{green}{0}) \quad f(e) = (\textcolor{red}{5}, \textcolor{blue}{2}, \textcolor{green}{2})$$

$$f(\textcolor{blue}{b}) = (\textcolor{red}{0}, \textcolor{blue}{9}, \textcolor{green}{0}) \quad f(g) = (\textcolor{red}{1}, \textcolor{blue}{2}, \textcolor{green}{6})$$

$$f(\textcolor{green}{c}) = (\textcolor{red}{0}, \textcolor{blue}{0}, \textcolor{green}{9})$$

Schnyder Drawing – Example





$$n = 7, 2n - 5 = 9 \quad f(d) = (\textcolor{red}{2}, \textcolor{blue}{6}, \textcolor{green}{1})$$

$$f(\textcolor{red}{a}) = (\textcolor{red}{9}, \textcolor{blue}{0}, \textcolor{green}{0}) \quad f(e) = (\textcolor{red}{5}, \textcolor{blue}{2}, \textcolor{green}{2})$$

$$f(\textcolor{blue}{b}) = (\textcolor{red}{0}, \textcolor{blue}{9}, \textcolor{green}{0}) \quad f(g) = (\textcolor{red}{1}, \textcolor{blue}{2}, \textcolor{green}{6})$$

$$f(\textcolor{green}{c}) = (\textcolor{red}{0}, \textcolor{blue}{0}, \textcolor{green}{9}) \quad f(h) = (\textcolor{red}{4}, \textcolor{blue}{1}, \textcolor{green}{4})$$

Schnyder Drawing – Example





$$n = 7, 2n - 5 = 9 \quad f(d) = (2, 6, 1)$$

$$f(a) = (9, 0, 0) \quad f(e) = (5, 2, 2)$$

$$f(b) = (0, 9, 0) \quad f(g) = (1, 2, 6)$$



$$f(c) = (0, 0, 9) \quad f(h) = (4, 1, 4)$$

Schnyder Drawing – Example



$$\begin{aligned}
 n &= 7, 2n - 5 = 9 & f(d) &= (2, 6, 1) \\
 f(a) &= (9, 0, 0) & f(e) &= (5, 2, 2) \\
 f(b) &= (0, 9, 0) & f(g) &= (1, 2, 6) \\
 f(c) &= (0, 0, 9) & f(h) &= (4, 1, 4)
 \end{aligned}$$



Schnyder Drawing – Example



$$n = 7, 2n - 5 = 9$$



$f(a) = (9, 0, 0)$	$f(d) = (2, 6, 1)$
$f(b) = (0, 9, 0)$	$f(e) = (5, 2, 2)$
$f(c) = (0, 0, 9)$	$f(g) = (1, 2, 6)$
	$f(h) = (4, 1, 4)$

Schnyder Drawing – Example





$$\begin{aligned}
 n &= 7, 2n - 5 = 9 & f(d) &= (2, 6, 1) \\
 f(a) &= (9, 0, 0) & f(e) &= (5, 2, 2) \\
 f(b) &= (0, 9, 0) & f(g) &= (1, 2, 6) \\
 f(c) &= (0, 0, 9) & f(h) &= (4, 1, 4)
 \end{aligned}$$

Schnyder Drawing – Example



$$\begin{aligned}
 n &= 7, 2n - 5 = 9 & f(d) &= (2, 6, 1) \\
 f(a) &= (9, 0, 0) & f(e) &= (5, 2, 2) \\
 f(b) &= (0, 9, 0) & f(g) &= (1, 2, 6) \\
 f(c) &= (0, 0, 9) & f(h) &= (4, 1, 4)
 \end{aligned}$$

Schnyder Drawing – Example





$$n = 7, 2n - 5 = 9 \quad f(d) = (2, 6, 1)$$

$$f(a) = (9, 0, 0) \quad f(e) = (5, 2, 2)$$

$$f(b) = (0, 9, 0) \quad f(g) = (1, 2, 6)$$

$$f(c) = (0, 0, 9) \quad f(h) = (4, 1, 4)$$

Schnyder Drawing – Example





$$n = 7, 2n - 5 = 9 \quad f(d) = (2, 6, 1)$$

$$f(a) = (9, 0, 0) \quad f(e) = (5, 2, 2)$$

$$f(b) = (0, 9, 0) \quad f(g) = (1, 2, 6)$$



$$f(c) = (0, 0, 9) \quad f(h) = (4, 1, 4)$$

Schnyder Drawing – Example



$$\begin{array}{ll}
 n = 7, 2n - 5 = 9 & f(d) = (2, 6, 1) \\
 f(a) = (9, 0, 0) & f(e) = (5, 2, 2) \\
 f(b) = (0, 9, 0) & f(g) = (1, 2, 6) \\
 f(c) = (0, 0, 9) & f(h) = (4, 1, 4)
 \end{array}$$

Schnyder Drawing – Example



$$n = 7, 2n - 5 = 9$$

$$f(a) = (9, 0, 0)$$

$$f(b) = (0, 9, 0)$$

$$f(c) = (0, 0, 9)$$



$$f(d) = (2, 6, 1)$$

$$f(e) = (5, 2, 2)$$

$$f(g) = (1, 2, 6)$$

$$f(h) = (4, 1, 4)$$

Schnyder Drawing – Example



$$n = 7, 2n - 5 = 9$$

$$f(a) = (9, 0, 0)$$

$$f(b) = (0, 9, 0)$$

$$f(c) = (0, 0, 9)$$



$$f(d) = (2, 6, 1)$$

$$f(e) = (5, 2, 2)$$

$$f(g) = (1, 2, 6)$$

$$f(h) = (4, 1, 4)$$



Schnyder Drawing – Example



$$n = 7, 2n - 5 = 9$$

$f(a) = (9, 0, 0)$	$f(d) = (2, 6, 1)$
$f(b) = (0, 9, 0)$	$f(e) = (5, 2, 2)$
$f(c) = (0, 0, 9)$	$f(g) = (1, 2, 6)$
	$f(h) = (4, 1, 4)$

Schnyder Drawing – Example





$$\begin{array}{ll}
 n = 7, 2n - 5 = 9 & f(d) = (2, 6, 1) \\
 f(a) = (9, 0, 0) & f(e) = (5, 2, 2) \\
 f(b) = (0, 9, 0) & f(g) = (1, 2, 6) \\
 f(c) = (0, 0, 9) & f(h) = (4, 1, 4)
 \end{array}$$

Schnyder Drawing – Example

Theorem.

Every n -vertex planar graph has a planar straight-line drawing of size $(2n - 5) \times (2n - 5)$.

[Schnyder '89]



$$\begin{array}{ll}
 n = 7, 2n - 5 = 9 & f(d) = (2, 6, 1) \\
 f(a) = (9, 0, 0) & f(e) = (5, 2, 2) \\
 f(b) = (0, 9, 0) & f(g) = (1, 2, 6) \\
 f(c) = (0, 0, 9) & f(h) = (4, 1, 4)
 \end{array}$$

Weak Barycentric Representation

A **weak barycentric representation** of a graph $G = (V, E)$ is an assignment of barycentric coordinates to V :

$$\phi: V \rightarrow \mathbb{R}_{\geq 0}^3, v \mapsto (\textcolor{red}{v}_1, \textcolor{blue}{v}_2, \textcolor{green}{v}_3)$$

with the following properties:


Weak Barycentric Representation

A **weak barycentric representation** of a graph $G = (V, E)$ is an assignment of barycentric coordinates to V :

$$\phi: V \rightarrow \mathbb{R}_{\geq 0}^3, v \mapsto (\textcolor{red}{v}_1, \textcolor{blue}{v}_2, \textcolor{green}{v}_3)$$

with the following properties:

(W1) $\textcolor{red}{v}_1 + \textcolor{blue}{v}_2 + \textcolor{green}{v}_3 = 1$ for all $v \in V$,



Weak Barycentric Representation


A **weak barycentric representation** of a graph $G = (V, E)$ is an assignment of barycentric coordinates to V :

$$\phi: V \rightarrow \mathbb{R}_{\geq 0}^3, v \mapsto (\textcolor{red}{v}_1, \textcolor{blue}{v}_2, \textcolor{green}{v}_3)$$

with the following properties:

(W1) $\textcolor{red}{v}_1 + \textcolor{blue}{v}_2 + \textcolor{green}{v}_3 = 1$ for all $v \in V$,

(W2) for each $xy \in E$



Weak Barycentric Representation


A **weak barycentric representation** of a graph $G = (V, E)$ is an assignment of barycentric coordinates to V :

$$\phi: V \rightarrow \mathbb{R}_{\geq 0}^3, v \mapsto (\textcolor{red}{v}_1, \textcolor{blue}{v}_2, \textcolor{green}{v}_3)$$

with the following properties:

(W1) $\textcolor{red}{v}_1 + \textcolor{blue}{v}_2 + \textcolor{green}{v}_3 = 1$ for all $v \in V$,

(W2) for each $xy \in E$ and each $z \in V \setminus \{x, y\}$



Weak Barycentric Representation


A **weak barycentric representation** of a graph $G = (V, E)$ is an assignment of barycentric coordinates to V :

$$\phi: V \rightarrow \mathbb{R}_{\geq 0}^3, v \mapsto (\textcolor{red}{v}_1, \textcolor{blue}{v}_2, \textcolor{green}{v}_3)$$

with the following properties:

(W1) $\textcolor{red}{v}_1 + \textcolor{blue}{v}_2 + \textcolor{green}{v}_3 = 1$ for all $v \in V$,

(W2) for each $xy \in E$ and each $z \in V \setminus \{x, y\}$
there exists $k \in \{\textcolor{red}{1}, \textcolor{blue}{2}, \textcolor{green}{3}\}$ with
 $(x_k, x_{k+1}) <_{\text{lex}} (z_k, z_{k+1})$ and $(y_k, y_{k+1}) <_{\text{lex}} (z_k, z_{k+1})$.



Weak Barycentric Representation

A **weak barycentric representation** of a graph $G = (V, E)$ is an assignment of barycentric coordinates to V :


$$\phi: V \rightarrow \mathbb{R}_{\geq 0}^3, v \mapsto (\color{red}{v_1}, \color{blue}{v_2}, \color{green}{v_3})$$

with the following properties:

(W1) $\color{red}{v_1} + \color{blue}{v_2} + \color{green}{v_3} = 1$ for all $v \in V$,

(W2) for each $xy \in E$ and each $z \in V \setminus \{x, y\}$ there exists $k \in \{\color{red}{1}, \color{blue}{2}, \color{green}{3}\}$ with

$$(\color{red}{x_k}, \color{red}{x_{k+1}}) <_{\text{lex}} (\color{red}{z_k}, \color{red}{z_{k+1}}) \text{ and } (\color{blue}{y_k}, \color{blue}{y_{k+1}}) <_{\text{lex}} (\color{blue}{z_k}, \color{blue}{z_{k+1}}).$$



i.e., either $y_k < z_k$ or
 $y_k = z_k$ and $y_{k+1} < z_{k+1}$

Weak Barycentric Representation


A **weak barycentric representation** of a graph $G = (V, E)$ is an assignment of barycentric coordinates to V :

$$\phi: V \rightarrow \mathbb{R}_{\geq 0}^3, v \mapsto (\textcolor{red}{v}_1, \textcolor{blue}{v}_2, \textcolor{green}{v}_3)$$

with the following properties:

(W1) $\textcolor{red}{v}_1 + \textcolor{blue}{v}_2 + \textcolor{green}{v}_3 = 1$ for all $v \in V$,

(W2) for each $xy \in E$ and each $z \in V \setminus \{x, y\}$ there exists $k \in \{1, 2, 3\}$ with
 $(x_k, x_{k+1}) <_{\text{lex}} (z_k, z_{k+1})$ and $(y_k, y_{k+1}) <_{\text{lex}} (z_k, z_{k+1})$.



i.e., either $y_k < z_k$ or
 $y_k = z_k$ and $y_{k+1} < z_{k+1}$

Weak Barycentric Representation


A **weak barycentric representation** of a graph $G = (V, E)$ is an assignment of barycentric coordinates to V :

$$\phi: V \rightarrow \mathbb{R}_{\geq 0}^3, v \mapsto (\textcolor{red}{v}_1, \textcolor{blue}{v}_2, \textcolor{green}{v}_3)$$

with the following properties:

(W1) $\textcolor{red}{v}_1 + \textcolor{blue}{v}_2 + \textcolor{green}{v}_3 = 1$ for all $v \in V$,

(W2) for each $xy \in E$ and each $z \in V \setminus \{x, y\}$ there exists $k \in \{1, 2, 3\}$ with
 $(x_k, x_{k+1}) <_{\text{lex}} (z_k, z_{k+1})$ and $(y_k, y_{k+1}) <_{\text{lex}} (z_k, z_{k+1})$.



i.e., either $y_k < z_k$ or
 $y_k = z_k$ and $y_{k+1} < z_{k+1}$

Weak Barycentric Representation


A **weak barycentric representation** of a graph $G = (V, E)$ is an assignment of barycentric coordinates to V :

$$\phi: V \rightarrow \mathbb{R}_{\geq 0}^3, v \mapsto (\textcolor{red}{v}_1, \textcolor{blue}{v}_2, \textcolor{green}{v}_3)$$

with the following properties:

(W1) $\textcolor{red}{v}_1 + \textcolor{blue}{v}_2 + \textcolor{green}{v}_3 = 1$ for all $v \in V$,

(W2) for each $xy \in E$ and each $z \in V \setminus \{x, y\}$ there exists $k \in \{1, 2, 3\}$ with
 $(x_k, x_{k+1}) <_{\text{lex}} (z_k, z_{k+1})$ and $(y_k, y_{k+1}) <_{\text{lex}} (z_k, z_{k+1})$.



i.e., either $y_k < z_k$ or
 $y_k = z_k$ and $y_{k+1} < z_{k+1}$

Weak Barycentric Representation

A **weak barycentric representation** of a graph $G = (V, E)$ is an assignment of barycentric coordinates to V :


$$\phi: V \rightarrow \mathbb{R}_{\geq 0}^3, v \mapsto (\color{red}{v}_1, \color{blue}{v}_2, \color{green}{v}_3)$$

with the following properties:

(W1) $\color{red}{v}_1 + \color{blue}{v}_2 + \color{green}{v}_3 = 1$ for all $v \in V$,

(W2) for each $xy \in E$ and each $z \in V \setminus \{x, y\}$ there exists $k \in \{1, 2, 3\}$ with

$$(\color{red}{x}_k, \color{red}{x}_{k+1}) <_{\text{lex}} (\color{red}{z}_k, \color{red}{z}_{k+1}) \text{ and } (\color{blue}{y}_k, \color{blue}{y}_{k+1}) <_{\text{lex}} (\color{blue}{z}_k, \color{blue}{z}_{k+1}).$$



i.e., either $y_k < z_k$ or $y_k = z_k$ and $y_{k+1} < z_{k+1}$

Weak Barycentric Representation

A **weak barycentric representation** of a graph $G = (V, E)$ is an assignment of barycentric coordinates to V :

$$\phi: V \rightarrow \mathbb{R}_{\geq 0}^3, v \mapsto (\color{red}{v_1}, \color{blue}{v_2}, \color{green}{v_3})$$

with the following properties:

(W1) $\color{red}{v_1} + \color{blue}{v_2} + \color{green}{v_3} = 1$ for all $v \in V$,

(W2) for each $xy \in E$ and each $z \in V \setminus \{x, y\}$ there exists $k \in \{1, 2, 3\}$ with


$$(\color{red}{x_k}, \color{red}{x_{k+1}}) <_{\text{lex}} (\color{red}{z_k}, \color{red}{z_{k+1}}) \text{ and } (\color{blue}{y_k}, \color{blue}{y_{k+1}}) <_{\text{lex}} (\color{blue}{z_k}, \color{blue}{z_{k+1}}).$$

Lemma.

For a weak barycentric representation $\phi: v \mapsto (\color{red}{v_1}, \color{blue}{v_2}, \color{green}{v_3})$ and a triangle $\color{red}{A}, \color{blue}{B}, \color{green}{C}$, the mapping


$$f: v \in V \mapsto \color{red}{v_1} \color{red}{A} + \color{blue}{v_2} \color{blue}{B} + \color{green}{v_3} \color{green}{C}$$

gives a **planar** drawing of G inside $\triangle \color{red}{ABC}$.




i.e., either $y_k < z_k$ or $y_k = z_k$ and $y_{k+1} < z_{k+1}$

Counting Vertices



Counting Vertices




$P_i(v)$: path from v to root of T_i .

$R_1(v)$: set of faces contained in P_2, bc, P_3 .

$R_2(v)$: set of faces contained in P_3, ca, P_1 .

$R_3(v)$: set of faces contained in P_1, ab, P_2 .

Counting Vertices



$P_i(v)$: path from v to root of T_i .


$R_1(v)$: set of faces contained in P_2, bc, P_3 .

$R_2(v)$: set of faces contained in P_3, ca, P_1 .

$R_3(v)$: set of faces contained in P_1, ab, P_2 .

$$v_i = |V(R_i(v))|$$

Counting Vertices



$P_i(v)$: path from v to root of T_i .


$R_1(v)$: set of faces contained in P_2, bc, P_3 .

$R_2(v)$: set of faces contained in P_3, ca, P_1 .

$R_3(v)$: set of faces contained in P_1, ab, P_2 .

$$v_i = |V(R_i(v))| - |P_{i-1}(v)|$$

Counting Vertices



$P_i(v)$: path from v to root of T_i .

$R_1(v)$: set of faces contained in P_2, bc, P_3 .


$R_2(v)$: set of faces contained in P_3, ca, P_1 .

$R_3(v)$: set of faces contained in P_1, ab, P_2 .

$$v_i = |V(R_i(v))| - |P_{i-1}(v)|$$

$$v_1 =$$

Counting Vertices



$P_i(v)$: path from v to root of T_i .

$R_1(v)$: set of faces contained in P_2, bc, P_3 .


$R_2(v)$: set of faces contained in P_3, ca, P_1 .

$R_3(v)$: set of faces contained in P_1, ab, P_2 .

$$v_i = |V(R_i(v))| - |P_{i-1}(v)|$$

$$v_1 =$$

Counting Vertices



$P_i(v)$: path from v to root of T_i .

$R_1(v)$: set of faces contained in P_2, bc, P_3 .


$R_2(v)$: set of faces contained in P_3, ca, P_1 .

$R_3(v)$: set of faces contained in P_1, ab, P_2 .

$$v_i = |V(R_i(v))| - |P_{i-1}(v)|$$

$$v_1 = 10 - 3 = 7$$

Counting Vertices



$P_i(v)$: path from v to root of T_i .

$R_1(v)$: set of faces contained in P_2, bc, P_3 .

$R_2(v)$: set of faces contained in P_3, ca, P_1 .


$R_3(v)$: set of faces contained in P_1, ab, P_2 .

$$v_i = |V(R_i(v))| - |P_{i-1}(v)|$$

$$v_1 = 10 - 3 = 7$$

$$v_2 =$$

Counting Vertices



$P_i(v)$: path from v to root of T_i .

$R_1(v)$: set of faces contained in P_2, bc, P_3 .

$R_2(v)$: set of faces contained in P_3, ca, P_1 .


$R_3(v)$: set of faces contained in P_1, ab, P_2 .

$$v_i = |V(R_i(v))| - |P_{i-1}(v)|$$

$$v_1 = 10 - 3 = 7$$

$$v_2 = 6 - 3 = 3$$

Counting Vertices



$P_i(v)$: path from v to root of T_i .

$R_1(v)$: set of faces contained in P_2, bc, P_3 .

$R_2(v)$: set of faces contained in P_3, ca, P_1 .

$R_3(v)$: set of faces contained in P_1, ab, P_2 .


$$v_i = |V(R_i(v))| - |P_{i-1}(v)|$$

$$v_1 = 10 - 3 = 7$$

$$v_2 = 6 - 3 = 3$$

$$v_3 =$$

Counting Vertices



$P_i(v)$: path from v to root of T_i .

$R_1(v)$: set of faces contained in P_2, bc, P_3 .

$R_2(v)$: set of faces contained in P_3, ca, P_1 .

$R_3(v)$: set of faces contained in P_1, ab, P_2 .


$$v_i = |V(R_i(v))| - |P_{i-1}(v)|$$

$$v_1 = 10 - 3 = 7$$

$$v_2 = 6 - 3 = 3$$

$$v_3 = 8 - 3 = 5$$

Counting Vertices



$P_i(v)$: path from v to root of T_i .

$R_1(v)$: set of faces contained in P_2, bc, P_3 .

$R_2(v)$: set of faces contained in P_3, ca, P_1 .

$R_3(v)$: set of faces contained in P_1, ab, P_2 .

$$v_i = |V(R_i(v))| - |P_{i-1}(v)|$$

$$v_1 = 10 - 3 = 7$$


$$v_2 = 6 - 3 = 3$$

$$v_3 = 8 - 3 = 5$$

Lemma.

- For inner vertices $u \neq v$ it holds that $u \in R_i(v) \Rightarrow (u_i, u_{i+1}) <_{\text{lex}} (v_i, v_{i+1})$.

Counting Vertices



$P_i(v)$: path from v to root of T_i .

$R_1(v)$: set of faces contained in P_2, bc, P_3 .

$R_2(v)$: set of faces contained in P_3, ca, P_1 .

$R_3(v)$: set of faces contained in P_1, ab, P_2 .

$$v_i = |V(R_i(v))| - |P_{i-1}(v)|$$

$$v_1 = 10 - 3 = 7$$


$$v_2 = 6 - 3 = 3$$

$$v_3 = 8 - 3 = 5$$

Lemma.

- For inner vertices $u \neq v$ it holds that $u \in R_i(v) \Rightarrow (u_i, u_{i+1}) <_{\text{lex}} (v_i, v_{i+1})$.

Counting Vertices



$P_i(v)$: path from v to root of T_i .

$R_1(v)$: set of faces contained in P_2, bc, P_3 .

$R_2(v)$: set of faces contained in P_3, ca, P_1 .

$R_3(v)$: set of faces contained in P_1, ab, P_2 .

$$v_i = |V(R_i(v))| - |P_{i-1}(v)|$$

$$v_1 = 10 - 3 = 7$$


$$v_2 = 6 - 3 = 3$$

$$v_3 = 8 - 3 = 5$$

Lemma.

- For inner vertices $u \neq v$ it holds that $u \in R_i(v) \Rightarrow (u_i, u_{i+1}) <_{\text{lex}} (v_i, v_{i+1})$.

Counting Vertices



$P_i(v)$: path from v to root of T_i .

$R_1(v)$: set of faces contained in P_2, bc, P_3 .

$R_2(v)$: set of faces contained in P_3, ca, P_1 .

$R_3(v)$: set of faces contained in P_1, ab, P_2 .

$$v_i = |V(R_i(v))| - |P_{i-1}(v)|$$

$$v_1 = 10 - 3 = 7$$


$$v_2 = 6 - 3 = 3$$

$$v_3 = 8 - 3 = 5$$

Lemma.

- For inner vertices $u \neq v$ it holds that $u \in R_i(v) \Rightarrow (u_i, u_{i+1}) <_{\text{lex}} (v_i, v_{i+1})$.

Counting Vertices



$P_i(v)$: path from v to root of T_i .

$R_1(v)$: set of faces contained in P_2, bc, P_3 .

$R_2(v)$: set of faces contained in P_3, ca, P_1 .

$R_3(v)$: set of faces contained in P_1, ab, P_2 .

$$v_i = |V(R_i(v))| - |P_{i-1}(v)|$$

$$v_1 = 10 - 3 = 7$$

$$v_2 = 6 - 3 = 3$$


$$v_3 = 8 - 3 = 5$$

Lemma.

■ For inner vertices $u \neq v$ it holds that $u \in R_i(v) \Rightarrow (u_i, u_{i+1}) <_{\text{lex}} (v_i, v_{i+1})$.

■ $v_1 + v_2 + v_3 =$

Counting Vertices



$P_i(v)$: path from v to root of T_i .

$R_1(v)$: set of faces contained in P_2, bc, P_3 .

$R_2(v)$: set of faces contained in P_3, ca, P_1 .

$R_3(v)$: set of faces contained in P_1, ab, P_2 .

$$v_i = |V(R_i(v))| - |P_{i-1}(v)|$$

$$v_1 = 10 - 3 = 7$$

$$v_2 = 6 - 3 = 3$$


$$v_3 = 8 - 3 = 5$$

Lemma.

■ For inner vertices $u \neq v$ it holds that $u \in R_i(v) \Rightarrow (u_i, u_{i+1}) <_{\text{lex}} (v_i, v_{i+1})$.

■ $v_1 + v_2 + v_3 = n - 1$

Counting Vertices



$P_i(v)$: path from v to root of T_i .

$R_1(v)$: set of faces contained in P_2, bc, P_3 .

$R_2(v)$: set of faces contained in P_3, ca, P_1 .

$R_3(v)$: set of faces contained in P_1, ab, P_2 .

$$v_i = |V(R_i(v))| - |P_{i-1}(v)|$$

$$v_1 = 10 - 3 = 7$$

$$v_2 = 6 - 3 = 3$$


$$v_3 = 8 - 3 = 5$$

Lemma.

■ For inner vertices $u \neq v$ it holds that
 $u \in R_i(v) \Rightarrow (u_i, u_{i+1}) <_{\text{lex}} (v_i, v_{i+1})$.

■ $v_1 + v_2 + v_3 = n - 1$

Counting Vertices



$P_i(v)$: path from v to root of T_i .

$R_1(v)$: set of faces contained in P_2, bc, P_3 .

$R_2(v)$: set of faces contained in P_3, ca, P_1 .

$R_3(v)$: set of faces contained in P_1, ab, P_2 .

$$v_i = |V(R_i(v))| - |P_{i-1}(v)|$$

$$v_1 = 10 - 3 = 7$$

$$v_2 = 6 - 3 = 3$$

$$v_3 = 8 - 3 = 5$$

Lemma.

■ For inner vertices $u \neq v$ it holds that $u \in R_i(v) \Rightarrow (u_i, u_{i+1}) <_{\text{lex}} (v_i, v_{i+1})$.

■ $v_1 + v_2 + v_3 = n - 1$

Schnyder Drawing^{*}

Set $A = (0, 0)$, $B = (n - 1, 0)$, and $C = (0, n - 1)$.

Theorem.


[Schnyder '90]

For a plane triangulation G , the mapping


$$f: v \mapsto \frac{1}{n-1} (v_1, v_2, v_3)$$

is a barycentric representation of G , which thus gives a planar straight-line drawing of G on the $(n - 2) \times (n - 2)$ grid.

Schnyder Drawing^{*} – Example





Schnyder Drawing^{*} – Example





$$n = 16, n - 2 = 14$$

Schnyder Drawing^{*} – Example





$$n = 16, n - 2 = 14$$

Schnyder Drawing^{*} – Example





$$n = 16, n - 2 = 14$$

Schnyder Drawing^{*} – Example



$$n = 16, n - 2 = 14$$



Schnyder Drawing^{*} – Example



$$n = 16, n - 2 = 14$$

$$f(a) = (\textcolor{red}{n} - \textcolor{red}{2}, \textcolor{blue}{1}, \textcolor{green}{0})$$



Schnyder Drawing^{*} – Example



$$n = 16, n - 2 = 14$$

$$f(a) = (\textcolor{red}{n} - \textcolor{red}{2}, \textcolor{blue}{1}, \textcolor{green}{0})$$

Schnyder Drawing^{*} – Example





$$n = 16, n - 2 = 14$$

$$f(a) = (n - 2, 1, 0)$$

$$f(b) = (0, n - 2, 1)$$

Schnyder Drawing^{*} – Example





$$n = 16, n - 2 = 14$$

$$f(a) = (\textcolor{red}{n} - \textcolor{red}{2}, \textcolor{blue}{1}, \textcolor{green}{0})$$

$$f(b) = (\textcolor{red}{0}, \textcolor{blue}{n} - \textcolor{blue}{2}, \textcolor{green}{1})$$

$$f(c) = (\textcolor{red}{1}, \textcolor{blue}{0}, \textcolor{green}{n} - \textcolor{green}{2})$$

Schnyder Drawing^{*} – Example





$$n = 16, n - 2 = 14$$

$$f(a) = (\textcolor{red}{n} - \textcolor{red}{2}, \textcolor{blue}{1}, \textcolor{green}{0})$$

$$f(b) = (\textcolor{red}{0}, \textcolor{blue}{n} - \textcolor{blue}{2}, \textcolor{green}{1})$$

$$f(c) = (\textcolor{red}{1}, \textcolor{blue}{0}, \textcolor{green}{n} - \textcolor{green}{2})$$

Schnyder Drawing^{*} – Example





$$n = 16, n - 2 = 14$$

$$f(a) = (\textcolor{red}{n} - \textcolor{red}{2}, \textcolor{blue}{1}, \textcolor{green}{0})$$

$$f(b) = (\textcolor{red}{0}, \textcolor{blue}{n} - \textcolor{blue}{2}, \textcolor{green}{1})$$

$$f(c) = (\textcolor{red}{1}, \textcolor{blue}{0}, \textcolor{green}{n} - \textcolor{green}{2})$$

Schnyder Drawing^{*} – Example




$$n = 16, n - 2 = 14$$

$$f(a) = (\textcolor{red}{n} - \textcolor{red}{2}, \textcolor{blue}{1}, \textcolor{green}{0})$$



$$f(b) = (\textcolor{red}{0}, \textcolor{blue}{n} - \textcolor{blue}{2}, \textcolor{green}{1})$$

$$f(c) = (\textcolor{red}{1}, \textcolor{blue}{0}, \textcolor{green}{n} - \textcolor{green}{2})$$



$$f(c) = (\textcolor{red}{1}, \textcolor{blue}{0}, \textcolor{green}{n} - \textcolor{green}{2})$$

Schnyder Drawing^{*} – Example




$$n = 16, n - 2 = 14$$


$$f(a) = (\textcolor{red}{n} - \textcolor{red}{2}, \textcolor{blue}{1}, \textcolor{green}{0})$$

$$f(b) = (\textcolor{red}{0}, \textcolor{blue}{n} - \textcolor{blue}{2}, \textcolor{green}{1})$$

$$f(c) = (\textcolor{red}{1}, \textcolor{blue}{0}, \textcolor{green}{n} - \textcolor{green}{2})$$





$$f(c) = (\textcolor{red}{1}, \textcolor{blue}{0}, \textcolor{green}{n} - \textcolor{green}{2})$$



$$f(c) = (\textcolor{red}{1}, \textcolor{blue}{0}, \textcolor{green}{n} - \textcolor{green}{2})$$

Schnyder Drawing^{*} – Example





$$n = 16, n - 2 = 14$$

$$f(a) = (\textcolor{red}{n} - \textcolor{red}{2}, \textcolor{blue}{1}, \textcolor{green}{0})$$

$$f(b) = (\textcolor{red}{0}, \textcolor{blue}{n} - \textcolor{blue}{2}, \textcolor{green}{1})$$

$$f(c) = (\textcolor{red}{1}, \textcolor{blue}{0}, \textcolor{green}{n} - \textcolor{green}{2})$$

Schnyder Drawing^{*} – Example





$$n = 16, n - 2 = 14$$

$$f(a) = (\textcolor{red}{n} - \textcolor{red}{2}, \textcolor{blue}{1}, \textcolor{green}{0})$$

$$f(b) = (\textcolor{red}{0}, \textcolor{blue}{n} - \textcolor{blue}{2}, \textcolor{green}{1})$$

$$f(c) = (\textcolor{red}{1}, \textcolor{blue}{0}, \textcolor{green}{n} - \textcolor{green}{2})$$

Schnyder Drawing^{*} – Example





$$n = 16, n - 2 = 14$$

$$f(a) = (\textcolor{red}{n} - \textcolor{red}{2}, \textcolor{blue}{1}, \textcolor{green}{0})$$

$$f(b) = (\textcolor{red}{0}, \textcolor{blue}{n} - \textcolor{blue}{2}, \textcolor{green}{1})$$

$$f(c) = (\textcolor{red}{1}, \textcolor{blue}{0}, \textcolor{green}{n} - \textcolor{green}{2})$$

Schnyder Drawing^{*} – Example




$$n = 16, n - 2 = 14$$


$$f(a) = (\textcolor{red}{n} - \textcolor{red}{2}, \textcolor{blue}{1}, \textcolor{green}{0})$$

$$f(b) = (\textcolor{red}{0}, \textcolor{blue}{n} - \textcolor{blue}{2}, \textcolor{green}{1})$$

$$f(c) = (\textcolor{red}{1}, \textcolor{blue}{0}, \textcolor{green}{n} - \textcolor{green}{2})$$





$$f(c) = (\textcolor{red}{1}, \textcolor{blue}{0}, \textcolor{green}{n} - \textcolor{green}{2})$$



$$f(c) = (\textcolor{red}{1}, \textcolor{blue}{0}, \textcolor{green}{n} - \textcolor{green}{2})$$

Schnyder Drawing^{*} – Example




$$n = 16, n - 2 = 14$$

$$f(a) = (\textcolor{red}{n} - \textcolor{red}{2}, \textcolor{blue}{1}, \textcolor{green}{0})$$



$$f(b) = (\textcolor{red}{0}, \textcolor{blue}{n} - \textcolor{blue}{2}, \textcolor{green}{1})$$

$$f(c) = (\textcolor{red}{1}, \textcolor{blue}{0}, \textcolor{green}{n} - \textcolor{green}{2})$$



$$f(c) = (\textcolor{red}{1}, \textcolor{blue}{0}, \textcolor{green}{n} - \textcolor{green}{2})$$

Schnyder Drawing^{*} – Example





$$n = 16, n - 2 = 14$$

$$f(a) = (\textcolor{red}{n} - \textcolor{red}{2}, \textcolor{blue}{1}, \textcolor{green}{0})$$

$$f(b) = (\textcolor{red}{0}, \textcolor{blue}{n} - \textcolor{blue}{2}, \textcolor{green}{1})$$

$$f(c) = (\textcolor{red}{1}, \textcolor{blue}{0}, \textcolor{green}{n} - \textcolor{green}{2})$$

Schnyder Drawing^{*} – Example





$$n = 16, n - 2 = 14$$

$$f(a) = (\textcolor{red}{n} - \textcolor{red}{2}, \textcolor{blue}{1}, \textcolor{green}{0})$$

$$f(b) = (\textcolor{red}{0}, \textcolor{blue}{n} - \textcolor{blue}{2}, \textcolor{green}{1})$$

$$f(c) = (\textcolor{red}{1}, \textcolor{blue}{0}, \textcolor{green}{n} - \textcolor{green}{2})$$

Schnyder Drawing^{*} – Example



$$n = 16, n - 2 = 14$$

$$f(a) = (\textcolor{red}{n} - \textcolor{red}{2}, \textcolor{blue}{1}, \textcolor{green}{0})$$

$$f(b) = (\textcolor{red}{0}, \textcolor{blue}{n} - \textcolor{blue}{2}, \textcolor{green}{1})$$

$$f(c) = (\textcolor{red}{1}, \textcolor{blue}{0}, \textcolor{green}{n} - \textcolor{green}{2})$$

Results & Variations

Theorem.



[De Fraysseix, Pach, Pollack '90]

Every n -vertex planar graph has a planar straight-line drawing of size $(2n - 4) \times (n - 2)$. Such a drawing can be computed in $O(n)$ time.

Theorem.

[Schnyder '90]

Every n -vertex planar graph has a planar straight-line drawing of size $(n - 2) \times (n - 2)$. Such a drawing can be computed in $O(n)$ time.



Results & Variations

Theorem.



[De Fraysseix, Pach, Pollack '90]

Every n -vertex planar graph has a planar straight-line drawing of size $(2n - 4) \times (n - 2)$. Such a drawing can be computed in $O(n)$ time.

Theorem.

[Schneider '90]

Every n -vertex planar graph has a planar straight-line drawing of size $(n - 2) \times (n - 2)$. Such a drawing can be computed in $O(n)$ time.



Results & Variations

Theorem.

[De Fraysseix, Pach, Pollack '90]

Every n -vertex planar graph has a planar straight-line drawing of size $(2n - 4) \times (n - 2)$. Such a drawing can be computed in $O(n)$ time.

Theorem.

[Schnyder '90]

Every n -vertex planar graph has a planar straight-line drawing of size $(n - 2) \times (n - 2)$. Such a drawing can be computed in $O(n)$ time.

Theorem.

[Chrobak & Kant '97]

Every n -vertex 3-connected planar graph has a planar straight-line drawing of size $(n - 2) \times (n - 2)$ where all faces are drawn convex. Such a drawing can be computed in $O(n)$ time.

Results & Variations

Theorem.

[De Fraysseix, Pach, Pollack '90]

Every n -vertex planar graph has a planar straight-line drawing of size $(2n - 4) \times (n - 2)$. Such a drawing can be computed in $O(n)$ time.

Theorem.

[Schnyder '90]

Every n -vertex planar graph has a planar straight-line drawing of size $(n - 2) \times (n - 2)$. Such a drawing can be computed in $O(n)$ time.

Theorem.

[Chrobak & Kant '97]


Every n -vertex 3-connected planar graph has a planar straight-line drawing of size $(n - 2) \times (n - 2)$ where all faces are drawn convex. Such a drawing can be computed in $O(n)$ time.

Theorem.


[Felsner '01]

Every 3-connected planar graph with f faces has a planar straight-line drawing of size $(f - 1) \times (f - 1)$ where all faces are drawn convex. Such a drawing can be computed in $O(n)$ time.


Schnyder Realizer \rightarrow Canonical Order




Schnyder Realizer \rightarrow Canonical Order




Schnyder Realizer \rightarrow Canonical Order




Schnyder Realizer \rightarrow Canonical Order




Schnyder Realizer \rightarrow Canonical Order




Schnyder Realizer \rightarrow Canonical Order




Schnyder Realizer \rightarrow Canonical Order




Schnyder Realizer \rightarrow Canonical Order




Schnyder Realizer \rightarrow Canonical Order




Schnyder Realizer \rightarrow Canonical Order




Schnyder Realizer \rightarrow Canonical Order




Schnyder Realizer \rightarrow Canonical Order




Schnyder Realizer \rightarrow Canonical Order




Schnyder Realizer \rightarrow Canonical Order




Schnyder Realizer \rightarrow Canonical Order




Schnyder Realizer \rightarrow Canonical Order




Schnyder Realizer \rightarrow Canonical Order




Schnyder Realizer \rightarrow Canonical Order



Schnyder Realizer \rightarrow Canonical Order




Schnyder Realizer \rightarrow Canonical Order



Theorem.

A ccw pre-order traversal on T_i induces a canonical order.


Schnyder Realizer \rightarrow Canonical Order



Theorem.

A ccw pre-order traversal on T_i induces a canonical order.


Schnyder Realizer \rightarrow Canonical Order



Theorem.

A ccw pre-order traversal on T_i induces a canonical order.


Schnyder Realizer \rightarrow Canonical Order




Theorem.

A ccw pre-order traversal on T_i induces a canonical order.


Canonical Order \rightarrow Schnyder Realizer




Canonical Order \rightarrow Schnyder Realizer




Canonical Order \rightarrow Schnyder Realizer




Canonical Order \rightarrow Schnyder Realizer




Canonical Order \rightarrow Schnyder Realizer




Canonical Order \rightarrow Schnyder Realizer




Canonical Order \rightarrow Schnyder Realizer




Canonical Order \rightarrow Schnyder Realizer




Canonical Order \rightarrow Schnyder Realizer




Canonical Order \rightarrow Schnyder Realizer




Canonical Order \rightarrow Schnyder Realizer




Canonical Order \rightarrow Schnyder Realizer




Canonical Order \rightarrow Schnyder Realizer




Canonical Order \rightarrow Schnyder Realizer




Canonical Order \rightarrow Schnyder Realizer




Canonical Order \rightarrow Schnyder Realizer




Canonical Order \rightarrow Schnyder Realizer




Canonical Order \rightarrow Schnyder Realizer



Canonical Order \rightarrow Schnyder Realizer




Canonical Order \rightarrow Schnyder Realizer




Canonical Order \rightarrow Schnyder Realizer

Theorem.

A canonical order induces a unique Schnyder Realizer.




Linear Time Computation




Linear Time Computation

- Compute Canonical Order




Linear Time Computation

- Compute Canonical Order
- Compute Schnyder Realizer




Linear Time Computation

- Compute Canonical Order
- Compute Schnyder Realizer




Linear Time Computation

- Compute Canonical Order
- Compute Schnyder Realizer
- Goal: $v_i = |V(R_i(v))| - |P_{i-1}(v)|$




Linear Time Computation

- Compute Canonical Order
- Compute Schnyder Realizer
- Goal: $v_i = |V(R_i(v))| - |P_{i-1}(v)|$
- With traversal of each T_i , compute:




Linear Time Computation

- Compute Canonical Order
- Compute Schnyder Realizer
- Goal: $v_i = |V(R_i(v))| - |P_{i-1}(v)|$
- With traversal of each T_i , compute:
 - The number of vertices in $P_i(v)$




Linear Time Computation

- Compute Canonical Order
- Compute Schnyder Realizer
- Goal: $v_i = |V(R_i(v))| - |P_{i-1}(v)|$
- With traversal of each T_i , compute:
 - The number of vertices in $P_i(v)$
 - The number of vertices in the subtree $T_i(v)$ of T_i rooted at v




Linear Time Computation

- Compute Canonical Order
- Compute Schnyder Realizer
- Goal: $v_i = |V(R_i(v))| - |P_{i-1}(v)|$
- With traversal of each T_i , compute:
 - The number of vertices in $P_i(v)$
 - The number of vertices in the subtree $T_i(v)$ of T_i rooted at v




Linear Time Computation

- Compute Canonical Order
- Compute Schnyder Realizer
- Goal: $v_i = |V(R_i(v))| - |P_{i-1}(v)|$
- With traversal of each T_i , compute:
 - The number of vertices in $P_i(v)$
 - The number of vertices in the subtree $T_i(v)$ of T_i rooted at v
- $|V(R_i(v))| =$




Linear Time Computation

- Compute Canonical Order
- Compute Schnyder Realizer
- Goal: $v_i = |V(R_i(v))| - |P_{i-1}(v)|$
- With traversal of each T_i , compute:
 - The number of vertices in $P_i(v)$
 - The number of vertices in the subtree $T_i(v)$ of T_i rooted at v
- $|V(R_i(v))| = \sum_{u \in P_{i+1}(v)} |T_i(u)|$




Linear Time Computation

- Compute Canonical Order
- Compute Schnyder Realizer
- Goal: $v_i = |V(R_i(v))| - |P_{i-1}(v)|$
- With traversal of each T_i , compute:
 - The number of vertices in $P_i(v)$
 - The number of vertices in the subtree $T_i(v)$ of T_i rooted at v
- $|V(R_i(v))| = \sum_{u \in P_{i+1}(v)} |T_i(u)| + \sum_{u \in P_{i-1}(v)} |T_i(u)|$




Linear Time Computation

- Compute Canonical Order
- Compute Schnyder Realizer
- Goal: $v_i = |V(R_i(v))| - |P_{i-1}(v)|$
- With traversal of each T_i , compute:
 - The number of vertices in $P_i(v)$
 - The number of vertices in the subtree $T_i(v)$ of T_i rooted at v
- $|V(R_i(v))| = \sum_{u \in P_{i+1}(v)} |T_i(u)| + \sum_{u \in P_{i-1}(v)} |T_i(u)| - T_i(v)$




Linear Time Computation

- Compute Canonical Order
- Compute Schnyder Realizer
- Goal: $v_i = |V(R_i(v))| - |P_{i-1}(v)|$
- With traversal of each T_i , compute:
 - The number of vertices in $P_i(v)$
 - The number of vertices in the subtree $T_i(v)$ of T_i rooted at v
- $|V(R_i(v))| = \sum_{u \in P_{i+1}(v)} |T_i(u)| + \sum_{u \in P_{i-1}(v)} |T_i(u)| - T_i(v)$
- Compute these sums in six tree traversals



Linear Time Computation

- Compute Canonical Order
- Compute Schnyder Realizer
- Goal: $v_i = |V(R_i(v))| - |P_{i-1}(v)|$
- With traversal of each T_i , compute:
 - The number of vertices in $P_i(v)$
 - The number of vertices in the subtree $T_i(v)$ of T_i rooted at v
- $|V(R_i(v))| = \sum_{u \in P_{i+1}(v)} |T_i(u)| + \sum_{u \in P_{i-1}(v)} |T_i(u)| - T_i(v)$
- Compute these sums in six tree traversals



Theorem.

[Schnyder '90]

Every n -vertex planar graph has a planar straight-line drawing of size $(n - 2) \times (n - 2)$. Such a drawing can be computed in $O(n)$ time.