

Pattern-avoidance in Binary Fillings of Grid Shapes

by

Alexey Spiridonov

A.B., Princeton University, 2004

Submitted to the Department of Mathematics
in partial fulfillment of the requirements for the degree of

Doctor of Philosophy

at the

MASSACHUSETTS INSTITUTE OF TECHNOLOGY

June 2009

© Alexey Spiridonov, 2009. All rights reserved.

The author hereby grants to MIT permission to reproduce and to
distribute publicly paper and electronic copies of this thesis document
in whole or in part in any medium now known or hereafter created.

Author
Department of Mathematics
May 8, 2009

Certified by
Alexander Postnikov
Associate Professor of Applied Mathematics
Thesis Supervisor

Accepted by
Michel X. Goemans
Chairman, Applied Mathematics Committee

Accepted by
David S. Jerison
Chairman, Department Committee on Graduate Students

Pattern-avoidance in Binary Fillings of Grid Shapes

by

Alexey Spiridonov

Submitted to the Department of Mathematics
on May 8, 2009 in partial fulfillment of the
requirements for the degree of
Doctor of Philosophy

Abstract

A *grid shape* is a set of boxes chosen from a square grid; any Young diagram is an example. We consider a notion of pattern-avoidance for 0-1 fillings of grid shapes, which generalizes permutation pattern-avoidance. A filling avoids a set of patterns if none of its sub-shapes, obtained by removing some rows and columns, equal any of the patterns. We focus on patterns that are *pairs* of 2×2 fillings.

Totally nonnegative Grassmann cells are in bijection with Young shape fillings that avoid particular 2×2 pair, which are, in turn, equinumerous with fillings avoiding another 2×2 pair. The latter ones correspond to acyclic orientations of the shape's bipartite graph. Motivated by this result, due to Postnikov and Williams, we prove a number of such analogs of Wilf-equivalence for these objects — that is, we show that, *in certain classes of shapes, some pattern-avoiding fillings are equinumerous with others*.

The equivalences in this paper follow from two very different bijections, and from a family of recurrences generalizing results of Postnikov and Williams. We used a computer to test each of the described equivalences on a diverse set of shapes. All our results are *nearly* tight, in the sense that we found no natural families of shapes, in which the equivalences hold, but the results' hypotheses do not.

One of these bijections gives rise to some new combinatorics on tilings of skew Young shapes with rectangles, which we name *Popeye diagrams*. In a special case, they are exactly Hugh Thomas's *snug partitions* for $d = 2$. We show that Popeye diagrams are a lattice, and, moreover, each diagram is a sublattice of the Tamari lattice. We also give a simple enumerative result.

Thesis Supervisor: Alexander Postnikov
Title: Associate Professor of Applied Mathematics