

Upper bounds for average Bayes accuracy in terms of mutual information

Charles Zheng and Yuval Benjamini

September 23, 2016

These are preliminary notes.

1 Introduction

Suppose X and Y are continuous random variables (or vectors) which have a joint distribution with density $p(x, y)$. Let $p(x) = \int p(x, y)dy$ and $p(y) = \int p(x, y)dx$ denote the respective marginal distributions, and $p(y|x) = p(x, y)/p(x)$ denote the conditional distribution.

Mutual information is defined

$$I[p(x, y)] = \int p(x, y) \log \frac{p(x, y)}{p(x)p(y)} dx dy.$$

ABE_k , or k -class Average Bayes accuracy is defined as follows. Let X_1, \dots, X_K be iid from $p(x)$, and draw Z uniformly from $1, \dots, k$. Draw $Y \sim p(y|X_Z)$. Then, the average Bayes accuracy is defined as

$$ABA_k[p(x, y)] = \sup_f \Pr[f(X_1, \dots, X_k, Y) = Z]$$

where the supremum is taken over all functions f . A function f which achieves the supremum is

$$f_{Bayes}(x_1, \dots, x_k, y) = \operatorname{argmax}_{z \in \{1, \dots, k\}} p(y|x_z),$$

where an arbitrary rule can be employed to break ties. Such a function f_{Bayes} is called a *Bayes classification rule*. It follows that ABA_k is given explicitly

by

$$\text{ABA}_k = \frac{1}{k} \int \left[\prod_{i=1}^k p(x_i) dx_i \right] \int dy \max_i p(y|x_i),$$

as stated in the following theorem.

Theorem 1.1 *For a joint distribution $p(x, y)$, define*

$$\text{ABA}_k[p(x, y)] = \sup_f \Pr[f(x_1, \dots, x_k, y) = Z]$$

where X_1, \dots, X_K are iid from $p(x)$, Z is uniform from $1, \dots, k$, and $Y \sim p(y|X_Z)$, and the supremum is taken over all functions $f : \mathcal{X}^k \times \mathcal{Y} \rightarrow \{1, \dots, k\}$. Then,

$$\text{ABA}_k = \frac{1}{k} \int \left[\prod_{i=1}^k p(x_i) dx_i \right] \int dy \max_i p(y|x_i).$$

Proof. First, we claim that the supremum is attained by choosing

$$f(x_1, \dots, x_k, y) = \operatorname{argmax}_{z \in \{1, \dots, k\}} p(y|x_z).$$

To show this claim, write

$$\sup_f \Pr[f(X_1, \dots, X_k, Y) = Z] = \sup_f \frac{1}{k} \int p_X(x_1) \dots p_X(x_k) p(y|x_{f(x_1, \dots, x_k, y)}) dx_1 \dots dx_k dy$$

We see that maximizing $\Pr[f(X_1, \dots, X_k, Y) = Z]$ over functions f additively decomposes into infinitely many subproblems, where in each subproblem we are given $\{x_1, \dots, x_k, y\} \in \mathcal{X}^k \times \mathcal{Y}$, and our goal is to choose $f(x_1, \dots, x_k, y)$ from the set $\{1, \dots, k\}$ in order to maximize the quantity $p(y|x_{f(x_1, \dots, x_k, y)})$. In each subproblem, the maximum is attained by setting $f(x_1, \dots, x_k, y) = \operatorname{argmax}_z p(y|x_z)$ —and the resulting function f attains the supremum to the functional optimization problem. This proves the claim.

We therefore have

$$p(y|x_{f(x_1, \dots, x_k, y)}) = \max_{i=1}^k p(y|x_i).$$

Therefore, we can write

$$\begin{aligned}
\text{ABA}_k[p(x, y)] &= \sup_f \Pr[f(X_1, \dots, X_k, Y) = Z] \\
&= \frac{1}{k} \int p_X(x_1) \dots p_X(x_k) p(y | x_{f(x_1, \dots, x_k, y)}) dx_1 \dots dx_k dy. \\
&= \frac{1}{k} \int p_X(x_1) \dots p_X(x_k) \max_{i=1}^k p(y | x_i) dx_1 \dots dx_k dy.
\end{aligned}$$

2 Problem formulation

Let \mathcal{P} denote the collection of all joint densities $p(x, y)$ on finite-dimensional Euclidean space. For $\iota \in [0, \infty)$ define $C_k(\iota)$ to be the largest k -class average Bayes error attained by any distribution $p(x, y)$ with mutual information not exceeding ι :

$$C_k(\iota) = \sup_{p \in \mathcal{P}: I[p(x, y)] \leq \iota} \text{ABA}_k[p(x, y)].$$

A priori, $C_k(\iota)$ exists since ABA_k is bounded between 0 and 1. Furthermore, C_k is nondecreasing since the domain of the supremum is monotonically increasing with ι .

It follows that for any density $p(x, y)$, we have

$$\text{ABA}_k[p(x, y)] \leq C_k(I[p(x, y)]).$$

Hence C_k provides an upper bound for average Bayes error in terms of mutual information.

Conversely we have

$$I[p(x, y)] \geq C_k^{-1}(\text{ABA}_k[p(x, y)])$$

so that C_k^{-1} provides a lower bound for mutual information in terms of average Bayes error.

On the other hand, there is no nontrivial *lower* bound for average Bayes error in terms of mutual information, nor upper bound for mutual information in terms of average Bayes error, since

$$\inf_{p \in \mathcal{P}: I[p(x, y)] \leq \iota} \text{ABA}_k[p(x, y)] = \frac{1}{k}.$$

regardless of ι .

The goal of this work is to attempt to compute or approximate the functions C_k and C_k^{-1} .

2.1 Notation

$|\cdot|$ denotes set cardinality.

3 Special case

We work out the special case where $p(x, y)$ lies on the unit square, and $p(x)$ and $p(y)$ are both the uniform distribution. Let \mathcal{P}^{unif} denote the set of such distributions, and

$$C_k^{unif}(\iota) = \sup_{p(x,y) \in \mathcal{P}^{unif} : I[p] \leq \iota} \text{ABA}_k[p].$$

We prove the following result:

Theorem 3.1 *For any $\iota > 0$, there exists $c_\iota \geq 0$ such that defining*

$$Q_c(t) = \frac{\exp[ct^{k-1}]}{\int_0^1 \exp[ct^{k-1}]},$$

we have

$$\int_0^1 Q_{c_\iota}(t) \log Q_{c_\iota}(t) dt = \iota.$$

Then,

$$C_k^{unif} = \int_0^1 Q_{c_\iota}(t) t^{k-1} dt.$$

The proof depends on the following lemmas.

Lemma 3.2 *Let $f(t)$ be an increasing function from $[a, b] \rightarrow \mathbb{R}$, where $a < b$, and let $g(t)$ be a bounded continuous function from $[a, b] \rightarrow \mathbb{R}$. Define the set*

$$A = \{t : f(t) \neq g(t)\}.$$

Then, we can write A as a countable union of intervals

$$A = \bigcup_{i=1}^{\infty} A_i$$

where A_i are mutually disjoint intervals, with $\inf A_i < \sup A_i$, and for each i , either $f(t) > g(t)$ for all $t \in A_i$ or $f(t) < g(t)$ for all $t \in A_i$.

Lemma 3.3 *Let $f(t)$ be a measurable function from $[a, b] \rightarrow \mathbb{R}$, where $a < b$. Then there exists sets \mathcal{B}_0 and \mathcal{B}_1 , satisfying the following properties:*

- $\mathcal{B} = \mathcal{B}_0 \cup \mathcal{B}_1$ is countable partition of $[a, b]$,
- $f(t)$ is constant on all $B \in \mathcal{B}_0$, but not constant on any proper super-interval $B' \supset B$, and
- $B \in \mathcal{B}_1$ contains no positive-length subinterval where $f(t)$ is constant.

Lemma 3.4 *For any measure G on $[0, \infty]$, let G^k denote the measure defined by*

$$G^k(A) = G(A)^k,$$

and define

$$E[G] = \int x dG(x).$$

$$I[G] = \int x \log x dG(x)$$

and

$$\psi_k[G] = \int x d(G^k)(x).$$

Then, defining Q_c and c_ι as in Theorem 1, we have

$$\sup_{G: E[G]=1, I[G] \leq \iota} \psi_k[G] = \int_0^1 Q_{c_\iota}(t) t^{k-1} dt.$$

Furthermore, the supremum is attained by a measure G that has cdf equal to Q_c^{-1} , and thus has a density g with respect to Lesbegue measure.

Lemma 3.5 *The map*

$$\iota \rightarrow \int_0^1 Q_{c_\iota}(t) t^{k-1} dt$$

is concave in $\iota > 0$.

Proof of Lemma 3.2. (This will appear in the appendix of the paper.)

The function $h(t) = f(t) - g(t)$ is measurable, since all increasing functions are measurable. Define $A^+ = \{t : f(t) > g(t)\}$ and $A^- = \{t : f(t) < g(t)\}$. Since A^+ and A^- are measurable subsets of \mathbb{R} , they both admit countable partitions consisting of open, closed, or half-open intervals. Let \mathcal{H}^+ be the

collection of all partitions of A^+ consisting of such intervals. There exists a least refined partition \mathcal{A}^+ within \mathcal{H}^+ . Define \mathcal{A}^- analogously, and let

$$\mathcal{A} = \mathcal{A}^+ \cup \mathcal{A}^-$$

and enumerate the elements

$$\mathcal{A} = \{A_i\}_{i=1}^\infty.$$

We claim that the partitions \mathcal{A}^+ and \mathcal{A}^- have the property that for all $t \in A^\pm$, the interval $I \in \mathcal{A}^\pm$ containing t has endpoints $l \leq u$ defined by

$$l = \inf_{x \in [a, b]} \{x : \text{Sign}(h([x, t])) = \{\text{Sign}(h(t))\}\}$$

and

$$u = \sup_{x \in [a, b]} \{x : \text{Sign}(h([t, x])) = \{\text{Sign}(h(t))\}\}.$$

We prove the claim for the partition \mathcal{A}^+ . Take $t \in A^+$ and define l and u as above. It is clear that $(l, u) \in A^+$, and furthermore, there is no $l' < l$ and $u' > u$ such that $(l', x) \in A^+$ or $(x, u') \in A^+$ for any $x \in I$. Let \mathcal{H} be any other partition of A^+ . Some disjoint union of intervals $H_i \in \mathcal{H}$ necessarily covers I for $i = 1, \dots$, and we can further require that none of the H_i are disjoint with I . Since each H_i has nonempty intersection with I , and I is an interval, this implies that $\cup_i H_i$ is also an interval. Let $l'' \leq u''$ be the endpoints of $\cup_i H_i$. Since $I \subseteq \cup_i H_i$, we have $l'' \leq l \leq u \leq u''$. However, since also $I \in A^+$, we must have $l \leq l'' \leq u'' \leq u$. This implies that $l'' = l$ and $u'' = u$. Since $\cup_i H_i = I$, and this holds for any $I \in \mathcal{A}^+$, we conclude that \mathcal{H} is a refinement of \mathcal{A}^+ . The proof of the claim for \mathcal{A}^- is similar.

It remains to show that there are not isolated points in \mathcal{A} , i.e. that for all $I \in \mathcal{A}$ with endpoints $l \leq u$, we have $l < u$. Take $I \in \mathcal{A}$ with endpoints $l \leq u$ and let $t = \frac{l+u}{2}$. By definition, we have $h(t) \neq 0$. Consider the two cases $h(t) > 0$ and $h(t) < 0$.

If $h(t) > 0$, then $t' = g^{-1}(h(t)) > t$, and for all $x \in [t, t']$ we have $h(x) > 0$. Therefore, it follows from definition that $[t, t'] \in I$, and since $l \leq t < t' \leq u$, this implies that $l < u$. The case $h(t) < 0$ is handled similarly. \square

Proof of Lemma 3.3. (This will appear in the appendix of the paper.) To construct the interval, define

$$l(t) = \inf\{x \in [0, 1] : f([x, t]) = \{f(t)\}\}$$

$$u(t) = \sup\{x \in [0, 1] : f([t, x]) = \{f(t)\}\},$$

Let B_0 be the set of all t such that $l(t) < u(t)$, and let B_1 be the set of all t such that $l(t) = t = u(t)$. For all $t \in B_0$, define

$$I(t) = (l(t), u(t)) \cup \{x \in \{l(t), u(t)\} : f(x) = f(t)\}.$$

Then we claim

$$\mathcal{B}_0 = \{I(t) : t \in B_0\}$$

is a countable partition of B_0 . The claim follows since the members of \mathcal{B}_0 are disjoint intervals of nonzero length, and B_0 has finite length. It follows from definition that for any $B \in \mathcal{B}_0$, that f is not constant on any proper superinterval $B' \supset B$.

Meanwhile, let \mathcal{B}_1 be a countable partition of B_1 into intervals.

Next, we show that for all $I \in \mathcal{B}_1$, I does not contain a subinterval I' of nonzero length such that f is constant on I' . Suppose to the contrary, we could find such an interval I and subinterval I' . Then for any $t \in I'$, we have $t \in B_0$. However, this implies that $t \notin B_1$, a contradiction.

Since $t \in [a, b]$ belongs to either B_0 or B_1 , letting $\mathcal{B} = \mathcal{B}_0 \cup \mathcal{B}_1$ yields the desired partition of $[a, b]$. \square .

Proof of Lemma 3.4. (This will appear in the appendix of the paper.)

Consider the quantile function $Q(t) = \inf_{x \in [0, 1]} : G((-\infty, x]) \geq t$. $Q(t)$ must be a monotonically increasing function from $[0, 1]$ to $[0, \infty)$. Let \mathcal{Q} denote the collection of all such quantile functions.

We have

$$E[G] = \int_0^1 Q(t) dt$$

$$\psi_k[G] = \int_0^1 Q(t) x^{k-1} dt.$$

and

$$I[G] = \int_0^1 Q(t) \log Q(t) dt.$$

For any given ι , let P_ι denote the class of probability distributions G on $[0, \infty]$ such that $E[G] = 1$ and $I[G] \leq \iota$. From Markov's inequality, for any $G \in P_\iota$ we have

$$G([x, \infty]) \leq x^{-1}$$

for any $x \geq 0$, hence P_ι is tight. From tightness, we conclude that P_ι is closed under limits with respect to weak convergence. Hence, since ψ_k is a

continuous function, there exists a distribution $G^* \in P_\iota$ which attains the supremum

$$\sup_{G \in P_\iota} \psi_k[G].$$

Let \mathcal{Q}_ι denote the collection of quantile functions of distributions in P_ι . Then, \mathcal{Q}_ι consists of monotonic functions $Q : [0, 1] \rightarrow [0, \infty]$ which satisfy

$$E[Q] = \int_0^1 Q(t) dt = 1,$$

and

$$I[Q] = \int_0^1 Q(t) \log Q(t) dt \leq \iota.$$

Let \mathcal{Q} denote the collection of *all* quantile functions from measures on $[0, \infty]$. And letting Q^* be the quantile function for G^* , we have that Q^* attains the supremum

$$\sup_{Q \in \mathcal{Q}_\iota} \phi_k[Q] = \sup_{Q \in \mathcal{Q}_\iota} \int_0^1 Q(t) t^{k-1} dt. \quad (1)$$

Therefore, there exist Lagrange multipliers $\lambda \geq 0$ and $\nu \leq 0$ such that defining

$$\mathcal{L}[Q] = E[Q] + \lambda \phi_k[Q] + \nu I[Q] = \int_0^1 Q(t) (1 + \lambda \log Q(t) + \nu t^{k-1}) dt,$$

Q^* attains the infimum of $\mathcal{L}[Q]$ over *all* quantile functions,

$$\mathcal{L}[Q^*] = \inf_{Q \in \mathcal{Q}} \mathcal{L}[Q].$$

We now claim that for such λ and ν , we have

$$1 + \lambda + \lambda \log Q(t) + \nu t^{k-1} = 0.$$

Consider a perturbation function $\xi : [0, 1] \rightarrow \mathbb{R}$. We have

$$\mathcal{L}[Q + \xi] \approx \mathcal{L}[Q] + \int_0^1 \xi(t) (1 + \lambda + \lambda \log Q(t) + \nu t^{k-1}) dt \quad (2)$$

for small ξ . Define

$$\nabla \mathcal{L}_{Q^*}(t) = 1 + \lambda + \lambda \log Q^*(t) + \nu t^{k-1}. \quad (3)$$

The function $\nabla \mathcal{L}_{Q^*}(t)$ is a *functional derivative* of the Lagrangian. Note that if we were able to show that $\nabla \mathcal{L}_{Q^*}(t) = 0$, this immediately yields

$$Q^*(t) = \exp[-\lambda^{-1} - 1 - \nu \lambda^{-1} t^{k-1}]. \quad (4)$$

At this point, we know that the right-hand side of (4) gives a stationary point of \mathcal{L} , but we cannot be sure that it gives the global minimizer. The reason is because the optimization occurs on a constrained space. We will show that (4) indeed gives the global minimizer Q^* , but we do so by showing that the set of points t where $\nabla \mathcal{L}_{Q^*}(t) \neq 0$ is of zero measure. Since sets of zero measure don't affect the integrals defining the optimization problem (1), we conclude there exists a global optimal solution with $\nabla \mathcal{L}_{Q^*}(t) = 0$ everywhere, which is therefore given explicitly by (4) for some $\lambda > 0$, $\nu < 0$.

The rest of the proof proceeds as follows. We will use Lemmas 3.2 and 3.3 to define a decomposition $A = D_0 \cup D_1 \cup D_2$, where D_2 is of measure zero. First, we show that assuming the existence of $t \in D_0$ yields a contradiction, and hence $D_0 = \emptyset$. Then, again using argument from contradiction we establish that $D_1 = \emptyset$. Finally, since D_2 is a set of zero measure, this allows us to conclude that the $Q^*(t) = 0$ on all but a set of zero measure.

We will now apply the Lemmas to obtain the necessary ingredients for constructing the sets D_i . Since $\nabla \mathcal{L}_{Q^*}(t)$ is a difference between an increasing function and a continuous strictly increasing function, we can apply Lemma 3.2 to conclude that there exists a countable partition \mathcal{A} of the set $A : \{t \in [0, 1] : \nabla \mathcal{L}_{Q^*}(t) \neq 0\}$ into intervals such that for all $J \in \mathcal{A}$, $|\text{Sign}(\nabla Q^*(J))| = 1$ and $\inf J < \sup J$. Applying Lemma 3.3 we get a countable partition $\mathcal{B} = \mathcal{B}_0 \cup \mathcal{B}_1$ of $[0, 1]$ so that each element $J \in \mathcal{B}_0$ is an interval such that $\nabla \mathcal{L}_{Q^*}(t)$ is constant on J , and furthermore is not properly contained in any interval with the same property, and each element $J \in \mathcal{B}_1$ is an interval, such that J contains no positive-length subinterval where $\nabla \mathcal{L}_{Q^*}(t)$ is constant. Also define B_i as the union of the sets in \mathcal{B}_i for $i = 0, 1$.

Note that B_0 is necessarily a subset of A . That is because if $\nabla \mathcal{L}_{Q^*}(t) = 0$ on any interval J , then that $Q^*(t)$ is necessarily not constant on the interval.

We will construct a new countable partition of A , called \mathcal{D} . The partition \mathcal{D} is constructed by taking the union of three families of intervals,

$$\mathcal{D} = \mathcal{D}_0 \cup \mathcal{D}_1 \cup \mathcal{D}_2.$$

Define D_i to be the union of intervals in \mathcal{D}_i for $i = 0, 1, 2$.

Define $\mathcal{D}_0 = \mathcal{B}_0$, Define a countable partition \mathcal{D}_1 by

$$\mathcal{D}_1 = \{J \cap L : J \in \mathcal{A}, L \in \mathcal{B}_1, \text{ and } |L| > 1\},$$

in order words, \mathcal{D}_1 consists of positive-length intervals where $\nabla Q^*(t)$ is entirely positive or negative and is not constant. Define

$$\mathcal{D}_2 = \{J \in \mathcal{B}_1 : J \subset A \text{ and } |J| = 1\},$$

i.e. \mathcal{D}_2 consists of isolated points in A .

One verifies that \mathcal{D} is indeed a partition of A by checking that $D_0 = B_0$, $D_1 \cup D_2 = B_1 \cap A$, so that $D_0 \cup D_1 \cup D_2 = A$: it is also easy to check that elements of \mathcal{D} are disjoint. Furthermore, as we mentioned earlier, the set D_2 is indeed of zero measure, since it consists of countably many isolated points.

Now we will show that the existence of $t \in D_0$ implies a contradiction. Take $t \in D$ for $D \in \mathcal{D}_0$, and let $a = \inf D$ and $b = \sup D$. Define

$$\xi^+ = I\{t \in D\}(Q^*(b) - Q^*(t))$$

and

$$\xi^- = I\{t \in D\}(Q^*(a) - Q^*(t)).$$

Observe that $Q + \epsilon \xi^+ \in \mathcal{Q}$ and $Q + \epsilon \xi^- \in \mathcal{Q}$ for any $\epsilon \in [0, 1]$. Now, if $\nabla \mathcal{L}_{Q^*}(t)$ is strictly positive on D , then for some $\epsilon > 0$ we would have $\mathcal{L}[Q^* + \epsilon \xi^-] < \mathcal{L}[Q^*]$, a contradiction. A similar argument with ξ^+ shows that $\nabla \mathcal{L}_{Q^*}(t)$ cannot be strictly negative on D either. From this perturbation argument, we conclude that $\nabla \mathcal{L}_{Q^*}(t) = 0$. Since this argument applies for all $t \in D_0$, we conclude that $D_0 = \emptyset$: therefore, on the set $[0, 1] \setminus (D_1 \cup D_2)$, we have $\nabla \mathcal{L}_{Q^*}(t) = 0$.

The following observation is needed for the next stage of the proof. If we look at the function $Q^*(t)$, then up to sets of negligible measure, it is given by the expression (4) on the set $[0, 1] \setminus D_1$, and it is piecewise constant in-between. But since (4) gives a strictly increasing function, and since Q^* is increasing, this implies that Q^* is discontinuous at the boundary of D_1 .

Now we are prepared to show that $\nabla \mathcal{L}_{Q^*}(t) = 0$ for $t \in D_1$. Take $t \in D$ for $D \in \mathcal{D}_1$, and let $a = \inf D$ and $b = \sup D$. From the previous argument, there is a discontinuity at both a and b , so that $\lim_{u \rightarrow a^-} Q(u) < Q(t) < \lim_{u \rightarrow b^+} Q(u)$. Therefore, for any $\xi(t)$ which is increasing on D and zero elsewhere, there exists $\epsilon > 0$ such that $\nabla Q^* + \epsilon \xi \in \mathcal{Q}$. It remains to find such a perturbation ξ such that $\mathcal{L}[Q + \epsilon \xi] < \mathcal{L}[Q]$.

Also, since by definition $\nabla \mathcal{L}_{Q^*}(t)$ is constant on D , follows from (3) that ∇Q^* is strictly decreasing, and thus either

- Case 1: $\nabla \mathcal{L}_{Q^*}(t) \geq 0$ on D ,
- Case 2: $\nabla \mathcal{L}_{Q^*}(t) \leq 0$ on D , or
- Case 3: $\nabla \mathcal{L}_{Q^*}(t) \geq 0$ for all $t \in D \cap [a, t_0]$ and $\nabla \mathcal{L}_{Q^*}(t) \leq 0$ for all $t \in D \cap [t_0, b]$.

Depending on the case, we construct a suitable perturbation ξ :

- Case 1: Construct $\xi(t) = -I\{t \in D\}$.
- Case 2: Construct $\xi(t) = I\{t \in D\}$
- Case 2: Construct

$$\xi(t) = \begin{cases} -1 & \text{for } t \in D \cap [a, t_0], \\ 0 & \text{otherwise.} \end{cases}$$

In all three cases, given the corresponding construction for $\xi(t)$ we get

$$\int_0^1 \xi(t) \nabla \mathcal{L}_{Q^*}(t) dt < 0.$$

Therefore, from (2), there exists some $\epsilon > 0$ such that $\mathcal{L}[Q + \epsilon\xi] < \mathcal{L}[Q]$, a contradiction. Again, since the contradiction applies for all $t \in D_1$, we conclude that $D_1 = \emptyset$.

By now we have established that a global optimum for (1) exists, and is given by (4) for some $\lambda > 0$, $\nu < 0$. It remains to determine the values of λ and ν .

Reparameterize $\alpha = \exp[-1 - \lambda^{-1}]$ and $\beta = -\nu\lambda t^{k-1}$. Therefore,

$$Q^*(t) = \alpha \exp[\beta t^{k-1}]$$

for $\alpha > 0$, $\beta > 0$. There is a one-to-one mapping from $(\alpha, \beta) \in (0, \infty)^2$ to $(\lambda, \nu) \in (0, \infty) \times (-\infty, 0)$.

Now, from the constraint

$$1 = E[Q^*] = \int_0^1 \alpha \exp[\beta t^{k-1}] dt.$$

we conclude that

$$\alpha = \frac{1}{\int_0^1 \exp[\beta t^{k-1}] dt}.$$

Therefore, we have reduced the set of possible solutions Q^* to a one-parameter family,

$$Q^*(t) = \frac{\exp[\beta t^{k-1}]}{Z(\beta)}.$$

where

$$Z(\beta) = \int_0^1 \exp[\beta t^{k-1}] dt.$$

Next, consider that

$$I[Q^*] = \int_0^1 Q^*(t) \log Q^*(t) = \frac{\beta}{Z(\beta)} \int_0^1 \exp[\beta t^{k-1}] t^{k-1} dt - \log Z(\beta).$$

Remark. More specifically, the supremum is attained by a distribution with density $p_\iota(x, y)$ where

$$p_\iota(x, y) = \begin{cases} g_\iota(y - x) & \text{for } x \geq y \\ g_\iota(1 + y - x) & \text{for } x < y \end{cases}$$

where

$$g_\iota(x) = \frac{d}{dx} G_\iota(x)$$

and G_ι is the inverse of Q_c .

In this case, letting $X_1, \dots, X_k \sim \text{Unif}[0, 1]$, and $Y \sim \text{Unif}[0, 1]$ define $Z_i(y) = p(y|X_i)$. We have $\mathbf{E}(Z(y)) = 1$ and,

$$I[p(x, y)] = \mathbf{E}(Z(Y) \log Z(Y))$$

while

$$\text{ABA}_k[p(x, y)] = k^{-1} \mathbf{E}(\max_i Z_i(Y)).$$

Letting g_y be the density of $Z(y)$, we have

$$I[p(x, y)] = \mathbf{E}(-H[g_Y])$$

and

$$\text{ABA}_k[p(x, y)] = \mathbf{E}(\psi_k[g_Y])$$

where

$$H[g] = - \int g(x) x \log x dx$$

and

$$\psi_k[g] = \int xg(x)G(x)^{k-1}dx$$

for $G(x) = \int_0^x g(t)dt$. Additionally g_y satisfies the constraint $\int xg(x)dx = 1$ since $\mathbf{E}[Z(y)] = 1$.

Define the set $D = \{(\alpha, \beta)\}$ as the set of possible values of $(-H[g], \psi_k[g])$ taken over all distributions g supported on $[0, \infty)$ with $\int xg(x)dx = 1$. Next, let $\mathcal{C}(D)$ denote the convex hull of D . It follows that $(I[p], \text{ABA}_k[p]) \in \mathcal{C}(D)$ since the pair is obtained via a convex average of points $(-H[g_y], \psi_k[g_y])$.

Define the upper envelope of D as the curve

$$d_k(\alpha) = \sup\{\beta : (\alpha, \beta) \in D\}.$$

We make the claim (to be shown in the following section) that $d_k(\alpha)$ is convex in α . As a result, the upper envelope of D is also the upper envelope of $\mathcal{C}(D)$. This in turn implies that $C_k^{unif}(\iota) = d_k(\iota)$. We establish these results, along with a open-form expression for C_k^{unif} , in the following section.

3.1 Variational methods

Consider the quantile function $Q(t) = G^{-1}(t)$. $Q(t)$ must be a continuous function from $[0, 1]$ to $[0, \infty)$. We can rewrite the moment constraint $\mathbf{E}[g] = 1$ as

$$\int_0^1 Q(t)dt = 1.$$

Meanwhile, $\beta = \psi_k[g]$ takes the form

$$\beta = \int_0^1 Q(t)x^{k-1}dt.$$

and $\alpha = -H[g]$ takes the form

$$\alpha = \int_0^1 Q(t) \log Q(t)dt.$$

To find the upper envelope, it will be useful to write the Langrangian

$$\begin{aligned} \mathcal{L}[g] &= \lambda \int_0^1 Q(t)dt + \mu \int_0^1 Q(t)x^{k-1}dt + \lambda \int_0^1 Q(t) \log Q(t)dt \\ &= \int_0^1 Q(t)(\lambda + \mu x^{k-1} + \nu \log Q(t))dt. \end{aligned}$$

In order for a quantile function $Q(t)$ to be on the upper envelope, it must be a local maximum of $-H$ with respect to small perturbations. Therefore, consider the functional derivative

$$D[\xi] = \lim_{\epsilon \rightarrow 0} \frac{\mathcal{L}[g + \epsilon\xi] - \mathcal{L}[g]}{\epsilon}.$$

We have

$$D[\xi] = \int_0^1 \xi(t)(\lambda + \nu + \mu x^{k-1} + \nu \log Q(t))dt.$$

Now consider the following three cases:

- $Q(t)$ is strictly monotonic, i.e. $Q'(t) > 0$.
- $Q(t)$ is differentiable but not strongly monotonic:
- $Q(t)$ is not strongly monotonic: there exist intervals $A_i = [a_i, b_i]$ such that $Q(t)$ is constant on A_i , and isolated points t_i where $Q'(t_i) = 0$.

Strictly monotonic case. Because Q is defined on a closed interval, strict monotonicity further implies the property of *strong monotonicity* where $\inf[0, 1]Q'(t) > 0$. Therefore, for any differentiable perturbation $\xi(t)$ with $\sup |\xi'(t)| < \infty$, and further imposing that $\xi(0) \geq 0$ in the case that $Q(0) = 0$, there exists some $\epsilon > 0$ such that $(Q + \epsilon\xi)(t)$ is still a valid quantile function. Therefore, in order for $Q(t)$ to be a local maximum, we must have

$$0 = \lambda + \nu + \mu x^{k-1} + \nu \log Q(t)$$

for $t \in [0, 1]$. This implies that

$$Q(t) = c_0 e^{-c_1 x^{k-1}}$$

for some $c_0, c_1 \geq 0$.

Other cases. (TODO) We have to show that these cannot be local maxima.

4 General case

We claim that the constants $C_k^{unif}(\iota)$ obtained for the special case also apply for the general case, i.e.

$$C_k(\iota) = C_k^{unif}(\iota).$$

We make use of the following Lemma:

Lemma. Suppose X, Y, W, Z are continuous random variables, and that $W \perp Y|Z$, $Z \perp X|Y$, and $W \perp Z|(X, Y)$. Then,

$$I[p(x, y)] = I[p((x, w), (y, z))]$$

and

$$ABA_k[p(x, y)] = ABA_k[p((x, w), (y, z))].$$

Proof. Due to conditional independence relationships, we have

$$p((x, w), (y, z)) = p(x, y)p(w|x)p(z|y).$$

It follows that

$$\begin{aligned} I[p((x, w), (y, z))] &= \int dx dw dy dz p(x, y)p(w|x)p(z|w) \log \frac{p((x, w), (y, z))}{p(x, w)p(y, z)} \\ &= \int dx dw dy dz p(x, y)p(w|x)p(z|w) \log \frac{p(x, y)p(w|x)p(z|y)}{p(x)p(y)p(w|x)p(z|y)} \\ &= \int dx dw dy dz p(x, y)p(w|x)p(z|w) \log \frac{p(x, y)}{p(x)p(y)} \\ &= \int dx dy p(x, y) \log \frac{p(x, y)}{p(x)p(y)} = I[p(x, y)]. \end{aligned}$$

Also,

$$\begin{aligned} ABA_k[p((x, w), (y, z))] &= \int \left[\prod_{i=1}^k p(x_i, w_i) dx_i dw_i \right] \int dy dz \max_i p(y, z|x_i, w_i). \\ &= \int \left[\prod_{i=1}^k p(x_i, w_i) dx_i dw_i \right] \int dy \max_i p(y|x_i) \int dz p(z|y). \\ &= \int \left[\prod_{i=1}^k p(x_i) dx_i \right] \left[\prod_{i=1}^k \int dw_i p(w_i|x_i) \right] \int dy \max_i p(y|x_i) \\ &= ABA_k[p(x, y)]. \end{aligned}$$

□

Next, we use the fact that for any $p(x, y)$ and $\epsilon > 0$, there exists a discrete distribution $p_\epsilon(\tilde{x}, \tilde{y})$ such that

$$|\mathbb{I}[p(x, y)] - \mathbb{I}[p_\epsilon(\tilde{x}, \tilde{y})]| < \epsilon,$$

where for discrete distributions, one defines

$$\mathbb{I}[p(x, y)] = \sum_x \sum_y p(x, y) \log \frac{p(x, y)}{p(x)p(y)}.$$

We require the additional condition that the marginals of the discrete distribution are close to uniform: that is, for some $\delta > 0$, we have

$$\sup_{x, x': p_\epsilon(x) > 0 \text{ and } p_\epsilon(x') > 0} \frac{p_\epsilon(x)}{p_\epsilon(x')} \leq 1 + \delta.$$

and likewise

$$\sup_{y, y': p_\epsilon(y) > 0 \text{ and } p_\epsilon(y') > 0} \frac{p_\epsilon(y)}{p_\epsilon(y')} \leq 1 + \delta.$$

To construct the discretization with the required properties, choose a regular rectangular grid Λ over the domain of $p(x, y)$ sufficiently fine so that partitioning X, Y into grid cells, we have

$$|\mathbb{I}[p(x, y)] - \mathbb{I}[\tilde{p}(\tilde{x}, \tilde{y})]| < \epsilon.$$

[NOTE: to be written more clearly] Next, define