

A NUMERICAL METHOD FOR SOLVING THE EIKONAL EQUATION ON SIMPLICIAL TESSELATIONS*

SACHIN NATESH[†]

Abstract. In this paper, we describe a method for solving the Eikonal equation on domains in $\mathbb{R}^2$ or $\mathbb{R}^3$ tessellated by triangles or tetrahedra. Our method generalizes to simplices in higher dimensions, owing to our choice of solution representation under the Jacobi family of orthogonal polynomials on the simplex. First, we describe a spectral decomposition for functions supported on the simplex in terms of their weighted L_2 expansion coefficients under the Jacobi basis. Computing this expansion requires an accurate and efficient quadrature, and we provide a novel optimization routine based on approximate joint-diagonalization of Jacobi matrices for generating such quadratures. Then, we consider differential operators acting on coefficients of a Jacobi expansion. Recent results due to Townsend, Oliver and Vasil enable the construction of sparse differential operators on the standard triangle, interoperating within the Jacobi polynomial family parameter space, and leading to low-storage, banded discrete differential operators. We generalize these results to the tetrahedra (laying the path for generalizing to $d > 3$), and use the resulting differential operators to construct the Eikonal operator on the simplex. Ultimately, we must solve a non-linear PDE, and working in coefficient space implies that our solution variables must be thought of as functions. Hence, we reformulate the Eikonal problem in terms of an infinite-dimensional L_2 minimization problem, which can be discretized with the aforementioned numerical quadrature. As with the README, this is more of a blog, but it's much easier to typeset math here.

Key words. multivariate orthogonal polynomials, spectral methods, partial differential equations, Eikonal equations, simplex

MSC codes. 68Q25, 68R10, 68U05

1. Solutions to the Eikonal Equation. Haven't really looked into the viscosity perspective, but..I think I converged on the same idea for solving the PDE. Consider the standard simplex

$$\mathcal{T}^d = \{\mathbf{x} = (x_1, \dots, x_d) \in \mathbb{R}^d \mid x_i \geq 0, 1 - |\mathbf{x}|_1 \geq 0\}$$

The *minimal* time $u(\mathbf{x})$ to reach the boundary $\partial\mathcal{T}^d$ from a point $x \in \mathcal{T}^d$ is computable through solving the Eikonal equation on $\mathcal{T}^d$:

$$(1.1) \quad \begin{cases} \sum_{i=1}^d (\partial_{x_i} u)^2 = \left(\frac{1}{f}\right)^2, & x \in \mathcal{T}^d, \\ u = q, & x \in \partial\mathcal{T}^d, \end{cases}$$

where $f : \overline{\mathcal{T}^d} \rightarrow \mathbb{R}_*^+$ is the speed in the medium, $q : \partial\mathcal{T}^d \rightarrow \mathbb{R}^+$ is the time to leave $\mathcal{T}^d$ from a point on $\partial\mathcal{T}^d$. We seek to represent u under an optimal, finite, weighted L_2 expansion, wherein

$$u(\mathbf{x}) \approx \sum_{j=1}^n u_j P_j(\mathbf{x}) = \mathbf{u} \cdot \mathbf{P}(\mathbf{x}),$$

and $\mathbf{u}$ is the vector in $\mathbb{R}^n$ of coefficients of u under the elements of the basis $\mathbf{P}(\mathbf{x}) : \mathbb{R}^d \rightarrow \mathbb{R}^n$. Here $\approx$ must be taken in the sense of weighted L_2 convergence of the series to u as $n \rightarrow \infty$. For the typical case of $q \equiv 0$ (i.e. time to boundary is 0 if on the boundary), we employ a weighted basis which evaluates to 0 on the boundary, thereby naturally satisfying homogeneous Dirichlet conditions.

*Submitted to the editors DATE.

Funding: This work was funded by nobody

[†]Department of Applied Mathematics, University of Colorado, Boulder, CO

1.1. Jacobi bases on the Triangle. Let us specialize for $d = 2$. By P_k^n , we denote the k^{th} Jacobi polynomial on the triangle of degree n with parameters (a, b, c) . When the parameters must be specified, we will refer to the same polynomial by $P_{k,n}^{(a,b,c)}$. The weight function is

$$(1.2) \quad \begin{aligned} W^{(a,b,c)}(x, y) &= x^{a-\frac{1}{2}} y^{b-\frac{1}{2}} (1-x-y)^{c-\frac{1}{2}}, \\ w_{(a,b,c)} &= \left(\int_{\mathcal{T}_{\text{ref}}^2} W(x, y) dx dy \right)^{-1} = \frac{\Gamma(|\kappa| + \frac{3}{2})}{\Gamma(a + \frac{1}{2}) \Gamma(b + \frac{1}{2}) \Gamma(c + \frac{1}{2})} \\ w^{(a,b,c)}(\mathbf{x}) &= w_{(a,b,c)} W^{(a,b,c)}(\mathbf{x}) \end{aligned}$$

with $a, b, c > -1/2$ the Jacobi parameters, and $|\kappa| = a + b + c$. The orthonormal polynomials are given in terms of the 1D Jacobi polynomials by

$$(1.3) \quad \begin{aligned} P_k^n(x, y) &= (h_{k,n})^{-1} P_{n-k}^{2k+b+c, a-1/2}(2x-1) (1-x)^k P_k^{c-1/2, b-1/2}\left(\frac{2y}{1-x} - 1\right), \\ (h_{k,n})^2 &= \frac{w_{a,b,c}}{(2n + |\kappa| + \frac{1}{2})(2k + b + c)} \\ &\times \frac{\Gamma(n + k + b + c + 1) \Gamma(n - k + a + \frac{1}{2}) \Gamma(k + b + \frac{1}{2}) \Gamma(k + c + \frac{1}{2})}{(n - k)! k! \Gamma(n + k + |\kappa| + \frac{1}{2}) \Gamma(k + b + c)}, \end{aligned}$$

and they satisfy a mutual orthonormality condition under the weighted inner product:

$$(1.4) \quad \langle P_{j,m}^{(a,b,c)}, P_{k,n}^{(a,b,c)} \rangle_{w^{(a,b,c)}} = \delta_{jk} \delta_{mn}.$$

A given basis polynomial is not only orthonormal to those with different total degree, but also to those others within the same-degree homogeneous subspace. In other words, P_k^n , $0 \leq k \leq n$ forms a basis for the space of homogeneous polynomials of degree n . By considering ordered collections of these subspaces (where ordering is in terms of the degree n of the space), instead of individual polynomials, we can more elegantly reformulate the Eikonal problem into a weak-type form. Let

$$\mathbb{P}_n(x_1, x_2) = (P_0^n(x_1, x_2), \dots, P_n^n(x_1, x_2))^T,$$

and let

$$\mathbf{P} = (\mathbb{P}_0^T, \mathbb{P}_1^T, \dots, \mathbb{P}_{n-1}^T)^T.$$

The above notation amounts to a vectorization of the basis for the general space of polynomials in two variables Π_{n-1}^2 defined of $\mathcal{T}^2$.

Now, consider the $(n-1)$ order Jacobi polynomial expansion of the solution u under the parameter family (a, b, c) . For $\mathbf{x} \in \mathcal{T}^d$, and $N = \dim \Pi_{n-1}^2$ coefficients $\mathbf{u} \in \mathbb{R}^N$, we can write it as

$$(1.5) \quad u(\mathbf{x}) = \mathbf{P}^{(a,b,c)}(\mathbf{x})^T \mathbf{u}.$$

There exists sparse discrete operators which map $\mathbf{u}$ to the coefficients of the derivative of u under a modified basis, as well as those which promote the basis parameters without taking derivatives:

$$(1.6) \quad \begin{aligned} \partial_{x_1} u(\mathbf{x}) &= \mathbf{P}^{(a+1, b+1, c+1)}(\mathbf{x})^T \mathcal{K}_{(a+1, b, c+1)}^{(a+1, b+1, c+1)} \mathcal{D}_{x_1} \mathbf{u} \\ \partial_{x_2} u(\mathbf{y}) &= \mathbf{P}^{(a+1, b+1, c+1)}(\mathbf{x})^T \mathcal{K}_{(a, b+1, c+1)}^{(a+1, b+1, c+1)} \mathcal{D}_{x_2} \mathbf{u} \end{aligned}$$

Inserting this into the fully constrained Eikonal equation gives

$$\begin{aligned}
 (1.7) \quad & \left(\mathbf{P}^{(a+1,b+1,c+1)}(\mathbf{x})^T \mathcal{K}_{(a,b,c)}^{(a+1,b+1,c+1)} \mathbf{c}_{\frac{1}{f}} \right)^2 = \\
 & \left(\mathbf{P}^{(a+1,b+1,c+1)}(\mathbf{x})^T \mathcal{K}_{(a+1,b,c+1)}^{(a+1,b+1,c+1)} \mathcal{D}_{x_1} \mathbf{u} \right)^2 + \\
 & \left(\mathbf{P}^{(a+1,b+1,c+1)}(\mathbf{x})^T \mathcal{K}_{(a,b+1,c+1)}^{(a+1,b+1,c+1)} \mathcal{D}_{x_2} \mathbf{u} \right)^2 = \\
 (1.8) \quad & \mathbf{P}^{(a+1,b+1,c+1)}(\mathbf{x})^T \left(\tilde{\mathcal{D}}_{x_1} \mathbf{u} \mathbf{u}^T \tilde{\mathcal{D}}_{x_1}^T + \tilde{\mathcal{D}}_{x_2} \mathbf{u} \mathbf{u}^T \tilde{\mathcal{D}}_{x_2}^T \right) \mathbf{P}^{(a+1,b,c+1)}(\mathbf{x}),
 \end{aligned}$$

where

$$\tilde{\mathcal{D}}_{x_1} = \mathcal{K}_{(a+1,b,c+1)}^{(a+1,b+1,c+1)} \mathcal{D}_{x_1}, \quad \tilde{\mathcal{D}}_{x_2} = \mathcal{K}_{(a,b+1,c+1)}^{(a+1,b+1,c+1)} \mathcal{D}_{x_2}$$

and, the coefficients of $(1/f)$ are represented as $\mathbf{c}_{\frac{1}{f}}$ but promoted to the $(a+1, b+1, c+1)$ Jacobi basis. We let $\tilde{\cdot}$ indicate promotion *to* $(a+1, b+1, c+1)$ from any other parameter set. The above equality holds for any point $\mathbf{x} \in \mathcal{T}^2$. So, for a solution $\mathbf{u}$, the coefficient operation must be such that:

$$(1.9) \quad (\tilde{\mathcal{D}}_{x_1} \mathbf{u} \mathbf{u}^T \tilde{\mathcal{D}}_{x_1}^T + \tilde{\mathcal{D}}_{x_2} \mathbf{u} \mathbf{u}^T \tilde{\mathcal{D}}_{x_2}^T) = \tilde{\mathbf{c}}_{\frac{1}{f}} \tilde{\mathbf{c}}_{\frac{1}{f}}^T.$$

Now, we seek a coefficient solution $\mathbf{u}$ which represents the solution *globally* in $\mathcal{T}^2$. However, polynomials are differentiable, and some of the problems we are interested in yield non-differentiable solutions. This is true even in the most basic case of $f \equiv 1$, in which u is the signed distance function from $\mathbf{x}$ to $\partial\mathcal{T}^2$, depicted below:




FIG. 1. A simple calculation shows that the minimum distance from $\mathbf{x} \in \mathcal{T}^2$ to the boundary $\partial\mathcal{T}^2$ is given by the minimum of $\left\{ x_1, x_2, \left\| \mathbf{x} - \begin{bmatrix} (x_1 - x_2 + 1)/2 \\ (x_2 - x_1 + 1)/2 \end{bmatrix} \right\|_2 \right\}$

We cannot hope to recover a solution $\mathbf{u}$ which satisfies (1.9) to machine precision, as that would imply the global solution has zero residual *pointwise* everywhere. Hence,

we must return to (1.7), and consider solving it weakly, in the sense of:

(1.10)

$$\begin{aligned} \int_{\mathcal{T}^2} w^{(a+1,b+1,c+1)}(\mathbf{x}) \mathbf{P}^{(a+1,b+1,c+1)}(\mathbf{x})^T \tilde{\mathbf{c}}_{\frac{1}{f}}^T \tilde{\mathbf{c}}_{\frac{1}{f}} \mathbf{P}^{(a+1,b+1,c+1)}(\mathbf{x}) d\mathbf{x} &\approx \|1/f\|_{w^{(a+1,b+1,c+1)}}^2 \\ &\approx \int_{\mathcal{T}^2} w^{(a+1,b+1,c+1)}(\mathbf{x}) \mathbf{P}^{(a+1,b+1,c+1)}(\mathbf{x})^T (\tilde{\mathcal{D}}_{x_1} \mathbf{u} \mathbf{u}^T \tilde{\mathcal{D}}_{x_1}^T + \tilde{\mathcal{D}}_{x_2} \mathbf{u} \mathbf{u}^T \tilde{\mathcal{D}}_{x_2}^T) \mathbf{P}^{(a+1,b+1,c+1)}(\mathbf{x}) d\mathbf{x} \\ &\approx \|\partial_x u\|_{w^{(a+1,b+1,c+1)}}^2 + \|\partial_y u\|_{w^{(a+1,b+1,c+1)}}^2. \end{aligned}$$

Here, the norm is viewed as that of a function in $L^2(\mathcal{T}^2, w^{(a+1,b+1,c+1)})$.

Parsing this equation out a bit further (condensing sub/super scripts), we have

$$\begin{aligned} \int_{\mathcal{T}^2} \mathbf{P}(\mathbf{x})^T (\mathbf{u}_x \mathbf{u}_x^T + \mathbf{u}_y \mathbf{u}_y^T) w(\mathbf{x}) \mathbf{P}(\mathbf{x}) d\mathbf{x} \quad (\text{Only } x \text{ part handled below}) \\ = \int_{\mathcal{T}^2} \mathbf{P}(\mathbf{x})^T \begin{bmatrix} u_{x,1}^2 & u_{x,1}u_{x,2} & \cdots & u_{x,1}u_{x,N} \\ u_{x,2}u_{x,1} & u_{x,2}^2 & \cdots & u_{x,2}u_{x,N} \\ \vdots & \ddots & \ddots & \\ u_{x,N}u_{x,1} & u_{x,N}u_{x,2} & \cdots & u_{x,N}^2 \end{bmatrix} \mathbf{P}(\mathbf{x}) w(\mathbf{x}) d\mathbf{x} \\ = \int_{\mathcal{T}^2} \begin{bmatrix} P_1(\mathbf{x}) & P_2(\mathbf{x}) & \cdots & P_N(\mathbf{x}) \end{bmatrix} \begin{bmatrix} u_{x,1}^2 & u_{x,1}u_{x,2} & \cdots & u_{x,1}u_{x,N} \\ u_{x,2}u_{x,1} & u_{x,2}^2 & \cdots & u_{x,2}u_{x,N} \\ \vdots & \ddots & \ddots & \\ u_{x,N}u_{x,1} & u_{x,N}u_{x,2} & \cdots & u_{x,N}^2 \end{bmatrix} \begin{bmatrix} P_1(\mathbf{x}) \\ P_2(\mathbf{x}) \\ \vdots \\ P_N(\mathbf{x}) \end{bmatrix} w(\mathbf{x}) d\mathbf{x} \\ = \int_{\mathcal{T}^2} \begin{bmatrix} P_1(\mathbf{x}) & P_2(\mathbf{x}) & \cdots & P_N(\mathbf{x}) \end{bmatrix} \begin{bmatrix} u_{x,1} \sum_{j=1}^N u_{x,j} P_j(\mathbf{x}) \\ u_{x,2} \sum_{j=1}^N u_{x,j} P_j(\mathbf{x}) \\ \vdots \\ u_{x,N} \sum_{j=1}^N u_{x,j} P_j(\mathbf{x}) \end{bmatrix} w(\mathbf{x}) d\mathbf{x} \\ = \int_{\mathcal{T}^2} w(\mathbf{x}) \sum_{k=1}^N P_k(\mathbf{x}) u_{x,k} \sum_{j=1}^N u_{x,j} P_j(\mathbf{x}) d\mathbf{x} = \int_{\mathcal{T}^2} w(\mathbf{x}) \sum_{k=1}^N \sum_{j=1}^N u_{x,k} u_{x,j} P_k(\mathbf{x}) P_j(\mathbf{x}) d\mathbf{x} \\ = \int_{\mathcal{T}^2} \sum_{k=1}^N u_{x,k}^2 P_k^2(\mathbf{x}) w(\mathbf{x}) d\mathbf{x} = \sum_{k=1}^N u_{x,k}^2 \int_{\mathcal{T}^2} P_k^2(\mathbf{x}) dw(\mathbf{x}) \quad (\text{now we add back in } y \text{ part}) \\ = \sum_{k=1}^N u_{x,k}^2 + \sum_{k=1}^N u_{y,k}^2 \\ = \mathbf{u}^T (\tilde{\mathcal{D}}_x^T \tilde{\mathcal{D}}_x + \tilde{\mathcal{D}}_y^T \tilde{\mathcal{D}}_y) \mathbf{u} = \tilde{\mathbf{c}}_{\frac{1}{f}}^T \tilde{\mathbf{c}}_{\frac{1}{f}}, \end{aligned}$$

where we've used the mutual orthonormality relations, which persist even within a subspace of homogeneous polynomials of degree m , to eliminate all off-diagonal terms in the sum. Note, not all orthonormal bases are created equal, and we normally would have to retain many outer product terms in the sum because orthonormality between vectors of the same subspace is not a given.

Thus, we want to solve the following quadratic program with equality constraints:

$$\begin{aligned} (1.11) \quad \min_{\mathbf{u} \in \mathbb{R}^N} F(\mathbf{u}) &= \frac{1}{2} (\mathbf{u}^T G \mathbf{u} - \tilde{\mathbf{c}}_{\frac{1}{f}}^T \tilde{\mathbf{c}}_{\frac{1}{f}}) \\ \text{subject to} \quad \mathbf{u} \cdot \mathbf{P}(\mathbf{x}) &= 0, \quad \mathbf{x} \in \partial\Omega, \end{aligned}$$

where $G = \tilde{D}_x^T \tilde{D}_x + \tilde{D}_y^T \tilde{D}_y \in \mathbb{R}^{N \times N}$ is symmetric and positive semi-definite ¹. The gradient and Hessian of F are given by

$$(1.12) \quad \nabla_{\mathbf{u}} F(\mathbf{u}) = G\mathbf{u},$$

$$(1.13) \quad \nabla_{\mathbf{u}}^2 F(\mathbf{u}) = G,$$

and we still need to consider the constraint equation. However, by making use of a weighted basis $\tilde{\mathbf{P}}(\mathbf{x})$, and expressing our solution coefficients and differential operators in terms of this weighted basis, we can eliminate the constraint equation and solve a completely unconstrained quadratic optimization problem This doesn't seem to work as well for the non-linear problem, but we will describe that approach and verify correctness in the next section anyway.

1.1.1. The weighted Jacobi basis.

$$(1.14) \quad \tilde{\mathbf{P}}^{(a,b,c)}(x, y) = x^{a-\frac{1}{2}} y^{b-\frac{1}{2}} (1-x-y)^{c-\frac{1}{2}} \mathbf{P}^{(a,b,c)}$$

will prove useful for incorporating boundary conditions to the discrete operator, while preserving its sparsity. Note, we are simply multiplying the original basis by the weight function $W^{(a,b,c)}(\mathbf{x})$ which forms the measure under which they are orthogonal. We need the following weighted ladder operators for differentiation in the weighted basis:

$$(1.15) \quad \begin{aligned} -(2k+b+c+1)h_{k,n}^{(a,b,c)} \partial_x \tilde{\mathbf{P}}_{k,n}^{(a,b,c)} &= (k+c)(n-k+1)h_{k,n+1}^{(a-1,b,c-1)} \tilde{\mathbf{P}}_{k,n+1}^{(a-1,b,c-1)} \\ &\quad + (k+1)(n-k+a)h_{k+1,n+1}^{(a-1,b,c-1)} \tilde{\mathbf{P}}_{k+1,n+1}^{(a-1,b,c-1)} \end{aligned}$$

$$(1.16) \quad h_{k,n}^{(a,b,c)} \partial_y \tilde{\mathbf{P}}_{k,n}^{(a,b,c)} = -(k+1)h_{k+1,n+1}^{(a,b-1,c-1)} \tilde{\mathbf{P}}_{k+1,n+1}^{(a,b-1,c-1)}.$$

If $\mathbf{f}$ are the coefficients of f under the (a,b,c) basis, then we have sparse banded matrices $\mathcal{W}_{x,(a,b,c)}^{(a-1,b,c-1)}, \mathcal{W}_{y,(a,b,c)}^{(a,b-1,c-1)}$ such that

$$(1.17) \quad \begin{aligned} \partial_x [x^{a-\frac{1}{2}} y^{b-\frac{1}{2}} (1-x-y)^{c-\frac{1}{2}} f(x, y)] &= \partial_x (\tilde{\mathbf{P}}^{(a,b,c)}(x, y)^T \mathbf{f}) \\ &= \tilde{\mathbf{P}}^{(a-1,b,c-1)}(x, y)^T \mathcal{W}_{x,(a,b,c)}^{(a-1,b,c-1)} \mathbf{f}, \end{aligned}$$

$$(1.18) \quad \begin{aligned} \partial_y [x^{a-\frac{1}{2}} y^{b-\frac{1}{2}} (1-x-y)^{c-\frac{1}{2}} f(x, y)] &= \partial_y (\tilde{\mathbf{P}}^{(a,b,c)}(x, y)^T \mathbf{f}) \\ &= \tilde{\mathbf{P}}^{(a,b-1,c-1)}(x, y)^T \mathcal{W}_{y,(a,b,c)}^{(a,b-1,c-1)} \mathbf{f} \end{aligned}$$

1.1.2. Lowering operators. Additionally, we have lowering operators which relate $xP_{k,n}^{(a,b,c)}$ to $P_{k,n}^{(a-1,b,c)}$ and $yP_{k,n}^{(a,b,c)}$ to $P_{k,n}^{(a,b-1,c)}$. These are given by the following recurrences:

$$(1.19)$$

$$(1.20) \quad \begin{aligned} (2n+a+b+c+2)h_{k,n}^{(a,b,c)} xP_{k,n}^{(a,b,c)} &= (n-k+a)h_{k,n}^{(a-1,b,c)} P_{k,n}^{(a-1,b,c)} \\ &\quad + (n-k+1)h_{k,n+1}^{(a-1,b,c)} P_{k,n+1}^{(a-1,b,c)}, \\ (2k+b+c+1)(2n+a+b+c+2)h_{k,n}^{(a,b,c)} yP_{k,n}^{(a,b,c)} &= (k+b)(n+k+b+c+1)h_{k,n}^{(a,b-1,c)} P_{k,n}^{(a,b-1,c)} \\ &\quad - (k+1)(n-k+a)h_{k+1,n}^{(a,b-1,c)} P_{k+1,n}^{(a,b-1,c)} \\ &\quad - (k+b)(n-k+1)h_{k,n+1}^{(a,b-1,c)} P_{k,n+1}^{(a,b-1,c)} \\ &\quad + (k+1)(n+k+a+b+c+2)h_{k+1,n+1}^{(a,b-1,c)} P_{k+1,n+1}^{(a,b-1,c)}. \end{aligned}$$

¹Note, $\forall \mathbf{x} \in \mathbb{R}_+^{n+1}, \mathbf{x}^T G \mathbf{x} = \mathbf{x}^T \tilde{D}_x^T \tilde{D}_x \mathbf{x} + \mathbf{x}^T \tilde{D}_y^T \tilde{D}_y \mathbf{x} = \|\tilde{D}_x \mathbf{x}\|^2 + \|\tilde{D}_y \mathbf{x}\|^2 \geq 0$. We have equality only for any $\mathbf{x} = \alpha \mathbf{e}_1$, as derivatives of the constant polynomial are 0.

So, there are sparse banded matrices $\mathcal{L}_{(a,b,c)}^{(a-1,b,c)}$ and $\mathcal{L}_{(a,b,c)}^{(a,b-1,c)}$ corresponding to multiplication by x and y , respectively, such that

$$(1.22) \quad xf(x, y) = x\mathbf{P}^{(a,b,c)}(x, y)^T \mathbf{f} = \mathbf{P}^{(a-1,b,c)}(x, y)^T \mathcal{L}_{(a,b,c)}^{(a-1,b,c)} \mathbf{f},$$

$$(1.23) \quad yf(x, y) = y\mathbf{P}^{(a,b,c)}(x, y)^T \mathbf{f} = \mathbf{P}^{(a,b-1,c)}(x, y)^T \mathcal{L}_{(a,b,c)}^{(a,b-1,c)} \mathbf{f}.$$

We will construct the modal Eikonal operator which operates on coefficients under the basis $\tilde{\mathbf{P}}^{(3/2,3/2,3/2)}$ and produces coefficients under $\mathbf{P}^{(3/2,3/2,3/2)}$. Note, the conversion from the weighted to regular Jacobi basis depends on our choice of parameters $a = b = c = 3/2$. With $\mathbf{f}$ the coefficients of $f(x, y)$ in the (a, b, c) basis,

$$\begin{aligned} \partial_x[W^{(a,b,c)}(x, y)f(x, y)] &= \tilde{\mathbf{P}}^{(a-1,b,c-1)}(x, y)^T \mathcal{W}_{x,(a,b,c)}^{(a-1,b,c-1)} \mathbf{f} \\ &= x^{a-\frac{3}{2}} y^{b-\frac{3}{2}} (1-x-y)^{c-\frac{3}{2}} (y\mathbf{P}^{(a-1,b,c-1)}(x, y))^T (\mathcal{W}_{x,(a,b,c)}^{(a-1,b,c-1)} \mathbf{f}) \\ &= x^{a-\frac{3}{2}} y^{b-\frac{3}{2}} (1-x-y)^{c-\frac{3}{2}} \mathbf{P}^{(a-1,b-1,c-1)}(x, y)^T (\mathcal{L}_{(a-1,b,c-1)}^{(a-1,b-1,c-1)} \mathcal{W}_{x,(a,b,c)}^{(a-1,b,c-1)} \mathbf{f}), \\ \partial_y[W^{(a,b,c)}(x, y)f(x, y)] &= \tilde{\mathbf{P}}^{(a,b-1,c-1)}(x, y)^T \mathcal{W}_{y,(a,b,c)}^{(a,b-1,c-1)} \mathbf{f} \\ &= x^{a-\frac{3}{2}} y^{b-\frac{3}{2}} (1-x-y)^{c-\frac{3}{2}} (x\mathbf{P}^{(a,b-1,c-1)}(x, y))^T (\mathcal{W}_{y,(a,b,c)}^{(a,b-1,c-1)} \mathbf{f}) \\ &= x^{a-\frac{3}{2}} y^{b-\frac{3}{2}} (1-x-y)^{c-\frac{3}{2}} \mathbf{P}^{(a-1,b-1,c-1)}(x, y)^T (\mathcal{L}_{(a,b-1,c-1)}^{(a-1,b-1,c-1)} \mathcal{W}_{y,(a,b,c)}^{(a,b-1,c-1)} \mathbf{f}) \blacksquare \end{aligned}$$

When $a = b = c = 3/2$, we have

$$\begin{aligned} \partial_x[W^{(a,b,c)}(x, y)f(x, y)] &= \mathbf{P}^{(a-1,b-1,c-1)}(x, y)^T (\mathcal{L}_{(a-1,b,c-1)}^{(a-1,b-1,c-1)} \mathcal{W}_{x,(a,b,c)}^{(a-1,b,c-1)} \mathbf{f}), \\ \partial_y[W^{(a,b,c)}(x, y)f(x, y)] &= \mathbf{P}^{(a-1,b-1,c-1)}(x, y)^T (\mathcal{L}_{(a,b-1,c-1)}^{(a-1,b-1,c-1)} \mathcal{W}_{y,(a,b,c)}^{(a,b-1,c-1)} \mathbf{f}), \end{aligned}$$

1.2. Numerical solution to the Eikonal equation. The process for solving (1.1) with zero Dirichlet conditions $q \equiv 0$ is as follows:

- Choose a high order quadrature rule $(\mathbf{x}, \mathbf{y}, \mathbf{w})$ on $\mathcal{T}^2$ and compute the Vandermonde matrix $\mathbf{P}^{(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})}(\mathbf{x}, \mathbf{y})$ up to a desired order (each column is a polynomial evaluated on abscissa), and there are $\mathcal{O}(n^2)$ columns for order n .
- Expand the right-hand side $f(x, y)$ in the basis $\mathbf{P}^{(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})}$ by computing

$$(1.24) \quad \mathbf{c}_{\frac{1}{2}} = \mathbf{P}^{(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})}(\mathbf{x}, \mathbf{y})^T (\mathbf{f}(\mathbf{x}, \mathbf{y}) \odot \mathbf{w})$$

- Assemble the operator matrix G as

$$(1.25) \quad G = \mathcal{L}_{(a-1,b,c-1)}^{(a-1,b-1,c-1)} \mathcal{W}_{x,(a,b,c)}^{(a-1,b,c-1)} + \mathcal{L}_{(a,b-1,c-1)}^{(a-1,b-1,c-1)} \mathcal{W}_{y,(a,b,c)}^{(a,b-1,c-1)}.$$

and solve for the coefficients $\mathbf{u}$ of $u(x, y)$ in the basis $(3/2, 3/2, 3/2)$ via (1.11).

- Evaluate the approximation $\hat{u}(x, y)$ using the weighted polynomials as

$$\hat{u}(\mathbf{x}, \mathbf{y}) = \tilde{\mathbf{P}}^{(\frac{3}{2}, \frac{3}{2}, \frac{3}{2})}(\mathbf{x}, \mathbf{y})^T \mathbf{u},$$

for any $(x, y) \in \mathcal{T}^2$

2. A variational approach. Let's try framing the problem from the perspective of infinite dimensional optimization. Thus far, my efforts have fallen under the discretize-then-optimize approach for solving non-linear PDEs. So, let's flip that and consider the optimize-then-discretize approach.

We want to solve

$$(2.1) \quad \begin{aligned} & \min_{u \in L_2(\mathcal{T}^2, w^{(a,b,c)})} \mathcal{L}(u), \\ & \text{s.t. } u = q \text{ on } \partial\mathcal{T}^2, \end{aligned}$$

where $\mathcal{L}$ is defined by

$$(2.2) \quad \mathcal{L}(u) := \int_{\mathcal{T}^2} (|\nabla_{\mathbf{x}} u|^2 - 1) d\mathbf{x}$$

The steps here are to derive the first variation of $\mathcal{L}$ as

$$\delta\mathcal{L}(u, \hat{u}) = \partial_\varepsilon \mathcal{L}(u + \varepsilon \hat{u})|_{\varepsilon=0} = 0,$$

where $\hat{u} \in \mathcal{H}_0^1(\mathcal{T}^2)$. This gives the necessary condition a minimizer of $\mathcal{L}$ must satisfy. Then, compute the second variation by considering another $\tilde{u} \in \mathcal{H}_0^1$, and deriving the newton step through

$$\delta^2\mathcal{L}(u, \hat{u}, \tilde{u}) = \partial_\varepsilon \delta\mathcal{L}(u + \varepsilon \tilde{u}, \hat{u})$$

and find the step direction by solving

$$\delta^2\mathcal{L}(u^k)(\hat{u}, \tilde{u}) = -\delta\mathcal{L}(u^k)(\hat{u}), \quad \forall \hat{u} \in \mathcal{H}_0^1$$

By choosing $u^0 = q$ on $\partial\mathcal{T}^2$, and setting $u^{k+1} = u^k + \tilde{u}$, we have an infinite-dimensional newton stepper for solving the Eikonal problem. I'm not sure I like where this goes in terms of complexity (mental and computational), so will stick to the DTO approach for now. There is also probably a correct energy functional which gives the Eikonal problem, which is likely not the one I started with.