

## Homework 3: Due at class on March 25

1. Let  $\varphi_t : M \rightarrow M$  be the flow generated by a vector field  $X \in \mathfrak{X}(M)$ . Then, show that for another vector field  $Y \in \mathfrak{X}(M)$

$$[X, Y] = \lim_{t \rightarrow 0} \frac{(\varphi_{-t})_* Y - Y}{t}$$

2. Given a vector field  $X \in \mathfrak{X}(M)$ , the Lie derivative

$$L_X : \Omega^k(M) \rightarrow \Omega^k(M)$$

can be defined by

$$L_X(\omega) = \lim_{t \rightarrow 0} \frac{\varphi_t^* \omega - \omega}{t}$$

where  $\varphi_t : M \rightarrow M$  be the flow as above. Show that it satisfies

$$L_X \omega(X_1, \dots, X_k) = X\omega(X_1, \dots, X_k) - \sum_{i=1}^k \omega(X_1, \dots, [X, X_i], \dots, X_k)$$

for  $X_1, \dots, X_k \in \mathfrak{X}(M)$ .

3. Define an  $n$ -form  $\omega$  on the space  $\mathbb{R}^{n+1} \setminus \{0\}$ , obtained from  $\mathbb{R}^{n+1}$  by removing the origin, by

$$\omega = \frac{1}{|x|^{n+1}} \sum_{i=0}^n (-1)^i x^i dx^0 \wedge \dots \wedge \widehat{dx^i} \wedge \dots \wedge dx^n.$$

Prove that  $d\omega = 0$ . Let  $\iota : S^n \hookrightarrow \mathbb{R}^{n+1}$  be the unit sphere in  $\mathbb{R}^{n+1}$ . Show that  $\iota^* \omega$  is the generator of  $n$ -th de Rham cohomology  $H_{dR}^n(S^n)$  of the  $n$ -sphere. In fact, the de Rham cohomology of  $S^n$  is

$$H_{dR}^k(S^n) = \begin{cases} \mathbb{R} & k = 0, n \\ 0 & \text{otherwise} \end{cases}.$$

In the case of  $n = 2$ , evaluate the integral

$$\int_{S^2} \iota^* \omega.$$

4. Let  $T^2 = S^1 \times S^1$  be the torus. Using the formula for de Rham cohomology of a product space,

$$H_{dR}^k(M \times N) \cong \bigoplus_{k=p+q} H_{dR}^p(M) \otimes H_{dR}^q(N),$$

find de Rham cohomology  $H_{dR}^*(T^2)$ . Find the Poincaré dual of each generator of  $H_{dR}^*(T^2)$ .

5. The Maxwell equations are written as

$$\nabla \cdot \mathbf{E} = \rho, \quad \nabla \times \mathbf{B} = \mathbf{J} + \frac{\partial \mathbf{E}}{\partial t},$$

$$\nabla \cdot \mathbf{B} = 0, \quad \nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}.$$

Let us write the gauge potential

$$A = A_\mu dx^\mu = \phi dt + A_1 dx^1 + A_2 dx^2 + A_3 dx^3$$

and the current

$$J = J_\mu dx^\mu = \rho dt + J_1 dx^1 + J_2 dx^2 + J_3 dx^3.$$

Then, the field strength can be written as  $F = dA$ . Show that the Maxwell equations are equivalent to the following equations

$$dF = 0, \quad \delta F = -j.$$

Find the equation of motion for the following action

$$S = -\frac{1}{2} \int F \wedge *F - \int A \wedge *J.$$

Discuss this for both a positive definite metric and a Lorentzian signature metrics.

6. (This is a bonus problem with extra 3 points which is NOT mandatory.) Find the preimage of a smooth map  $f : \text{SO}(3) \rightarrow \text{SO}(3); g \mapsto g^2$  and the rank of  $f_*$  at every point of  $\text{SO}(3)$ . [Hint: An element  $g \neq 1 \in \text{SO}(3)$  is an rotation along a certain axis in  $\mathbb{R}^3$ .]