

Homework 12: Due at class on June 3

1 Chern classes of $\mathbb{C}P^n$

An arbitrary point $\ell = [z_0; \cdots; z_n] \in \mathbb{C}P^n$ is a complex line \mathbb{C} through the origin of \mathbb{C}^{n+1} . So we can construct a complex line bundle L over the complex projective space $\mathbb{C}P^n$ as

$$\begin{aligned} L &= \{(\ell, w) \in \mathbb{C}P^n \times \mathbb{C}^{n+1} \mid w \in \ell\}, \\ &= \{([z_0; \cdots; z_n], w) \in \mathbb{C}P^n \times \mathbb{C}^{n+1} \mid w = \lambda \cdot (z_0, \cdots, z_n), \lambda \in \mathbb{C}\}. \end{aligned} \quad (1.1)$$

(The notation is the same as in Problem 4 of Homework 10.) This is called the **Hopf line bundle**. If we consider points with unit length in each fiber of the Hopf line bundle, then we obtain a sphere

$$S^{2n+1} = \{(\ell, z) \in \mathbb{C}P^n \times \mathbb{C}^{n+1} \mid z \in \ell, |z| = 1\}.$$

Therefore, the Hopf line bundle is the associated line bundle of the Hopf S^1 -bundle.

In fact, the Whitney sum of the tangent bundle $T\mathbb{C}P^n$ and the trivial bundle ϵ is the Whitney sum of $(n+1)$ copies of L^*

$$T\mathbb{C}P^n \oplus \epsilon \cong L^* \oplus \cdots \oplus L^*.$$

where L^* is the dual line bundle of L . (See Morita Proposition 5.13 for the derivation.)

Let us first consider the case of $n = 1$. We know that cohomology $H^2(\mathbb{C}P^1; \mathbb{Z}) \cong \mathbb{Z}$ and its generator x is the volume form. Show that $c_1(L) = -x$ and $c_1(T\mathbb{C}P^1) = 2x$. In addition, find the curvature of the Ehresmann connection

$$A = i \operatorname{Im}(\bar{z}_0 dz_0 + \bar{z}_1 dz_1)$$

given in Problem 4 of Homework 10, and compare it with the volume form. Actually, show that over $\mathbb{C}P^1$ one can find an infinite family of non-equivalent complex line bundles.

Find the total Chern class of $T\mathbb{C}P^n$. In fact, the $2i$ -th de Rham cohomology group of $\mathbb{C}P^n$ is isomorphic to \mathbb{R} generated by $x^i = x \wedge \cdots \wedge x$:

$$H_{dR}^{2i}(\mathbb{C}P^n) \cong \mathbb{R}\langle x^i \rangle \quad \text{for } i \leq n.$$

Moreover, the de Rham cohomology groups of $\mathbb{C}P^n$ has a ring structure where the multiplication is given by the wedge product:

$$H_{dR}^*(\mathbb{C}P^n) \cong \mathbb{R}[x]/(x^{n+1}).$$

2 Chern-Simons and Wess-Zumino-Witten

Let us define the Chern-Simons action over a 3-manifold M

$$S_{CS} = \frac{k}{4\pi} \int_M \operatorname{Tr} \left(A \wedge dA + \frac{2}{3} A \wedge A \wedge A \right).$$

Under the gauge transformation

$$A \rightarrow g^{-1}dg + g^{-1}Ag ,$$

show that the action is transformed as

$$S_{CS} \rightarrow S_{CS} + \frac{k}{4\pi} \int_M d \operatorname{Tr} \left(g^{-1}Ag \wedge g^{-1}dg \right) - \frac{k}{12\pi} \int_M \operatorname{Tr} \left((g^{-1}dg)^3 \right) .$$

Let us consider the case $M = S^3$ and $G = \mathrm{SU}(2)$. Then, the second term vanishes because S^3 is closed. Show that the third term is

$$\frac{k}{12\pi} \int_M \operatorname{Tr} \left((g^{-1}dg)^3 \right) = 2\pi k \deg(g) ,$$

where $\deg(g)$ is the mapping degree of

$$g : S^3 \rightarrow \mathrm{SU}(2) \cong S^3 .$$

(Hint: $\mu = g^{-1}dg$ is the Maurer-Cartan form and $\frac{1}{12} \operatorname{Tr}(\mu \wedge \mu \wedge \mu)$ is the left-invariant volume form of $\mathrm{SU}(2)$.) Argue that the level k must be an integer in order for the partition function to be well-defined.

3 Chern-Simons from 4-dimension

Show the following identity

$$\operatorname{Tr}(F \wedge F) = d \operatorname{Tr} \left(A \wedge dA + \frac{2}{3} A \wedge A \wedge A \right) .$$

Therefore, if there exists a four-manifold B whose boundary is $\partial B = M$, then we can indeed define Chern-Simons action by

$$S_{CS} = \frac{k}{4\pi} \int_B \operatorname{Tr}(F \wedge F) ,$$

when G is compact and semi-simple.