

# Homework 4: Due at class on Oct 19

## 1 Modular invariant

Show that (4.98) is modular  $S$ -invariant by using the Poisson resummation formula (4.99).

## 2 Modular transformation

The modular group is defined by

$$\mathrm{SL}(2, \mathbb{Z}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a, b, c, d \in \mathbb{Z}, \quad ad - bc = 1 \right\} .$$

Show that

$$\mathrm{Im}(\gamma \cdot \tau) = \frac{\mathrm{Im} \tau}{|c\tau + d|^2} ,$$

where


$$\gamma : \tau \mapsto \gamma \cdot \tau = \frac{a\tau + b}{c\tau + d} .$$

Therefore, if  $\mathrm{Im} \tau > 0$ , then  $\mathrm{Im}(\gamma \cdot \tau) > 0$  so that the upper half-plane

$$\mathbb{H} = \{x + iy \mid y > 0; x, y \in \mathbb{R}\}$$

receives the action of the modular group. Express elements of modular group that maps the gray region to blue, red and green region in terms of modular  $S$  and  $T$  transformations

$$T : \tau \rightarrow \tau + 1 , \quad S : \tau \rightarrow -1/\tau .$$



### 3 Boson-Fermion correspondence

We have learnt that the OPE of the free boson is

$$\varphi(z)\varphi(0) \sim -\ln z .$$

Let us also consider two Majorana-Weyl fermions  $\psi^1, \psi^2$  with OPE

$$\psi^i(z)\psi^j(0) \sim \frac{\delta^{ij}}{z} .$$

We can define the complex fermion

$$\psi(z) = 2^{-1/2}(\psi^1(z) + i\psi^2(z)), \quad \bar{\psi}(z) = 2^{-1/2}(\psi^1(z) - i\psi^2(z)) .$$

Show the equivalence of operators in boson and fermion

$$:e^{i\varphi}: \cong \psi, \quad :e^{-i\varphi}: \cong \bar{\psi}, \quad i\partial\varphi \cong \psi\bar{\psi}, \quad T_\varphi \cong T_\psi ,$$

by calculating the OPEs of operators in both theories and comparing. Note that the energy-momentum tensor of the complex fermion is

$$T_\psi := -\frac{1}{2} : \psi \partial \bar{\psi} : - \frac{1}{2} : \bar{\psi} \partial \psi : .$$