

IHES - June 2026

Mysterious relations between geometry of Hitchin moduli spaces and VOA modules from 4d $N=2$ theories

Based on joint work with Gukov-Koroteev-Pei-Saberi [\[arXiv:2206.03565\]](#)
& Junkang Huang-Yutai Zhang-Shutong Zhuang [\[arXiv:2412.19647\]](#)

Related works by Peng Shan-Wenbin Yan-Dan Xie, Wenbin Pan-Yiwen Yan

Satoshi Nawata Fudan University


Motivation

4d N=2 Class S theory $\mathcal{T}$ associated to Hitchin system

Higgs branch

assoc. var.

VOA $\mathbb{V}(\mathcal{T})$



chiral
quantization

Hyper-Kähler manifolds



Relation ???

Coulomb branch

Hitchin moduli space


def'n
quantization

Skein algebras
Spherical DAHA



Class S theory

6d (2,0) SCFT of type $\mathfrak{g}$ ADE



$$\mathbb{R}^3 \times S^1 \times C_{g,n}$$

Gaiotto

Gaiotto-Moore-Neitzke

4d N=2 Class S theory $\mathcal{T}_{g,n}$ associated to Hitchin system


- Regular class S theory ($g \geq 0, n \geq 0$ s.t. $2g - 2 + n > 0$)



$$\phi|_{z_i} \sim \frac{f_i^\vee}{z - z_i} + \dots, \quad f_i \in \mathcal{O}_{f_i}$$

Higgs branch = Star-shaped quiver variety

- Argyres-Douglas type (a sphere with one irregular and one regular puncture)



$$\phi|_0 \sim \frac{f^\vee}{z} + \dots,$$

$$\phi|_\infty \sim \frac{s}{w^{2+\frac{u}{m}}} + \dots, \quad s: \text{ semisimple regular}$$

Dan Xie

SECTION 1

4-punctured sphere and brane quantization

4-punctured sphere and Betti moduli space

- Moduli space of flat $\text{SL}(2, \mathbb{C})$ -connections on a four-punctured sphere $C_{0,4}$

$$\mathfrak{X} = \mathcal{M}_{\text{flat}}(C_{0,4}, \text{SL}(2, \mathbb{C})) = \langle M_1, M_2, M_3, M_4 \in \text{SL}(2, \mathbb{C}) \mid M_1 M_2 M_3 M_4 = \text{Id} \rangle / \text{SL}(2, \mathbb{C}) ,$$

- These variables are subject to the **cubic equation**

$$f(x, y, z) = -xyz + x^2 + y^2 + z^2 + \theta_1 x + \theta_2 y + \theta_3 z + \theta_4 = 0 ,$$


where, using the notation $\bar{f} \equiv f + f^{-1}$, θ_i are expressed by

$$\theta_1 = \chi_{D_4}(\mathbf{8}_V) = \bar{t}_1 \bar{t}_2 + \bar{t}_3 \bar{t}_4 ,$$

$$\theta_2 = \chi_{D_4}(\mathbf{8}_S) = \bar{t}_1 \bar{t}_3 + \bar{t}_2 \bar{t}_4 ,$$

$$\theta_3 = \chi_{D_4}(\mathbf{8}_C) = \bar{t}_1 \bar{t}_4 + \bar{t}_2 \bar{t}_3 ,$$

$$\theta_4 = \chi_{D_4}(\mathbf{adj}) = \bar{t}_1^2 + \bar{t}_2^2 + \bar{t}_3^2 + \bar{t}_4^2 + \bar{t}_1 \bar{t}_2 \bar{t}_3 \bar{t}_4 - 4 .$$



- This space is the Coulomb branch of 4d $\mathcal{N} = 2$ $\text{SU}(2)$ $N_f = 4$ Seiberg-Witten theory on $S^1 \times \mathbb{R}^3$
- Flavor symmetry is $\text{SO}(8)$

Dolbeault moduli space

- Moduli space of parabolic $SU(2)$ -Higgs bundles on a four-punctured sphere $C_{0,4}$

$$\mathcal{M}_H(C_{0,4}, SU(2)) = \{(E, \varphi) \mid E \text{ holom. bdle}, \varphi \in H^0(C_{0,4}, \text{ad}(E) \otimes K)\}_{\text{stab. cond.}},$$

- Tame ramifications $(\alpha_i, \beta_i, \gamma_i) \in (\mathfrak{t}_{A_1})^3$ at punctures $i = (1, 2, 3, 4)$

$$A = \alpha_j d\vartheta + \dots$$

$$\varphi = \frac{1}{2}(\beta_j + i\gamma_j) \frac{dz}{z} + \dots$$



- Ramification data are enhanced to the Cartan subalgebra of D_4

$$\prod_{j=1}^4 \mathfrak{t}_j^{(\alpha)} \times \mathfrak{t}_j^{(\beta)} \times \mathfrak{t}_j^{(\gamma)} \cong \mathfrak{t}_{D_4}^{(\alpha)} \times \mathfrak{t}_{D_4}^{(\beta)} \times \mathfrak{t}_{D_4}^{(\gamma)}$$

- Non-Abelian Hodge correspondence: $\mathfrak{X} = \mathcal{M}_{\text{flat}}(C_{0,4}, SL(2, \mathbb{C})) \cong \mathcal{M}_H(C_{0,4}, SU(2))$

- Complex structures as a hyper-Kähler manifold

$$t_j = \exp(-2\pi(\gamma_j + i\alpha_j)), \quad (j = 1, 2, 3, 4)$$




Hitchin fibration

- In **complex structure** I , the **Hitchin moduli space** admits the **Hitchin fibration** over $\mathcal{B}_H$:

$$h : \mathcal{M}_H(C_{0,4}, \mathrm{SU}(2)) \rightarrow \mathcal{B}_H = H^0(C_{0,4}, K_C^{\otimes 2}) .$$


$$(E, \varphi) \mapsto \mathrm{Tr}(\varphi^2)$$

- In our case, the **Hitchin fibration** is an **elliptic fibration** with a **singular fiber**.
- When $\beta_i = 0 = \gamma_i$, the **singular fiber** is located at the **origin**.
- **fiber class relation**: $[\mathbf{F}] = 2[\mathbf{D}_4] + \sum_{i=0}^3 [\mathbf{D}_i]$



$$\dot{W}(D_4) \subset H_2(\mathfrak{X}, \mathbb{Z}) \cong \dot{Q}(D_4) .$$

Afine D_4 Dynkin diagram



Deformation quantization and spherical DAHA

- Deformation quantization of coordinate ring $\mathcal{O}(\mathfrak{X})$ with respect to Ω_J

Oblomkov

$$\mathcal{O}(\mathfrak{X}) \rightsquigarrow \mathcal{O}^q(\mathfrak{X}) \cong S\ddot{H}_{q,t} , \quad [\widehat{f}, \widehat{g}] = -i\hbar \widehat{\{f, g\}} .$$

- It is spherical double affine Hecke algebra $S\ddot{H}_{q,t}$ of type $C^\vee C_1$:

$$\begin{aligned} [x, y]_q &= (q - q^{-1})z + (q^{\frac{1}{2}} - q^{-\frac{1}{2}})\theta_3 , \\ [y, z]_q &= (q - q^{-1})x + (q^{\frac{1}{2}} - q^{-\frac{1}{2}})\theta_1 , \\ [z, x]_q &= (q - q^{-1})y + (q^{\frac{1}{2}} - q^{-\frac{1}{2}})\theta_2 , \\ -q^{\frac{1}{2}}xyz + qx^2 + q^{-1}y^2 + qz^2 + q^{\frac{1}{2}}\theta_1x + q^{-\frac{1}{2}}\theta_2y + q^{\frac{1}{2}}\theta_3z + \theta_4(q) &= 0 , \end{aligned}$$

where, using the notation $\overline{f} \equiv f + f^{-1}$, θ_i are expressed by

$$\begin{aligned} \theta_1 &= \chi_{D_4}(\mathbf{8}_V) = \overline{t_1} \overline{t_2} + \overline{t_3} \overline{t_4} , \\ \theta_2 &= \chi_{D_4}(\mathbf{8}_S) = \overline{t_1} \overline{t_3} + \overline{t_2} \overline{t_4} , \\ \theta_3 &= \chi_{D_4}(\mathbf{8}_C) = \overline{t_1} \overline{t_4} + \overline{t_2} \overline{t_3} , \\ \theta_4(q) &= \overline{t_1}^2 + \overline{t_2}^2 + \overline{t_3}^2 + \overline{t_4}^2 + \overline{t_1} \overline{t_2} \overline{t_3} \overline{t_4} - q - q^{-1} - 2 . \end{aligned}$$

A-model and Brane Quantization

- Brane quantization is geometric approach to representation theory of $\mathcal{O}^q(\mathfrak{X})$ for affine hyper-Kähler manifold $\mathfrak{X}$
- Consider A-model $\Sigma \hookrightarrow (\mathfrak{X}, \omega_{\mathfrak{X}})$
- The distinguished “big” A-brane called “canonical coisotropic brane” $\mathfrak{B}_{cc}$

Gukov-Witten

Kontsevich-Soibelman

$$\mathfrak{B}_{cc} : \begin{array}{c} \mathcal{L} \\ \downarrow \\ \mathfrak{X} \end{array} \quad c_1(\mathcal{L}) = [F/2\pi] \in H^2(\mathfrak{X}, \mathbb{Z}) .$$

where $q = \exp(2\pi i \hbar)$ and

$$\frac{\Omega_J}{2\pi i \hbar} = F + B + i\omega_{\mathfrak{X}}$$

- M is a Lagrangian $\omega_{\mathfrak{X}}|_M = 0$ with respect to $\omega_{\mathfrak{X}}$, so can consider an ordinary A-brane on $\mathfrak{X}$

$$\mathfrak{B}_M : \begin{array}{c} \mathcal{L}' \\ \downarrow \\ M \end{array} \quad \text{local system}$$

Derived equivalence from brane quantization



- Joining of strings provides **representation theory** of $\mathcal{O}^q(\mathfrak{X})$ via **A-model** on $\mathfrak{X}$:

$$\begin{aligned}\mathcal{O}^q(\mathfrak{X}) &= \mathrm{Hom}(\mathfrak{B}_{cc}, \mathfrak{B}_{cc}) \\ \bigcirc_Q V_M &= \mathrm{Hom}(\mathfrak{B}_M, \mathfrak{B}_{cc}) .\end{aligned}$$

- This functor is expected to provide derived equivalence

$$\begin{aligned}\mathrm{RHom}(-, \mathfrak{B}_{cc}) : D^b A\text{-Brane}(\mathfrak{X}, \omega_{\mathfrak{X}}) &\rightarrow D^b \mathrm{Rep}(S\ddot{H}), \\ \mathfrak{B}_M &\mapsto \mathcal{H} = \mathrm{Hom}(\mathfrak{B}_M, \mathfrak{B}_{cc}) .\end{aligned}$$

Geometric category $\longleftrightarrow$ Representation category




compact A-branes $\longleftrightarrow$ finite-dimensional $\mathcal{O}^q(\mathfrak{X})$ -modules
 non-compact A-branes $\longleftrightarrow$ infinite-dimensional $\mathcal{O}^q(\mathfrak{X})$ -modules

(A,B,A)-branes and polynomial representation

Claim. The (A, B, A) -branes supported on $27 - 3 = 24$ lines in the affine cubic surface $\mathfrak{X}$ correspond to the polynomial representation of $S\ddot{H}_{q,t}$ and its images under certain symmetries.

The polynomial representation acts on the ring of symmetric functions $\mathcal{P} = \mathbb{C}_{q,t}[X^\pm]^{\mathbb{Z}_2}$:

$$\begin{aligned} \text{pol} : S\ddot{H}_{q,t} &\rightarrow \text{End}(\mathcal{P}) , \\ x &\mapsto X + X^{-1} , \\ y &\mapsto A(X; \mathbf{t})(\varpi - 1) + A(X^{-1}; \mathbf{t})(\varpi^{-1} - 1) - q^{\frac{1}{2}} t_1 t_3 - q^{-\frac{1}{2}} (t_1 t_3)^{-1} , \\ z &\mapsto q^{\frac{1}{2}} [X A(X) \varpi + X^{-1} A(X^{-1}) \varpi^{-1}] \\ &\quad - \frac{X + X^{-1}}{q^{\frac{1}{2}} + q^{-\frac{1}{2}}} \left[A(X) + A(X^{-1}) + q^{\frac{1}{2}} t_1 t_3 + q^{-\frac{1}{2}} t_1^{-1} t_3^{-1} \right] - \frac{\theta_3}{q^{\frac{1}{2}} + q^{-\frac{1}{2}}} . \end{aligned}$$



Askey-Wilson polynomial is the eigenfunction of y under polynomial representation

$$\text{pol}(y) \cdot P_n(x; q, \mathbf{t}) = (-q^{n+\frac{1}{2}} t_1 t_3 - q^{-n-\frac{1}{2}} t_1^{-1} t_3^{-1}) P_n(x; q, \mathbf{t})$$

This corresponds to an (A, B, A) -branes supported on

$$\mathbf{P}_{(1,0,1,0)} = \{y = -t_1 t_3 - t_1^{-1} t_3^{-1}, z = -t_1 t_3 x - t_1 t_4 - t_1 t_4^{-1} - t_2 t_3 - t_2^{-1} t_3\}$$

Compact A-branes and finite-dimensional modules


Irreducible components at a Hitchin fiber are (B, A, A) -branes which is Lagrangian when $\hbar \in \mathbb{R}$

$$\ell_i = \dim \mathcal{D}_i = \int_{\mathbf{D}_i} \frac{\Omega_J}{2\pi i \hbar} = \frac{\alpha \cdot e^i}{\hbar} \, .$$

Claim. The functor $\mathrm{Hom}(-, \mathfrak{B}_{cc})$ restricts to a derived equivalence of the full subcategory of compact Lagrangian A -branes of $\mathfrak{X}$ and the category of finite-dimensional $S\ddot{H}_{q,t}$ -modules.

finite-dim rep	shortening condition	A -brane	A -brane condition
$\mathcal{F}_m^{(x_m)}$	$q^m = 1$	$\mathfrak{B}_{\mathbf{F}}^{(x_m)}$	$m = \frac{1}{\hbar}$
$\mathcal{V}_k$	$q^k = t_1^{-2}$	$\mathfrak{B}_{\mathbf{V}}$	$k = \frac{2\alpha_1}{\hbar}$
$\mathcal{D}_{\ell_1}^{(1)}$	$q^{\ell_1} = t_1 t_2 t_3 t_4$	$\mathfrak{B}_{\mathbf{D}_1}$	$\ell_1 = \frac{1}{\hbar} - \frac{\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4}{\hbar}$
$\mathcal{D}_{\ell_2}^{(2)}$	$q^{\ell_2} = t_1 t_2 t_3^{-1} t_4^{-1}$	$\mathfrak{B}_{\mathbf{D}_2}$	$\ell_2 = \frac{-\alpha_1 - \alpha_2 + \alpha_3 + \alpha_4}{\hbar}$
$\mathcal{D}_{\ell_3}^{(3)}$	$q^{\ell_3} = t_1 t_2^{-1} t_3 t_4^{-1}$	$\mathfrak{B}_{\mathbf{D}_3}$	$\ell_3 = \frac{-\alpha_1 + \alpha_2 - \alpha_3 + \alpha_4}{\hbar}$
$\mathcal{D}_{\ell_4}^{(4)}$	$q^{\ell_4} = t_1 t_2^{-1} t_3^{-1} t_4$	$\mathfrak{B}_{\mathbf{D}_4}$	$\ell_4 = \frac{-\alpha_1 + \alpha_2 + \alpha_3 - \alpha_4}{\hbar}$

Compact A-branes and finite-dimensional modules




Claim. The functor $\mathrm{Hom}(-, \mathfrak{B}_{cc})$ restricts to a **derived equivalence** of the full subcategory of **compact Lagrangian A-branes** of $\mathfrak{X}$ and the category of **finite-dimensional $S\ddot{H}_{q,t}$ -modules**.

finite-dim rep	shortening condition	A-brane	A-brane condition
$\mathcal{F}_m^{(x_m)}$	$q^m = 1$	$\mathfrak{B}_{\mathbf{F}}^{(x_m)}$	$m = \frac{1}{\hbar}$
$\mathcal{V}_k$	$q^k = t_1^{-2}$	$\mathfrak{B}_{\mathbf{V}}$	$k = \frac{2\alpha_1}{\hbar}$
$\mathcal{D}_{\ell_1}^{(1)}$	$q^{\ell_1} = t_1 t_2 t_3 t_4$	$\mathfrak{B}_{\mathbf{D}_1}$	$\ell_1 = \frac{1}{\hbar} - \frac{\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4}{\hbar}$
$\mathcal{D}_{\ell_2}^{(2)}$	$q^{\ell_2} = t_1 t_2 t_3^{-1} t_4^{-1}$	$\mathfrak{B}_{\mathbf{D}_2}$	$\ell_2 = \frac{-\alpha_1 - \alpha_2 + \alpha_3 + \alpha_4}{\hbar}$
$\mathcal{D}_{\ell_3}^{(3)}$	$q^{\ell_3} = t_1 t_2^{-1} t_3 t_4^{-1}$	$\mathfrak{B}_{\mathbf{D}_3}$	$\ell_3 = \frac{-\alpha_1 + \alpha_2 - \alpha_3 + \alpha_4}{\hbar}$
$\mathcal{D}_{\ell_4}^{(4)}$	$q^{\ell_4} = t_1 t_2^{-1} t_3^{-1} t_4$	$\mathfrak{B}_{\mathbf{D}_4}$	$\ell_4 = \frac{-\alpha_1 + \alpha_2 + \alpha_3 - \alpha_4}{\hbar}$

Affine braid group action on categories

- The A -model depends only on complexified Kähler moduli, but is independent of complex structure.
- If $\hbar \in \mathbb{R}$, then (α, β) is the complex structure, while $\gamma + i\eta$ is the complexified Kähler moduli.
- Variations along a loop in the complex structure moduli space (avoiding singularities) induce non-trivial transformations of the A -brane category.




P. Seidel

- When $\gamma + i\eta = 0$, the affine braid group $\dot{\text{Br}}(D_4)$ acts on the category:

$$\begin{array}{ccc} & \dot{\text{Br}}(D_4) & \\ \circlearrowleft & & \circlearrowright \\ D^b A\text{-Brane}(\mathfrak{X}, \omega_{\mathfrak{X}})_{\gamma=0=\eta} & \cong & D^b \text{Rep}(S\ddot{H}_{q,t})_{|q|=1=|t_j|} \end{array}$$

where $T_a([\mathbf{D}_b]) = [\mathbf{D}_b] + ([\mathbf{D}_a] \cdot [\mathbf{D}_b])[\mathbf{D}_a]$ are subject to

$$\begin{aligned} T_j T_k &= T_k T_j \\ T_j T_4 T_j &= T_4 T_j T_4 \end{aligned}$$



SECTION 2

VOA modules vs $\mathbb{C}^\times$ -fixed components of $\mathcal{M}_H$

SCFT/VOA Correspondence

Beem-Lemos-Liendo-Peelaers-Rastelli-van Rees,

$$4d \mathcal{N} = 2 \text{ SCFT } \mathcal{T} \longrightarrow \text{VOA } \mathbb{V}(\mathcal{T})$$

- The 2d central charge is related to the 4d central charge by

$$c_{2d} = -12c_{4d}$$

- The vacuum character of the VOA equals the Schur index of the 4d theory:

$$\chi_{\mathbb{V}(\mathcal{T})} = I_{\text{Schur}}(\mathcal{T})$$

- The flavor symmetry algebra decomposes into simple factors

$$\mathfrak{g}_F = \bigoplus_i \mathfrak{g}_i \oplus \mathfrak{u}(1)^{\oplus s},$$

and each simple factor gives an affine Lie subalgebra $L_{k^{2d}}(\mathfrak{g}_i) \subset \mathbb{V}(\mathcal{T})$, with negative level

$$k_i^{2d} = -k_i^{4d}$$


- The associated variety $X_{\mathbb{V}(\mathcal{T})}$ is the Higgs branch.

Fixed components vs simple modules

- For a 4-punctured sphere, VOA is affine D_4 Lie algebra $\mathbb{V}(\mathcal{T}_{0,4}) = L_{-2}(\mathfrak{d}_4)$ with level -2
- There are 5 simple modules:

$$0, -2\omega_1, -2\omega_3, -2\omega_4, -\omega_2$$

- Fixed components under the $U(1)_r$ action consists of 4 points and $\text{Bun}_G \cong \mathbb{P}^1$:



Fixed components vs simple modules

- The $\mathbb{V}(\mathcal{T}_{0,4})$ -module characters $\{\chi_\lambda\}$ satisfy a set of flavor modular differential equations (FMDEs) These form a set of partial differential equations.
- The quasi-modularity of FMDEs implies that characters belong to a representation $\mathbb{S}_{0,4}^{mod}$ of the modular group
- Generally, the dimension of $\mathbb{S}_{0,4}^{mod}$ is larger than the number of SMs, where the extra solutions are usually logarithmic modules, and T is not semisimple.

$$SL(2, \mathbb{Z}) \curvearrowright \mathbb{S}_{0,4}^{mod} = \text{Span}_{\mathbb{C}} \{ 5 \text{ characters} + \log\text{-module} \} , \quad \text{Jord}(T) = [2, 1^4]$$

- Another perspective comes from the mixed Hodge polynomial of character variety We need its pure part:

$$PH(\mathcal{M}_{0,4}^{\text{Betti}}; q) = 4 + q$$

- $\#$ of simple modules = $\#$ fixed manifolds = $\#$ Jordan block $(T) = PH(q = 1)$

Fixed components vs simple modules


Yiwen Pan-Wenbin Yan

- $SL(2, \mathbb{Z}) \curvearrowright \mathbb{S}_{0,4}^{mod} = \text{Span}_{\mathbb{C}} \{ 5\text{characters} + \text{log-module} \} , \quad \text{Jord}(T) = [2, 1^4]$
- Another perspective comes from the mixed Hodge polynomial of character variety We need its pure part:

$$\text{PH}(\mathcal{M}_{0,4}^{\text{Betti}}; q) = 4 + q$$

- # of simple modules = # fixed manifolds = # Jordan block $(T) = PH(q = 1)$

$L(\omega_i)$	h	$\mathfrak{sl}(2)^{\otimes 4}$ rep	$\dim M$	$\mu - \mu_{max} _{\alpha_i \mapsto -\omega^{(i)}}$
$-\omega_2$	-1	$[-2, 0, 0, 0]$	1	$-1 - 2\omega^{(1)}$
$-2\omega_1$	-1	$[-2, -2, 0, 0]$	0	$-1 - 2\omega^{(1)} - 2\omega^{(2)}$
$-2\omega_3$	-1	$[-2, 0, -2, 0]$	0	$-1 - 2\omega^{(1)} - 2\omega^{(3)}$
$-2\omega_4$	-1	$[-2, 0, 0, -2]$	0	$-1 - 2\omega^{(1)} - 2\omega^{(4)}$
$-2\omega_0$	0	$[0, 0, 0, 0]$	0	0



SECTION 3

Time for wine? I bought a new decanter!

1-punctured torus

- Moduli space of flat $\mathrm{SL}(2, \mathbb{C})$ -connections on a once-punctured torus C_p

$$\pi_1(C_p) = \langle \mathfrak{m}, \mathfrak{l}, \mathfrak{c} \mid \mathfrak{m} \mathfrak{l} \mathfrak{m}^{-1} \mathfrak{l}^{-1} = \mathfrak{c} \rangle .$$

- These variables are subject to the cubic equation

$$\mathcal{M}_{\mathrm{flat}}(C_p, \mathrm{SL}(2, \mathbb{C})) = \{(x, y, z) \in \mathbb{C}^3 \mid x^2 + y^2 + z^2 - xyz - 2 = t^2 + t^{-2}\} .$$

From the 4-punctured sphere, we can obtain it by setting $\theta_{1,2,3} = 0$.

- The deformation quantization of $\mathcal{O}(\mathfrak{X})$ with respect to Ω_J is the spherical DAHA $S\ddot{H}_{q,t}$ of type A_1

$$\mathcal{O}(\mathfrak{X}) \rightsquigarrow \mathcal{O}^q(\mathfrak{X}) \cong S\ddot{H}_{q,t}$$

- Relations are given by:

$$[x, y]_q = (q^{-1} - q)z$$

$$[y, z]_q = (q^{-1} - q)x$$

$$[z, x]_q = (q^{-1} - q)y$$

$$q^{-1}x^2 + qy^2 + q^{-1}z^2 - q^{-\frac{1}{2}}xyz = (q^{-\frac{1}{2}}t - q^{\frac{1}{2}}t^{-1})^2 + (q^{\frac{1}{2}} + q^{-\frac{1}{2}})^2 ,$$



Symmetry

- **Modular group** $\mathrm{PSL}(2, \mathbb{Z})$: Since the algebra is constructed from a torus, it enjoys $\mathrm{PSL}(2, \mathbb{Z})$ -automorphism generated by

$$\tau_+ \mapsto \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, \quad \tau_- \mapsto \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

and

$$\sigma := \tau_+ \tau_-^{-1} \tau_+ = \tau_-^{-1} \tau_+ \tau_-^{-1} \mapsto \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad \tau_+ \tau_-^{-1} \mapsto \begin{pmatrix} 1 & -1 \\ 1 & 0 \end{pmatrix}.$$

The actions on the generators are given by

$$\tau_+ : (x, y, z) \mapsto \left(x, \frac{xy + yx}{q^{1/2} + q^{-1/2}} - z, y \right),$$

$$\tau_- : (x, y, z) \mapsto \left(\frac{xy + yx}{q^{1/2} + q^{-1/2}} - z, y, x \right),$$

$$\sigma : (x, y, z) \mapsto \left(y, x, \frac{xy + yx}{q^{1/2} + q^{-1/2}} - z \right).$$

- “**sign changes**” $\Xi = \mathbb{Z}_2 \times \mathbb{Z}_2$ are holomorphic involutions, and its generators ξ_1, ξ_2 and their combination $\xi_3 := \xi_1 \circ \xi_2$ act as

$$\xi_1 : (x, y, z) \mapsto (-x, y, -z),$$


$$\xi_2 : (x, y, z) \mapsto (x, -y, -z),$$

$$\xi_3 : (x, y, z) \mapsto (-x, -y, z),$$



Modularity

$PSL(2, \mathbb{Z})$ -actions on $\mathfrak{B}_{D_0}$ and $\mathfrak{B}_{D_4}$ are categorified by modular tensor categories



Affine Lie algebra $L_k(\mathfrak{a}_1)$


$PSL(2, \mathbb{Z}) \curvearrowright \text{Hom}(\mathfrak{B}_{D_4}, \mathfrak{B}_{cc})$

Virasoro minimal model

$PSL(2, \mathbb{Z}) \curvearrowright \text{Hom}(\mathfrak{B}_{D_0}, \mathfrak{B}_{cc})$

$(2, 2\ell + 1)$ Virasoro minimal model is VOA of $(A_1, A_{2(\ell-1)})$ Argyres-Douglas theory

(A_1, A_2) Argyres-Douglas theory and Lee-Yang model



$$\phi(z) \sim \frac{\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}}{z^{2+\frac{3}{2}}} + \dots$$

$$\mathbb{V}(A_1, A_2) = \text{Lee-Yang minimal model}$$

$$2 \text{ primaries } \{\varphi_0, \varphi_{-\frac{2}{5}}\} \curvearrowright PSL(2, \mathbb{Z})$$

$$S = \frac{2}{\sqrt{5}} \begin{pmatrix} -\sin \frac{2\pi}{5} & \sin \frac{\pi}{5} \\ \sin \frac{\pi}{5} & \sin \frac{2\pi}{5} \end{pmatrix}, \quad T = \begin{pmatrix} e^{\frac{11\pi i}{30}} & 0 \\ 0 & e^{-\frac{\pi i}{30}} \end{pmatrix}$$


Bezrukavnikov-Boixeda Alvarez-McBreen-Yun

2 fixed points in the central fiber (Kodaira type II)

In wild Hitchin moduli space

the central fiber is affine Springer fiber


$$\mathbb{C}^\times \curvearrowright Fl_\gamma = \{g \in G((z))/G[[z]] \mid Ad_{g^{-1}}(\gamma) \in \mathfrak{g}[[z]]\}$$




Web of relations

Simple modules of affine Lie algebras
with boundary admissible level


Peng Shan-Wenbin Yan-Dan Xie



$\mathbb{C}^\times$ -fixed points of
Affine springer fiber




$PSL(2, \mathbb{Z})$




Finite-dimensional module
of DAHA

$$t^{h^\vee} q^n = 1$$
$$q^m = 1$$





Bezrukavnikov-Boixeda Alvarez-McBreen-Yun

$\mathbb{C}^\times$ -fixed points of wild Hitchin moduli space




Hitchin moduli space and Coulomb branch

Coulomb branch of 4d $\mathcal{N} = 4$ theory is a $\mathbb{Z}_2$ quotient of $SU(2)$ Hitchin moduli space.



Simple modules vs fixed points via Langlands duality

- For a **once-punctured torus**, **VOA** $\mathbb{V}(\mathcal{T}_{0,1})$ is **small** $\mathcal{N} = 4$ **superconformal Virasoro algebra** with $c = -9$.
- There are **2 simple modules** of $\mathbb{V}(\mathcal{T}_{0,1})$: $0, -\omega_1$.
- $\mathbb{S}_{1,1}^{\text{mod}} = \text{Span}_{\mathbb{C}}(\text{2 simple} + \text{log})$, **Jordan type** of T is $[2,1]$
- **Fixed manifolds** of **SO(3) Hitchin moduli space** $\mathcal{M}_H(C_p, \text{SO}(3))$ consist of a point $+ \mathbb{P}^1 / (\mathbb{Z}_2 \times \mathbb{Z}_2)$
- $\text{SU}(2) \leftrightarrow \text{SO}(3)$ Langlands duality **Gukov-Pei-Hsin**



Mysterious relationship

$$\text{Simple modules of } \mathbb{V}(\mathcal{T}(C_{g,n}, G)) \overset{1-1}{\longleftrightarrow} \mathbb{C}^\times\text{-fixed components of } \mathcal{M}_H(C_{g,n}, G^\vee)$$

- More precise correspondence has been formulated for $G=\text{SU}(2)$ Yiwen Pan-Wenbin Yan
- Very little is understood for higher-rank
- Koszul type duality? What is the underlying intrinsic reason?
- Very mysterious even for physicists
- Any insights from mathematicians will be appreciated

EPILOGUE
Cheers!



Fudan University

Math dept



SIMIS




Shanghai Center for Mathematical Science

Fudan University



Many
Activities



We bring four professors which give five lectures a week
我们邀请了四位教授，他们每周讲授五场讲座



Beijing-Shanghai Summer
School in Mathematics

水暑 结



Jun 22 - July 3, 2026 at SIMIS

Lectures by:


- Tobias Ekholm
- Ko Honda
- Mikhail Khovanov
- Vivek Shende

Organized by:

- 李骏 (Jun Li)
- Andrei Okounkov
- Anton Dzhamay
- 单凡 (Peng Shan)
- 周业浩 (Yehao Zhou)



Every room is equipped with air conditioners!!!



Opportunities at the Xuemin Institute, Fudan University

Independent PI and Visiting Fellow positions



Xuemin Investigator Program (Independent PI)

For outstanding early-career scholars with bold vision and exceptional research potential. The program offers generous long-term funding, support to build an independent research group, academic freedom, tenure-track to tenured appointments, and the possibility of dual positions across Fudan University.



Eligibility

All nationalities welcome.
A doctoral degree is required (ideally within 7–8 years).
Exceptional candidates with a new doctoral degree or advanced doctoral candidates in a relevant field will also be considered.



Application materials

- Cover Letter
- Curriculum Vitae
- Research Statement
- Milestone Publications
- List of References



Xuemin Visiting Fellow Program

For scientists, engineers, clinicians, and scholars from around the world seeking a stimulating and collaborative environment. Fellows may be at any career stage and in various disciplines. Most Visiting Fellows stay from a few weeks up to a year.



How to apply

Early applications are encouraged (at least 3–4 months before the intended visit date) to allow sufficient time for review and arrangements.



Application materials

- Cover Letter
- Curriculum Vitae
- Milestone Publications
- Recommendation Letters
- Visiting Proposal



Submit application materials in PDF format to: xuemin_ias@fudan.edu.cn




More information: www.xuemin.org



Please feel free to share this with interested colleagues and collaborators.

Every room is equipped with air conditioners!!!





深化友谊

approfondir les liens d'amitié

IHES



復旦大學

