

Finite-word contraction and short descent certificates for the Juggler map

Publication draft. Not submitted.

Philippe Cochin

28 August 2026

Contents

Abstract	1
1. Introduction	2
1.1 Evidence discipline	2
1.2 Related maps	3
2. Finite words, envelope, and defect	3
3. Inverse cells and cycles	5
4. Short descent certificates	7
5. Ambient discrepancy and the short-certificate class	8
6. The remaining gap	10
7. Software archive	10
Acknowledgments	11
References	11

Abstract

The Juggler map is the nonlinear integer map

$$J(n) = \begin{cases} \lfloor \sqrt{n} \rfloor, & n \text{ even,} \\ \lfloor n^{3/2} \rfloor, & n \text{ odd.} \end{cases}$$

It is conjectured that every positive integer eventually reaches 1. We do not prove that conjecture.

For every finite parity word w realized at n , the iterate satisfies the power envelope

$$J^{|w|}(n)^{2^{|w|}} \leq n^{3^{\#(w)}}.$$

Hence $3^{\#O(w)} < 2^{|w|}$ forces $J^{|w|}(n) < n$ whenever $n \geq 2$. Local floor remainders lift through later exponents to an exact global defect: concatenation is a two-term power-gap, not an additive sum, and zero defect is the rigid monochrome towers. These statements are conditional on a realized word: they do not say that every orbit meets a contracting word, and the defect is not a state-independent tax.

A start $n \geq 2$ has a *descent certificate* if some realized finite word sends it strictly below n . Every even start, and every odd start whose first image is even, has a uniform one- or two-step certificate. A classical discrepancy bound

$$|S_O(N)| \ll N^{5/6}, \quad S_O(N) = \sum_{\substack{n \leq N \\ n \text{ odd}}} (-1)^{\lfloor n^{3/2} \rfloor},$$

implies that the complementary odd-to-odd starts have density $1/4$, so the uniform short-certificate class has density $3/4$. This is not a density of all descent certificates, and it is not a density of starts that reach 1. Odd-to-odd starts expand on the first step and may still descend later. The bound itself is ambient: it does not transfer to sparse Juggler-generated image sets.

The remaining question, analogue of Terras’s almost-all stopping-time theorem, is whether almost every odd-to-odd start has some finite descent. No such theorem is proved here.

1. Introduction

The Juggler sequence was introduced by Pickover [1]; see also OEIS A094683 [2]. Universal arrival at 1 remains open. An exact check in the software archive shows every start $n \leq 4000$ reaches 1. That check is not a proof.

The map combines a contracting even branch with an expanding odd branch,

$$E(n) = \lfloor n^{1/2} \rfloor, \quad O(n) = \lfloor n^{3/2} \rfloor.$$

A word of length k with o odd letters has ideal exponent $3^o/2^k$. Floors are applied after every letter, and a word is available only when the orbit realizes those parities. The paper records the exact finite-word comparison, the compositional slack, the uniform short certificates, and the density of the class those certificates cover.

The contribution is a short exact-arithmetic note, not a survey of failed compressions and not a description of the computational atlas used in the surrounding laboratory. We do not prove a Collatz theorem, and we do not transfer Collatz stopping-time results to J .

1.1 Evidence discipline

- **EXACT — LEAN VERIFIED:** a theorem checked by Lean in the software archive named in Section 7.
- **EXACT — HUMAN PROOF:** a mathematical argument not packaged in Lean. Theorem 5.1 is of this kind.
- **COMPUTATIONALLY VERIFIED:** an exact finite computation under stated bounds.
- **OBSERVATION:** a descriptive pattern in finite data.
- **REFUTED:** a universal candidate killed by a certificate.

A finite check is never called a termination proof.

1.2 Related maps

The nearest published comparison is the Collatz problem. Lagarias [3,4] surveys parity words and almost-all statements short of totality. Terras [5] and Everett [6] proved that almost every positive integer has a finite Collatz stopping time; Tao [7] later showed that almost all Collatz orbits attain almost bounded values. Those results are methodological cousins, not theorems about J . The Juggler analogue of Terras would be an almost-all descent theorem on the odd-to-odd class.

Crandall [8] and Matthews–Watts [9] treat piecewise-affine Hasse–Syracuse maps; the Juggler branches are floor powers, not affine. Prasad and Prasad [11] estimate excursion and stopping constants for juggler-like random walks; Section 6 keeps that comparison descriptive.

2. Finite words, envelope, and defect

Let $\mathcal{B} = \{E, O\}$. A finite word $w \in \mathcal{B}^*$ is *realized* at $n \in \mathbb{N}$ when the successive parities of the orbit of n are exactly the letters of w . Write $J^{|w|}(n)$ for the endpoint after those letters, and $\#O(w)$ for the number of odd letters.

The identities in this section are formalized in Lean (`follows_iff_word`, `image_eq_iterate`, `image_append`, `image_monotone_of_follows`, `power_bound_word`, `power_bound_contracts`, `global_defect_identity`, `global_defect_eq_zero_iff_localsTight`, `global_defect_append`, `power_bound_eq_iff_extremal`). The proofs below are the ordinary integer arguments.

Theorem 2.1 (fixed-word monotonicity; EXACT — LEAN VERIFIED). If $n \leq m$ and both realize w , then $J^{|w|}(n) \leq J^{|w|}(m)$.

The realizing set of a fixed word need not be an interval.

Theorem 2.2 (finite-word power envelope; EXACT — LEAN VERIFIED). If w is realized at n and $m = J^{|w|}(n)$, then

$$m^{2^{|w|}} \leq n^{3^{\#O(w)}}.$$

Proof. Write $k = |w|$ and $o = \#O(w)$. The empty word is the equality $n \leq n$. Suppose the bound holds at a realized prefix ending at x , and the next letter is realized.

If the next letter is even, then $J(x) = \lfloor \sqrt{x} \rfloor$, so $J(x)^2 \leq x$. Raising the inductive bound to the second power gives

$$J(x)^{2^{k+1}} = (J(x)^2)^{2^k} \leq x^{2^k} \leq n^{3^o}.$$

The odd count is unchanged.

If the next letter is odd, then $J(x) = \lfloor x^{3/2} \rfloor$, so $J(x)^2 \leq x^3$. Therefore

$$J(x)^{2^{k+1}} = (J(x)^2)^{2^k} \leq x^{3 \cdot 2^k} = (x^{2^k})^3 \leq (n^{3^o})^3 = n^{3^{o+1}}.$$

This is the claimed bound after one more odd letter. $\square$

Corollary 2.3 (exponent-gap contraction; EXACT — LEAN VERIFIED). If $n \geq 2$, w is realized at n , and $3^{\#O(w)} < 2^{|w|}$, then $J^{|w|}(n) < n$.

Proof. Let $m = J^{|w|}(n)$ and $k = |w|$. Theorem 2.2 gives $m^{2^k} \leq n^{3^{\#O(w)}}$. The exponent gap and $n \geq 2$ give $n^{3^{\#O(w)}} < n^{2^k}$, so $m^{2^k} < n^{2^k}$. Since $m \geq 1$, one has $m < n$. $\square$

The corollary includes familiar contracting blocks such as $OOOEE$. It does not prove that every start realizes some contracting word.

The floor slack is exact, and it is not an additive path sum. For a single branch,

$$x^e = J(x)^2 + \rho(x), \quad e = \begin{cases} 1, & x \text{ even}, \\ 3, & x \text{ odd}, \end{cases}$$

with $0 \leq \rho(x) < 2J(x) + 1$. Write

$$\text{gap}(a, \rho, e) = (a + \rho)^e - a^e,$$

so $a^e + \text{gap}(a, \rho, e) = (a + \rho)^e$. If $e \geq 1$, then $\text{gap}(a, \rho, e) = 0$ if and only if $\rho = 0$, and $\rho^e \leq \text{gap}(a, \rho, e)$.

Along a realized word w , write $x_0 = n$ and $x_{j+1} = J(x_j)$, and let $\rho_j = \rho(x_j)$. Define a running slack by $D_0 = 0$ and

$$D_{j+1} = \begin{cases} D_j + \text{gap}(x_{j+1}^2, \rho_j, 2^j), & \text{next letter even,} \\ \text{gap}(x_{j+1}^2, \rho_j, 2^j) + \text{gap}(x_j^{2^j}, D_j, 3), & \text{next letter odd.} \end{cases}$$

The *global defect* is $\Delta_w(n) = D_{|w|}$. An even letter keeps the old slack and lifts the new remainder through 2^j . An odd letter first cubes the running slack and then lifts the new remainder. In particular Δ is not the sum of the local remainders.

Theorem 2.4 (global defect identity; EXACT — LEAN VERIFIED). If w is realized at n and $m = J^{|w|}(n)$, then

$$n^{3^{\#O(w)}} = m^{2^{|w|}} + \Delta_w(n), \quad \Delta_w(n) \geq 0.$$

Theorem 2.2 is the inequality $\Delta_w(n) \geq 0$.

Proof. The empty word is $n = n + 0$. Suppose a realized prefix of length k with odd count o ends at x and satisfies $n^{3^o} = x^{2^k} + D$.

If the next letter is even, then $x = J(x)^2 + \rho(x)$, so

$$n^{3^o} = (J(x)^2 + \rho(x))^{2^k} + D = J(x)^{2^{k+1}} + D + \text{gap}(J(x)^2, \rho(x), 2^k).$$

The odd count is unchanged, and the new slack is the even update.

If the next letter is odd, then $x^3 = J(x)^2 + \rho(x)$. Cubing the inductive identity gives

$$n^{3^{o+1}} = (x^{2^k} + D)^3 = x^{3 \cdot 2^k} + \text{gap}(x^{2^k}, D, 3).$$

The leading term is

$$x^{3 \cdot 2^k} = (J(x)^2 + \rho(x))^{2^k} = J(x)^{2^{k+1}} + \text{gap}(J(x)^2, \rho(x), 2^k),$$

which is the odd update. $\square$

Theorem 2.5 (vanishing; EXACT — LEAN VERIFIED). If w is realized at n , the following are equivalent.

1. $\Delta_w(n) = 0$.
2. Every local remainder along w vanishes.
3. $J^{|w|}(n)^{2^{|w|}} = n^{3^{\#O(w)}}$.

In that case w is monochrome. More precisely, either $w = E^k$ and $n = a^{2^k}$ for an even a , or $w = O^k$ and $n = a^{2^k}$ for an odd a . A realized mixed word therefore has $\Delta_w(n) > 0$.

Proof. The identity of Theorem 2.4 gives $(1) \Leftrightarrow (3)$. For $(1) \Leftrightarrow (2)$, use the recurrence. If $e \geq 1$, a power-gap vanishes if and only if its addend vanishes. The even update is a sum of two nonnegative terms, and the odd update is a sum of two power-gaps; either vanishes only when the incoming slack and the new remainder both vanish. Thus $D_{|w|} = 0$ forces $D_0 = 0$ and every $\rho_j = 0$. The converse is immediate.

If every remainder vanishes, each step is an exact square or an exact cube-to-square, so the current parity is preserved and the itinerary cannot use both letters. The exact towers are the saturation of the envelope (`power_bound_eq_iff_extremal`). $\square$

Theorem 2.6 (composition; EXACT — LEAN VERIFIED). If u is realized at n and v is realized at $m = J^{|u|}(n)$, then

$$\Delta_{uv}(n) = \text{gap}(m^{2^{|u|}}, \Delta_u(n), 3^{\#O(v)}) + \text{gap}(J^{|v|}(m)^{2^{|v|}}, \Delta_v(m), 2^{|u|}).$$

In particular $\Delta_v(m)^{2^{|u|}} \leq \Delta_{uv}(n)$. Every local remainder satisfies $\rho_j^{2^j} \leq \Delta_w(n)$.

Proof. Write $o_u = \#O(u)$ and $o_v = \#O(v)$. Theorem 2.4 on u and on uv gives

$$n^{3^{o_u+o_v}} = (m^{2^{|u|}} + \Delta_u(n))^{3^{o_v}} = J^{|uv|}(n)^{2^{|uv|}} + \Delta_{uv}(n).$$

The first power expands as

$$m^{2^{|u|} \cdot 3^{o_v}} + \text{gap}(m^{2^{|u|}}, \Delta_u(n), 3^{o_v}).$$

Theorem 2.4 on v at m rewrites the leading term as

$$(J^{|v|}(m)^{2^{|v|}} + \Delta_v(m))^{2^{|u|}},$$

which expands as the second gap plus $J^{|uv|}(n)^{2^{|uv|}}$. Comparing the two expressions for $n^{3^{o_u+o_v}}$ yields the identity. The suffix inequality is $\rho^e \leq \text{gap}(a, \rho, e)$ on the second summand. For a local remainder at index j , split w after j letters: the suffix defect is at least ρ_j , and raising through 2^j gives the stated bound. $\square$

Composition is polynomial, not additive: later odd letters raise the prefix slack to the third power, and later letters of either parity raise a suffix slack through $2^{|u|}$. The naive recurrence $\Delta \leftarrow \Delta + \rho$ is false.

The identity does not supply a state-independent positive tax. Persistent expanding blocks can have arbitrarily small observed normalized slack $\Delta/n^{3^{\#O(w)}}$ at large scale. The inequality $\Delta_w(n) > n^{3^{\#O(w)}} - n^{2^{|w|}}$ on a formally expanding word is exactly $J^{|w|}(n) < n$, so it does not forbid mixed expanding prefixes.

3. Inverse cells and cycles

The exact one-step fibers are

$$J(n) = q \iff q^2 \leq n < (q+1)^2 \quad (n \text{ even})$$

and

$$J(n) = m \iff m^2 \leq n^3 < (m+1)^2 \quad (n \text{ odd}).$$

An odd fiber contains at most one integer (`odd_cell_unique`; EXACT — LEAN VERIFIED). An even fiber is a parity-restricted square interval and may contain many predecessors.

A nonempty realized word w with $J^{|w|}(n) = n$ is a cycle word.

The identities in this section are formalized in Lean (`cycle_word_formally_expanding`, `cycleMin_not_end_odd`, `square_scale_superquadratic`, `cycleMin_to_even_superquadratic`, `no_cycle_word_oooeoe`, `no_cycle_word_ooooee`). The proofs below are the ordinary integer arguments.

Theorem 3.1 (cycle restrictions; EXACT — LEAN VERIFIED). Let w be a cycle word at $n \geq 2$.

(i) The word is formally expanding:

$$2^{|w|} < 3^{\#O(w)}$$

(`cycle_word_formally_expanding`). A contracting word cannot close a nontrivial cycle.

(ii) The cycle minimum is odd and the cycle maximum is even. A minimum-based orientation cannot end in an odd letter (`cycleMin_not_end_odd`).

(iii) A realized word v is *superquadratic* if

$$3^{\#O(v)} \geq 2^{|v|+1}.$$

Any realized path from a start $n \geq 2$ to a state at least n^2 is superquadratic (`square_scale_superquadratic`). On a cycle minimum the path to any later even state is superquadratic (`cycleMin_to_even_superquadratic`). The prefix *OOE* is expanding ($9 > 8$) but not superquadratic ($9 < 16$), so it cannot carry the minimum to square scale.

These are necessary restrictions, not an exclusion of every nontrivial cycle.

The two remaining legal minimum-based length-six orientations are nevertheless impossible. The argument uses the last-even cell together with a coarse lower envelope. For $n \geq 1$,

$$n < 4 \lfloor \sqrt{n} \rfloor^2,$$

so an even step satisfies $n \leq 4J(n)^2$ and an odd step satisfies $n^3 \leq 4J(n)^2$. Composing along a realized word v of length k with odd count o gives

$$n^{3^o} \leq D_v J^k(n)^{2^k}.$$

The denominator starts at 1 and updates by $D \mapsto D \cdot 4^{2^j}$ on an even letter and $D \mapsto D^3 \cdot 4^{2^j}$ on an odd letter, at step j . In particular $D_{OOO} = 2^{38}$ and $D_{OOOO} = 2^{130}$. If a cycle word ends in an even letter, the last-even cell is $J^{|w|-1}(n) < (n+1)^2$.

Theorem 3.2 (leftover length-six orientations; EXACT — LEAN VERIFIED). Neither *OOOEOE* nor *OOOOEE* is a cycle word at any $n \geq 2$. This is not an exclusion of every length-six word, nor of every cycle.

Proof. First, if $n \geq 256$, then

$$n^{81} > 2^{130}(n+1)^{64}.$$

Indeed $257^{64} < 2 \cdot 256^{64}$ and $256(n+1) \leq 257n$, so $(n+1)^{64} < 2n^{64}$. Then $2^{130}(n+1)^{64} < 2^{131}n^{64}$. Since $n \geq 256 = 2^8$, one has $n^{17} \geq 2^{136}$, and therefore $2^{131}n^{64} < 2^{136}n^{64} \leq n^{81}$.

For $n < 256$, neither word returns to its start. Lean checks both itineraries by `native_decide` on `Fin 256`.

Now suppose $n \geq 256$ realizes $OOOOEE$, and write $z = J^4(n)$ for the image after the prefix $OOOO$. The last two even letters give $J(z) < (n+1)^2$ and $z < (J(z)+1)^2$, hence $z < (n+1)^4$. The lower envelope on $OOOO$ is $n^{81} \leq 2^{130}z^{16}$, so $n^{81} < 2^{130}(n+1)^{64}$, contradicting the tail inequality.

Finally suppose $n \geq 256$ realizes $OOOEOE$. Write $z_3 = J^3(n)$ and $y = J(z_3) = \lfloor \sqrt{z_3} \rfloor$, so $z_3 < (y+1)^2$. The lower envelope on OOO is $n^{27} \leq 2^{38}z_3^8 < 2^{38}(y+1)^{16}$. Cubing yields $n^{81} < 2^{114}(y+1)^{48}$. The last letters OE give the odd-cell bound $y^3 < (n+1)^4$. Write $A = n+1 \geq 257$. Then $(y+1)^3 < 2A^4$: if $y \leq A$, this follows from $(A+1)^3 < 2A^4$; if $y > A$, then $y \geq 4$ and $(y+1)^3 = y^3 + 3y^2 + 3y + 1 < A^4 + 4y^2$, while $4y^2 < A^4$ because $4y^3 < 4A^4 \leq yA^4$. Raising $(y+1)^3 < 2A^4$ to the sixteenth power gives $(y+1)^{48} < 2^{16}(n+1)^{64}$. Combining with the cubed lower envelope produces again $n^{81} < 2^{130}(n+1)^{64}$. $\square$

4. Short descent certificates

A start $n \geq 2$ has a *descent certificate* if there exists a realized finite word w with $J^{|w|}(n) < n$, or with image 1. Lean packages this predicate as `FiniteProgress`, an abbreviation of `DescentCertificate`. The four constructors of that type are proof forms of the same predicate, not four different claims. The predicate is existential over all finite words, not only over words of length one or two.

Theorem 4.1 (uniform short certificates; EXACT — LEAN VERIFIED). Let $n \geq 2$.

1. If n is even, then the one-letter word E is a descent certificate: $J(n) = \lfloor \sqrt{n} \rfloor < n$.
2. If n is odd and $J(n)$ is even, then the two-letter word OE is a descent certificate.

Theorem 4.2 (unresolved starts are odd-to-odd; EXACT — LEAN VERIFIED). If $n \geq 2$ has no descent certificate, then n is odd and $J(n)$ is odd.

The converse is false: many odd-to-odd starts descend after a longer word.

If every start above 1 has some descent certificate, ordinary strong induction yields arrival at 1 (`reachesOne_of_all_finiteProgress`). The hypothesis is not proved. A certificate at n only reduces the problem to a strictly smaller positive integer, which may itself be odd-to-odd.

Lean also certifies a finite landing class (`reachesOne_of_lt_twelve, even_lt_sq_twelve_reachesOne`):

Theorem 4.3 (small residuals; EXACT — LEAN VERIFIED). Every $y \in \{1, \dots, 11\}$ reaches 1. Consequently every even residual strictly below $144 = 12^2$ reaches 1.

This enlarges the set of fatal landings for a hypothetical minimal counterexample. It does not prove that every even start reaches 1.

Independently, an exact computation gives:

Proposition 4.4 (window totality; COMPUTATIONALLY VERIFIED). Every start $1 \leq n \leq 4000$ reaches 1.

This is a finite check, not a bound on all starts.

On the complementary odd-to-odd class, first return below the start is frequent at a short horizon, but not automatic.

Proposition 4.5 (odd-to-odd first return; OBSERVATION). For starts $2 \leq n \leq N$ and horizon 20, write OO for the odd-to-odd starts in that range. The exact census is:

N	#OO	OO first return ≤ 20	all starts, first return ≤ 20
10^3	252	0.877	0.969
10^4	2504	0.887	0.972
10^5	24984	0.896	0.974
10^6	249926	0.895	0.974

At $N = 10^6$, 26,243 odd-to-odd starts have no return below the start in 20 steps. None of these rows is a Lean descent certificate, and none is an almost-all theorem.

5. Ambient discrepancy and the short-certificate class

For odd n write $s(n) = (-1)^{\lfloor n^{3/2} \rfloor}$ and

$$S_O(N) = \sum_{\substack{n \leq N \\ n \text{ odd}}} s(n).$$

Let $M(N)$ be the number of odd integers $n \leq N$, and let

$$\text{OO}(N) = \#\{n \leq N : n \text{ odd}, J(n) \text{ odd}\}.$$

Then $s(n) = -1$ if and only if $J(n)$ is odd, and

$$S_O(N) = M(N) - 2 \text{OO}(N).$$

Also $M(N) = N/2 + O(1)$.

The floor sign has an exact fractional-part form. Write $x = \lfloor x \rfloor + \{x\}$. If $\lfloor x \rfloor$ is even then $\{x/2\} < 1/2$; if $\lfloor x \rfloor$ is odd then $\{x/2\} \geq 1/2$. Thus

$$\lfloor x \rfloor \text{ is odd} \iff \{x/2\} \geq \frac{1}{2}.$$

For odd $n = 2r + 1$ set

$$g(r) = \frac{(2r+1)^{3/2}}{2}.$$

Then $s(n) = -1$ if and only if $\{g(r)\} \geq 1/2$, and $S_O(N)$ is twice the interval discrepancy of the sequence $\{g(r)\}$ against $[1/2, 1)$.

The following two lemmas are classical; see Kuipers–Niederreiter [10, Ch. 1–2].

Lemma 5.A (van der Corput, second-derivative form). Let f be twice differentiable on an interval of length M , with $\lambda \leq |f''| \leq \alpha\lambda$ for some $\alpha \geq 1$. Then

$$\left| \sum e(f) \right| \ll_{\alpha} M\lambda^{1/2} + \lambda^{-1/2},$$

where the sum runs over the integers in the interval and $e(t) = e^{2\pi it}$.

Lemma 5.B (Erdős–Turán). The discrepancy of R points $x_j \in \mathbb{R}/\mathbb{Z}$ against an interval satisfies

$$D \ll \frac{R}{H} + \sum_{h=1}^H \frac{1}{h} \left| \sum_{j=1}^R e(hx_j) \right|$$

for every cutoff $H \geq 1$.

Theorem 5.1 (ambient odd-input discrepancy; EXACT — HUMAN PROOF).

$$|S_O(N)| \ll N^{5/6}.$$

This argument is not packaged in Lean.

Proof. The second derivative is

$$g''(r) = \frac{3}{2}(2r+1)^{-1/2}.$$

It is positive and decreasing. On a dyadic block $r \asymp M$ one has $g''(r) \asymp M^{-1/2}$. For the h -th Fourier mode take $f = hg$, so $\lambda \asymp hM^{-1/2}$. Lemma 5.A gives

$$\left| \sum_{r \asymp M} e(hg(r)) \right| \ll h^{1/2} M^{3/4} + h^{-1/2} M^{1/4}.$$

Lemma 5.B, with $R \asymp M$ and the relation between S_O and interval discrepancy, yields

$$|S_O| \ll \frac{M}{H} + M^{3/4} H^{1/2} + M^{1/4}$$

on that block: the middle term is $\sum_{h \leq H} h^{-1} \cdot h^{1/2} M^{3/4} \ll M^{3/4} H^{1/2}$, and the last term is $\sum_{h \leq H} h^{-1} \cdot h^{-1/2} M^{1/4} \ll M^{1/4}$. The choice $H \asymp M^{1/6}$ balances the first two terms at $M^{5/6}$. Summing dyadic blocks up to N preserves the exponent. The floor sign is not replaced by a single exponential; the exponential sums are those of the sequence $\{g(r)\}$. $\square$

Corollary 5.2 (short-certificate class; EXACT — HUMAN PROOF). The odd-to-odd starts have natural density $1/4$:

$$|\text{OO}(N) - N/4| \ll N^{5/6}.$$

Equivalently, the starts that admit the uniform one- or two-step certificates of Theorem 4.1 — every even start together with every odd-to-even start — have natural density $3/4$.

This is a density of the uniform short-certificate class of Theorem 4.1, not of all descent certificates and not of starts that reach 1.

The exact census through $N = 10^7$ has $\text{OO}(N)/N = 0.2499896$. Through $N = 10^6$ one has $S_O(N) = 146$ and running maximum 256, at $n = 985351$. A spot computation at 10^7 has running maximum 459. The observed growth is much smaller than $N^{5/6}$. The descriptive $N^{1/3}$ -scale envelope is not promoted to a theorem.

The bound does not transfer from intervals to Juggler images. For a general interval $I = [A, B]$ the variation of S_O depends on the right endpoint, not only on $|I|$. The expanding image $Y = J_O(O(I))$ is a sparse gap set. At source bound 10^6 , recorded consecutive gaps in Y range from 4 to 3000. A certified finite concentration is the run of 52 consecutive odd sources of equal sign on $[952525, 952627]$. Therefore ambient interval cancellation does not imply orbit or image-set cancellation. This is a boundary for the present argument, not a proof that no sparse-sequence estimate can exist.

6. The remaining gap

Theorem 4.1 and Corollary 5.2 together say that a uniform one- or two-step argument covers a set of density $3/4$. The first odd-to-odd image expands, so ordinary strong induction cannot fire on that class. Proposition 4.5 shows that most odd-to-odd starts in a finite window still return below the start inside twenty steps. That is an observation, not Terras’s theorem for J .

A mixed-parity heuristic, ignoring floors, gives mean log-log drift $\frac{1}{2} \log(3/4) < 0$. Finite ensembles sit near this value; hard paths are more odd-rich. This agrees qualitatively with the juggler-like random-walk model of Prasad and Prasad [11]. Fair parity is an assumption, not a dynamical theorem, and typical negative drift is not pointwise contraction.

The gap is therefore:

No theorem forces every exact integer state into a contracting prefix. In particular, it is open whether almost every odd-to-odd start has a finite descent certificate.




Figure 1: Even and odd-to-even starts have uniform short certificates. The complementary odd-to-odd class is the remaining pointwise gap. This figure is not a theorem.

That is the Juggler form of the Terras question. A sharper ambient discrepancy exponent does not answer it.

7. Software archive

Lean proofs of the exact-arithmetic theorems live at https://github.com/sneakyweasel/balanced_ternary/. The archive is not required to read the arguments above. Theorem 5.1 is a human proof; Lean does not certify it.

From a clone, the review object is

```
cd formal && lake build Problems.JugglerPaper
```

A Word Atlas used in the laboratory records bounded realizers through length 20 and starts $n \leq 10^8$. It is infrastructure, not a theorem of this note.

Acknowledgments

I used large language models extensively while drafting and revising the text, organizing companion notes, and as an interactive assistant for Lean statements, tests, and literature records. The models are not authors. Lean theorems and named computations are the certificates for those claims. The discrepancy bound (Theorem 5.1) is a human proof using classical analytic inequalities; it is not Lean-certified. I take full responsibility for the contents.

References

1. C. A. Pickover, *Computers and the Imagination: Visual Adventures Beyond the Edge*, St. Martin's Press, New York, 1991, ch. 40, p. 232.
2. OEIS Foundation Inc., “Juggler sequence: if n even then $\lfloor \sqrt{n} \rfloor$ else $\lfloor n^{3/2} \rfloor$,” Sequence A094683 in *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org/A094683> (accessed 28 August 2026).
3. J. C. Lagarias, “The $3x+1$ problem and its generalizations,” *Amer. Math. Monthly* 92 (1985), 3–23. doi:10.1080/00029890.1985.11971528.
4. J. C. Lagarias (ed.), *The Ultimate Challenge: The $3x+1$ Problem*, American Mathematical Society, Providence, RI, 2010.
5. R. Terras, “A stopping time problem on the positive integers,” *Acta Arith.* 30 (1976), 241–252. doi:10.4064/aa-30-3-241-252.
6. C. J. Everett, “Iteration of the number-theoretic function $f(2n) = n$, $f(2n+1) = 3n+2$,” *Adv. Math.* 25 (1977), 42–45. doi:10.1016/0001-8708(77)90087-1.
7. T. Tao, “Almost all orbits of the Collatz map attain almost bounded values,” *Forum Math. Pi* 10 (2022), e12. doi:10.1017/fmp.2022.8.
8. R. E. Crandall, “On the “ $3x+1$ ” problem,” *Math. Comp.* 32 (1978), 1281–1292. doi:10.1090/S0025-5718-1978-0480321-3.
9. K. R. Matthews and A. M. Watts, “A generalization of Hasse’s generalization of the Syracuse algorithm,” *Acta Arith.* 43 (1984), 167–175. doi:10.4064/aa-43-2-167-175.
10. L. Kuipers and H. Niederreiter, *Uniform Distribution of Sequences*, Wiley-Interscience, New York, 1974.
11. V. Prasad and M. A. Prasad, “Estimates of the maximum excursion constant and stopping constant of juggler-like sequences,” ResearchGate preprint, 2025. doi:10.13140/RG.2.2.14110.04168.