

Small cycles of the Juggler map

Philippe Cochin
philippe@cochin.fr

29 August 2026

Abstract

The Juggler map is the nonlinear integer map

$$J(n) = \begin{cases} \lfloor \sqrt{n} \rfloor, & n \text{ even,} \\ \lfloor n^{3/2} \rfloor, & n \text{ odd.} \end{cases}$$

It is conjectured that every positive integer eventually reaches 1. We do not prove that conjecture.

The main theorem is a small-cycle census: the Juggler map has no nontrivial cycle of length at most six. The argument is elementary. A realized parity word w obeys the power envelope $(J^{|w|}(n))^{2^{|w|}} \leq n^{3^{\#O(w)}}$, so a cycle word is formally expanding. Inverse cells and two next-square thresholds exclude every even-terminating expanding word of length at most six except two leftover shapes, *OOOEOE* and *OOOOEE*. Those two are excluded by a tail inequality for $n \geq 256$ and a check of the 254 starts $2 \leq n < 256$. Length seven is the first combinatorial shape the thresholds do not reach.

The same envelope gives an exact global defect by keeping the floor remainders. Zero defect characterizes the monochrome power towers; concatenation is a two-term power-gap. The defect is not used as a uniform tax — none exists — and is recorded because it makes the envelope identity exact.

Even starts and odd-to-even starts have a uniform one- or two-step descent. The complementary class is the odd-to-odd starts; many of those still descend after a longer word. No density result is stated or used here.

2020 Mathematics Subject Classification. 11B83, 37P99, 11Y55.

1. Introduction

The Juggler sequence was introduced by Pickover [1,2]. The one-step map is OEIS A094683 [3]; the stopping-time table is A007320 [4]. The orbit of 3 is 3, 5, 11, 36, 6, 2, 1. The orbit of 37 already peaks at 24906114455136. Universal arrival at 1 remains the open conjecture of the subject.

The map combines a contracting even branch with an expanding odd branch,

$$E(n) = \lfloor n^{1/2} \rfloor, \quad O(n) = \lfloor n^{3/2} \rfloor.$$

A word of length k with o odd letters has ideal exponent $3^o/2^k$. Floors are applied after every letter, and a word is available only when the orbit realizes those parities.

The main theorem of the note is the census: there is no nontrivial cycle of period at most six (Theorem 3.6). The tool is the finite-word envelope (Theorem 2.2), together with the inverse cells and two next-square thresholds. The exact defect (Theorems 2.4–2.6) is the same recurrence with remainders kept rather than dropped. It classifies the rigid zero cases and shows that a uniform per-step slack tax is impossible; it is not needed for the length-six exclusion beyond a short alternative to the exponent comparison on mixed words.

Throughout, $\mathbb{N} = \{1, 2, 3, \dots\}$. Write J^k for the k -fold iterate. A nonempty realized word w with $J^{|w|}(n) = n$ is a *cycle word*. The unique fixed point is 1; a cycle is *nontrivial* when it contains some $n \geq 2$.

1.1 Related work

Pickover’s later exposition is Chapter 45 of [2]. Weisstein [5] records the map, the stopping-time sequences A094670, A094679, A095908, and a verification of arrival at 1 through 10^6 . OEIS A094716 [6] records extreme heights, including the start 48443 whose peak has 972 463 digits. Those computations do not bound the period and are not used in the proofs below.

We know of no published exclusion of nontrivial cycles for the exact floor-power map J . Prasad–Prasad [7] estimate excursion and stopping constants for juggler-like maps by a random-walk large-deviation model; those estimates do not apply here. Small-cycle censuses are a standard first layer for Collatz-like maps, surveyed by Lagarias [8,9]. Those results do not transfer: the branches of J are floor powers rather than affine maps (Crandall [10], Matthews–Watts [11]).

Itinerary-class densities are the subject of a companion manuscript [12] and are not used here.

1.2 Verification

The arguments of Sections 2–4 may be read without machine assistance. An independent Lean check of those arguments lives in the repository named in Section 4; the corresponding theorem names are collected in Appendix A. Lemma 3.5 uses a table of 254 six-step evaluations for $2 \leq n < 256$, one table for each of the two leftover words. That table is a finite computation, not a termination proof.

2. Envelope and defect

Let $\mathcal{B} = \{E, O\}$. A finite word $w \in \mathcal{B}^*$ is *realized* at $n \in \mathbb{N}$ when the successive parities of the orbit of n are exactly the letters of w . Write $J^{|w|}(n)$ for the endpoint after those letters, and $\#O(w)$ for the number of odd letters. When w is realized, this endpoint is the $|w|$ -fold iterate.

Theorem 2.1 (fixed-word monotonicity). If $n \leq m$ and both realize w , then $J^{|w|}(n) \leq J^{|w|}(m)$.

Proof. Induct on w . The empty word is immediate. The images after a common realized prefix remain ordered, and both current states realize the same next letter. The even branch is the monotone integer square root; the odd branch is $x \mapsto x^3$ followed by the integer square root. $\square$

The realizing set of a fixed word need not be an interval: OE is realized at 7 and at 11 but not at 9.

Theorem 2.2 (finite-word power envelope). If w is realized at n and $m = J^{|w|}(n)$, then

$$m^{2^{|w|}} \leq n^{3^{\#O(w)}}.$$

Proof. The empty word is the equality $n \leq n$. Suppose a realized prefix of length ℓ with odd count o ends at x and satisfies $x^{2^\ell} \leq n^{3^o}$, and the next letter is realized.

If the next letter is even, then $J(x)^2 \leq x$. Raising the inductive bound to the second power gives

$$J(x)^{2^{\ell+1}} = (J(x)^2)^{2^\ell} \leq x^{2^\ell} \leq n^{3^o}.$$

The odd count is unchanged.

If the next letter is odd, then $J(x)^2 \leq x^3$, so

$$J(x)^{2^{\ell+1}} = (J(x)^2)^{2^\ell} \leq x^{3 \cdot 2^\ell} = (x^{2^\ell})^3 \leq (n^{3^o})^3 = n^{3^{o+1}}.$$

This is the claimed bound after one more odd letter. $\square$

Corollary 2.3 (exponent-gap contraction). If $n \geq 2$, w is realized at n , and $3^{\#O(w)} < 2^{|w|}$, then $J^{|w|}(n) < n$.

Proof. Let $m = J^{|w|}(n)$ and $k = |w|$. Theorem 2.2 gives $m^{2^k} \leq n^{3^{\#O(w)}}$. The exponent gap and $n \geq 2$ give $n^{3^{\#O(w)}} < n^{2^k}$, so $m^{2^k} < n^{2^k}$. Since $m \geq 1$, one has $m < n$. $\square$

The corollary includes familiar contracting blocks such as $OOOEE$. It does not prove that every start realizes some contracting word.

The floor slack is exact. For a single branch,

$$x^e = J(x)^2 + \rho(x), \quad e = \begin{cases} 1, & x \text{ even}, \\ 3, & x \text{ odd}, \end{cases}$$

with $0 \leq \rho(x) < 2J(x) + 1$. Write

$$\text{gap}(a, \rho, e) = (a + \rho)^e - a^e,$$

so $a^e + \text{gap}(a, \rho, e) = (a + \rho)^e$. If $e \geq 1$, then $\text{gap}(a, \rho, e) = 0$ if and only if $\rho = 0$, and $\rho^e \leq \text{gap}(a, \rho, e)$.

Along a realized word w , write $x_0 = n$ and $x_{j+1} = J(x_j)$, and let $\rho_j = \rho(x_j)$. Define a running slack by $D_0 = 0$ and

$$D_{j+1} = \begin{cases} D_j + \text{gap}((x_{j+1})^2, \rho_j, 2^j), & \text{next letter even,} \\ \text{gap}((x_{j+1})^2, \rho_j, 2^j) + \text{gap}(x_j^{2^j}, D_j, 3), & \text{next letter odd.} \end{cases}$$

The *global defect* is $\Delta_w(n) = D_{|w|}$. An even letter keeps the old slack and lifts the new remainder through 2^j . An odd letter first cubes the running slack and then lifts the new remainder. In particular Δ is not the sum of the local remainders.

Theorem 2.4 (global defect identity). If w is realized at n and $m = J^{|w|}(n)$, then

$$n^{3^{\#O(w)}} = m^{2^{|w|}} + \Delta_w(n), \quad \Delta_w(n) \geq 0.$$

Theorem 2.2 is the inequality $\Delta_w(n) \geq 0$.

Proof. The empty word is $n = n + 0$. Suppose a realized prefix of length ℓ with odd count o ends at x and satisfies $n^{3^o} = x^{2^\ell} + D$.

If the next letter is even, then $x = J(x)^2 + \rho(x)$, so

$$n^{3^o} = (J(x)^2 + \rho(x))^{2^\ell} + D = J(x)^{2^{\ell+1}} + D + \text{gap}(J(x)^2, \rho(x), 2^\ell).$$

The odd count is unchanged, and the new slack is the even update.

If the next letter is odd, then $x^3 = J(x)^2 + \rho(x)$. Cubing the inductive identity gives

$$n^{3^{o+1}} = (x^{2^\ell} + D)^3 = x^{3 \cdot 2^\ell} + \text{gap}(x^{2^\ell}, D, 3).$$

The leading term is

$$x^{3 \cdot 2^\ell} = (J(x)^2 + \rho(x))^{2^\ell} = J(x)^{2^{\ell+1}} + \text{gap}(J(x)^2, \rho(x), 2^\ell),$$

which is the odd update. $\square$

For a one-letter illustration, take $n = 3$ and $w = O$. Then $J(3) = 5$ and $3^3 = 27 = 5^2 + 2$, so $\Delta_O(3) = 2$. After a later letter the lift is a power-gap, not a sum of remainders.

Theorem 2.5 (vanishing). If w is realized at n , the following are equivalent.

1. $\Delta_w(n) = 0$.
2. Every local remainder along w vanishes.
3. $(J^{|w|}(n))^{2^{|w|}} = n^{3^{\#O(w)}}$.

In that case w is monochrome. More precisely, either $w = E^k$ and $n = a^{2^k}$ for an even a , or $w = O^k$ and $n = a^{2^k}$ for an odd a . A realized mixed word therefore has $\Delta_w(n) > 0$.

Proof. The identity of Theorem 2.4 gives (1) $\Leftrightarrow$ (3). For (1) $\Leftrightarrow$ (2), use the recurrence. If $e \geq 1$, a power-gap vanishes if and only if its addend vanishes. The even update is a sum of two nonnegative terms, and the odd update is a sum of two power-gaps; either vanishes only when the incoming slack and the new remainder both vanish. Thus $D_{|w|} = 0$ forces $D_0 = 0$ and every $\rho_j = 0$. The converse is immediate.

If every remainder vanishes, an even step satisfies $x_j = x_{j+1}^2$, while an odd step satisfies $x_j^3 = x_{j+1}^2$. In either case x_j and x_{j+1} have the same parity, so the itinerary is monochrome. If $w = E^k$, repeated substitution gives $n = x_k^{2^k}$; writing $a = x_k$, its parity is even. If $w = O^k$, unique factorization applied to $x_j^3 = x_{j+1}^2$ shows that every prime valuation of x_j is even. Iterating this valuation relation shows that every prime valuation of $n = x_0$ is divisible by 2^k , hence $n = a^{2^k}$ for an odd a . Conversely, these even and odd towers make every local remainder zero. $\square$

Theorem 2.6 (composition). If u is realized at n and v is realized at $m = J^{|u|}(n)$, then

$$\Delta_{uv}(n) = \text{gap}(m^{2^{|u|}}, \Delta_u(n), 3^{\#O(v)}) + \text{gap}((J^{|v|}(m))^{2^{|v|}}, \Delta_v(m), 2^{|u|}).$$

In particular $\Delta_v(m)^{2^{|u|}} \leq \Delta_{uv}(n)$. Every local remainder satisfies $\rho_j^{2^j} \leq \Delta_w(n)$.

Proof. Write $o_u = \#O(u)$ and $o_v = \#O(v)$. Theorem 2.4 on u and on uv gives

$$n^{3^{o_u+o_v}} = (m^{2^{|u|}} + \Delta_u(n))^{3^{o_v}} = (J^{|uv|}(n))^{2^{|uv|}} + \Delta_{uv}(n).$$

The first power expands as

$$m^{2^{|u|} \cdot 3^{o_v}} + \text{gap}(m^{2^{|u|}}, \Delta_u(n), 3^{o_v}).$$

Theorem 2.4 on v at m rewrites the leading term as

$$\left((J^{|v|}(m))^{2^{|v|}} + \Delta_v(m) \right)^{2^{|u|}},$$

which expands as the second gap plus $(J^{|uv|}(n))^{2^{|uv|}}$. Comparing the two expressions for $n^{3^{o_u+o_v}}$ yields the identity. The suffix inequality is $\rho^e \leq \text{gap}(a, \rho, e)$ on the second summand. For a local remainder at index j , split w after j letters: the suffix defect is at least ρ_j , and raising through 2^j gives the stated bound. $\square$

Composition is polynomial, not additive: later odd letters raise the prefix slack to the third power. The naive recurrence $\Delta \leftarrow \Delta + \rho$ is false.

Corollary 2.7 (cycle surplus). If w is a cycle word at n , then

$$\Delta_w(n) = n^{3^{\#O(w)}} - n^{2^{|w|}}$$

exactly.

Proof. Theorem 2.4 with $m = n$. $\square$

The identity does not supply a state-independent positive tax, and none exists. For one realized letter at x with branch exponent $e \in \{1, 3\}$,

$$x^e < (J(x) + 1)^2,$$

so the relative slack $1 + \eta = x^e / J(x)^2$ satisfies

$$\eta < \frac{2}{J(x)} + \frac{1}{J(x)^2},$$

which tends to 0 as the state grows. The recorded extreme is an *OOE* block at $n = 180370579261640036336071806107777 \approx 1.80 \cdot 10^{32}$ whose word-relative slack

$$q_w(n) = \frac{n^{3^{\#O(w)}}}{(J^{|w|}(n))^{2^{|w|}}} - 1$$

satisfies $0 < q_{OOE}(n) < 10^{-30}$, by the exact integer comparison $0 < (n^9 - J^3(n)^8) 10^{30} < J^3(n)^8$.

3. Inverse cells and the census

The exact one-step fibers are

$$J(n) = q \iff q^2 \leq n < (q+1)^2 \quad (n \text{ even})$$

and

$$J(n) = m \iff m^2 \leq n^3 < (m+1)^2 \quad (n \text{ odd}).$$

Lemma 3.1 (odd cells are unique). An odd fiber contains at most one integer. An even fiber is a parity-restricted square interval and may contain many predecessors.

Proof. Suppose $a < b$ lie in the same odd cell indexed by m . Then $(a+1)^3 \leq b^3 < (m+1)^2$ and $m^2 \leq a^3$. Subtracting the latter lower bound from the former upper bound gives

$$3a^2 + 3a + 1 = (a+1)^3 - a^3 < (m+1)^2 - m^2 = 2m + 1,$$

hence $3a^2 + 3a < 2m$ and $3a^2 < 2m$. If $a = 0$, then $m = 0$ and the displayed strict inequality is already impossible. If $a > 0$, squaring and using $m^2 \leq a^3$ gives $9a^4 < 4m^2 \leq 4a^3$, contradicting $4a^3 < 9a^4$. $\square$

Theorem 3.2 (cycle restrictions). Let w be a cycle word at $n \geq 2$.

- (i) The word is formally expanding:

$$2^{|w|} < 3^{\#O(w)}.$$

A contracting word cannot close a nontrivial cycle.

- (ii) The cycle minimum is odd and the cycle maximum is even. A minimum-based orientation cannot end in an odd letter.

- (iii) A realized word v is *superquadratic* if

$$3^{\#O(v)} \geq 2^{|v|+1}.$$

Any realized path from a start $n \geq 2$ to a state at least n^2 is superquadratic. On a cycle minimum the path to any later even state is superquadratic. The prefix OOE is expanding ($9 > 8$) but not superquadratic ($9 < 16$), so it cannot carry the minimum to square scale.

Proof. For (i), Theorem 2.2 applied to a cycle endpoint gives $n^{2^{|w|}} \leq n^{3^{\#O(w)}}$. Since $n \geq 2$, comparison of exponents gives $2^{|w|} \leq 3^{\#O(w)}$; equality is impossible because the two sides have different prime divisors and $|w| \geq 1$. Alternatively: a mixed cycle word has $\Delta_w(n) > 0$ by Theorem 2.5, so the envelope is strict; a monochrome tower cannot return for $n \geq 2$.

Every state on such a cycle is at least 2: once an orbit reaches 1, it remains there and cannot return to a start $n \geq 2$. An even state $x \geq 2$ satisfies $J(x) < x$, so a cycle minimum cannot be even. An odd state $x \geq 3$ satisfies $J(x) > x$, so a cycle maximum cannot be odd. This proves the first assertion of (ii). For the final-letter assertion, let x be the predecessor of a minimum-oriented return n . If the last letter were odd, the odd return cell would give $n^2 \leq x^3 < (n+1)^2$. Minimality gives $x \geq n$, hence $n^3 < (n+1)^2$, impossible for $n \geq 3$; and the minimum is odd, so $n \geq 3$.

For (iii), let a realized word v send n to $y \geq n^2$. Theorem 2.2 gives

$$n^{2^{|v|+1}} = (n^2)^{2^{|v|}} \leq y^{2^{|v|}} \leq n^{3^{\#O(v)}}.$$

Since $n \geq 2$, comparison of exponents proves the superquadratic inequality. On a cycle minimum, if a later state y is even, its successor is at least n . Thus $n \leq J(y) = \lfloor \sqrt{y} \rfloor$, so $n^2 \leq y$, and the preceding argument applies to that prefix. $\square$

Lemma 3.3 (coarse lower envelope). For $n \geq 1$, write $q = \lfloor \sqrt{n} \rfloor$. Then $q^2 \leq n < (q+1)^2$. For $q \geq 1$ one has $(q+1)^2 \leq 4q^2$, and the second inequality is strict for $q \geq 2$, while for $q = 1$ one has $n < 4$. In all cases

$$n < 4 \lfloor \sqrt{n} \rfloor^2.$$

Thus an even step satisfies $n \leq 4J(n)^2$ and an odd step satisfies $n^3 \leq 4J(n)^2$. Composing along a realized word v of length k with odd count o gives

$$n^{3^o} \leq C_v J^k(n)^{2^k}.$$

The constant starts at 1 and updates by $C \mapsto C \cdot 4^{2^j}$ on an even letter and $C \mapsto C^3 \cdot 4^{2^j}$ on an odd letter, at step j . In particular $C_{OOO} = 2^{38}$ and $C_{OOOO} = 2^{130}$.

Proof. The comparison $n < 4q^2$ is the paragraph above. The composition law is the same recurrence as the proof of Theorem 2.2, with a factor 4^{2j} inserted at each letter. The values C_{OOO} and C_{OOOO} are the result of that recurrence on those two words. $\square$

Lemma 3.4 (next-square thresholds). (i) If $q \geq 5$ realizes OO , then $J^2(q) \geq (q+1)^2$. (ii) If $q \geq 3$ realizes OOO , then $J^3(q) \geq (q+1)^2$. (iii) If a realized word v satisfies $J^{|v|}(q) \geq (q+1)^2$ and the next realized letter is odd, then $J^{|v|+1}(q) \geq (q+1)^2$. (iv) If vE is a cycle word at n , then $J^{|v|}(n) < (n+1)^2$. (v) If $a \geq 3$, then O^aE is not a cycle word at any $n \geq 2$.

Proof. For (i), write $m = \lfloor q^{3/2} \rfloor$. The bound $J^2(q) \geq (q+1)^2$ follows from $m^3 \geq (q+1)^4$, because then $\lfloor m^{3/2} \rfloor \geq (q+1)^2$. If $q = 5$, then $11^2 = 121 \leq 125 = 5^3 < 144 = 12^2$, so $m = 11$, and $11^3 = 1331 \geq 6^4 = 1296$. Since q realizes OO , it is odd, so the only remaining case is $q \geq 7$. Then $m \geq q^{3/2} - 1$, so it is enough that $(q^{3/2} - 1)^3 \geq (q+1)^4$. Expanding the left side and dropping the positive remainder $3q^{3/2} - 1$ reduces this to $q^{9/2} - 3q^3 \geq (q+1)^4$, or equivalently $\sqrt{q} - 3/q \geq (1 + 1/q)^4$. The left side increases for $q > 0$ and the right side decreases, so the case $q = 7$ suffices:

$$\sqrt{7} - \frac{3}{7} > \frac{5}{2} - \frac{3}{7} = \frac{29}{14} > \frac{4096}{2401} = \left(\frac{8}{7}\right)^4,$$

where $\sqrt{7} > 5/2$ follows from $25/4 < 7$.

For (ii), the orbit $3 \rightarrow 5 \rightarrow 11 \rightarrow 36$ gives $J^3(3) = 36 \geq 16$. If $q \geq 5$ realizes OOO , then it realizes OO , so (i) gives $J^2(q) \geq (q+1)^2$. The third letter is odd, hence $J^3(q) = J(J^2(q)) \geq J^2(q)$.

For (iii), the image after v is odd and at least $(q+1)^2 \geq 4$, so the odd branch does not decrease it.

For (iv), the last letter is even, so the preimage $z = J^{|v|}(n)$ is even and satisfies $n^2 \leq z < (n+1)^2$.

For (v), suppose O^aE is a cycle word at $n \geq 2$. The start is odd, hence at least 3, and realizes O^a . Parts (ii) and (iii) give $J^a(n) \geq (n+1)^2$, contradicting (iv). $\square$

Lemma 3.5 (two length-six exclusions). Neither $OOOEEOE$ nor $OOOOEEE$ is a cycle word at any $n \geq 2$.

Proof. First, if $n \geq 256$, then

$$n^{81} > 2^{130}(n+1)^{64}.$$

Indeed $257^{64} < 2 \cdot 256^{64}$ because $(1 + 1/256)^{64} < e^{64/256} = e^{1/4} < 2$, and $256(n+1) \leq 257n$, so $(n+1)^{64} < 2n^{64}$. Then $2^{130}(n+1)^{64} < 2^{131}n^{64}$. Since $n \geq 256 = 2^8$, one has $n^{17} \geq 2^{136}$, and therefore $2^{131}n^{64} < 2^{136}n^{64} \leq n^{81}$.

For $2 \leq n < 256$, neither word returns to its start. This is a table of 254 evaluations of each word: at every start, either some letter fails to match the current parity, or the six-step image differs from the start. The same finite check is the Lean `native_decide` evaluation behind `no_cycle_word_ooooee` and `no_cycle_word_ooooee` (Appendix A).

Now suppose $OOOOEEE$ is a cycle word at $n \geq 256$, and write $z = J^4(n)$ for the image after the prefix $OOOO$. Lemma 3.4(iv) gives $J(z) < (n+1)^2$, and the preceding even letter gives $z < (J(z)+1)^2$, hence $z < (n+1)^4$. Lemma 3.3 on $OOOO$ gives $n^{81} \leq 2^{130}z^{16}$, so $n^{81} < 2^{130}(n+1)^{64}$, contradicting the tail inequality.

Finally suppose $OOOEEOE$ is a cycle word at $n \geq 256$. Write $z_3 = J^3(n)$ and $y = J(z_3) = \lfloor \sqrt{z_3} \rfloor$, so $z_3 < (y+1)^2$. Lemma 3.3 on OOO gives $n^{27} \leq 2^{38}z_3^8 < 2^{38}(y+1)^{16}$. Cubing yields $n^{81} < 2^{114}(y+1)^{48}$. The last letters OE give the odd-cell bound $y^3 < (n+1)^4$. Write $A = n+1 \geq 257$.

We claim $(y+1)^3 < 2A^4$. If $y \leq A$, this is $(A+1)^3 < 2A^4$. If $y > A$, then $y \geq 258$, so $3y+1 < y^2$ and hence $(y+1)^3 = y^3 + 3y^2 + 3y + 1 < A^4 + 4y^2$, while $4y^2 < A^4$ because $4y^3 < 4A^4 \leq yA^4$. Raising $(y+1)^3 < 2A^4$ to the sixteenth power gives $(y+1)^{48} < 2^{16}(n+1)^{64}$. Combining with the cubed lower envelope produces again $n^{81} < 2^{130}(n+1)^{64}$. $\square$

Theorem 3.6 (small-cycle census). No word of length at most six is a cycle word at any $n \geq 2$. Equivalently, a nontrivial Juggler cycle, if one exists, has period at least seven.

Proof. Rotating a cycle word by one letter moves the base point one step along the orbit and yields another cycle word; every state of the cycle is at least 2, because an orbit that reaches 1 stays at 1 and cannot return to a start $n \geq 2$.

If every letter is odd, the start is odd, hence $n \geq 3$, and the odd branch strictly increases there: $J(x) > x$ for odd $x \geq 3$, since $x^3 \geq (x+1)^2$. The orbit ascends strictly and never returns. Otherwise some letter is even, and a rotation ending just after that letter produces an even-terminating cycle word vE of the same length based at a cycle state $m \geq 2$. It therefore suffices to exclude even-terminating cycle words of length at most six.

By Theorem 3.2(i) a cycle word is formally expanding. No even-terminating word of length one or two is expanding ($2 > 3^0$ and $4 > 3$).

Length three: the only expanding candidate is OOE . If $m \geq 5$ realizes OO , Lemma 3.4(i) gives $J^2(m) \geq (m+1)^2$, contradicting Lemma 3.4(iv). The only smaller odd start is $m = 3$, where $J^2(3) = 11$ is odd, so the final even letter is not realized.

Length four: the expanding filter requires three odd letters among the first three, leaving only O^3E , excluded by Lemma 3.4(v).

Length five: the filter requires four odd letters among the first four, leaving only O^4E , excluded by Lemma 3.4(v).

Length six: the filter requires at least four odd letters among the first five, leaving O^5E , $EOOOOE$, $OEEOOE$, $OOEOOE$, $OOOEEOE$, and $OOOOEE$. Lemma 3.4(v) excludes O^5E . The word $EOOOOE$ rotates one step onto $OOOOEE$, and $OEEOOE$ rotates two steps onto $OOOEEOE$; both are excluded by Lemma 3.5.

It remains to exclude $OOEOOE$. Let this word be a cycle word at some start, and rotate to a cycle minimum $m \geq 2$. The three alignments of the two even letters are $OOEOOE$, $OEOOEO$, and $EOOEEO$. The last starts even, so it cannot be a minimum, by Theorem 3.2(ii). The middle starts OE : the first image is even and strictly below m^2 , whereas every even state after a cycle minimum is at least m^2 , as proved in Theorem 3.2(iii). Thus the minimum orientation is $OOEOOE$. In particular m is odd, so $m \geq 3$. The prefix OOE must be realized. If $m = 3$, then $3 \rightarrow 5 \rightarrow 11$ and 11 is odd, so OOE is not realized. Hence $m \geq 5$. Write $y = J^3(m)$ for the state after OOE . Then $y \geq m$ by minimality, so $y \geq 5$. The suffix OO is realized at y , and Lemma 3.4(i) gives $J^2(y) \geq (y+1)^2 \geq (m+1)^2$. The last letter is even, so Lemma 3.4(iv) gives $J^2(y) < (m+1)^2$. $\square$

Lemma 3.4(v) excludes every odd-run-then-even word O^aE with $a \geq 3$, of any length. Length seven is the first even-terminating expanding length whose surviving candidates have an internal even letter (five odd letters among the first six). Those shapes are not reached by the recorded thresholds. No exclusion of cycles of length seven or more is claimed. The census stops at length six because length seven admits even-terminating expanding words with an internal even letter that none of the recorded thresholds reach; no exclusion at length seven is claimed.

4. Remarks

A start $n \geq 2$ has a *descent certificate* if there exists a realized finite word w with $J^{|w|}(n) < n$.

Theorem 4.1 (uniform short certificates). Let $n \geq 2$.

1. If n is even, then the one-letter word E is a descent certificate: $J(n) = \lfloor \sqrt{n} \rfloor < n$.
2. If n is odd and $J(n)$ is even, then the two-letter word OE is a descent certificate.

Proof. For (1), $\lfloor \sqrt{n} \rfloor \leq \sqrt{n} < n$ for $n \geq 2$. For (2), the word OE is realized by hypothesis, and

$$J^2(n) = \lfloor \sqrt{\lfloor n^{3/2} \rfloor} \rfloor \leq n^{3/4} < n$$

for $n \geq 2$. $\square$

The starts not covered by Theorem 4.1 are exactly the odd-to-odd starts. In particular, if $n \geq 2$ has no descent certificate of any length, then n is odd-to-odd. The converse is false: many odd-to-odd starts descend after a longer word. If every start above 1 has some descent certificate, ordinary strong induction yields arrival at 1. That hypothesis is not proved.

The first odd-to-odd image expands, so ordinary strong induction cannot fire on the complement of Theorem 4.1. A uniform run bound on expanding blocks is likewise unavailable: four consecutive expanding blocks occur already at

$$1999 \xrightarrow{OOE} 5169 \xrightarrow{OOOOEE} 50093 \xrightarrow{OOE} 193753 \xrightarrow{OOE} 887471,$$

and the relative slack of a single letter tends to 0 with the state (Section 2).

No theorem forces every exact integer state into a contracting prefix. In particular, it is open whether every start reaches 1, and open whether a nontrivial cycle of length seven or more exists.

Lean proofs of the theorems of Sections 2–4 are in the project repository. From a clone, `lake build Problems.JugglerPaper` builds only the modules named in Appendix A.

Appendix A. Lean names

The proofs in the text are ordinary integer arguments. The following names are the corresponding Lean theorems in `formal/Problems/Juggler/`, imported by `Problems.JugglerPaper`.

Text	Lean
itinerary semantics	<code>follows_iff_word</code> , <code>image_eq_iterate</code> , <code>image_append</code>
Theorem 2.1	<code>image_monotone_of_follows</code>
Theorem 2.2	<code>power_bound_word</code>
Corollary 2.3	<code>power_bound_contracts</code>
Theorem 2.4	<code>global_defect_identity</code>
Theorem 2.5	<code>global_defect_eq_zero_iff_localsTight</code> , <code>global_defect_eq_zero_implies_monochrome</code> , <code>power_bound_eq_iff_extremal</code>
Theorem 2.6	<code>global_defect_append</code>

Text	Lean
Corollary 2.7	<code>image_eq_start_defectRatio</code>
per-step slack	<code>one_plus_eta_lt_succ_sq</code>
Lemma 3.1	<code>odd_cell_unique</code>
Theorem 3.2	<code>cycle_word_formally_expanding,</code> <code>cycleMin_start_odd, cycleMax_start_even,</code> <code>cycleMin_not_end_odd,</code> <code>square_scale_superquadratic,</code> <code>cycleMin_to_even_superquadratic</code>
Lemma 3.3	<code>lower_growth_word</code>
Lemma 3.4	<code>oo_suffix_threshold,</code> <code>ooo_suffix_threshold,</code> <code>threshold_inherits_odd_append,</code> <code>cycle_last_even_interval,</code> <code>no_cycle_odd_run_append_even</code>
Lemma 3.5	<code>no_cycle_word_oooeoe,</code> <code>no_cycle_word_ooooee</code>
Theorem 3.6	<code>no_cycle_word_length_le_six</code>
Theorem 4.1	<code>even_finiteProgress,</code> <code>odd_even_finiteProgress</code>
no certificate $\Rightarrow$ odd-to-odd	<code>no_finiteProgress_implies_odd_odd</code>
induction to 1	<code>reachesOne_of_all_finiteProgress</code>
four-block chain	<code>four_block_pe_1999</code>

Acknowledgments

I used large language models extensively while drafting and revising the text, organizing companion notes, and as an interactive assistant for Lean statements, tests, and literature records. The models are not authors. Lean theorems and named computations are the certificates for the claims of Sections 2–4. I take full responsibility for the contents.

References

1. C. A. Pickover, *Computers and the Imagination: Visual Adventures Beyond the Edge*, St. Martin's Press, New York, 1991, ch. 40, p. 232.
2. C. A. Pickover, *The Mathematics of Oz: Mental Gymnastics from Beyond the Edge*, Cambridge University Press, Cambridge, 2002, ch. 45, pp. 102–106.
3. OEIS Foundation Inc., “Juggler sequence: if n even then $\lfloor \sqrt{n} \rfloor$ else $\lfloor n^{3/2} \rfloor$,” Sequence A094683 in *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org/A094683> (accessed 28 August 2026).
4. OEIS Foundation Inc., “Number of steps needed for n to reach 1 in the juggler sequence,” Sequence A007320 in *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org/A007320> (accessed 29 August 2026).
5. E. W. Weisstein, “Juggler Sequence,” *MathWorld*, <https://mathworld.wolfram.com/JugglerSequence.html> (accessed 29 August 2026).
6. OEIS Foundation Inc., “Largest value in trajectory of n under the juggler map of A094683,” Sequence A094716 in *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org/A094716>

(accessed 29 August 2026).

7. V. Prasad and M. A. Prasad, “Estimates of the maximum excursion constant and stopping constant of juggler-like sequences,” ResearchGate preprint, 2025. <https://doi.org/10.13140/RG.2.2.14110.04168>.
8. J. C. Lagarias, “The $3x+1$ problem and its generalizations,” *Amer. Math. Monthly* 92 (1985), 3–23. doi:10.1080/00029890.1985.11971528.
9. J. C. Lagarias (ed.), *The Ultimate Challenge: The $3x+1$ Problem*, American Mathematical Society, Providence, RI, 2010.
10. R. E. Crandall, “On the “ $3x+1$ ” problem,” *Math. Comp.* 32 (1978), 1281–1292. doi:10.1090/S0025-5718-1978-0480321-3.
11. K. R. Matthews and A. M. Watts, “A generalization of Hasse’s generalization of the Syracuse algorithm,” *Acta Arith.* 43 (1984), 167–175. doi:10.4064/aa-43-2-167-175.
12. P. Cochin, “Parity equidistribution of nested floor powers, with descent applications to the Juggler map,” companion manuscript, in preparation, 2026.