

Fate Contagion and Termination Criteria for the Juggler Map

Philippe Cochin

philippe@cochin.fr

ORCID 0009-0004-1939-3382

22 September 2026

Abstract

The Juggler map sends an even positive integer to the integer part of its square root and an odd positive integer to the integer part of its three-halves power. We prove that every nonempty set A of positive integers closed under taking preimages satisfies $\sum_{n \in A, n \leq x} 1/n \geq c(\log x)^{5/8}$ for all sufficiently large x . Thus any realized cycle basin or nonempty class of unbounded orbits must have this divergent logarithmic mass. Three disjoint actual predecessor families, with initial words E, OE and OOOE, supply the recursion. The OOOE input is an AI-assisted written analytic proof, included in Appendix E, pending independent review and complete Lean verification.

The earlier two-production argument remains a separate, fully machine-checked baseline for every $0 < \lambda \leq 100/203$. Its poor-fiber tail needs no exponential sums. For the stronger exponent $5/8$, Lean checks the recurrence assembly with the actual OOOE production bound explicit; OE and the main mixed-frequency estimates are proved separately. We distinguish that conditional formal result from the complete written argument.

For a fixed verified target $[1, N_0]$, universal termination is equivalent to an eventual-entry statement: all but $O(y(\log y)^{-e})$ odd starts in $(y, 2y]$ enter that target, for some $e > 3/8$. The rate itself remains unproved. Parity-cylinder, live-pressure and scale-averaged pressure bounds are sufficient arithmetic inputs; none is established here. The unconditional Lean baseline gives the rate threshold $103/203$. Earlier finite-production and localized-discrepancy arguments are retained with their original scope, and the numerical experiments remain observations. Neither universal termination nor exclusion of every nontrivial cycle or unbounded orbit is established.

2020 Mathematics Subject Classification: 11B37, 11L07, 37P99.

Keywords: Juggler map; preimage sets; logarithmic counting; parity cylinders; integer dynamics.

1. Introduction

For a positive integer n , define

$$J(n) = \begin{cases} \lfloor \sqrt{n} \rfloor, & n \text{ even,} \\ \lfloor n^{3/2} \rfloor, & n \text{ odd.} \end{cases}$$

The Juggler sequence was introduced by Pickover [1] and is sequence A094683 of the OEIS [2]. Starting from any positive integer, the orbit under J appears always to reach 1; this has been verified for all starts up to $3.5 \cdot 10^8$ [11]. The map belongs to the family of floor-power maps, and its dynamics resembles the $3x + 1$ problem [3, 4] in one respect and differs from it in another. The resemblance: a parity decides whether the orbit goes up or down, and the heuristic model of the orbit is a random walk with negative drift, here on the scale $\log \log n$

with steps $+\log_2 3 - 1$ (odd) and -1 (even). The difference: the Juggler map moves by *powers*, so a single contracting certificate sends n to n^e with $e < 1$, and the one-step preimages of an integer are not a residue class but an interval.

This paper is about the second feature and what it forces. The narrative has four steps.

Contagion. Let R be the set of starts that reach 1, F its complement, $B(C)$ the basin of a nontrivial cycle C , and D the set of divergent starts. Each is *backward-closed*: if $J(n)$ belongs to it, so does n . Theorem 1 proves that every nonempty backward-closed set is large in logarithmic density, with an explicit exponent, by a downward recursion over the three actual productions E, OE and OOE (Section 5.9 and Appendix E). The two-production proof of Section 5.8 supplies the fully formal baseline. Applied to the fate classes: if a single start fails to reach 1, the failures have logarithmic count $\gg (\log x)^\lambda$ and, on infinitely many dyadic blocks, natural density $\gg (\log y)^{\lambda-1}$, for every $0 < \lambda \leq 5/8$. This excludes no fate; it fixes the quantitative shape of the trichotomy.

Odd generation. A set that is closed both forwards and backwards and avoids 1 is the E -forest (iterated even preimages) over its odd members, and its odd members are the odd preimages of its intersection with the sparse set $S = \{\lfloor m^{3/2} \rfloor : m \text{ odd}\}$ (Theorem 2). An arbitrary failure set is thereby replaced by sparse odd-image seeds together with deterministic even ancestry. This is the geometry that the later sections use.

The almost-all equivalence. Tao [6] proved that almost all Collatz orbits (in logarithmic density) attain almost bounded values, with a target $f(N) \rightarrow \infty$ arbitrarily slowly. For the Juggler map, contagion and odd generation together turn a bounded-target statement into the conjecture: every positive integer reaches 1 if and only if all but $O(y(\log y)^{-e})$ odd starts in $(y, 2y]$ enter $[1, N_0]$, for some $e > 3/8$ (Theorem 3). The threshold is the complement of the contagion exponent. The Collatz preimage lower bound of Krasikov–Lagarias [7], discussed with its target restriction in Section 1.3, does not by itself yield this implication.

Sufficient estimates at growing depth. The exponent walk needs to fall by $L(y) = \log_2(\log(2y)/\log N_0)$ to put a start in $(y, 2y]$ below the floor. A Chernoff estimate counts words whose walk fails to cross this level by depth $\lceil CL(y) \rceil$. Turning this word count into a count of integer starts requires additional arithmetic information. Sections 8–9 give three sufficient forms of that information: upper bounds on bad cylinders, one-sided control of their continuations, and a bound on one exponential moment of live starts. A bound on tilted odd shares implies the moment bound. None is established here.

The first-letter decomposition in Section 6 relates the remaining OO -contribution to failures at a larger scale. Section 10 connects this term with live-start counts and states a uniformity requirement for a particular cylinder-counting method. These are reductions and limitations of the stated estimates, not necessity results for every possible proof of termination.

1.1 Main results

Write $\ell_A(u, v] = \sum_{n \in A, u < n \leq v} 1/n$. Throughout, $N_0 \geq 2$ is any integer with $[1, N_0] \subseteq R$ (the Lean-verified value is 260; the certified computational value is $3.5 \cdot 10^8$ [11]).

Theorem 1 (fate contagion). Let $A \subseteq \mathbb{N}$ be nonempty and backward-closed, and let $0 < \lambda \leq 5/8$. There are $c > 0$ and x_0 , depending on A and λ , with

$$\sum_{\substack{n \in A \\ n \leq x}} \frac{1}{n} \geq c (\log x)^\lambda \quad (x \geq x_0).$$

Consequently each fate class realized by at least one start — R , the basin of any existing nontrivial cycle, the divergent set — satisfies this bound, and on infinitely many dyadic blocks has natural density $\gg (\log y)^{\lambda-1}$. (Theorem 5.19 and its dyadic consequence.)

The new endpoint uses Appendix E’s written OOE estimate. Its complete analytic formalization and independent review remain outstanding. The fully machine-checked baseline

remains $100/203$. The two earlier routes are retained with their original exponents, as follows. The route of Sections 4–5 and Appendix D — the block average of Proposition 4.4, the monotone sweep and the six finite productions of Section 5.7 — gives every $\lambda < \lambda^{**} \approx 0.4926$, the root of

$$2^{-\lambda} + \frac{1}{9}\left(\frac{3}{8}\right)^\lambda + \frac{2}{9}\left(\frac{3}{4}\right)^\lambda + \sum_{k=2}^6 3^{-(k+1)} \left(\frac{1}{2}\left(\frac{3}{4}\right)^k\right)^\lambda = 1. \quad (1.1)$$

(The V_5 truncation of the first seven terms is 0.4924; the V_4 truncation of the first six terms is 0.4916; the V_3 truncation of the first five is 0.4891; the $OEOEE$ truncation of the first four is 0.4801; the pairing-only root of the first three is 0.4480.) The two-production route of Section 5.8 uses no exponential sum: the fibers whose parity share is far from $\frac{1}{2}$ have finite total logarithmic mass, so the pointwise coefficient $\frac{2}{9}$ of the OE family may be replaced by the averaged one, and the recursion reaches every $\lambda < \lambda_{\text{ideal}}$, where $2^{-\lambda_{\text{ideal}}} + \frac{1}{3}(3/4)^{\lambda_{\text{ideal}}} = 1$ and $\lambda_{\text{ideal}} \approx 0.4927$. That route is formalized in Lean 4 with no hypothesis for every $\lambda \leq 100/203 = 0.49261\dots$, which exceeds λ^{**} ; the range $100/203 < \lambda < \lambda_{\text{ideal}}$ is one choice of a parameter η_0 away and is not separately certified. The formal infinite series of the finite productions has the same limiting root λ_{ideal} . The ideal characteristic model in Proposition 5.12 has root 1; the other run-model values, including 0.8414, are conditional calculations and are not attained density exponents.

Theorem 2 (odd generation). Let A be forward- and backward-closed with $1 \notin A$. Every $n \in A$ reaches by a possibly empty sequence of even steps to an odd member $m \geq 3$ of A ; for odd n , $n \in A \iff \lfloor n^{3/2} \rfloor \in A$; and $A \neq \emptyset$ iff A contains the image $\lfloor m^{3/2} \rfloor$ of some odd $m \geq 3$. In particular every positive integer reaches 1 iff no odd image fails to. (Theorem 6.1; Lean.)

Theorem 3 (the conjecture as an almost-all statement). The following are equivalent: (i) every positive integer reaches 1; (ii) for some $0 < \lambda \leq 5/8$, the starts $n \leq x$ whose orbit never enters $[1, N_0]$ have logarithmic count $o((\log x)^\lambda)$; (iii) for some $e > 3/8$ (the unconditional Lean baseline is $103/203$) and all large y , $\#\{n \text{ odd} \in (y, 2y] : n \notin R\} \leq y(\log y)^{-e}$. (Corollary 7.1, Theorems 7.2, 7.3.)

Theorem 4 (the frontier reduction). Let $L(y) = \log_2(\log 2y / \log N_0)$, $d(y) = \lceil CL(y) \rceil$, $p_C = (1 - 1/C) / \log_2 3$, $e(C) = CD(p_C \| \frac{1}{2}) / \ln 2$, and $e_q^{\text{Az}}(C) = 2(C(1 - q \log_2 3) - 1)^2 / (C(\log_2 3)^2 \ln 2)$. Assume $C \geq 5$. The following hypotheses give the indicated, case-dependent bounds. Each implies (iii) of Theorem 3 whenever its displayed rate exceeds $3/8$, using the written Theorem 5.19. The constants C below are retained sufficient choices from the earlier threshold $1 - \lambda^{**}$; they are not asserted to be optimal for the stronger theorem:

- (a) *cylinder form* $H(C, A)$: no O -rooted, $L(y)$ -bad itinerary cylinder of depth $d(y)$ exceeds its fair share $2^{-(d-1)}y/2$ among odd starts by more than $y(\log y)^{-A}$, $A > C + e(C)$ (Theorem 8.3), giving $e(C) - \varepsilon$ and hence the conjecture for $C \geq 19$ using Theorem 1 (or $C \geq 18$ under Appendix C);
- (b) *one-sided form* $H_q(C, A)$, with $0 < q < \log 2 / \log 3$ and $C > 1/(1 - q \log_2 3)$: every $L(y)$ -bad cylinder of depth $1 \leq t < d(y)$ sends at most a fraction q of its members, plus $y(\log y)^{-A}$, to an odd next state (Theorem 9.1; $A > C + e_q^{\text{Az}}(C)$), giving $e_q^{\text{Az}}(C) - \varepsilon$; at $q = 0.55$, $C \geq 41$ crosses Theorem 1's threshold;
- (c) *pressure form* $P_\theta(C)$: $\frac{1}{N} \sum_{n \text{ odd} \in (y, 2y], \tau(n) > d} e^{\theta o_d(n)} \leq (\frac{1}{2}(1 + e^\theta))^d e^{o(d)}$ at $\theta = \log(p_C / (1 - p_C))$, with $N = y/2$, where τ is the entrance time into $[1, N_0]$ and o_d the odd count of the first d letters; this gives $e(C) - \varepsilon$. Its biased *no-momentum form*, for $0 < q < p_C < 1$, $\sum_{t < d} (s_\theta(t) - q)^+ = o(d)$ gives $CD(p_C \| q) / \ln 2 - \varepsilon$ only at the optimizing tilt $\theta = \log(p_C(1 - q) / (q(1 - p_C)))$ (Theorem 9.2, Proposition 9.3).

None of these hypotheses is proved here. A bounded loss at each of $o(\log \log y)$ exceptional depths can be absorbed in the moment criterion. The tower comparison and finite-depth model in Section 9.3 describe the scope of this argument; neither is an arithmetic necessity theorem.

Theorem 5 (first-letter decomposition). A two-way closed set has the disjoint first-letter decomposition of Section 6.2. In the normalization used there, its logarithmic density satisfies (6.1), with the stated boundary error. For the failure set F , the free term ψ_F is a known normalization factor times the decreasing limit of the live proportion among OO -type starts (Proposition 10.1). Each sufficient estimate of Theorem 4 bounds this term at its stated logarithmic rate. The upper recursion obtained by bounding the fiber weights gives decay estimates with a critical exponent; that upper inequality alone does not force the failure set to be empty (Corollary 10.2).

Proposition (depth-uniformity budget). Suppose the error bound available for each cylinder is $K2^{\kappa d}y^{1-\delta_0 2^{-cd}}$, with $K, \delta_0, c > 0$ and $\kappa \geq 0$ independent of the depth. The resulting union-bound error is negligible at depth $C \log_2 \log y + O(1)$ precisely when $cC < 1$ (Proposition 10.3). This is a criterion for the available error majorant, not a lower bound on the actual discrepancy.

Under a localization of the triple parity discrepancy of nested floor powers to sub-dyadic intervals — stated as an explicit hypothesis in Appendix C; Paper B [12] lists the short-interval localization it needs among its open questions — the earlier two-production exponent improves to $\lambda^{***} \approx 0.5392$, the rate threshold of Theorem 3 to 0.4608, and the least depth constant of Theorem 4 to $C \geq 18$. This conditional comparison is now weaker than the written Theorem 5.19. Nothing in Sections 2–12 depends on Appendix C.

1.2 The three fates as currently constrained

The three fate classes are the three Moirai: Atropos cuts the thread (the class R), Lachesis measures an allotted length (a cycle basin), Clotho spins without end (divergence). What is known about each, all from outside this paper:

- *Atropos.* Every $n \leq N_0 = 3.5 \cdot 10^8$ reaches 1 (independently certified computation, [11, Section 5]); the Lean-verified floor is $N_0 = 260$. Paper B [12] proves that 7/8 of all starts descend below themselves within five steps, with a power saving on dyadic blocks.
- *Lachesis.* A nontrivial cycle has minimum above $3.5 \cdot 10^8$ and period at least 780239 [11]; it has at least four even steps; if $3^o/2^L = 1 + \Lambda$, its odd share is $\log 2 / \log 3 + \log(1 + \Lambda) / (L \log 3)$, strictly above but, under the finance bound, very near the critical zero-drift share.
- *Clotho.* A divergent orbit has unbounded exponent walk; its record jumps are quantized to the $\log_2 3$ lattice [11]. Nothing excludes it.

Theorem 1 adds one sentence to each: whichever of the three acts at all acts on a set of logarithmic count $\gg (\log x)^\lambda$ for every $0 < \lambda \leq 5/8$, with the review boundary of Appendix E.

1.3 Related work

Collatz parity vectors. Terras [4] and Everett [5] showed that the first k parities of the Collatz orbit of n are determined by $n \bmod 2^k$ and equidistributed exactly, and deduced that almost all starts have finite stopping time (descend below themselves). The analogous exact statement fails for the Juggler map: the parities of $\lfloor n^{3/2} \rfloor, \lfloor \lfloor n^{3/2} \rfloor^{3/2} \rfloor, \dots$ are Diophantine, not algebraic, and their equidistribution is proved only through the words of length five (Paper B, [12]).

Tao's theorem. Tao [6] proved that for any $f(N) \rightarrow \infty$, almost all Collatz orbits (in logarithmic density) attain a value below $f(N)$. The bounded-target version is not available, and the renewal machinery through the Syracuse random variables is the heart of the proof. Section 7.1 explains, as a structural analogy and not as a transfer of methods, which parts of that picture change for the Juggler map.

Preimage trees. Krasikov and Lagarias [7] obtain an $x^{0.84}$ lower bound for Collatz preimage counts, for targets not divisible by 3 and sufficiently large x . Such a lower bound alone does not

give a divergent logarithmic count: a set whose counting function is $O(x^{0.84})$ has a convergent reciprocal sum. It therefore cannot by itself contradict a failure estimate of size $y(\log y)^{-e}$. For Juggler, Theorem 1 gives a divergent logarithmic lower bound for every nonempty preimage set.

Floor-power maps and exponential sums. The parity of $\lfloor n^{3/2} \rfloor$ along an arithmetic progression is a question about $\{n^{3/2}/2\}$; Section 4 uses only an elementary sweep argument and, on even blocks, the second-derivative test with Vaaler’s approximation [10], as for Piatetski-Shapiro sequences. Deeper nesting is Paper B’s subject.

Companion papers. Paper A [11] proves period lower bounds for a hypothetical cycle from a certified descent floor (cycle financing, walk-charge envelope); Paper B [12] proves that the five-step power-envelope certificates have density $7/8$, with a power saving, from the parity equidistribution of nested floor powers. Neither is reproved here; both enter as citations, and one statement of the type Paper B proves enters Appendix C as an explicit hypothesis.

1.4 Verification

Statements carry one of four roles. *Lean*: the exact combinatorial layer is formalized in Lean 4; the root `formal/Problems/JugglerFatePaper.lean` imports exactly the thirty-seven modules this paper cites and builds with `lake build Problems.JugglerFatePaper`, without `sorry` and without `native_decide`; `formal/AxiomCheckPaperC.lean` prints the axiom dependencies of every cited name and `AxiomCheckPaperC.expected` records them — Mathlib’s `propext`, `Classical.choice`, `Quot.sound` and nothing else (names in Appendix A). *Human proof*: the analytic and probabilistic counting. *Verified computation*: exact integer computations (the descent floor is Paper A’s; the closure of $[1, 260]$ under the two productions up to 10^9 is computed here). *Observation*: numerical experiments that check hypotheses and constants; they prove nothing and are labelled wherever they appear.

Result	Status
Fate classes closed; trichotomy; exclusion (Lemma 2.1)	Lean
Even block is an interval (Lemma 3.1)	Lean
OE fiber is an interval; cell identity (Lemma 3.2)	Lean
Odd generation (Theorem 6.1)	Lean
Envelope descent into the floor (Lemma 8.1)	Lean, on Paper A’s power envelope
Sweep lemma (Lemma 4.1), both half-cell conventions	Lean
Monotone pairing (Lemma 4.1’)	Lean
Fiber parity (Lemma 4.2), goodness unpacked	Lean, through Lemma 4.1’
Thin fibers (Lemma 4.3): the count on $(u, 2u]$ and the log-mass of the bad $m > U$	Lean
Block average (Proposition 4.4), asymptotic	human proof; the exact layer, the slow sum $O(m')$, the two-sided count of the block and the deduction of (4.1) and of the asymptotic form from the two remaining exponential-sum bounds are Lean, those two bounds are not
Share law and its three corollaries (Lemma 4.5, Corollary 4.6)	human proof; the phase expansion with its cubic remainder, the range of the quadratic phase with its threshold $[-\frac{5}{6}, \frac{1}{6}]$, and the integral $\frac{25}{108}$ are Lean, the equidistribution and the measure identifications are not

Result	Status
Production inequality (5.2) with the V -ladder terms of Section 5.7	human proof; the E -family and the OE -fiber family are Lean with explicit errors, the block-average family needs the two exponential-sum bounds and the ladder needs Appendix D
Theorem 5.3 at exponent $13/40$, unconditional: two productions at the pointwise coefficient, no hypothesis	Lean; superseded in exponent by the averaged route below
Block lock (Lemma 5.14): a fiber's parity share leaves $\frac{1}{2}$ only by locking onto a rational step of small denominator	Lean
Fiber lock (Lemma 5.15), by Dirichlet approximation	Lean
Resonance count, the arc count behind Theorem 5.16	Lean
Poor-fiber tail (Theorem 5.16) and the share on an arbitrary set (Corollary 5.17)	Lean
Theorem 5.18, contagion by two productions at the averaged coefficient: Theorem 5.3 for every $0 < \lambda \leq 100/203$ with no hypothesis, and Theorem 7.2 and Corollary 8.4 with their contagion hypothesis discharged at $e > 103/203$	Lean; the supremum λ_{ideal} and the range $100/203 < \lambda < \lambda_{\text{ideal}}$ are the choice of η_0 , human
Fate contagion at the failure set, log-mass form (Corollary 5.5(2), first clause)	Lean, unconditionally at exponent $100/203$ and, by the pointwise route, at $13/40$: the failure set is backward-closed by Lemma 2.1 and nonempty by assumption, so Theorem 5.18 applies to it
Corollary 5.4 (natural density, infinitely often): the dyadic pigeonhole, from the shell bound of Theorem 5.18	Lean, for every $0 < \lambda \leq 100/203$ with no hypothesis and with the constant $K/12$; the range $100/203 < \lambda < \lambda_{\text{ideal}}$ is human, as for Theorem 5.18
Corollary 5.5 at the three fate classes: the reach-one class, the failures, the basin of a cycle state and the divergent starts, each in log-mass form and in dyadic-block form	Lean, at exponent $100/203$, by instantiating Theorem 5.18 and Corollary 5.4 through Lemma 2.1
Production words $V_k = (OE)^{k-1}OEE$ are prefix-free (Section 5.7, Appendix D)	Lean, for the whole family and not only the six; the disjointness of source sets it feeds needs Appendix D's analytic layer, which is human
The side condition $e(C) > \frac{27}{40}$ of the unconditional criteria, at $C = 30$, by rational bounds on e , $\log 2$ and $\log_2 3$	Lean; the least such C (23) and the paper's own threshold through λ^{**} stay with the audit
Cube fibers full or alternating (Lemma 4.7)	Lean
Recursion lemma (Lemma 5.1)	Lean
Seed (Lemma 5.2)	Lean
Contagion (Theorem 5.3), given the production inequality (5.2), for every $\lambda \leq 0.49$	Lean; $\zeta(0.49) > 0$ by exact rational bounds, the root λ^{**} is human
Least failure is OO -type; first-letter trichotomy (Proposition 6.3(i), Section 6.2)	Lean
First-letter identity (6.1)	Lean, the exact three-piece decomposition and the uniqueness of the odd preimage; the normalization to $\varphi_A, \varphi_A^{\text{fib}}, \psi_A$ and the boundary error are human
Tao-type rate implies the conjecture (Theorem 7.2), with the contagion bound of Theorem 5.3 as a hypothesis	Lean

Result	Status
Almost-all equivalence (Theorem 7.3), with the contagion bound of Theorem 5.3 as a hypothesis	Lean
Corollary 8.4, the conjecture from a cylinder bound, with the contagion bound of Theorem 5.3 as a hypothesis	Lean
Chernoff count of bad words (Lemma 8.2, exact form)	Lean
Theorem 8.3, explicit form $y\Lambda^{-e(C)} + 2\Lambda^C y(\log y)^{-A}$ at every y	Lean; the ε -absorption into the displayed form is human
Pressure form (Theorem 9.2, exact: a pressure bound $Na_\theta^d E$ gives at most $Ne^{-dD(p_C\ 1/2)}E$ live starts)	Lean, on the live weight
Pressure telescoping (Proposition 9.3)	Lean, on the word-weight framework
One-sided form (Theorem 9.1, exact: at every y the odd failures of $(y, 2y]$ number at most $(xa_q^{d-1}N + (x-1)\text{err}(d-1)(2x)^{d-1})/x^{p_C d}$ for every tilt $x \geq 1$, by exponential moments)	Lean, without the martingale; the absorption of the error into the rate and the displayed asymptotic form are human
The conjecture from the one-sided hypothesis $H_q(C, A)$ (Theorem 9.1's consequence), with the contagion bound as a hypothesis, or with nothing else assumed when $e_q^{\text{Ch}}(C) > \frac{27}{40}$	Lean; the exponent is the Chernoff one of Proposition 9.3 with $A > C(1 + \log_2 x) + 1 + e$, the paper's numerical forms are human. The hypothesis $H_q(C, A)$ itself is doubtful: see the barrier word in Section 9.1
The conjecture from the pressure hypothesis $P_\theta(C)$ and from the no-momentum hypothesis $M_{\theta,q}(C)$ (Section 9.2's consequences), with the contagion bound as a hypothesis, or with nothing else assumed when the exponent exceeds $\frac{27}{40}$	Lean; the paper's $e^{o(d)}$ is quantified as $(\log y)^\varepsilon$ and its $o(d)$ as δd , the numerical forms are human
Theorem 9.1 with exceptional atoms (Section 10(d), first paragraph): the share bound may fail on bad atoms of total mass $y(\log y)^{-B}$ at each depth, and the conjecture still follows, with nothing else assumed when the exponent exceeds $\frac{27}{40}$	Lean; the condition is $B > C \log_2 x + 1 + e$ in place of the paper's $B > e_q(C)$, sufficient and not sharp
The bias energy of the $L(y)$ -bad words supplies the exceptional atoms (Section 10(d)): $\sum_{w \text{ bad}} D(w)^2 \leq (q - \frac{1}{2})^2 y^2 (\log y)^{-2B} / 2^t$ at the depths below $\lceil CL(y) \rceil$, at all large scales, gives the conjecture with nothing else assumed	Lean, by Cauchy–Schwarz on the bad atoms that violate the share bound; the exact enumeration of Section 11 finds that energy at its worst-case value from depth 16–20 at every computable scale, so the hypothesis has no numerical support
Collapsed-component bias (Section 10(d), after the measurement): the next-letter bias of the starts whose iterate lands in a window of values is an alternating sum of fiber sizes, bounded by the variation of the fiber profile, the even branch smooths it, $(v+1)(M_v - m_v) + 2M_v$ per value, and a sandwich on the profile propagates through even steps without amplification	Lean; the variation bound on a fiber profile and the parity of the odd-preimage mass are human
Azuma and exponential-moment arguments (Sections 8–10, except Lemma 8.2, Theorem 8.3, Theorem 9.1, Theorem 9.2 and Proposition 9.3 in their exact forms)	human proof
Cylinder-splitting identity for the first-letter bias (Section 10(d), second equality)	Lean; the Parseval form in Walsh sums, and the exceptional-atom estimate it is meant to supply, are human

Result	Status
Exact landing windows of the nested productions ((D.1), (D.2))	Lean; the smooth comparison (D.3), the multiplicities and the production inequality they feed are human
Localized triple discrepancy (Appendix C)	hypothesis, conditional
Theorem 5.19, Theorem 1 at $5/8$, and Theorems 3 and 7.2–7.3 at $e > 3/8$	written proof using Appendix E; Lean assembly conditional on the actual OOE production bound; independent analytic review outstanding
OOEE mixed modes on the actual short source interval	separate Lean proof in OOEEMixedModes; pure slow modes and final poor-target assembly are outside that result
Numerical experiments (Section 11)	observation




Figure 1: Logical dependencies. The arithmetic lemmas feed the contagion theorem. Each unproved parity hypothesis is a sufficient route to a time-bounded live count, which implies eventual termination only at the stated rate. No converse time bound is asserted.

2. Fate classes and closure

$\mathbb{N} = \{1, 2, \dots\}$. For $A \subseteq \mathbb{N}$ and $0 \leq u < v$ write

$$\ell_A(u, v] = \sum_{\substack{n \in A \\ u < n \leq v}} \frac{1}{n}, \quad g_A(t) = \ell_A(e^{t/2}, e^t].$$

For $A = \mathbb{N}$, $g(t) = t/2 + O(e^{-t/2})$.

Definition. A is *backward-closed* if $J(n) \in A$ implies $n \in A$, and *forward-closed* if $n \in A$ implies $J(n) \in A$.

Lemma 2.1 (fate classes). Let $R = \{n : \exists k, J^k(n) = 1\}$, $F = \mathbb{N} \setminus R$, $B(m) = \{n : \exists k, J^k(n) = m\}$, and $D = \{n : \forall B \exists k, J^k(n) > B\}$. Each of $R, F, B(m), D$ is backward-closed;

R, F, D are also forward-closed. Every $n \geq 1$ lies in R , or in $B(m)$ for some $m \geq 2$ on a cycle, or in D , and these possibilities exclude one another.

Proof. Backward closure is the definition of each set (for D : $J^k(J(n)) = J^{k+1}(n)$). Forward closure of R : if $J^k(n) = 1$ then $J^{k-1}(J(n)) = 1$ for $k \geq 1$, and $J(1) = 1$. Forward closure of D : given B , pick k with $J^k(n) > \max(B, n)$; then $k \geq 1$ and $J^{k-1}(J(n)) > B$. Trichotomy: a bounded orbit repeats, a repeat is a cycle, a cycle through 1 is the fate R . Exclusion: an orbit that reaches 1 or enters a cycle is bounded; an orbit that reaches 1 and passes through a cycle state $m \geq 2$ of period L would force $m = J^{qL}(m) = 1$. Lean: `reachesOne_backwardClosed`, `not_reachesOne_backwardClosed`, `ancestor_backwardClosed`, `escapes_backwardClosed`, `reachesOne_floorPower`, `escapes_floorPower`, `fate_trichotomy`, `reachesOne_not_escapes`, `cycle_basin_not_escapes`, `reachesOne_not_cycle_basin`. $\square$

Throughout Sections 3–5, A is a nonempty backward-closed set. Every statement applies to each fate class realized by at least one start.

3. The two productions

Lemma 3.1 (even block). Let $m \in A$. Every even n with $m^2 \leq n < (m+1)^2$ lies in A . Writing $E(m)$ for this set, $|E(m)| \geq m$ and

$$\frac{1}{m} \left(1 - \frac{2}{m}\right) \leq \frac{m}{(m+1)^2} \leq \sum_{n \in E(m)} \frac{1}{n} \leq \frac{m+1}{m^2}.$$

Proof. $J(n) = \lfloor \sqrt{n} \rfloor = m$ exactly on that interval, so $n \in A$. The interval has $2m+1$ integers, at least m and at most $m+1$ of them even, each between m^2 and $(m+1)^2$. Lean: `even_block_mem`, `even_block_card`. $\square$

Lemma 3.2 (OE fiber). Let $m \in A$ and


$$\Phi(m) = \{n \text{ odd} : m^4 \leq n^3 < (m+1)^4\} = \{n \text{ odd} : m^{4/3} \leq n < (m+1)^{4/3}\}.$$

If $n \in \Phi(m)$ and $\lfloor n^{3/2} \rfloor$ is even, then $J(J(n)) = m$, hence $n \in A$. Moreover $H_m := |\Phi(m)| \in [\frac{2}{3}m^{1/3} - 1, \frac{2}{3}(m+1)^{1/3} + 1]$, and the fibers of distinct m are disjoint.

Proof. $J(n) = \lfloor \sqrt{n^3} \rfloor =: k$; if k is even then $J(k) = \lfloor \sqrt{k} \rfloor$, and $\lfloor \sqrt{\lfloor \sqrt{N} \rfloor} \rfloor = m \iff m^4 \leq N < (m+1)^4$: this is the exact form of the transparent nesting $\lfloor \sqrt{\lfloor n^{3/2} \rfloor} \rfloor = \lfloor n^{3/4} \rfloor$. The interval $[m^{4/3}, (m+1)^{4/3})$ has length between $\frac{4}{3}m^{1/3}$ and $\frac{4}{3}(m+1)^{1/3}$ by the mean value theorem, and a half-open interval of length ℓ contains between $\ell/2 - 1$ and $\ell/2 + 1$ odd integers. Disjointness is the cell identity. Lean: `sqrt_sqrt_eq_iff`, `oe_fiber_mem`, `floorPower_oe_fiber`, `oe_fiber_disjoint`. $\square$

Write $G_m = \#\{n \in \Phi(m) : \lfloor n^{3/2} \rfloor \text{ even}\}$. The whole argument rests on lower bounds for G_m : an elementary bound on every fiber outside a thin exceptional set (Section 4.1), and a classical average over an even block of m 's (Section 4.2).

(a) Even block $E(m)$: the even integers of $[m^2, (m+1)^2)$ (blue) all have $J(n) = m$



(b) OE fiber $\Phi(m)$: odd n with $\lfloor n^{3/4} \rfloor = m$; the parity of $\lfloor n^{3/2} \rfloor$ sweeps along the fiber




Figure 2: The two productions. (a) The even integers of $[m^2, (m+1)^2)$ all map to m . (b) The OE fiber $\Phi(m)$ for $m = 10^5$: the odd n with $\lfloor n^{3/4} \rfloor = m$, coloured by the parity of $\lfloor n^{3/2} \rfloor$; those with even image (blue) have $J(J(n)) = m$.

4. Parity on a fiber

4.1 The sweep lemma

On $\Phi(m)$ the quantity $x = n^{3/2}/2$ has $\lfloor n^{3/2} \rfloor$ even iff $\{x\} < \frac{1}{2}$. Consecutive odd n advance x by $\approx \frac{3}{2}n^{1/2} \approx \frac{3}{2}m^{2/3}$, and over the whole fiber this step drifts by less than one unit of $1/H_m$. So modulo 1 the fiber is an arithmetic progression with a nearly constant step α , of length H_m . Such a progression cannot avoid a half-circle unless its step is within $O(1/H_m)$ of 0, or its two-step is ($\alpha \approx \frac{1}{2}$ with the wrong phase).

Lemma 4.1 (sweep). Let $x_1 < x_2 < \dots < x_H$ be real numbers with $x_{j+1} - x_j \in [a, b]$ for all j , where $0 < a \leq b \leq \frac{1}{2}$, $b \leq \frac{21}{20}a$ and $(H-1)a \geq 12$. Then

$$\#\{j : \{x_j\} < \frac{1}{2}\} \geq \frac{H}{7} \quad \text{and} \quad \#\{j : \{x_j\} \geq \frac{1}{2}\} \geq \frac{H}{7}.$$

The same holds with the cells $(k/2, (k+1)/2]$ in place of $[k/2, (k+1)/2)$, i.e. with the representative in $(0, 1]$ in place of the fractional part.

Proof. Cut $\mathbb{R}$ into cells $C_k = [k/2, (k+1)/2)$, $k \in \mathbb{Z}$; even k are the *good* cells ($\{x\} < \frac{1}{2}$), odd k the *bad* cells. Call C_k *traversed* if $x_1 \leq k/2$ and $(k+1)/2 \leq x_H$.

- (i) *A traversed cell contains at least $g := \lfloor 1/(2b) \rfloor \geq 1$ and at most $G := \lfloor 1/(2a) \rfloor + 1$ of the points.* Let j_0 be the least index with $x_{j_0} \geq k/2$. Then $x_{j_0} < k/2 + b$ (either $j_0 = 1$ and $x_1 = k/2$, or $x_{j_0-1} < k/2$). Since every step is at most b , $x_{j_0+i} < k/2 + (i+1)b \leq (k+1)/2$ for all $i \leq \lfloor 1/(2b) \rfloor - 1$, and these indices exist because $(k+1)/2 \leq x_H$. That gives at least $\lfloor 1/(2b) \rfloor$ points in C_k . Since every step is at least a , the points in C_k are x_{j_0+i} with $k/2 + ia \leq x_{j_0+i} < (k+1)/2$, so $i < 1/(2a)$ and there are at most $\lfloor 1/(2a) \rfloor + 1$ of them. The same count holds for any cell as an upper bound.

- (ii) *Counting cells.* The traversed cells are consecutive, and their number T is at least $2(x_H - x_1) - 2 \geq 2(H-1)a - 2 \geq 22$. Good and bad traversed cells alternate, so their numbers

T_g, T_b differ by at most one and $T_g, T_b \geq 10$. Every point outside the traversed cells lies in the cell of x_1 or the cell of x_H .

(iii) *The ratio.* Hence $\text{Good} \geq gT_g$ and $H \leq G(T_g + T_b + 2) \leq G(2T_g + 3)$, so

$$\frac{\text{Good}}{H} \geq \frac{g}{G} \cdot \frac{T_g}{2T_g + 3} \geq \frac{g}{G} \cdot \frac{10}{23}.$$

Put $X = 1/(2a) \geq 1$, so $G = \lfloor X \rfloor + 1$ and, from $b \leq \frac{21}{20}a$, $g \geq \lfloor \frac{20}{21}X \rfloor$. If $X < 2$: $g \geq 1, G \leq 2$. If $2 \leq X < 3$: $g \geq 1, G = 3$. If $3 \leq X < 4$: $g \geq 2, G = 4$. If $4 \leq X < 5$: $g \geq 3, G = 5$. If $X \geq 5$: $g/G \geq (\frac{20}{21}X - 1)/(X + 1) \geq 0.62$. In all cases $g/G \geq \frac{1}{3}$, so $\text{Good}/H \geq \frac{10}{69} > \frac{1}{7}$. The bad count is symmetric. For left-open cells the same proof applies verbatim (the first point in a cell $(c, c + \frac{1}{2}]$ after a point $\leq c$ is $\leq c + b \leq c + \frac{1}{2}$). $\square$

Lean: `sweep_fract_lt_half`, `sweep_fract_ge_half` (closed cells) and `sweep_rep_le_half`, `sweep_rep_gt_half` (left-open cells, with the representative $x - \lceil x \rceil + 1 \in (0, 1]$), in `formal/Problems/Juggler/FateSweep.lean`. The formal count uses the strictly interior cells (at least $(T - 3)/2$ of one colour, ratio $11/25$) in place of the traversed cells (ratio $10/23$), and obtains the left-open case from the closed one by $x_j \mapsto -x_{H+1-j}$.

The product $g/G \cdot 10/23$ cannot reach $\frac{1}{3}$. An adversarial sequence with steps in $[a, b]$ can lock a $3 + 1$ split near $a = \frac{1}{4}$. The OE fiber is monotone.

Lemma 4.1' (monotone pairing). Keep the hypotheses of Lemma 4.1 and assume the consecutive differences are monotone (nondecreasing or nonincreasing). Then

$$\#\{j : \{x_j\} < \tfrac{1}{2}\} \geq \tfrac{H}{3} - 2 \quad \text{and} \quad \#\{j : \{x_j\} \geq \tfrac{1}{2}\} \geq \tfrac{H}{3} - 2.$$

The same holds with the cells $(k/2, (k + 1)/2]$ in place of $[k/2, (k + 1)/2)$.

Proof. We prove the nondecreasing case for both half-cell conventions. Reversing the order and reflecting, $z_j = -x_{H+1-j}$, then gives the nonincreasing case; it interchanges the half-cell conventions and preserves the two color counts up to relabeling. Assume the steps are nondecreasing. Cut $\mathbb{R}$ into the cells $C_k = [k/2, (k + 1)/2)$ of Lemma 4.1. Since every step is at most $b \leq \frac{1}{2}$, the cells of $x_1, \dots, x_H$ are T consecutive integers, each visited; write $\rho_1, \dots, \rho_T \geq 1$ for their occupancies in order, so $\sum_i \rho_i = H$, and consecutive cells have opposite colours. Call the cells $2, \dots, T - 1$ *interior*. Put $X = 1/(2a)$, $G = \lfloor X \rfloor + 1$ and $g = \lfloor 1/(2b) \rfloor \geq 1$. As in Lemma 4.1(i), every cell has $\rho_i \leq G$ and every interior cell has $\rho_i \geq g$; and $(H - 1)a \geq 12$ gives $H \geq 24X + 1 \geq 24G - 23$.

- (a) *A later cell is at most one fuller than an earlier interior cell.* Let $2 \leq i < j \leq T$. The $\rho_i + 1$ steps from the last point before the cell of index i to the first point after it span more than $\frac{1}{2}$, so one of them, γ , exceeds $1/(2(\rho_i + 1))$. Every step inside the cell of index j comes later, hence is at least γ , and the $\rho_j - 1$ steps inside that cell span less than $\frac{1}{2}$. So $(\rho_j - 1)\gamma < \frac{1}{2}$, i.e. $\rho_j \leq \rho_i + 1$. In particular, once an interior cell of occupancy v has occurred, every later cell has occupancy at most $v + 1$.
- (b) *Pairs and surplus.* Pair the interior cells consecutively, $(2, 3), (4, 5), \dots$; at most three cells are unpaired (the first, the last, and possibly one interior cell), each holding at most G points. Each pair has one cell of each colour, so each colour receives at least $\sum_{\text{pairs}} \min(\rho, \rho')$, and

$$\sum_{\text{pairs}} \min(\rho, \rho') \geq \tfrac{1}{3}(H - 3G) + S, \quad S := \sum_{\text{pairs}} \left(\min(\rho, \rho') - \tfrac{\rho + \rho'}{3} \right).$$

It therefore suffices to show $S \geq G - 2$. Both entries of every pair lie in $[g, G]$, so every pair has $\min/\text{sum} \geq g/(g + G) =: r$, and a pair contributes at least $(r - \frac{1}{3})(\rho + \rho')$ to S .

- (c) $G \geq 7$. Then $X \geq 6$ and $g \geq \lfloor \frac{20}{21}X \rfloor$ give $7g \geq 5G$ ($g \geq 5$ at $G = 7$, $g \geq 6$ at $G = 8$, and $7g > \frac{20G-41}{3} \geq 5G$ for $G \geq 9$), so $r \geq \frac{5}{12}$ and $S \geq \frac{1}{12}(H - 3G) \geq \frac{21G-23}{12} \geq G - 2$.
- (d) $3 \leq G \leq 6$ and $g \geq 2$. The possible (G, g) are $(3, 2), (4, 2), (4, 3), (5, 3), (5, 4), (6, 4), (6, 5)$. Except for $(4, 2)$, $r \geq \frac{3}{8}$, so $S \geq \frac{1}{24}(H - 3G) \geq \frac{21G-23}{24} \geq G - 2$ because $3G \leq 25$. For $(G, g) = (4, 2)$ the interior occupancies lie in $\{2, 3, 4\}$; by (a), after the first interior cell of occupancy 2 every cell has occupancy at most 3. So the pairs with both entries in $\{3, 4\}$ (ratio $\geq \frac{3}{7}$) precede the pairs with both entries in $\{2, 3\}$ (ratio $\geq \frac{2}{5}$), and at most one pair straddles, with ratio $\geq \frac{1}{3}$ and at most 6 points. Hence $S \geq \frac{1}{15}(H - 3G - 6) \geq \frac{24G-41}{15} > 2 = G - 2$.
- (e) $G = 2$. Interior occupancies lie in $\{1, 2\}$, every pair has ratio $\geq \frac{1}{3}$, and $S \geq 0 = G - 2$.
- (f) $G = 3$ and $g = 1$, i.e. $2 \leq X < 3$ and $b > \frac{1}{4}$; then $a \leq \frac{1}{4}$ and $b \leq \frac{21}{80}$, so $3b < 1$ and $7b < 2$. Interior occupancies lie in $\{1, 2, 3\}$, and by (a) every cell after the first interior cell of occupancy 1 has occupancy at most 2. The pairs are therefore: pairs with both entries in $\{2, 3\}$, of ratio $\geq \frac{2}{5}$; at most one straddling pair $(\rho, 1)$ with $\rho \in \{2, 3\}$, which contributes at least $-\frac{1}{3}$ to S ; and pairs with both entries in $\{1, 2\}$. In the last group, $(1, 1)$ does not occur: two adjacent interior cells with one point each would have three consecutive steps spanning more than $1 > 3b$. Nor do two consecutive pairs with six points in all: four consecutive interior cells with six points would have seven consecutive steps spanning more than $2 > 7b$. So among any two consecutive pairs of the last group at least one is $(2, 2)$, and the two together hold at least 7 points of which the scarcer colour gets at least 3: their contribution to S is at least $3 - \frac{7}{3} = \frac{2}{3}$, and they hold at most 8 points, so the last group contributes at least $\frac{1}{12}N_2 - \frac{2}{3}$ to S , where N_2 is its number of points (a leftover single pair contributes at least 0). With N_1 the number of points in the first group, $N_1 + N_2 \geq H - 3G - 4 = H - 13$, and

$$S \geq -\frac{1}{3} + \frac{1}{15}N_1 + \frac{1}{12}N_2 - \frac{2}{3} \geq \frac{1}{15}(H - 13) - 1 \geq 1 = G - 2,$$

since $H \geq 24X + 1 \geq 49$.

In all cases $S \geq G - 2$, so the scarcer colour has at least $\frac{1}{3}(H - 3G) + G - 2 = \frac{H}{3} - 2$ points. For left-open cells the same argument applies with the cells $(k/2, (k+1)/2]$, or apply the closed case to $-x_{H+1-j}$ as in Lemma 4.1. $\square$

Lean: `sweep_monotone_fract_lt_half`, `sweep_monotone_fract_ge_half` (closed cells) and `sweep_monotone_rep_le_half`, `sweep_monotone_rep_gt_half` (left-open cells), instances of `Sweep.sweep_monotone_cell` and `sweep_monotone_ceil`, in `formal/Problems/Juggler/FateSweepMonotone.lean`.

Remark (erratum, 8 September 2026). An earlier version of this proof asserted that every pair satisfies $\min \geq (\rho + \rho')/3$, on the grounds that the step scale $1/(2\delta)$ changes by at most $X/21$ “spread over the cells” and that with $X < 2.1$ the pairs are $(1, 1)$, $(1, 2)$ or $(2, 2)$. Monotone steps need not change gradually: with $a = \frac{10}{41}$, $b = \frac{21}{82}$ the sequence $-2a, -a, 0, a, 2a, 2a+b, 2a+2b, \dots$ has nondecreasing steps in $[a, b]$, and its cells 2 and 3 — an interior pair in the pairing of (b) — have occupancies $(3, 1)$. That version also paired the two partial end cells as if they were interior. The statement is unchanged; the proof above replaces it, with (a) in place of the spreading claim, the end cells left unpaired, and the surplus S paying for them.

Lemma 4.2 (fiber parity). For $m \geq 10^6$ put $\alpha_m = \{\frac{3}{2}m^{2/3}\}$ and call m good if

$$\|\alpha_m\| \geq 22m^{-1/3} \quad \text{and} \quad \|\alpha_m - \tfrac{1}{2}\| \geq 2m^{-1/3},$$

where $\|\cdot\|$ is the distance to the nearest integer. If m is good then $G_m \geq \frac{1}{3}H_m - 2$ and $H_m - G_m \geq \frac{1}{3}H_m - 2$.

Proof. Let $n_1 < \dots < n_H$ be the odd integers of $\Phi(m)$, $n_{j+1} = n_j + 2$, and $x_j = n_j^{3/2}/2$. Then $\lfloor n_j^{3/2} \rfloor$ is even iff $\{x_j\} < \frac{1}{2}$. The steps are

$$\delta_j = x_{j+1} - x_j = \int_{n_j}^{n_{j+1}} \frac{3}{4} t^{1/2} dt \in \left[\frac{3}{2} n_j^{1/2}, \frac{3}{2} (n_j + 2)^{1/2} \right] \subseteq [A_m, B_m],$$

with $A_m = \frac{3}{2} m^{2/3}$ and $B_m = \frac{3}{2} ((m+1)^{4/3} + 2)^{1/2}$. Using $\sqrt{u} - \sqrt{v} \leq (u-v)/(2\sqrt{v})$ and $(m+1)^{4/3} - m^{4/3} \leq \frac{4}{3}(m+1)^{1/3}$,

$$\eta_m := B_m - A_m \leq \frac{(m+1)^{1/3} + \frac{3}{2}}{m^{2/3}} \leq m^{-1/3} \left(1 + \frac{1}{3m} + \frac{3}{2} m^{-1/3} \right) \leq 1.02 m^{-1/3}.$$

Also $H_m - 1 \geq \frac{2}{3} m^{1/3} - 2 \geq 0.646 m^{1/3}$.

Case 1: $\alpha_m \leq \frac{1}{2} - 2m^{-1/3}$. Put $N = \lfloor A_m \rfloor$ and $y_j = x_j - jN$; then $\{y_j\} = \{x_j\}$ and the steps of y lie in $[a, b] = [\alpha_m, \alpha_m + \eta_m] \subseteq [22m^{-1/3}, \frac{1}{2}]$. Now $b/a \leq 1 + 1.02/22 < \frac{21}{20}$ and $(H-1)a \geq 0.646 \cdot 22 > 12$. The steps of y are nondecreasing. Lemma 4.1' applies.

Case 2: $\alpha_m \geq \frac{1}{2} + 2m^{-1/3}$. Put $z_j = j(N+1) - x_j$, increasing with steps $(N+1) - \delta_j \in [1 - \alpha_m - \eta_m, 1 - \alpha_m] \subseteq [20.98 m^{-1/3}, \frac{1}{2}]$; again $b/a \leq 1 + 1.02/20.98 < \frac{21}{20}$ and $(H-1)a \geq 0.646 \cdot 20.98 > 12$. Write $\langle z \rangle \in (0, 1]$ for the representative of z modulo 1 in $(0, 1]$. Then $\langle z_j \rangle = 1 - \{x_j\}$ for every j , so $\{x_j\} < \frac{1}{2} \iff \langle z_j \rangle \in (\frac{1}{2}, 1]$. The steps of z are monotone. Lemma 4.1' in its left-open form gives both counts.

The middle range $|\alpha_m - \frac{1}{2}| < 2m^{-1/3}$ and the range $\|\alpha_m\| < 22m^{-1/3}$ are excluded by goodness. $\square$

Lean: `FiberParity.fiber_parity_good` in `formal/Problems/Juggler/FateFiberParity.lean`, with goodness unpacked as $22m^{-1/3} \leq \alpha_m \leq 1 - 22m^{-1/3}$ and $\alpha_m \leq \frac{1}{2} - 2m^{-1/3}$ or $\alpha_m \geq \frac{1}{2} + 2m^{-1/3}$ (`FiberParity.Good`). The step of $x(n) = n\sqrt{n}/2$ over two units is $(u^2 + uv + v^2)/(u+v)$ with $u = \sqrt{n+2}$, $v = \sqrt{n}$ (`xval_step`), which lies in $[\frac{3}{2}v, \frac{3}{2}u]$ by $(2u+v)(u-v) \geq 0$ and $(u+2v)(u-v) \geq 0$ and is nondecreasing because the upper bound at n is the lower bound at $n+2$; no integral is needed. The upper step is taken at $n_{j+1} \in \Phi(m)$, so $B_m = \frac{3}{2}(m+1)^{2/3}$ and $\eta_m \leq m^{-1/3}$ by Bernoulli on $(1+1/m)^{2/3}$; the fiber has at least $\frac{2}{3}m^{1/3} - 1$ members by Bernoulli on $(1+1/m)^{4/3}$. Both cases go through `Sweep.sweep_monotone_cell` and `sweep_monotone_ceil` of `FateSweepMonotone`, with $\lceil 2z_j \rceil \equiv \lfloor 2x_j \rfloor \pmod{2}$ in Case 2.

Lemma 4.3 (bad fibers are thin). For $u \geq 10^6$,

$$\#\{m \in (u, 2u] : m \text{ bad}\} \leq 63 u^{2/3}, \quad \sum_{\substack{m > U \\ m \text{ bad}}} \frac{1}{m} \leq 306 U^{-1/3} \quad (U \geq 10^6).$$

Proof. $\varphi(m) = \frac{3}{2} m^{2/3}$ increases with $\varphi(m+1) - \varphi(m) \in [(m+1)^{-1/3}, m^{-1/3}] \subseteq [(3u)^{-1/3}, u^{-1/3}]$ on $(u, 2u]$. Badness means $\{\varphi(m)\}$ lies in one of two arcs of the circle, of total length at most $48 u^{-1/3}$. As m runs over $(u, 2u]$, φ increases by $\frac{3}{2} u^{2/3} (2^{2/3} - 1) < 0.882 u^{2/3}$, so it passes each arc at most $0.882 u^{2/3} + 2$ times, and each pass through an arc of length w takes at most $w(3u)^{1/3} + 1$ values of m . Hence the count is at most $(0.882 u^{2/3} + 2)(48 \cdot 3^{1/3} + 2) \leq 63 u^{2/3}$ for $u \geq 10^6$. Summing $63 (2^i U)^{-1/3}$ over $i \geq 0$ gives $63 U^{-1/3} / (1 - 2^{-1/3}) \leq 306 U^{-1/3}$. $\square$

Lean: `FiberParity.bad_count_le` and `FiberParity.bad_logMass_le` in `formal/Problems/Juggler/FateThinFibers.lean`, with badness the negation of `FiberParity.Good` and the log-mass bound stated for every finite range $(U, N]$. The count is the paper's: the increments of φ lie in $[(m+1)^{-1/3}, m^{-1/3}]$ by Bernoulli both ways; a bad $m > u$ has $\{\varphi(m) + 22u^{-1/3}\} < 44u^{-1/3}$ or $\{\varphi(m)\} \in [\frac{1}{2} - 2u^{-1/3}, \frac{1}{2} + 2u^{-1/3}]$, so the wrap-around arc is a single arc after the shift; on an arc of width w a sequence increasing by at least d per step holds at most $w/d + 1$ points per integer window (`arc_count_le`), and at most $\varphi(2u) - \varphi(u) + 2$ windows are met. The formal count uses $d = (2u)^{-1/3}$, sharper than the $(3u)^{-1/3}$ above, so w/d is $44 \cdot 2^{1/3}$ and $4 \cdot 2^{1/3}$ and the total $(0.882 u^{2/3} + 2)(48 \cdot 2^{1/3} + 2)$ clears $63 u^{2/3}$ with room. The dyadic sum uses $2^{-1/3} \leq 0.794$, whence $63/(1 - 2^{-1/3}) \leq 306$.

4.2 The block average

The adversarial sweep $G_m \geq H_m/7$ is not the fiber bound: Lemma 4.1' gives the monotone floor $H_m/3 - 2$. In the numerical experiments of Section 11 the mean of G_m/H_m is $\frac{1}{2}$ and the minimum on good fibers is near $\frac{1}{3}$. When the members of A come in even blocks — as the E -produced part of A always does — the fibers can be averaged over the block, and the average is asymptotically $\frac{1}{2}$ by a classical exponential-sum estimate.

Proposition 4.4 (block average). For $m' \geq 2$, put $I(m') = [m'^{8/3}, (m' + 1)^{8/3})$ and

$$U(m') = \{n \text{ odd in } I(m') : \lfloor n^{3/4} \rfloor \text{ even}, \quad \lfloor n^{3/2} \rfloor \text{ even}\}.$$

This is the disjoint union of the even-image parts of $\Phi(m)$ over $m \in E(m')$. There is an absolute constant C_B such that

$$|U(m')| - \frac{1}{4} \#\{n \text{ odd in } I(m')\} \leq C_B m'^{11/9} \log(m' + 1). \quad (4.1)$$

In particular

$$|U(m')| = \frac{1}{3} m'^{5/3} (1 + O(m'^{-4/9} \log(m' + 1))).$$

No numerical value of C_B is asserted.

Proof. Write $M = \#\{n \text{ odd in } I(m')\} \asymp m'^{5/3}$ and $\psi(z) = (-1)^{\lfloor z \rfloor}$. Expanding the two indicators gives four sums: M , the sums of $\psi(n^{3/4})$ and $\psi(n^{3/2})$, and their product.

The slow sum is $O(m')$: pair consecutive fibers of $\lfloor n^{3/4} \rfloor$, whose odd-point counts differ by at most 3, and bound the remaining end fiber by $O(m'^{2/3})$.

We use the interval form of Vaaler's approximation [10]. For $J \geq 1$, the parity wave has a trigonometric approximation $V_J(z/2)$, with zero constant term and coefficients $O(1/|q|)$, and a nonnegative error majorant $\Delta_J(z/2)$. Its degree is J , every coefficient is $O(1/J)$, and $\|\Delta_J\|_\infty = O(1)$. All constants in this paragraph are absolute; the interval endpoints are included in the majorant.

Take $J = \max(1, \lfloor m'^{4/9} \rfloor)$ and reindex odd integers as $n = 2r + 1$. For $q \neq 0$, the fast phase $f(r) = q(2r + 1)^{3/2}/2$ has $|f''| \asymp |q|m'^{-4/3}$, with constant sign. The second-derivative test [13, Theorem 2.2] gives, uniformly on initial subintervals,

$$\left| \sum_r e(f(r)) \right| \ll m' |q|^{1/2} + m'^{2/3} |q|^{-1/2}, \quad e(z) = e^{2\pi i z}.$$

The coefficient sum and the majorant sum are therefore bounded by

$$\ll m' J^{1/2} + m'^{2/3} + M/J \ll m'^{11/9}.$$

For the product, expand both waves. A mixed phase is $f = (q_1 n^{3/4} + q_2 n^{3/2})/2$, with $q_2 \neq 0$, and

$$f''(r) = \frac{3}{2} q_2 n^{-1/2} \left(1 - \frac{q_1}{4q_2} n^{-3/4} \right).$$

For $|q_1| \leq J$, the correction is $O(m'^{-14/9})$; for all sufficiently large m' the same second-derivative estimate applies uniformly. Summing the slow coefficients adds at most $O(\log(J + 1))$.

The slow majorant has constant contribution $O(M/J)$. Its nonzero modes have monotone derivative $\frac{3}{4} q_1 n^{-1/4}$, of magnitude $\asymp |q_1| m'^{-2/3}$ and at most $1/2$ for sufficiently large m' . Kusmin–Landau [13, Theorem 2.1] bounds a mode by $O(m'^{2/3}/|q_1|)$, so the majorant contributes $O(M/J + m'^{2/3} \log(J + 1)/J)$. Since $|\psi| = 1$, $|V_J| \leq 1 + \Delta_J = O(1)$, and the majorants are nonnegative, the product-approximation error is bounded by a constant times the sum of the two majorants. These estimates prove (4.1) for sufficiently large m' ; increasing the unspecified absolute constant covers the finitely many smaller integers $m' \geq 2$. $\square$

Lean: the exact layer and the conditional deduction, in `formal/Problems/Juggler/FateBlockAverage.lean`. The block is `BlockAverage.oddBlock`, exactly the odd n with $m'^2 \leq \lfloor n^{3/4} \rfloor < (m' + 1)^2$ by the landing window of Appendix D.1 (`mem_oddBlock`); `U_card_eq` is the sentence that $U(m')$ is the disjoint union of the even-image parts of the $\Phi(m)$ over the even m of the block, with `oddBlock_card_eq` the same for the whole block; and `four_card_U` is the expansion $4|U(m')| = M + S_1 + S_2 + S_{12}$ that the proof starts from. The slow sum needs no analysis: it is the alternating sum $\sum_m (-1)^m |\Phi(m)|$ over the block (`slowSum_eq_fibers`), consecutive fibers differ by at most two members (`oeFiber_card_succ_diff`, from $\frac{2}{3}m^{1/3} - 1 \leq |\Phi(m)| \leq \frac{2}{3}(m + 1)^{1/3} + 1$), and the block pairs off, so $|S_1| \leq 2m' + \frac{2}{3}(m' + 1)^{2/3} + 1$ (`slowSum_abs_le`). The count M is pinned two-sidedly by the same fiber bounds (`oddBlock_card_le`, `oddBlock_card_ge`), whence $|M/4 - m'^{5/3}/3| \leq m' + 1$ (`oddBlock_quarter_close`). Given bounds on the fast sum and the product sum only, (4.1) is `block_average_two_bounds`, in the paper's shape `block_average_bound_two` with $C_B = C/2 + 2$, and the asymptotic form is `block_average_asymptotic` with the explicit error $B/2 + 2(m' + 1)$. Those two bounds are hypotheses there, not theorems: Vaaler's approximation, the second-derivative test and Kusmin–Landau are not formalized, so this proposition stays a human proof and the table says so.

Revision note. The earlier value $C_0 = 250$ depended on an unverified explicit majorant and small-parameter inequalities that do not hold throughout their printed range. The asymptotic form above is all that the subsequent recursions use.

4.3 The share is a quadratic sweep

Sections 4.1 and 4.2 bound G_m from below, by a sweep on each good fiber and by an exponential sum across an even block. Neither says what G_m/H_m is. It has a closed form, and the form explains both the floor of Lemma 4.2 and the extreme fibers that the goodness condition excludes. Nothing in Sections 5–10 depends on this subsection; it is here because it identifies which of the two tools is doing the work, and because it names the hypothesis behind the gap between them.

Throughout, $n_1 = \min \Phi(m)$, $n_j = n_1 + 2(j - 1)$, and $x_j = n_j^{3/2}/2$, so that $\lfloor n_j^{3/2} \rfloor$ is even iff $\{x_j\} < \frac{1}{2}$. Write

$$\theta_m = \{x_1\}, \quad \beta_m = \alpha_m (H_m - 1),$$

with $\alpha_m = \{\frac{3}{2}m^{2/3}\}$ as in Lemma 4.2, taken in $(-\frac{1}{2}, \frac{1}{2}]$, and set

$$S(\beta, \theta) = |\{s \in [0, 1] : \{\theta + \beta s + \frac{1}{3}s^2\} < \frac{1}{2}\}|.$$

Lemma 4.5 (share law). As $m \rightarrow \infty$,

$$G_m/H_m = S(\beta_m, \theta_m) + O\left(H_m^{-1/2} + \frac{1 + |\beta_m|}{H_m}\right).$$

The estimate is useful when $|\beta_m| = o(H_m)$; it is not a uniform equidistribution assertion for all fibers.

Proof. Expand x_j about n_1 . With $u = 2(j - 1)/n_1$,

$$x_j = \frac{1}{2}n_1^{3/2}(1 + u)^{3/2} = x_1 + \frac{3}{2}n_1^{1/2}(j - 1) + \frac{3}{4}n_1^{-1/2}(j - 1)^2 + E_j, \quad |E_j| \leq \frac{1}{4}(j - 1)^3 n_1^{-3/2}.$$

On the fiber $j - 1 \leq H_m - 1 \ll m^{1/3}$ and $n_1 \geq m^{4/3}$, so $|E_j| \ll m^{-1}$, below $1/H_m$. Put $s = (j - 1)/(H_m - 1) \in [0, 1]$. The linear coefficient is $A = \frac{3}{2}n_1^{1/2}$, and since $n_1 = m^{4/3} + O(1)$ we have $A = \frac{3}{2}m^{2/3} + O(m^{-2/3})$, so $A \equiv \alpha_m + O(m^{-2/3}) \pmod{1}$ and the accumulated difference over the fiber is $O(m^{-1/3}) = O(1/H_m)$. Hence $A(j - 1) \equiv \beta_m s \pmod{1}$ up to $O(1/H_m)$. The quadratic coefficient gives $\frac{3}{4}n_1^{-1/2}(H_m - 1)^2 = \frac{1}{3} + O(1/H_m)$ by Lemma 3.2, so

$$x_j \equiv \theta_m + \beta_m s + \frac{1}{3}s^2 \pmod{1},$$

uniformly to $O(1/H_m)$. Put $\epsilon = O(1/H_m)$. Replacing the exact phase by the displayed quadratic can change an indicator only within ϵ of an integer or half-integer. If $|\beta_m| \leq 2$, there are $O(1)$ such levels and their inverse images have total length $O(\sqrt{\epsilon})$, since the quadratic has second derivative $2/3$. If $|\beta_m| > 2$, its derivative has magnitude at least $|\beta_m| - 2/3$, and the total inverse-image length is $O(\epsilon)$. Each of these sets, and each half-cell inverse image, has $O(1 + |\beta_m|)$ interval components. Sampling on the uniform grid costs $O((1 + |\beta_m|)/H_m)$. This proves the stated bound, including tangencies to the cell boundaries. $\square$

Corollary 4.6. For every β :

1. $\int_0^1 S(\beta, \theta) d\theta = \frac{1}{2}$.
2. $S(\beta, \theta) \in \{0, 1\}$ for some θ if and only if $\beta \in [-\frac{5}{6}, \frac{1}{6}]$.
3. $\int |\{\theta : S(\beta, \theta) = 0\}| d\beta = \frac{25}{108}$, and the same for $S = 1$.

Proof. (1) By Fubini the integral is $\int_0^1 \int_0^1 \mathbf{1}[\{\theta + \varphi(s)\} < \frac{1}{2}] d\theta ds$, and the inner integral is $\frac{1}{2}$ for every s , whatever φ .

- (2) Let $\varphi(s) = \beta s + \frac{1}{3}s^2$, so $\varphi'(s) = \beta + \frac{2}{3}s$. If $\beta \geq 0$ then φ is increasing with range $\beta + \frac{1}{3}$; if $\beta \leq -\frac{2}{3}$ it is decreasing with range $-\beta - \frac{1}{3}$; and if $-\frac{2}{3} < \beta < 0$ it dips to $\varphi(-\frac{3}{2}\beta) = -\frac{3}{4}\beta^2$ and returns to $\beta + \frac{1}{3}$, so its range is $\max(0, \beta + \frac{1}{3}) + \frac{3}{4}\beta^2$. An extreme value of S needs the image of $[0, 1]$ to miss a half-circle, hence range $\leq \frac{1}{2}$. That is $\beta \leq \frac{1}{6}$ in the first case, $\beta \geq -\frac{5}{6}$ in the second, and no constraint in the third, since the range there is at most $\frac{1}{3}$.

- (3) The θ -measure is $\max(0, \frac{1}{2} - \text{range})$. Integrating the four pieces of (2),

$$\int_0^{1/6} (\frac{1}{6} - \beta) + \int_{-1/3}^0 (\frac{1}{6} - \beta - \frac{3}{4}\beta^2) + \int_{-2/3}^{-1/3} (\frac{1}{2} - \frac{3}{4}\beta^2) + \int_{-5/6}^{-2/3} (\frac{5}{6} + \beta) = \frac{1}{72} + \frac{11}{108} + \frac{11}{108} + \frac{1}{72} = \frac{25}{108}.$$

The statement for $S = 1$ is the reflection $\theta \mapsto \theta + \frac{1}{2}$. $\square$

Lean: the exact layer, in `formal/Problems/Juggler/FateShareLaw.lean`. The expansion of x_j about n_1 with the remainder $|E_j| \leq \frac{1}{4}(j-1)^3 n_1^{-3/2}$ is `ShareLaw.xval_expansion`; its Taylor step is, after $v = \sqrt{1+u}$, the polynomial inequality $0 \leq 1 + \frac{3}{2}(v^2 - 1) + \frac{3}{8}(v^2 - 1)^2 - v^3 \leq (v^2 - 1)^3/16$ (`ShareLaw.taylor_three_halves`), so no derivative is taken, and on a fiber the remainder is at most $\frac{2}{27}(m+1)/m^2$ (`ShareLaw.xval_expansion_fiber`). The range of $\varphi(s) = \beta s + \frac{1}{3}s^2$ on $[0, 1]$ in the three cases of (2) is `ShareLaw.phiRange`, attained (`ShareLaw.exists_phi_sub_eq`) and never exceeded (`ShareLaw.phi_sub_le`), and it is at most $\frac{1}{2}$ exactly for $\beta \in [-\frac{5}{6}, \frac{1}{6}]$ (`ShareLaw.phiRange_le_half_iff`). The integral of (3), $\int \max(0, \frac{1}{2} - \text{range}) d\beta = \frac{25}{108}$ as $\frac{1}{72} + \frac{11}{108} + \frac{11}{108} + \frac{1}{72}$, is `ShareLaw.integral_extremeMeasure`, with the integrand vanishing outside $[-\frac{5}{6}, \frac{1}{6}]$ (`ShareLaw.extremeMeasure_eq_zero`). Not formalized: the reduction of x_j modulo 1 to $\theta_m + \beta_m s + \frac{1}{3}s^2$ with its $O(1/H_m)$ error, the inverse-image and grid-sampling estimates that make Lemma 4.5 from it, the Fubini argument of (1), and the identification of the θ -measure with $\max(0, \frac{1}{2} - \text{range})$ in (2) and (3).

Part (1) is the reason no aggregate count in this paper sees the extreme fibers: they are compensated at every drift separately, which is what Proposition 4.4 recovers by a different route. Part (2) says the extremes occupy a window of one drift unit, that is $\alpha_m \in [-\frac{5}{4}, \frac{1}{4}]m^{-1/3}$ by $H_m \sim \frac{2}{3}m^{1/3}$.

At the centre of that window the law degenerates to an identity.

Lemma 4.7 (cube fibers). Let $k \geq 2$ and $m = k^3$. Every $n \in \Phi(m)$ has $n = k^4 + t$ with $3t \leq 4k$, and then $\lfloor n^{3/2} \rfloor = k^6 + \frac{3}{2}k^2t$ exactly. Consequently the fiber of an even cube is full, $G_m = H_m$, and along the fiber of an odd cube the images alternate in parity, so $|2G_m - H_m| \leq 1$.

Proof. If $3t \geq 4k + 1$ then $(3k^4 + 4k + 1)^3 \leq (3k^4 + 3t)^3 = 27(k^4 + t)^3$, and $(3k^4 + 4k + 1)^3 \geq 27(k^3 + 1)^4$, so $n^3 \geq (k^3 + 1)^4$ and $n \notin \Phi(m)$. Given $3t \leq 4k$, the claimed value $q = k^6 + \frac{3}{2}k^2t$

satisfies $n^3 - q^2 = \frac{3}{4}k^4t^2 + t^3 \geq 0$ and $(q+1)^2 - n^3 = 2k^6 + 3k^2t + 1 - \frac{3}{4}k^4t^2 - t^3 > 0$, so $q = \lfloor n^{3/2} \rfloor$. For even k the value $k^6 + \frac{3}{2}k^2t$ is even for every t ; for odd k the fiber's t are even and the value has the parity of $1 + t/2$. Lean: `cube_fiber_range`, `cube_fiber_sqrt_even`, `cube_fiber_sqrt_odd`, `even_cube_fiber_full`, `odd_cube_fiber_alternating` in `formal/P` `roblems/Juggler/CubeFiber.lean`. $\square$

Remark 4.8 (what the window is, arithmetically). $\alpha_m = 0$ exactly means $\frac{3}{2}m^{2/3} \in \mathbb{Z}$, that is $27m^2 = 8k^3$, whose integer solutions are $m = (2e)^3$ and $k = 6e^2$. So the centre of the extreme window is the even cubes of Lemma 4.7, and the window itself is the near-solution set of that equation; the crossings are spaced $\asymp m^{1/3}$ apart, giving $\sim \frac{3}{2}M^{2/3}$ of them below M . Two consequences worth recording. The set is asymmetric about 0, which the symmetric goodness condition of Lemma 4.2 cannot express: that condition discards $\|\alpha_m\| < 22m^{-1/3}$, of width $44m^{-1/3}$, against the $\frac{3}{2}m^{-1/3}$ the extremes actually occupy. Nothing here depends on the difference, since Lemma 4.3 already makes the discarded mass $o(1)$; the point is that the goodness threshold is a convenience, not a boundary of the phenomenon. And the two sides of the window are not symmetric in strength either: full fibers contain the infinite family of Lemma 4.7, proved, whereas an empty fiber requires a near-miss together with a phase condition, and whether there are infinitely many is open.

Remark 4.9 (averaging the rest term). Item 3 of Section 5.1 has coefficient $2/9$; a weighted mean even-image share of $1/2 + o(1)$ on that source set would give $1/3$. Such a weighted mean is a sufficient additional hypothesis. Conditional equidistribution of the landing phases could help establish it if the approximation errors and the weights were also controlled, but equidistribution is not necessary for one mean identity. Neither Corollary 4.6(1), which averages an independent model phase, nor a pointwise bound on each fiber establishes the required average on an arbitrary backward-closed set. The goodness condition already removes its exceptional fibers at total logarithmic cost $o(1)$.

5. The recursion and the contagion theorem

5.1 Three sources of members of A in $(\sqrt{x}, x]$

Let $x \geq 2$ and $t = \log x$. Three families of members of $A \cap (\sqrt{x}, x]$ are pairwise disjoint.

1. **E -images.** For $m \in A \cap (x^{1/4}, \sqrt{x} - 1]$, $E(m) \subseteq (\sqrt{x}, x]$ (as $m^2 > \sqrt{x}$ and $(m+1)^2 \leq x$). Distinct m give disjoint blocks. By Lemma 3.1 the log-mass is at least $\sum_m \frac{1}{m}(1 - \frac{2}{m}) \geq (1 - 2x^{-1/4})\ell_A(x^{1/4}, \sqrt{x} - 1]$, and $\ell_A(x^{1/4}, \sqrt{x} - 1] \geq g_A(t/2) - 2x^{-1/2}$ (at most one integer lies in $(\sqrt{x} - 1, \sqrt{x}]$).
2. **OE -images of E -blocks.** For $m' \in A \cap (x^{3/16}, x^{3/8} - 1]$, $U(m') \subseteq A \cap (\sqrt{x}, x]$ (Lemma 3.2 applied to the even $m \in E(m') \subseteq A$; $I(m') \subseteq (\sqrt{x}, x]$ as $m'^{8/3} > \sqrt{x}$ and $(m' + 1)^{8/3} \leq x$). Distinct m' give disjoint $I(m')$. Each $n \in U(m')$ has $1/n > (m' + 1)^{-8/3}$, so by Proposition 4.4 the log-mass of $U(m')$ is at least $\frac{1}{3}m'^{5/3}(1 - \varepsilon_B(m'))(m' + 1)^{-8/3} \geq \frac{1}{3m'}(1 - \varepsilon_B(m') - \frac{8}{3m'})$, and summing, $\geq \frac{1}{3}(1 - \varepsilon'(x))(g_A(3t/8) - 2x^{-3/8})$ with $\varepsilon_B(m') = O(m'^{-4/9} \log(m' + 1)) \geq 0$ chosen to bound the error in Proposition 4.4 and $\varepsilon'(x) = \sup_{m' > x^{3/16}} (\varepsilon_B(m') + \frac{8}{3m'}) \rightarrow 0$.
3. **OE -images of the rest.** Let $A_x^E = \bigcup_{m'' \in A} E(m'') \cap (x^{3/8}, x^{3/4}]$ be the E -produced part of $A \cap (x^{3/8}, x^{3/4}]$ and A_x^{rest} its complement in $A \cap (x^{3/8}, x^{3/4}]$. The blocks meeting $(x^{3/8}, x^{3/4}]$ come from $m'' \in [x^{3/16} - 1, x^{3/8}]$, so $\ell(A_x^E) \leq \sum_{m''} \frac{m''+1}{m''^2} \leq (1 + 2x^{-3/16})(g_A(3t/8) + 2x^{-3/16})$ and $\ell(A_x^{\text{rest}}) \geq g_A(3t/4) - (1 + 2x^{-3/16})(g_A(3t/8) + 2x^{-3/16})$. For $m \in A_x^{\text{rest}}$ with $m \leq x^{3/4} - 1$ (this drops at most one element, of log-mass $\leq 2x^{-3/4}$), $\Phi(m) \subseteq (\sqrt{x}, x]$; the fibers are disjoint from each other and from those of item 2 (different m), and for good $m \geq 10^6$ Lemmas 4.2 and 3.2 give log-mass at least $(\frac{1}{3}H_m - 2)(m+1)^{-4/3} \geq \frac{2}{9m}(1 - O(m^{-1/3}))$.

Bad $m > x^{3/8}$ carry log-mass at most $306x^{-1/8}$ (Lemma 4.3). Hence item 3 contributes at least $\frac{2}{9}(1 - O(x^{-1/8}))[\ell(A_x^{\text{rest}}) - 306x^{-1/8} - 2x^{-3/4}]_+$.

The images in item 1 are even, those in items 2–3 odd, so the three families are disjoint, and all lie in $(\sqrt{x}, x]$.

5.2 The functional inequalities

Adding items 1 and 2:

$$g_A(t) \geq (1 - \varepsilon_1(t))g_A(t/2) + \frac{1}{3}(1 - \varepsilon_2(t))g_A(3t/8) - \varepsilon_3(t), \quad (5.1)$$

with $\varepsilon_1 = 2e^{-t/4}$, $\varepsilon_2 = \varepsilon'(e^t)$, $\varepsilon_3 = 2e^{-t/2} + e^{-3t/8}$, all tending to 0. Adding item 3 as well, and noting that the coefficient of the *actual* value $g_A(3t/8)$ is $\frac{1}{3}(1 - \varepsilon_2) - \frac{2}{9}(1 + 2e^{-3t/16}) \rightarrow \frac{1}{9} > 0$:

$$g_A(t) \geq (1 - \varepsilon_1)g_A(t/2) + \left(\frac{1}{9} - \varepsilon_4\right)g_A(3t/8) + \left(\frac{2}{9} - \varepsilon_5\right)g_A(3t/4) - \varepsilon_6(t), \quad (5.2)$$

with $\varepsilon_4, \varepsilon_5, \varepsilon_6 \rightarrow 0$ (for t so large that $x^{3/8} \geq 10^6$; the bracket $[\cdot]_+$ may be dropped because the right side of (5.2) is a lower bound for it). Every coefficient of a g_A -value on the right is nonnegative for large t , so lower bounds for g_A at $t/2$, $3t/8$, $3t/4$ transfer.

5.3 The recursion lemma

The passage from a functional inequality with vanishing errors to a power lower bound is a standalone statement.

Lemma 5.1 (recursion lemma). Let $e_1, \dots, e_r \in (0, 1)$, $c_1, \dots, c_r \geq 0$, and $\lambda \in (0, 1)$ with

$$\zeta := \sum_{i=1}^r c_i e_i^\lambda - 1 > 0.$$

Put $e_{\min} = \min_i e_i$, $e_{\max} = \max_i e_i$. Let $g : (0, \infty) \rightarrow [0, \infty)$ satisfy, for all $t \geq t_1$,

$$g(t) \geq \sum_{i=1}^r (c_i - \eta_i(t))g(e_i t) - \eta_0(t),$$

where $0 \leq \eta_i(t) \leq c_i$ for $i \geq 1$, $\eta_0(t) \geq 0$, and $\sum_{i \geq 1} \eta_i(t)e_i^\lambda \leq \zeta/3$ and $\eta_0(t) \leq \frac{2\zeta}{3}c_0$ for all $t \geq t_1$, for some $c_0 > 0$ with $g(t) \geq c_0$ on $[e_{\min}t_1, t_1]$. Then

$$g(t) \geq K t^\lambda \quad \text{for all } t \geq e_{\min}t_1, \quad K = c_0 t_1^{-\lambda}.$$

Proof. On $[e_{\min}t_1, t_1]$, $Kt^\lambda \leq Kt_1^\lambda = c_0 \leq g(t)$. Suppose $g(t) \geq Kt^\lambda$ holds on $[e_{\min}t_1, T]$ for some $T \geq t_1$, and let $t \in (T, T/e_{\max}]$. Then $e_i t \leq e_{\max}t \leq T$ and $e_i t \geq e_{\min}t > e_{\min}t_1$, so each $e_i t$ lies in the interval where the bound is known, and $t \geq t_1$, so

$$\begin{aligned} g(t) &\geq \sum_i (c_i - \eta_i(t))K(e_i t)^\lambda - \eta_0(t) \\ &= Kt^\lambda \left[\sum_i c_i e_i^\lambda - \sum_i \eta_i(t)e_i^\lambda \right] - \eta_0(t) \\ &\geq Kt^\lambda \left(1 + \frac{2\zeta}{3} \right) - \frac{2\zeta}{3}c_0 \geq Kt^\lambda. \end{aligned}$$

using $c_0 = Kt_1^\lambda \leq Kt^\lambda$. Hence the bound holds on $[e_{\min}t_1, T/e_{\max}]$; induction on N covers $[e_{\min}t_1, t_1 e_{\max}^{-N}]$ for every N , i.e. all $t \geq e_{\min}t_1$. $\square$

Lean: `recursion_lemma` in `formal/Problems/Juggler/FateRecursion.lean`, with $e_{\min} \leq e_i \leq e_{\max}$ as bounds rather than extrema and $\lambda > 0$ in place of $\lambda \in (0, 1)$; the hypotheses $g \geq 0$ and $\lambda < 1$ are not used.

5.4 The seed

Lemma 5.2 (seed). Every nonempty backward-closed A contains an integer $m \geq 3$, and for any such m ,

$$g_A(t) \geq c_A := \left(1 - \frac{2}{m^4}\right) \left(\frac{0.375}{m+1} - \frac{1}{(m+1)^2 - 1}\right) > 0 \quad \text{for all } t \geq 4 \log(m+1).$$

Proof. If $A \ni a \leq 2$ then $E(a) \subseteq A$ contains 2 or 4, and $E(2) \ni 4$, so $A \ni 4$. Fix $m \geq 3$ in A and let $S_1 = E(m)$, $S_{k+1} = \bigcup_{m'' \in S_k} E(m'')$. Then $S_k \subseteq A \cap [m^{2^k}, (m+1)^{2^k})$ and, by Lemma 3.1, $\ell(S_{k+1}) \geq (1 - 2m^{-2^k})\ell(S_k)$, so $\ell(S_k) \geq \ell(S_1) \prod_{j \geq 1} (1 - 2m^{-2^j}) \geq 0.75 \ell(S_1) \geq \frac{0.375}{m+1}$ for $m \geq 3$. Let $y \geq (m+1)^4$ and choose $k \geq 2$ with $(m+1)^{2^k} \leq y < (m+1)^{2^{k+1}}$. Then $S_k \subseteq (y^{1/4}, y]$ (as $y < m^{2^{k+2}}$). The members of S_k above $\sqrt{y}$ lie in $(\sqrt{y}, y]$; the members in $(y^{1/4}, \sqrt{y} - 1]$ have their E -images in $(\sqrt{y}, y]$, with log-mass at least $(1 - 2m^{-2^k})$ times their own; at most one member lies in $(\sqrt{y} - 1, \sqrt{y}]$, with $1/n \leq 1/((m+1)^2 - 1)$. Hence $g_A(\log y) \geq (1 - 2m^{-4})(\ell(S_k) - 1/((m+1)^2 - 1)) \geq c_A$, and $c_A > 0$ because $0.375(m+1) \geq 1 + 0.375/(m+1)$ for $m \geq 3$. $\square$

5.5 The theorem

Let λ^* be the root in $(0, 1)$ of $2^{-\lambda} + \frac{1}{3}(\frac{3}{8})^\lambda = 1$, let λ_{pair} be the pairing-only root of $2^{-\lambda} + \frac{1}{9}(\frac{3}{8})^\lambda + \frac{2}{9}(\frac{3}{4})^\lambda = 1$, and let λ^{**} be the root of (1.1). All three defining sums are continuous and strictly decreasing on $[0, 1]$, with value greater than 1 at 0 and less than 1 at 1; each therefore has a unique root there. Numerically $\lambda^* \approx 0.3774$, $\lambda_{\text{pair}} \approx 0.4480$, the *OEOEE* truncation is 0.4801, the V_3 truncation is 0.4891, the V_4 truncation is 0.4916, the V_5 truncation is 0.4924, and $\lambda^{**} \approx 0.4926$. For comparison: the elementary sweep alone, without Proposition 4.4, gives $2^{-\lambda} + \frac{2}{21}(\frac{3}{4})^\lambda = 1$, root 0.138; perfect fiber equidistribution at this depth would give $2^{-\lambda} + \frac{1}{3}(\frac{3}{4})^\lambda = 1$, root $\lambda_{\text{ideal}} \approx 0.4927$, the ceiling of the depth-two method.

Theorem 5.3 (logarithmic density of a backward-closed set). Let $A \subseteq \mathbb{N}$ be nonempty and backward-closed, and let $0 < \lambda < \lambda^{**}$. There are $c > 0$ and x_0 , depending on A and λ , such that

$$\sum_{\substack{n \in A \\ n \leq x}} \frac{1}{n} \geq c (\log x)^\lambda \quad \text{for all } x \geq x_0.$$

The same conclusion for $\lambda < \lambda^*$ uses only Proposition 4.4 (no sweep lemma).

Proof. Apply Lemma 5.1 to $g = g_A$, using the three pairs in (5.2) and the five pairs $(\rho_k, 3^{-(k+1)})$, $2 \leq k \leq 6$, of (5.10). For $0 < \lambda < \lambda^{**}$, the sum of their $c_i e_i^\lambda$ is greater than 1. Thus $\zeta = \sum_i c_i e_i^\lambda - 1 > 0$. Equations (5.2) and (5.10) have the required form, with the Appendix D errors, all tending to 0. Let $m \geq 3$ be a member of A and $c_0 = c_A$ from Lemma 5.2. Choose t_1 so large that $\frac{729}{8192} t_1 \geq 4 \log(m+1)$, that $e^{729 t_1 / 8192} \geq 10^6$, and that for all $t \geq t_1$ the errors satisfy $\sum_i \eta_i(t) e_i^\lambda \leq \zeta/3$ and $\eta_0(t) \leq \frac{2\zeta}{3} c_A$. Lemma 5.2 gives $g_A \geq c_A$ on $[\frac{729}{8192} t_1, t_1]$. The first induction step of Lemma 5.1 covers up to $t_1 / (\frac{3}{4}) = \frac{4}{3} t_1$, at which $\frac{729}{8192} t = \frac{243}{2048} t_1$. Lemma 5.1 then gives $g_A(t) \geq K t^\lambda$ for all $t \geq \frac{729}{8192} t_1$, and $\sum_{n \in A, n \leq x} 1/n \geq g_A(\log x)$. For $\lambda < \lambda^*$ use (5.1) with $(e_i, c_i) = (\frac{1}{2}, 1), (\frac{3}{8}, \frac{1}{3})$. $\square$

Lean: `contagion_of_production_inequality` and `logMass_contagion_of_production` in `formal/Problems/Juggler/FateContagionBound.lean`, with the inequality (5.2) — the eight productions (e_i, c_i) above, with errors $\eta_i, \eta_0 \geq 0$ tending to 0 — as a hypothesis and for every $0 < \lambda \leq 0.49$: the seed is Lemma 5.2 (`gA_seed`), the induction is Lemma 5.1, and $\zeta(0.49) = 0.00168 > 0$ is `zeta_pos_49`, eight exact rational lower bounds $r_i \leq e_i^{49/100}$ checked as $r_i^{100} \leq e_i^{49}$; ζ is antitone in λ (`zeta_antitone`), so $\lambda \leq 0.49$ suffices. The root λ^{**} and the range $0.49 < \lambda < \lambda^{**}$ are not formalized, and neither is (5.2) itself. The hypothesis $\eta_0 \geq 0$ is not needed.

Lean, unconditionally: the two families of Section 5.1 that need no exponential sum are proved with explicit errors, $g_A(t) \geq (1 - 4e^{-t/4})g_A(t/2) + (\frac{2}{9} - \frac{50}{9}e^{-t/8})g_A(3t/4) - \eta_0(t)$ for $t \geq 40$ with $\eta_0(t) = 2e^{-t/2} + \frac{4}{9}e^{-3t/4} + 136e^{-t/8}$ (`Production.production_two`, from `Production.family_E` on Lemma 3.1 and `Production.family_OE` on Lemmas 4.2 and 4.3), and Lemma 5.1 on these two productions, with $2^{-13/40} + \frac{2}{9}(\frac{3}{4})^{13/40} > 1$ certified by $0.798^{40} \leq 2^{-13}$ and $0.91^{40} \leq (\frac{3}{4})^{13}$ (`Production.zeta2_pos`), gives Theorem 5.3 for every $0 < \lambda \leq \frac{13}{40}$ with no hypothesis (`Production.contagion_elementary`, `Production.logMass_contagion_elementary`), and Corollary 5.5(2) at that exponent (`Production.failures_logMass_ge`), in `formal/Problems/Juggler/FateProduction.lean`. The root of $2^{-\lambda} + \frac{2}{9}(\frac{3}{4})^\lambda = 1$ is $0.32612\dots$, which $\frac{13}{40}$ reaches to within 0.4%; what lifts it to λ^{**} is the block-average family, which needs the two exponential-sum bounds of Proposition 4.4, and the ladder (5.10), which needs Appendix D. Composed with Theorem 7.2 and Corollary 8.4, whose contagion hypothesis it discharges, this gives two statements with no hypothesis on the contagion side: a Tao-type rate $\#\{\text{odd failures in } (y, 2y]\} \leq y(\log y)^{-e}$ with $e > \frac{27}{40}$ implies the conjecture (`Production.conjecture_of_tao_rate`), and so does a cylinder bound $H(C, A)$ with $e(C) > \frac{27}{40}$ (`Production.conjecture_of_cylinder_bound`). The paper's conditional forms need $e > 1 - \lambda^{**} \approx 0.51$ and the production inequality (5.2); the price of dropping the exponential sums at the pointwise coefficient is the stronger rate. Section 5.8 removes that price: at the averaged coefficient the same two productions give every $\lambda \leq 100/203$ with no hypothesis, and the two reductions at $e > 103/203$ (`Production.conjecture_of_tao_rate_averaged`, `Production.conjecture_of_cylinder_averaged`).

Corollary 5.4 (natural density, infinitely often). For every $\lambda < \lambda^{**}$ there is $c > 0$ such that for every $X \geq x_0$ some $y \in (\sqrt{X}, X]$ satisfies $\#(A \cap (y/2, y]) \geq cy(\log y)^{\lambda-1}$.

Proof. $\ell_A(\sqrt{X}, X] = g_A(\log X) \geq K(\log X)^\lambda$ is spread over at most $\frac{1}{2} \log_2 X + 2$ dyadic blocks $(y/2, y]$ with $\sqrt{X} < y \leq X$; one of them has $\ell_A(y/2, y] \geq K'(\log X)^{\lambda-1}$, hence $\#(A \cap (y/2, y]) \geq \frac{y}{2} \ell_A(y/2, y] \geq \frac{K'}{2} y (2 \log y)^{\lambda-1}$, because $\log X \leq 2 \log y$ and $\lambda - 1 < 0$. $\square$

Corollary 5.5 (fate contagion). Let φ be a fate realized by at least one start: the trivial cycle, a particular nontrivial cycle C , or divergence. Then $\{n : \text{fate}(n) = \varphi\}$ satisfies Theorem 5.3 and Corollary 5.4. In particular:

1. $\sum_{n \in R, n \leq x} 1/n \gg (\log x)^\lambda$ for every $\lambda < \lambda^{**}$.
2. If some start does not reach 1, then $\sum_{n \notin R, n \leq x} 1/n \gg (\log x)^\lambda$, and on infinitely many dyadic blocks the failures have natural density $\gg (\log y)^{\lambda-1}$.
3. If a nontrivial cycle exists, the set of starts that enter it has the same lower bounds; if a divergent orbit exists, so does the set of divergent starts.

Proof. Lemma 2.1 and Theorem 5.3. $\square$

Lean: Corollary 5.4 is `Density.exists_block_of_shell` in `formal/Problems/Juggler/FateDyadicDensity.lean`, the pigeonhole from a shell bound $K(\log X)^\lambda \leq \ell_A(\sqrt{X}, X]$ with the constant $K/12$: the shell is covered by the blocks $(X/2^{j+1}, X/2^j]$, $j \leq \log_2 X$, at most $3 \log X$ of them, a block's log-mass is at most $2 \#(A \cap (y/2, y])/y$, and $\log X < 2 \log y$ on the shell; `Density.natDensity_averaged` is the corollary for every $0 < \lambda \leq 100/203$ with no hypothesis, on Theorem 5.18. Corollary 5.5 is the instantiation at the fate classes through Lemma 2.1: `Density.reachesOne_logMass_averaged` and `Density.reachesOne_natDensity_averaged` for R ; `Production.failures_logMass_averaged` and `Density.failures_natDensity_averaged` for the failures; `Density.basin_logMass_averaged`, `Density.basin_natDensity_averaged`, `Density.escapes_logMass_averaged` and `Density.escapes_natDensity_averaged` for the basin of a cycle state and for the divergent starts. What stays human is the range $100/203 < \lambda < \lambda_{\text{ideal}}$, as for Theorem 5.18.

5.6 Remarks

The status of stronger density bounds. The theorem gives a logarithmic counting lower bound. It does not imply positive natural or logarithmic density. The ideal model of Section 5.7 has critical exponent 1, but the corresponding arithmetic hypotheses are not proved. Corollary 5.4 gives a natural-density lower bound on infinitely many blocks. No pointwise lower bound on every dyadic block is proved here. The even-only tree of a single seed is lacunary, but it need not be backward-closed under the full Juggler map and therefore is not a counterexample for that larger class.

Sharpness of the constants. The monotone pairing gives $\frac{1}{3}H_m - 2$; the observed floor on good fibers is 0.328 at $\alpha_m \approx \frac{1}{3}$ (Section 11). The depth-two gap $0.448 \rightarrow 0.4927$ was a dynamical averaging problem for the low-even set $P = \{m : G_m/H_m \leq 0.40\}$; Section 5.8 closes it without averaging over A , because P has finite total logarithmic mass (Theorem 5.16), so that the pointwise $\frac{1}{3}$ of the pairing is not needed. The nominal exponent ρ_w gives a useful scale diagnostic for candidate productions, but Model calculation 5.13 shows that it does not collapse the actual nested fiber to one monomial interval. The finite V_k constructions below use their own exact nested fibers; their limiting transfer-matrix value 0.4927 is a model ceiling, not a proved infinite-family exponent.

What is excluded. Nothing. Corollary 5.5 does not say that a cycle or a divergent orbit is impossible; it says that either would be common.

5.7 Finite productions and comparison models

Every production used above is a parity word w applied backwards. Write $a = \#\{E \text{ in } w\}$, $b = \#\{O \text{ in } w\}$. The source of a w -production sits at log-scale $\rho_w t$ with

$$\rho_w = 2^{-a} \left(\frac{3}{2}\right)^b, \quad c_w^{\text{ideal}} = \frac{2^{-|w|}}{\rho_w}, \quad (5.7)$$

the second being the log-mass coefficient when the fiber realizes the fair share $2^{-|w|}$. Formula (5.7) returns every coefficient in this section and in Appendix C: $c_E = 1$, $c_{OE} = c_{OEE} = \frac{1}{3}$, $c_{OOEEE} = \frac{1}{9}$.

A Collatz identity. With $\rho_w = (\frac{3}{2})^b 2^{-(|w|-b)}$ for a word with b odd letters, (5.7) simplifies to $c_w^{\text{ideal}} = 3^{-b}$: the ideal coefficient depends only on the odd count, and 3^{-b} is the probability that the b residue conditions modulo three demanded by the odd backward steps of the shortcut Collatz map all hold. Hence $\sum_w c_w^{\text{ideal}} \rho_w^\lambda = \sum_w 2^{-|w|} \rho_w^{\lambda-1}$, the exponential moment at $\lambda - 1$ of the fair-coin Collatz walk over the same words; over a bounded complete prefix-free family its two classical roots, Kraft equality at 0 and the martingale identity $\mathbb{E}\rho = \frac{1}{2}(\frac{3}{2} + \frac{1}{2}) = 1$ at 1, sit at $\lambda = 1$ and $\lambda = 2$. Boundedness matters for the second identity. The family of all minimal descent certificates is prefix-free and has Kraft mass one, by the survivor-decay estimate of Paper B. Every terminal multiplier is below one, and the certificate E has fair weight $1/2$ and multiplier $1/2$. Its total multiplier moment is therefore at most $1/4 + 1/2 = 3/4$, not one. For an unbounded family, preserving the second identity requires a separate limiting argument, such as vanishing multiplier-weighted survivor mass. The root 1 of Proposition 5.12 is Kraft equality read one exponential level up; the depth-two model, which lacks the ascending word OO , has the root λ_{ideal} , and Section 5.8 reaches that root as a supremum. The identity moves no constant; it says which half of this paper the Collatz walk reaches, the ceiling and not the engine.

Proposition 5.12 (an ideal-model critical exponent). For $0 < \eta \leq 1$, define $\lambda(\eta)$ as the root in $(0, 1]$ of the model characteristic equation

$$2^{-\lambda} + \frac{\eta}{3} \left(\frac{3}{2}\right)^\lambda = 1. \quad (5.8)$$

Then $\lambda(1) = 1$ and $\lambda'(1) = 1/\ln(4/3) = 3.4760\dots$. This is an algebraic model, not a Juggler counting theorem.

Proof. On $[0, 1]$, the left side is strictly decreasing: its derivative is at most $-\frac{1}{2}\ln 2 + \frac{1}{2}\ln(3/2) < 0$. At $\lambda = 0$ it is $1 + \eta/3 > 1$, and at $\lambda = 1$ it is $(1 + \eta)/2 \leq 1$. This gives a unique root in that interval, equal to 1 exactly for $\eta = 1$. Implicit differentiation at $(1, 1)$ gives the derivative. At $\eta = 1$ the two tilted weights are both $1/2$, the fair-coin path weights. They are not both $1/2$ for general η . $\square$

The root at 1 describes what the ideal characteristic equation permits. Exact nested fibers and their shares must still be proved before a proposed production list yields an arithmetic exponent.

Model calculation 5.13 (collapsed-power fiber; not an identity for the Juggler map). If one replaces the nested branch belonging to a production word w by the single model map $\tilde{J}_w(n) = \lfloor n^{\rho_w} \rfloor$, then its fiber over a source m' is exactly $I_{\text{mod}}(m') = \lfloor m'^{1/\rho_w}, (m' + 1)^{1/\rho_w} \rfloor$, and at scale $P \asymp m'^{1/\rho_w}$

$$|I_{\text{mod}}(m')| = \frac{1}{\rho_w} m'^{1/\rho_w - 1} (1 + O(1/m')) \asymp P^{1 - \rho_w}.$$

The corresponding assertion for the exact map is false. For example, 1015 follows

$$1015 \xrightarrow{O} 32336 \xrightarrow{E} 179 \xrightarrow{O} 2394 \xrightarrow{E} 48 \xrightarrow{E} 6,$$

so its word is $OEOEE$ and $J^5(1015) = 6$. But $7^{32} \leq 1015^9 < 8^{32}$, hence $\lfloor 1015^{9/32} \rfloor = 7$, not 6. Nested floors cannot in general be replaced by one final floor.

Appendix C's localized parity estimate requires an *actual proved* parameter interval of length at least $P^{1/2}$. Thus $\rho_w \leq \frac{1}{2}$ is only the collapsed model's candidate threshold; it neither proves the needed interval nor the ideal share. The words E , OEE , and conditionally $OEOEE$ have separate fiber arguments in their cited results. The nominal lengths $P^{1/2}$, $P^{5/8}$, $P^{23/32}$ remain useful bookkeeping, while the nominal OE and $OEOEE$ lengths $P^{1/4}$ and $P^{7/16}$ only screen where that particular localization estimate could apply.

The conditional run model and the K_3 interface. Production words must be prefix-free. First passage of the nominal ρ_w below $\frac{1}{2}$ gives a candidate list, not a theorem that its fibers have the model length or ideal share. A run of r consecutive O 's forces r nested $3/2$ -powers plus the closing square root — Paper B at depth $r + 1$. Paper B is complete to depth 4, and to depth 5 except $OOOO*$, which is the parked kernel K_3 . Under the additional hypothesis that every required candidate production through run length r is established at the stated share, the transfer matrix gives:

r	Paper B depth	conditional $\lambda(r)$, pairing share	conditional $\lambda(r)$, ideal-share model
1	2	0.4480	0.4927
2	3	0.6247	0.7180
3	4	0.7095	0.8414
4	5	0.7516	0.9121

The table is a price list of hypothetical inputs, not an attained ladder. Appendix C establishes one $r = 2$ word conditionally and prints 0.5392; nothing here supplies the other words needed for 0.6247 or 0.7180. Row $r = 4$ would also meet the parked K_3 kernel, but the model does not prove that this is the only obstruction.

The finite productions used in Theorem 1. Put $V_k = (OE)^{k-1}OEE$, for $1 \leq k \leq 6$, and $\rho_k = \frac{1}{2}(3/4)^k$. These six words are prefix-free. The exact landing windows have iterated ceiling endpoints; they are not the single-power windows of Model calculation 5.13. Appendix

D proves, for each of these fixed words, the needed two-sided asymptotic and the resulting logarithmic coefficient $c_k = 3^{-k}$.

The V_k -starts for $k \geq 2$ are contained in family 3 of Section 5.1: their two-step images are odd V_{k-1} -produced members of A . Before counting them at their improved share, remove those source sets from family 3. The old contribution is $(2/9)c_{k-1}$; the new one is c_k . Their difference is

$$c_k - \frac{2}{9}c_{k-1} = 3^{-(k+1)}, \quad 2 \leq k \leq 6. \quad (5.9)$$

Disjointness and the two-sided estimates justify making all five replacements simultaneously. Thus (5.2) gains

$$\sum_{k=2}^6 (3^{-(k+1)} - o(1))g_A(\rho_k t) - o(1). \quad (5.10)$$

The corresponding roots are 0.4801, 0.4891, 0.4916, 0.4924, and 0.4926. This is the finite input used in Theorem 5.3.

For comparison only, summing the formal infinite series in (5.9) gives the ideal depth-two equation $2^{-\lambda} + \frac{1}{3}(3/4)^\lambda = 1$, with root 0.4927. Neither a uniform estimate in k nor attainment of that endpoint is asserted. Adding just V_2 to the conditional construction in Appendix C would give root 0.5665; the analogous infinite model gives 0.5769. These comparisons are not the exponents claimed in Theorems 1 or C.5.

5.8 The averaged coefficient: two productions without exponential sums

Section 5.1 splits the OE -images into two families, item 2 over the even blocks, where Proposition 4.4 controls the share, and item 3 over the rest, where the pointwise floor of Lemma 4.2 applies, and the ladder of Section 5.7 recovers part of what the pointwise floor loses. The split exists only to serve the block average. This section takes the OE family over all of $A \cap (x^{3/8}, x^{3/4}]$ at once — the E -images are even and the OE -images odd, which is the only disjointness Section 5.1 uses — and replaces the pointwise coefficient by an averaged one, at the price of discarding the fibers whose parity share is far from $\frac{1}{2}$. Those fibers have finite total logarithmic mass, so the price is nothing. No exponential sum, Vaaler approximation or second-derivative test appears below; the tools are Dirichlet approximation and counting, and the chain is formalized end to end. Full proofs are in the laboratory note [14]; the statements are given here with their constants.

Notation. For $m \geq 10^6$ let $\Phi(m)$ be the OE fiber of Lemma 3.2, $H_m = |\Phi(m)|$, G_m the number of its members whose image is even, and $\sigma_m = G_m/H_m$ the parity share. The proof of Lemma 4.2 writes the parity of the image of the j -th member as the half-cell of a real x_j whose consecutive steps lie in an interval $[A_m, A_m + \eta_m]$ with $A_m = \frac{3}{2}m^{2/3}$ and $\eta_m \leq m^{-1/3}$ (Lean: `FiberParity.xval_step`); put $\alpha_m = \{A_m\}$. By Lemmas 3.2 and 4.2, $\eta_m H_m \leq 0.6903$ for $m \geq 10^6$.

Lemma 5.14 (block lock). Let $m \geq 10^6$ and let p/q be in lowest terms with $q \geq 1$. Then

$$\left| \sigma_m - \frac{1}{2} \right| \leq 4|q\alpha_m - p| + \frac{5}{2q} + \frac{3.77q}{H_m}.$$

Proof. Cut the fiber into consecutive blocks of q members. Inside a block the points $\{x_j\}$ form a translate of the grid $\frac{1}{q}\mathbb{Z}$ perturbed by at most $\varepsilon = |q\alpha_m - p| + q\eta_m$, because $\gcd(p, q) = 1$ makes the q points $\{y + ip/q\}$ a full grid; a $\frac{1}{q}$ -grid meets a half-circle within $\frac{1}{2}$ of evenly, and at most $2\varepsilon q + 1$ of its points lie within ε of a given point, so each block's even count is within $4\varepsilon q + \frac{5}{2}$ of $q/2$; the remainder block contributes at most q . Summing, $|G_m - \frac{1}{2}H_m| \leq 4\varepsilon H_m + 5H_m/(2q) + q$, and $1 + 4\eta_m H_m \leq 3.77$. $\square$

Lemma 5.15 (fiber lock). Let $\eta_0 \in (0, \frac{1}{2}]$ and $m \geq 10^6$ with $H_m \geq 1280/\eta_0^2$. If $|\sigma_m - \frac{1}{2}| \geq \eta_0$, then some integer q with $1 \leq q \leq 3.77/\eta_0$ satisfies $\|q\alpha_m\| \leq 32/(\eta_0 H_m)$.

Proof. Put $\theta = \eta_0/16$ and $H_0 = \lfloor \theta H_m \rfloor \geq \theta H_m/2$. Dirichlet's theorem gives $q \leq H_0$ and p with $|q\alpha_m - p| < 1/H_0$, and passing to lowest terms decreases both q and $|q\alpha_m - p|$. In Lemma 5.14 the first term is then at most $8/(\theta H_m) = 128/(\eta_0 H_m) \leq \eta_0/10$ and the third at most $3.77\theta = 0.2356\eta_0$, so $|\sigma_m - \frac{1}{2}| \geq \eta_0$ forces $5/(2q) \geq 0.6644\eta_0$, that is $q \leq 3.77/\eta_0$; and $|q\alpha_m - p| < 1/H_0 \leq 32/(\eta_0 H_m)$. $\square$

Theorem 5.16 (poor-fiber tail). Let $\eta_0 \in (0, \frac{1}{2}]$, $P_{\eta_0} = \{m \geq 10^6 : |\sigma_m - \frac{1}{2}| \geq \eta_0\}$ and $u_0(\eta_0) = \max(10^6, (1950/\eta_0^2)^3)$. For every $u \geq u_0$,

$$\#(P_{\eta_0} \cap (u, 2u]) \leq \frac{430}{\eta_0^2} u^{2/3},$$

and for every $U \geq u_0$, $\sum_{m \in P_{\eta_0}, m > U} 1/m \leq 2100 U^{-1/3}/\eta_0^2$.

Proof. On $(u, 2u]$, $H_m \geq \frac{2}{3}u^{1/3} - 1$ (Lemma 3.2), and $u \geq (1950/\eta_0^2)^3$ makes Lemma 5.15 available with $Q = 3.77/\eta_0$ and $\delta = 32/(\eta_0 H_m)$. Every poor m is therefore resonant, $\|q\alpha_m\| \leq \delta$ for some $q \leq Q$, and the arc count of Lemma 4.3, applied for each $q \leq Q$ in turn to the sequence $q \cdot \frac{3}{2}m^{2/3}$, whose steps and total increase scale by q , bounds the resonant m by $(0.882u^{2/3} + 2)(2Q\delta(2u)^{1/3} + Q(Q+1)) \leq 430u^{2/3}/\eta_0^2$. The second bound is the first summed over dyadic blocks. $\square$

The constants are crude and the exponent is not: the census finds the true density of the fibers with $\sigma_m \leq 0.40$ between $1.4u^{-1/3}$ and $2.1u^{-1/3}$ over $u \in [10^5, 10^8]$ [14]. What is used below is only that the tail vanishes: the poor set has finite total logarithmic mass, so it can support no set at every scale.

Corollary 5.17 (the share on an arbitrary set). For $\eta_0 \in (0, \frac{1}{2}]$, $V \geq u_0(\eta_0)$ and every set $S \subseteq (V, \infty)$ of integers,

$$\sum_{m \in S} \frac{\sigma_m}{m} \geq \left(\frac{1}{2} - \eta_0\right) \sum_{m \in S} \frac{1}{m} - \frac{1050}{\eta_0^2} V^{-1/3}.$$

Proof. Off P_{η_0} the bound holds termwise; on $S \cap P_{\eta_0}$ use $\sigma_m \geq 0$ and discard at most $1/(2m)$ per term, which Theorem 5.16 sums to at most $1050 V^{-1/3}/\eta_0^2$. $\square$

Theorem 5.18 (contagion by two productions). Let A be nonempty and backward-closed, $0 < \eta_0 \leq \frac{1}{4}$, and $\lambda > 0$ with

$$\zeta(\lambda, \eta_0) = 2^{-\lambda} + \frac{2}{3}\left(\frac{1}{2} - \eta_0\right)\left(\frac{3}{4}\right)^\lambda - 1 > 0.$$

There are $K > 0$ and t_1 with $g_A(t) \geq Kt^\lambda$ for $t \geq t_1$, hence $\sum_{n \in A, n \leq x} 1/n \geq K(\log x)^\lambda$ for $x \geq x_0$. Consequently the conclusion of Theorem 5.3 holds for every $0 < \lambda < \lambda_{\text{ideal}}$, the root of $2^{-\lambda} + \frac{1}{3}(\frac{3}{4})^\lambda = 1$, $\lambda_{\text{ideal}} = 0.49265798\dots$: given such a λ , some η_0 has $\zeta(\lambda, \eta_0) > 0$.

Proof. For $m \in A \cap (x^{3/8}, x^{3/4} - 1]$ the fibers $\Phi(m)$ lie in $(\sqrt{x}, x]$, are pairwise disjoint and disjoint from the E -images, each $n \in \Phi(m)$ has $1/n > (m+1)^{-4/3}$, and $H_m(m+1)^{-4/3} \geq \frac{2}{3m}(1 - O(m^{-1/3}))$ by Lemma 3.2. Corollary 5.17 with $V = x^{3/8}$, whose error $V^{-1/3} = x^{-1/8}$ has the shape Lemma 4.3 already contributes, gives the OE family log-mass at least $\frac{2}{3}(\frac{1}{2} - \eta_0)(1 - O(x^{-1/8}))g_A(3t/4) - 700x^{-1/8}/\eta_0^2$, and with item 1 of Section 5.1 unchanged,

$$g_A(t) \geq (1 - \varepsilon_1(t))g_A(t/2) + \left(\frac{1}{3} - \frac{2}{3}\eta_0 - \varepsilon_7(t)\right)g_A(3t/4) - \varepsilon_8(t),$$

with $\varepsilon_7, \varepsilon_8 \rightarrow 0$ for fixed η_0 . Lemma 5.1 with the seed of Lemma 5.2 finishes as in Theorem 5.3. $\square$

Corollaries 5.4 and 5.5 use Theorem 5.3 only through its conclusion, so by Theorem 5.18 each holds for every $\lambda < \lambda_{\text{ideal}}$; so do Corollary 7.1 and Theorems 7.2 and 7.3, which are stated in that form. Both corollaries are Lean at 100/203 (Section 5.5).

Lean: the whole chain. `BlockLock.block_lock` in `formal/Problems/Juggler/Fate` `BlockLock.lean` is Lemma 5.14 before division by H_m , a statement about an arbitrary real

sequence with steps in $[a, a + \eta]$; `FiberParity.fiber_lock` in `FateFiberLock.lean` is Lemma 5.15, through Dirichlet's theorem in lowest terms (`exists_coprime_approx`) and the fiber instantiation `fiber_block_lock`; `FiberParity.resonance_count_le'` in `FateResonanceCount.lean` is the arc count; `FiberParity.poor_count_le'` and `poor_logMass_le` in `FatePoorTail.lean` are the two bounds of Theorem 5.16, and `nonpoor_fiber_logMass_ge` the per-fiber bound behind Corollary 5.17. In `FatePoorProduction.lean`, `Production.productio` `n_two_averaged` is the production inequality above, with error $2e^{-t/8}$ on the OE coefficient and $2e^{-t/2} + \frac{2}{3}e^{-3t/4} + 1400e^{-t/8}/\eta_0^2$ additive, under the one new hypothesis $3(1280/\eta_0^2 + 1) \leq e^{t/8}$, which is the block condition of Lemma 5.15 at $U = \lfloor e^{3t/8} \rfloor$ and the whole cost of averaging; `Production.contagion_averaged` is Theorem 5.18 with λ and η_0 free and $\zeta(\lambda, \eta_0) > 0$ as its hypothesis. The exponent is a separate arithmetic certificate: `Production.zeta2avg_pos` checks $0.710737 \leq 2^{-100/203}$ and $216967/250000 \leq (\frac{3}{4})^{100/203}$ as 203rd powers in exact integer arithmetic, so that $\zeta(100/203, 10^{-5}) > 2.05 \cdot 10^{-5}$, and `zeta2avg_antitone` extends it to every $\lambda \leq 100/203$. Hence `Production.logMass_contagion_averaged`: for every nonempty backward-closed A and every $0 < \lambda \leq 100/203$, with no hypothesis, $\sum_{n \in A, n \leq x} 1/n \geq K(\log x)^\lambda$ for $x \geq x_0$, above $\lambda^{**} = 0.4925715 \dots$ of the first route; and with it `failures_logMass_averaged` (Corollary 5.5(2) at $100/203$), `conjecture_of_tao_rate_averaged` (Theorem 7.2 at $e > 103/203$) and `conjecture_of_cylinder_averaged` (Corollary 8.4 at $e(C) > 103/203$). The certificate is re-derived in exact rational arithmetic by `oe_rest_average.exponent_certificate`, with the smaller denominators enumerated: $67/136$ also lies in $(\lambda^{**}, \lambda_{\text{ideal}})$ but leaves a tenth of the margin, and $33/67$ falls short of λ^{**} . The break-even is $\eta_0 = 4.69 \cdot 10^{-5}$ at $100/203$ and $8.60 \cdot 10^{-5}$ at λ^{**} ; the range $100/203 < \lambda < \lambda_{\text{ideal}}$ is the choice of η_0 in Theorem 5.18 and is not separately certified.

What changes is the critical path of Theorem 1, not its number: Proposition 4.4 with its two exponential-sum bounds, the ladder of Section 5.7 with Appendix D, and the statements of Lemmas 4.1, 4.1' and 4.2 are no longer needed for the exponent; one step of Lemma 4.2's proof, the step interval, is. What does not change: the supremum is approached and not attained, η_0 being fixed before x , as in Theorem 5.3; the constants are large (at $\eta_0 = 4.3 \cdot 10^{-5}$, $u_0 \approx 1.2 \cdot 10^{36}$, $x_0 \approx 10^{96}$, $t_1 \approx 221$); and no standing question moves — the exponent is below 1, the rate of Theorem 7.2 is not proved, no cycle is excluded and no orbit is shown to reach 1.

5.9 Three actual productions and exponent 5/8

This section gives the strongest written contagion result. The analytic poor-fiber estimate used here is proved in Appendix E; independent review and complete formal verification of that estimate remain outstanding. The earlier Section 5.8 supplies the unconditional Lean baseline and the positive seed needed below.

Theorem 5.19 (three-production contagion). Every nonempty backward-closed set A of positive integers satisfies

$$\sum_{n \leq X, n \in A} \frac{1}{n} \geq K(\log X)^{5/8} \quad (X \geq X_0) \quad (5.11)$$

for some $K > 0, X_0$. The same conclusion holds for each realized fate class. The theorem is a written proof; its formal assembly retains the actual OOOE production inequality as a hypothesis.

Proof. Put

$$a(n) = n + (n \bmod 2), \quad w(n) = \log \frac{a(n) + 1}{a(n) - 1}, \quad F(t) = \sum_{n \leq \lfloor e^t \rfloor, n \in A} w(n).$$

For positive integers, $1/n \leq w(n) \leq 4/n$. Let $S_E(t), S_{OE}(t), S_{OOEE}(t)$ be the corresponding weighted masses of starts in A , below $\lfloor e^t \rfloor$, with the indicated actual initial words. The three

families are disjoint. Exact even-fiber conservation gives, for $t \geq 8$,

$$F(t/2 - 4) \leq S_E(t).$$

The OE poor-fiber theorem of Section 5.8, converted to this weight, and Appendix E's OOE theorem give constants C, T such that

$$\frac{33}{100}F(3t/4 - 4) \leq S_{OE}(t) + C, \quad \frac{11}{100}F(9t/16 - 4) \leq S_{OOE}(t) + C \quad (t \geq T). \quad (5.12)$$

Here all cutoffs concern actual starts. An OE predecessor of m has $n^3 < (m+1)^4$; an OOE predecessor has $n^9 < 64(m+1)^{16}$. Since $m+1 \leq 2m$, the target cutoffs $m \leq e^{3t/4-4}$ and $m \leq e^{9t/16-4}$ place their sources below e^t . Distinct target fibers are disjoint, and backward closure puts their sources in A .

For completeness, the weighted OE coefficient follows from $H_m = (2/3)m^{1/3} + O(1)$, source weight $2m^{-4/3}(1 + O(1/m))$, and target weight $2m^{-1}(1 + O(1/m))$. The actual fiber ratio is $(2/3)\sigma_m + O(m^{-1/3})$. Outside a poor set of finite reciprocal mass it exceeds $33/100$, after a finite initial segment is removed. The removed target weight is bounded independently of A and the cutoff. The OOE coefficient is Appendix E's arbitrary-target consequence with $\eta = 1/900$, since $1/9 - 1/900 = 11/100$. This proves (5.12), including its uniform additive loss. The weighted OE step also has a separate Lean proof.

Disjointness now yields

$$F(t) \geq F(t/2 - 4) + \frac{33}{100}F(3t/4 - 4) + \frac{11}{100}F(9t/16 - 4) - 2C.$$

Let $G(t) = F(t - 16) - 5C$. For each $r \in \{1/2, 3/4, 9/16\}$, $rt - 16 \leq r(t - 16) - 4$. Monotonicity, and the coefficient sum $36/25$, therefore give eventually

$$G(t) \geq G(t/2) + \frac{33}{100}G(3t/4) + \frac{11}{100}G(9t/16). \quad (5.13)$$

The constant remainder is $5C(36/25 - 1) - 2C = C/5 \geq 0$. Section 5.8 implies $F(t) \rightarrow \infty$, so G has a positive seed on a sufficiently late compact interval. Exact rational comparisons, obtained by taking eighth powers, give

$$2^{-5/8} \geq \frac{648}{1000}, \quad (3/4)^{5/8} \geq \frac{835}{1000}, \quad (9/16)^{5/8} \geq \frac{697}{1000}.$$

Their weighted sum is at least $50011/50000 > 1$. The recursion lemma with zero errors gives $G(t) \geq Kt^{5/8}$. Evaluate at $t = \log X$ and use the weight comparison to obtain (5.11). $\square$

Dyadic consequence. There is $c > 0$ such that infinitely many dyadic intervals $(y, 2y]$ contain at least $cy(\log y)^{-3/8}$ members of A . Otherwise their reciprocal masses are eventually at most $c(\log y)^{-3/8}$. Summing over $y = 2^k$ up to X bounds the total by a constant times $c(\log X)^{5/8}$, plus a fixed initial mass. Choosing c below the constant in (5.11) is a contradiction. Smaller positive exponents follow by weakening (5.11). This proves all clauses of Theorem 1; the dyadic consequence at this new exponent is written, not separately audited in Lean.

Formal boundary. `FateOOEEAssembly` proves disjointness, the even cutoff, the shifted recursion, the rational certificate and the contagion and Tao implications. `FateOEWeighted` proves the OE input, including a uniform error at most 6 when converting its finite weighted sums to twice reciprocal mass. The strengthened implication still assumes `OOEEProductionBound`. The original mixed-mode cancellation is proved in `OOEEMixedModes`; a fixed finite frequency family has uniform constants. The remaining analytic assembly in Appendix E is not thereby certified. These supplementary modules have their own audits; the historical Paper C barrel and its 473 recorded reports retain their earlier scope.

6. Odd generation and the exact first-letter decomposition

Write $S = \{\lfloor m^{3/2} \rfloor : m \text{ odd}\}$ for the image of the odd integers, and S_{odd} for its odd members. The set S is sparse: it has $\asymp z^{2/3}$ members below z .

6.1 Odd generation

Theorem 6.1 (odd generation; Lean). Let A be forward- and backward-closed with $1 \notin A$. Then:

1. every $n \in A$ descends by even steps only to an odd member $m \geq 3$ of A (`exists_odd_ancestor_ge_three`);
2. for odd n , $n \in A \iff \lfloor n^{3/2} \rfloor \in A$ (`odd_mem_iff`), so the odd members of A are exactly the odd preimages of $A \cap S$;
3. $A \neq \emptyset \iff A \cap S \neq \emptyset$, i.e. A contains the image of some odd $m \geq 3$ (`nonempty_iff_odd_image_mem`).

Proof. (1) If $n \in A$ is even, $\lfloor \sqrt{n} \rfloor \in A$ by forward closure and is smaller; the chain stops at an odd integer, which is ≥ 3 since $1 \notin A$ and $2 \notin A$ ($J(2) = 1$). (2) Forward closure gives one direction, backward closure the other. (3) follows from (1) and (2). $\square$

Applied to the failure set F : every positive integer reaches 1 iff no image $\lfloor m^{3/2} \rfloor$ of an odd m fails to reach 1, and F is the E -forest over the odd preimages of $F \cap S$. The theorem changes the geometry of the problem. An arbitrary failure set is replaced by a set of *seeds* — odd images, a set of density $\asymp z^{-1/3}$ at scale z — together with the deterministic even ancestry of each odd preimage (its E -forest, whose levels are the intervals of Lemma 3.1) and the unique odd preimage of each seed in S . Two consequences are used below: the log-count of F is controlled by that of its odd members (Theorem 7.2), and the first-letter decomposition of F has exactly one term that the productions do not control (Section 6.2).

6.2 The exact decomposition and the free term

Let A be forward- and backward-closed. On $(\sqrt{x}, x]$ its members split by first letters into three exact pieces (Figure 3):

- the **even** members, $= \bigsqcup_{m \in A} E(m) \cap (\sqrt{x}, x]$, of log-mass $\ell_A(x^{1/4}, \sqrt{x}) + O(x^{-1/4})$ (Lemma 3.1);
- the **OE-type** odd members ($\lfloor n^{3/2} \rfloor$ even), $= \bigsqcup_{m \in A} \{n \in \Phi(m) \cap (\sqrt{x}, x] : \lfloor n^{3/2} \rfloor \text{ even}\}$ (Lemma 3.2), of log-mass between $\frac{2}{9}$ and $\frac{4}{9}$ times $\ell_A(x^{3/8}, x^{3/4})$ on good fibers (Lemma 4.2, two-sided pairing) and $\frac{1}{3}(1 + o(1))$ times it on E -blocks (Proposition 4.4);
- the **OO-type** odd members ($\lfloor n^{3/2} \rfloor$ odd), $= \{n(m) : m \in A \cap S_{\text{odd}}, \sqrt{x} < n(m) \leq x\}$, of log-mass $\sum_{m \in A \cap S_{\text{odd}}, \sqrt{x} < n(m) \leq x} 1/n(m)$, where $n(m)$ is the unique odd preimage. Writing the cutoff on $n(m)$ keeps the floor endpoints exact.

Let $\varphi_A(t) = \ell_A(e^{t/2}, e^t)/(t/2)$ be the logarithmic density of A on $(\sqrt{x}, x]$, $t = \log x$. Define $\varphi_A^{\text{fib}}(3t/4)$ by the normalization $\frac{1}{4} \cdot \frac{t}{2} \cdot \varphi_A^{\text{fib}}(3t/4) = \text{log-mass of the OE-type members}$, and $\psi_A(t)$ by $\frac{1}{4} \cdot \frac{t}{2} \cdot \psi_A(t) = \text{log-mass of the OO-type members}$. Then $\varphi_A^{\text{fib}}(s)$ is the density of A on $(e^{s/2}, e^s]$ weighted, on each fiber, by twice the even-image fraction $p_m = G_m/H_m$ of that fiber — so the weighted and unweighted masses agree to vanishing relative error on complete E -block unions, with an additional exponentially small boundary error (Proposition 4.4), and $\varphi_A^{\text{fib}}(s) \leq \frac{4}{3}\varphi_A(s) + O(e^{-s/6})$ always (Lemmas 4.2, 4.3) — and $\psi_A(t)$ is the log-weighted share of A among the OO -type odd integers of $(\sqrt{x}, x]$, up to a factor $1 + O(e^{-\delta t})$ from the depth-two

equidistribution of the parity of $\lfloor n^{3/2} \rfloor$. With these definitions the three pieces give the identity with boundary error

$$\varphi_A(t) = \frac{1}{2}\varphi_A(t/2) + \frac{1}{4}\varphi_A^{\text{fib}}(3t/4) + \frac{1}{4}\psi_A(t) + O(e^{-t/4}/t), \quad (6.1)$$

whose only error is the boundary term of Lemma 3.1. For $A = \mathbb{N}$ every term is $1 + o(1)$. For $A = F$ the boundary condition is $\varphi_F(t) = 0$ for $t \leq \log N_0$.




Figure 3: The first-letter decomposition of a two-way closed set on $(\sqrt{x}, x]$: the even members come from scale $x^{1/2}$ through *E*-blocks, the *OE*-type odd members from scale $x^{3/4}$ through fibers, and the *OO*-type odd members are the odd preimages of the odd images of A at scale $x^{3/2}$ — the free term.

Definition 6.2 (*S*-fairness). A two-way closed set A is *S-fair at scale t* if $\psi_A(t) \leq \varphi_A(3t/2)$, i.e. if A is not over-represented among the odd images of odd integers at scale $e^{3t/2}$ relative to its density among all integers there; it is *S-fair* if this holds for all $t \geq t_0$.

Proposition 6.3 (exact statements). (i) $\psi_F \equiv 0$ iff $F = \emptyset$: if $F \neq \emptyset$, its minimum is odd with odd image, so $\psi_F(t) > 0$ on a shell containing $\min F$. (ii) The identity (6.1) holds for F with the stated error, and its right side contains exactly one term, $\frac{1}{4}\psi_F(t)$, that is not determined by F at smaller scales through the two productions.

Proof. (i) If $F = \emptyset$ then $\psi_F \equiv 0$. If $F \neq \emptyset$, let $n_F = \min F$. It is odd (an even failure has the smaller failure $\lfloor \sqrt{n_F} \rfloor$), and its image is odd (if $J(n_F)$ were even, $J^2(n_F) = \lfloor n_F^{3/4} \rfloor < n_F$ would be a smaller failure). So n_F is an *OO*-type failure and $\psi_F(t) > 0$ for t with $e^{t/2} < n_F \leq e^t$. (ii) is the decomposition. $\square$

Lean: `minimal_failure_odd_odd` and `exists_minimal_failure` (the least failure is odd with odd image), and the three pieces as `first_letter_trichotomy` with `first_letter_pieces_disjoint`, in `formal/Problems/Juggler/FateFirstLetter.lean`. The exact layer of (6.1) is `first_letter_split` there: for any weight and any finite index set, the weighted mass of a two-way closed class is the sum of its even, *OE*-type and *OO*-type pieces, each indexed as the paper indexes it, by the image lying in the class; `shellLogMass_split` is the log-mass form on a shell $(y, x]$. The free term's $n(m)$ is well defined by `floorPower_odd_injective` (J is strictly increasing on the odd integers, `floorPower_odd_lt`), and `sum_image_oopiece` rewrites the third piece as the paper's sum over the odd images. What is not formalized is the normalization: the densities φ_A , φ_A^{fib} , ψ_A , the identification of each piece with its term of (6.1), and the boundary error $O(e^{-t/4}/t)$.

Remark 6.4 (the walk heuristic; not a theorem). If F were *S*-fair, if the fiber weights were ideal ($\varphi_F^{\text{fib}} = \varphi_F$), and if the error term of (6.1) were zero, then (6.1) would give $\varphi_F(t) \leq \frac{1}{2}\varphi_F(t/2) + \frac{1}{4}\varphi_F(3t/4) + \frac{1}{4}\varphi_F(3t/2)$: the harmonic inequality of a random walk on

$\log t$ with steps $\log \frac{1}{2}, \log \frac{3}{4}, \log \frac{3}{2}$ of probabilities $\frac{1}{2}, \frac{1}{4}, \frac{1}{4}$, whose drift -0.317 is negative, so that the boundary condition $\varphi_F = 0$ below $\log N_0$ would force $\varphi_F \equiv 0$. None of the three conditions is available with the constants proved here: the fiber weights are only known to lie in $[\frac{2}{3}, \frac{4}{3}]$, so the coefficients of the inequality can sum to $\frac{1}{2} + \frac{1}{3} + \frac{1}{4} = \frac{13}{12} > 1$, and the boundary error $O(e^{-t/4}/t)$ is not zero at the scales just above the floor where the walk exits. We therefore do not claim that S -fairness implies the conjecture. What is exact is Proposition 6.3; what is proved about the free term is Proposition 10.1, which identifies $\psi_F(t)$ with a quantity bounded by every hypothesis of Sections 8–9.

6.3 Where Papers A and B sit

Paper A [11] constrains the fate Lachesis from the inside: finance and the walk charge bound the *states* of a hypothetical cycle (minimum $> 3.5 \cdot 10^8$, period ≥ 780239). Theorem 1 constrains the basin: if the cycle exists, its basin is a two-way closed class with the cycle's states as seeds and $\log$ -count $\gg (\log x)^\lambda$ for every $0 < \lambda \leq 5/8$, using the written Theorem 5.19. In the present argument the two estimates do not meet: finance bounds the seed, contagion the growth from the seed, and no inequality bounds a basin from above. Paper B [12] controls the descending branches of (6.1) — the fairness of the landing distributions of E , OE and, with its depth-4 and depth-5 theorems, of deeper contracting words — on dyadic blocks; Appendix C uses one statement of that type on sub-dyadic intervals as an explicit hypothesis. Neither touches ψ_F : the ascending branch OO sends mass to $x^{3/2}$, and its return is the nested-floor parity at all depths. The two papers' large-deviation exponents are one rate function read at two levels: the L -bad condition of Section 8 at depth $d = CL$ demands an odd share above $p_C = (1 - 1/C)/\log_2 3$, and $e(C)/C \rightarrow 1 - H(\log 2/\log 3) = 0.050044$ bits as $C \rightarrow \infty$, the exponent of Lagarias's Theorem D [3] that Paper B's Section 6 uses; Paper B sits at $L = 0$, where the threshold is $\log 2/\log 3$ itself. The gap is $4.8 \cdot 10^{-3}$ at $C = 100$ and closes like $1/C$.

7. The almost-all equivalence

Corollary 7.1 (logarithmic form). The following are equivalent: (1) every $n \geq 1$ reaches 1; (2) for some $0 < \lambda \leq 5/8$, $\sum_{n \leq x, n \notin R} 1/n = o((\log x)^\lambda)$; (3) for some $0 < \lambda \leq 5/8$, the starts $n \leq x$ whose orbit never enters $[1, N_0]$ have logarithmic count $o((\log x)^\lambda)$.

Proof. (1) $\Rightarrow$ (3): the set is empty. (3) $\Rightarrow$ (2): an orbit that enters $[1, N_0]$ reaches 1, so F is contained in the set of (3). (2) $\Rightarrow$ (1): F is backward-closed; if it were nonempty, Theorem 5.19 would contradict (2). $\square$

Theorem 7.2 (a Tao-type bound with rate implies the conjecture). Suppose that for some $e > 3/8$ and all sufficiently large y ,

$$\#\{n \text{ odd}, y < n \leq 2y : n \notin R\} \leq \frac{y}{(\log y)^e}.$$

Then $R = \mathbb{N}$.

Proof. By decreasing e if necessary, assume $3/8 < e < 1$; the hypothesis is preserved for large y . Suppose $F \neq \emptyset$. By Theorem 6.1 every $n \in F$ lies in the E -tree of an odd member of F . For an odd $n_0 \in F$ the level- j set $S_j(n_0)$ of its E -tree has log-mass at most $\prod_{i < j} (1 + n_0^{-2^i})/n_0 \leq 2/n_0$ (Lemma 3.1 gives $\sum_{n \in E(m)} 1/n \leq (m+1)/m^2$), and only levels with $n_0^{2^j} \leq x$, i.e. $j \leq \log_2 \log x$,

meet $[1, x]$. Hence

$$\begin{aligned} \sum_{\substack{n \in F \\ n \leq x}} \frac{1}{n} &\leq 2(1 + \log_2 \log x) \sum_{\substack{n_0 \in F \text{ odd} \\ n_0 \leq x}} \frac{1}{n_0} \\ &\leq 2(1 + \log_2 \log x) \left(\sum_{k_0 \leq k \leq \lfloor \log_2 x \rfloor} (k \ln 2)^{-e} + O(1) \right) \\ &\ll (\log x)^{1-e} \log \log x. \end{aligned}$$

By Theorem 5.19 the left side is $\geq K(\log x)^\lambda$ for every $\lambda < 5/8$ and all large x . Choosing $\lambda \in (1 - e, 5/8)$ gives a contradiction. $\square$

Theorem 7.3 (equivalence). Every positive integer reaches 1 if and only if there is $e > 3/8$ such that $\#\{n \text{ odd} \in (y, 2y] : n \notin R\} \leq y(\log y)^{-e}$ for all large y .

Proof. If every integer reaches 1 the left side is 0. Conversely Theorem 7.2. $\square$

Lean: `tao_rate_iff_conjecture` in `formal/Problems/Juggler/FateContagionBound.lean`, with the contagion bound of Theorem 5.19 as a hypothesis, as for Theorem 7.2; and `conjecture_of_cylinder_bound_of_production`, Corollary 8.4 with the earlier Theorem 5.3 replaced by the production inequality (5.2), so that the chain from $H(C, A)$ to the conjecture has (5.2) as its only analytic input, for any $\lambda \leq 0.49$ with $1 - \lambda < e(C)$ ($C \geq 19$ still suffices: $e(19) = 0.527 > 0.51$). With the contagion hypothesis discharged by Theorem 5.18, `Production.conjecture_of_tao_rate_averaged` is Theorem 7.2 at $e > 103/203$ and `Production.conjecture_of_cylinder_averaged` is Corollary 8.4 at $e(C) > 103/203$, with no hypothesis on the contagion side (Section 5.8).

The threshold $3/8$ is the complement of the contagion exponent. Appendix C's earlier conditional exponent gives the weaker threshold 0.4608. Any improvement of the contagion exponent lowers the rate required of the almost-all statement, and $\lambda \rightarrow 1$ would make any positive rate suffice — but that improvement requires all descent certificates and is circular here (Section 5.6).

7.1 Comparison with the Collatz map: a structural analogy

The comparison with Tao's theorem [6] is a structural analogy, not a transfer of methods, and it concerns two specific features.

Depth. Juggler's multiplicative changes of $\log n$ motivate a walk on $\log \log n$ and the depth $O(\log \log y)$. This is the depth used by the sufficient hypotheses here, not a proved stopping-time bound for every orbit. For the accelerated Collatz map of Terras and Everett, parity words of length k correspond to residue classes modulo 2^k . Their counts in an interval of length y have an additive $O(1)$ boundary error; relative control by that fact alone becomes ineffective once 2^k is comparable to or larger than y . Saying that every depth of order $\log y$ has singleton cylinders would be too strong: the multiplicative constant in the depth matters.

Preimage counts. The result of Krasikov–Lagarias [7] cited in Section 1.3 has a different counting scale from Theorem 1. An $x^{0.84}$ lower bound is compatible with a bounded reciprocal sum, and by itself cannot contradict a logarithmic error bound. This comparison explains why the particular argument of Theorem 7.2 does not transfer. It does not exclude other almost-all-to-all arguments for Collatz.

The analogue of Theorem 1 fails for Collatz, by exhibit. For the shortcut map $n \mapsto n/2$, $n \mapsto (3n + 1)/2$, the set $A = \{3 \cdot 2^k : k \geq 0\}$ is backward-closed: a multiple of three has no odd preimage, since $2 \cdot 3 \cdot 2^k - 1 \equiv 2 \pmod{3}$, so the fiber of $3 \cdot 2^k$ is $\{3 \cdot 2^{k+1}\}$; the same holds for the standard map. A is infinite, its counting function is $\lfloor \log_2(x/3) \rfloor + 1$, and its reciprocal sum is exactly $\frac{2}{3}$. What separates the two maps is not fiber size but worst-case backward log-mass: the even block of m returns log-mass at least $\frac{1}{m}(1 - \frac{2}{m})$ for every m (Lemma 3.1, item 1 of

Section 5.1), whereas the Collatz fiber of m returns $\frac{1}{m}$ in the mean over m and exactly $\frac{1}{2m}$ on every multiple of three. Theorem 1 quantifies over every backward-closed set, so the worst case governs and the mean is irrelevant.

Tao's theorem [6] has an arbitrarily slowly growing target and logarithmic-density exceptional set. Our eventual-entry equivalence has a fixed verified target and a specified dyadic rate; obtaining that rate remains open. No part of Tao's renewal argument is claimed to have been supplied for Juggler.

The strengthened conclusions of this section use Appendix E's written analytic proof. The unconditional Lean specialization remains at 100/203 for contagion and at rates greater than 103/203; the 5/8 assembly retains its actual OOE production hypothesis.

8. From parity control to the almost-all bound

8.1 Notation

For a word $w \in \{O, E\}^d$ and $t \leq d$ write $o_t(w)$ for the number of odd letters among the first t and

$$u_t(w) = o_t(w) \log_2 3 - t$$

for the *exponent walk*, so that the ideal image after t letters is $n^{2^{u_t}}$. For $y > N_0$ put

$$L(y) = \log_2 \frac{\log 2y}{\log N_0}, \quad d(y) = \lceil C L(y) \rceil,$$

with a constant $C \geq 5$ fixed below. A word w of length d is *L-bad* if $u_t(w) > -L$ for every $1 \leq t \leq d$ (its walk never descends to $-L$). $\text{word}_d(n)$ is the realized itinerary of the first d steps of n (letter s is the parity of $J^{s-1}(n)$); the *cylinder* $[w]_y$ is the set of odd $n \in (y, 2y]$ with $\text{word}_{|w|}(n) = w$. Odd starts have first letter O , so the fair share of an O -rooted cylinder of depth d relative to the reference mass $N = y/2$. The actual number of odd starts in $(y, 2y]$ is $N + O(1)$; this bounded endpoint error is absorbed in the asymptotic bounds below is $2^{-(d-1)}N$.

8.2 Envelope descent and the rarity of bad words

Lemma 8.1 (envelope descent). Let $n \in (y, 2y]$ and $d \geq 1$. If $\text{word}_d(n)$ is not $L(y)$ -bad, then $n \in R$.

Proof. Some $t \leq d$ has $u_t \leq -L(y)$, i.e. $3^{o_t}/2^t \leq \log N_0 / \log 2y \leq \log N_0 / \log n$, i.e. $n^{3^{o_t}} \leq N_0^{2^t}$ as integers. The power envelope $J^t(n)^{2^t} \leq n^{3^{o_t}}$ (Paper A, Theorem 2.2; `power_bound_word`) gives $J^t(n) \leq N_0$, so $J^t(n) \in R$ and $n \in R$ by backward closure. Lean: `iterate_le_of_envelope`, `mem_of_envelope_floor`, `reachesOne_of_itinerary_envelope`. $\square$

Lemma 8.2 (Chernoff). For $C \geq 5$ put

$$p_C = \frac{1 - 1/C}{\log_2 3}, \quad e(C) = \frac{C D(p_C \| \frac{1}{2})}{\ln 2}, \quad D(p \| \frac{1}{2}) = p \ln(2p) + (1 - p) \ln(2(1 - p)).$$

For $L > 0$ and $d \geq CL$, the number of L -bad words of length d is at most $2^d \cdot 2^{-e(C)L}$. Moreover, for every $\varepsilon > 0$ there is L_0 such that for $L \geq L_0$ and $d = \lceil CL \rceil$ the number of L -bad words of length d beginning with O is at most $2^{d-1} \cdot 2^{-(e(C)-\varepsilon)L}$.

Proof. An L -bad word has $u_d > -L$, i.e. $o_d > (d - L)/\log_2 3 \geq p_C d$ (as $L/d \leq 1/C$). For a fair coin, $\Pr[\text{Bin}(d, \frac{1}{2}) \geq pd] \leq e^{-dD(p \| 1/2)}$ for $p \geq \frac{1}{2}$, and $D(\cdot \| \frac{1}{2})$ is increasing on $[\frac{1}{2}, 1]$; $p_C \geq \frac{1}{2}$ for $C \geq 5$. Hence the count is at most $2^d e^{-dD(p_C \| 1/2)} \leq 2^d e^{-CL D(p_C \| 1/2)} = 2^d 2^{-e(C)L}$. For the O -rooted count, the remaining $d - 1$ letters contain $o_d - 1 > (d - L)/\log_2 3 - 1 = p'(d - 1)$ odd letters with $p' = p_C - O(1/L)$; the same bound with $d - 1$ trials gives $2^{d-1} e^{-(d-1)D(p' \| 1/2)}$, and $(d - 1)D(p' \| 1/2) \geq (e(C) - \varepsilon)L \ln 2$ for L large. $\square$

Lean: the first statement, in the exact form $\#\{w \text{ } L\text{-bad}, |w| = d\} \leq 2^d 2^{-e(C)L}$ for $C \geq 5$, $d \geq CL$, $d \geq 1$, is `LBad_count_le` in `formal/Problems/Juggler/FateChernoff.lean`: the Markov tilt `weight_markov` on the constant weight `(weightGen_one, (1+x)^d)`, the value $\exp(dh(p))$ at $x = p/(1-p)$ (`tilt_value`, `count_oddCount_ge_le_exp`), Gibbs' inequality $D(p \parallel \frac{1}{2}) \geq 0$ (`klHalf_nonneg`), and $p_C \in [\frac{1}{2}, 1)$ from $\log_2 3 \leq 8/5$ (`half_le_pC`, `pC_lt_one`). The O -rooted statement with ε stays a human proof.

Numerically $e(18) \approx 0.480$, $e(19) \approx 0.527$, $e(20) \approx 0.574$, $e(21) = 0.621$, $e(25) = 0.812$, $e(30) = 1.054$; $e(C)$ grows linearly in C with slope $D(1/\log_2 3 \parallel \frac{1}{2})/\ln 2 = 0.050$.

8.3 The cylinder hypothesis and the bound

Hypothesis $H(C, A)$ (log-log-depth cylinder bound). For all sufficiently large y , with $d = d(y)$, every word $w \in \{O, E\}^d$ that begins with O and is $L(y)$ -bad satisfies

$$\#[w]_y \leq 2^{-(d-1)} \cdot \frac{y}{2} + \frac{y}{(\log y)^A}.$$

Only an upper bound is asked, only at one depth per scale, and only for the $L(y)$ -bad words. Since $2^{-(d-1)}y/2 \asymp y(\log y)^{-C}$, $H(C, A)$ with $A > C$ says that no O -rooted cylinder of depth $d(y)$ whose envelope has not crossed the floor exceeds its fair share by more than a relative $O((\log y)^{C-A})$. The restriction is essential: the older all-word version is false because already absorbed starts can share a long terminating tail and overpopulate a cylinder without contributing any live mass; see the absorbed-cylinder obstruction. This obstruction does not refute $H(C, A)$ as stated here.

Theorem 8.3 (Tao-type bound from the cylinder hypothesis). Assume $H(C, A)$ with $C \geq 5$ and $A > C + e(C)$. Then for every $\varepsilon > 0$ and all sufficiently large y ,

$$\begin{aligned} \#\{n \text{ odd in } (y, 2y] : n \notin R\} &\leq \#\{n \text{ odd in } (y, 2y] : J^t(n) > N_0 \ \forall t \leq d(y)\} \\ &\leq \frac{y}{2} \left(\frac{\log(2y)}{\log N_0} \right)^{-(e(C)-\varepsilon)}. \end{aligned}$$

In words: all but $O(y(\log y)^{-e(C)+\varepsilon})$ odd starts in $(y, 2y]$ enter $[1, N_0]$ within $d(y) = O(\log \log y)$ steps.

Proof. A failure never enters the verified floor, giving the first inequality. A live start has a bad word by Lemma 8.1. Apply Lemma 8.2 with $\varepsilon/2$ in place of ε . Every odd start has an O -rooted word; by Lemma 8.2 the O -rooted bad words number at most $2^{d-1}2^{-(e(C)-\varepsilon/2)L(y)}$, and by $H(C, A)$ each carries at most $2^{-(d-1)}y/2 + y(\log y)^{-A}$ odd starts. Hence the count is at most

$$\frac{y}{2} 2^{-(e(C)-\varepsilon/2)L(y)} + 2^{d-1} \frac{y}{(\log y)^A} \leq \frac{y}{2} \left(\frac{\log 2y}{\log N_0} \right)^{-(e(C)-\varepsilon/2)} + \left(\frac{\log 2y}{\log N_0} \right)^C \frac{y}{(\log y)^A},$$

using $2^{d-1} \leq 2^{CL}$. The second term is $O(y(\log y)^{C-A}) = o(y(\log y)^{-e(C)})$ because $A > C + e(C)$. For large y , both terms fit inside the displayed bound with exponent $e(C) - \varepsilon$, using the remaining $\varepsilon/2$ slack. $\square$

Lean: the exact skeleton, in `formal/Problems/Juggler/FateChernoff.lean`. `oddFailure_s_subset_bad_cylinders` is Lemma 8.1 in covering form: with every start up to N_0 reaching 1, an odd $n \in (y, 2y]$ that does not reach 1 lies in the cylinder of a word whose every prefix fails the integer envelope comparison $N_0^{2^t} < (2y)^{3^{o_t}}$ (`EnvelopeBad`); `LBad_of_envelopeBad` identifies those words as $L(y)$ -bad with $L(y) = \log_2(\log 2y / \log N_0)$; `oddFailures_card_le` is the union bound, and `oddFailures_card_le_chernoff` composes it with Lemma 8.2: if every $L(y)$ -bad cylinder of depth $d \geq CL(y)$ holds at most M starts, the odd failures in $(y, 2y]$ number at most $2^d 2^{-e(C)L(y)} M$. `oddFailures_card_le_explicit` substitutes $d = \lceil CL(y) \rceil$ and the

bound of $H(C, A)$ at the scale y , for O -rooted words only (`cylinder_even_root_empty`): with $\Lambda = \log 2y / \log N_0$, the odd failures in $(y, 2y]$ number at most $y\Lambda^{-e(C)} + 2\Lambda^C y(\log y)^{-A}$, at every $y \geq 2$ and without ε . The displayed form follows by absorbing the factor 2 into Λ^ε and the second term into the first for $A > C + e(C)$; that absorption is not formalized. Corollary 8.4 uses only the exponent.

Corollary 8.4 (the conjecture from a cylinder bound). If $H(C, A)$ holds for some $C \geq 19$ and $A > C + e(C)$, then every positive integer reaches 1. Under the conditional exponent λ^{**} of Appendix C the same conclusion holds for $C \geq 18$.

Proof. Theorem 8.3 gives the hypothesis of Theorem 7.2 with $e = e(C) - \varepsilon \geq e(19) - \varepsilon > 1 - \lambda^{**}$; with λ^{**} the threshold is $0.4608 < e(18) \approx 0.480$. The pairing-only intermediate still needed $C \geq 20$ ($e(20) = 0.574 > 0.5520$). $\square$

Lean: `cylinder_bound_implies_conjecture` in `formal/Problems/Juggler/FateCylinderCorollary.lean`: if $H(C, A)$ holds at all large scales (`CylinderBound`, the displayed bound on every O -rooted $L(y)$ -bad cylinder of depth $d(y)$) with $C \geq 5$ and $A > C + e(C)$, and the contagion bound of Theorem 5.3 holds for the failure set with an exponent λ satisfying $1 - \lambda < e(C)$, then every positive integer reaches 1. The step from the explicit bound of Theorem 8.3 to the rate $y(\log y)^{-e}$ of Theorem 7.2, for any $e < e(C)$, is `oddFailures_eventually_le`: $\Lambda \geq \log y / \log N_0$ turns the first term into $(\log N_0)^{e(C)} y(\log y)^{-e(C)}$, $\Lambda \leq 2 \log y / \log N_0$ turns the second into $2(2 / \log N_0)^C y(\log y)^{C-A}$, and both exponent gaps are positive. The numerical threshold $C \geq 19$ is a statement about the root λ^{**} and stays with the audit.

8.4 Constants

The least integer depth constant in this sufficient criterion is $C = 19$ ($e(19) \approx 0.527 > 1 - \lambda^{**}$). With the certified floor $N_0 = 3.5 \cdot 10^8$, the table below keeps $C = 20$ as a conservative a-fortiori display:

y	$L(y)$	$d(y)$	exact fair-coin bad probability	$(\log y)^{-0.6}$	least depth for rate 0.6
10^{20}	1.25	25	0.065	0.100	19
10^{100}	3.55	72	0.017	0.038	56
10^{1000}	6.87	138	0.0038	0.0096	117

The bad probability is the fair-coin probability that a walk of length $d(y)$ never reaches $-L(y)$, by dynamic programming over the walk (unconditioned first letter; conditioning on $u_1 = \log_2 3 - 1$ roughly doubles the finite-depth values and leaves the exponent unchanged); the last column is the least depth at which it drops below $(\log y)^{-0.6}$ — about $17 L(y)$. With the Lean floor $N_0 = 260$ the depths grow by about 38 letters. For comparison, Paper B controls depth 4 (all words) and two words of depth 5, with relative error $y^{-1/96}$ on dyadic blocks; the hypothesis asks for the bad cylinders at depth $d(y) \rightarrow \infty$ with relative error $o(1)$ — weaker in rate than a power saving, unbounded in depth.

Lean: a certified instance of the side condition, in `formal/Problems/Juggler/FateCertified.lean`. The constants above are an audit script; for one concrete C the script can be removed from the chain entirely. A logarithm is certified by an integer power comparison — if $E \leq e$ and $x^n \leq E^m$ then $n \log x \leq m$ (`Certified.log_le_of_pow_le`, and three siblings) — so with Mathlib’s $2.718 < e < 2.719$, six comparisons ($2^{10} \leq 2.718^7$, $2.719^2 \leq 2^3$, $2^{84} \leq 3^{53}$, $3^{94} \leq 2^{149}$, $2.719^{19} \leq (3049/2500)^{100}$, $(50/39)^4 \leq 2.718$) give $\log 2 \leq \frac{7}{10}$ and $\frac{84}{53} \leq \log_2 3 \leq \frac{149}{94}$, hence $p_{30} \in [0.6098, 0.61]$, hence $D(p_{30} \| \frac{1}{2}) \geq 0.018312$ and $e(30) > \frac{39}{50}$ (`Certified.bernoulliExponent_thirty_gt`), with $e(30) < \frac{13}{10}$ in the other direction. So `Certified.cylinder_bound_thirty` is Corollary 8.4 at $C = 30$ with every constant a numeral: a cylinder bound $H(30, A)$ at all large scales with $A \geq 32$, above a certified floor, gives the conjecture, and nothing in its proof reads

a table. `Certified.pressure_thirty` and `Certified.one_sided_thirty` do the same for Theorem 9.2's and Theorem 9.1's criteria (the latter at the fair share $q = \frac{1}{2}$, where the one-sided exponent is the Chernoff one, `Certified.oneSidedExponent_half`, and $A \geq 52$ suffices by `Certified.logb_tilt_thirty_le`). The least C with $e(C) > \frac{27}{40}$ is 23, unchanged by the drop from $\frac{7}{10}$: $e(22) = 0.668392$ misses the lower threshold too, and $e(23) = 0.716046$ clears both. The paper's own threshold $e(C) > 1 - \lambda^{**}$ at $C = 19$ is a statement about λ^{**} ; both stay with the audit.

9. Weaker forms of the hypothesis

Theorem 8.3 asks for a fair-share upper bound on every envelope-bad cylinder at one depth. Lower bounds and control of cylinders after their envelope has crossed the floor are not needed. This section gives the sufficient conditions, including a single moment bound, and records what that form does not need.

9.1 One-sided odd-share control

Hypothesis $H_q(C, A)$. For all sufficiently large y , every $1 \leq t < d(y)$ and every $L(y)$ -bad word $w \in \{O, E\}^t$,

$$\#\{n \in [w]_y : \text{word}_{t+1}(n) = wO\} \leq q \# [w]_y + \frac{y}{(\log y)^A}.$$

(The depth-0 cylinder is all odd starts, whose next letter is O with share 1; the hypothesis starts at depth 1. As in Section 8.3, no condition is imposed after an envelope crossing.)

Theorem 9.1 (one-sided form). Let $0 < q < \log 2 / \log 3 = 0.6309$, $\mu = 1 - q \log_2 3 > 0$, $C > 1/\mu$, and $e_q(C) = 2(C\mu - 1)^2 / (C(\log_2 3)^2 \ln 2)$. Under $H_q(C, A)$ with $A > C + e_q(C)$, for every $\varepsilon > 0$ and all large y ,

$$\#\{n \text{ odd} \in (y, 2y] : n \notin R\} \leq \frac{y}{2} \left(\frac{\log 2y}{\log N_0} \right)^{-(e_q(C) - \varepsilon)}.$$

Hence $e_q(C) > 1 - \lambda^{**}$ implies the conjecture; the least C is 19 at $q = \frac{1}{2}$, 41 at 0.55, 223 at 0.60, 1586 at 0.62 (with λ^{***} : 18, 39, 206, 1451).

Proof. Let n be uniform on the $N_y = y/2 + O(1)$ odd integers of $(y, 2y]$. Bounded endpoint factors are absorbed using ε -slack. Follow its actual word until the first prefix that is not $L(y)$ -bad, and after that crossing append synthetic letters E . Let $\hat{\mathcal{F}}_t$ be the filtration of these stopped words, $\hat{X}_t$ the indicator that the synthetic next letter is O , and $\hat{u}_t = (\log_2 3 - 1) + \sum_{1 \leq s < t} (\hat{X}_s \log_2 3 - 1)$. On a bad atom the stopped word is the actual word, so H_q applies; on a crossed atom $\hat{X}_t = 0$. Thus, with $\eta_t = (\mathbb{P}(\hat{X}_t = 1 \mid \hat{\mathcal{F}}_t) - q)^+$, each bad atom $[w]$ has $\eta_t \leq y(\log y)^{-A} / \# [w]$, while every crossed atom has $\eta_t = 0$. There are at most 2^t atoms, so $\mathbb{E}[\eta_t] \ll 2^t (\log y)^{-A}$ and $\mathbb{E}[\sum_{1 \leq t < d} \eta_t] \ll (\log 2y / \log N_0)^C (\log y)^{-A}$. Since $\mathbb{E}[\hat{u}_{t+1} - \hat{u}_t \mid \hat{\mathcal{F}}_t] \leq -\mu + \eta_t \log_2 3$, the process $M_t = \hat{u}_t - \hat{u}_1 - \sum_{1 \leq s < t} \mathbb{E}[\hat{u}_{s+1} - \hat{u}_s \mid \hat{\mathcal{F}}_s]$ is a martingale with increments in an interval of length $\log_2 3$, and $\hat{u}_d \leq \hat{u}_1 + M_d - (d-1)\mu + \log_2 3 \sum_{s < d} \eta_s$. Fix $\kappa \in (0, 1)$. If $n \notin R$, the contrapositive of Lemma 8.1 gives $u_t > -L$ for every $t \leq d$. Thus no crossing occurred, $\hat{u}_d = u_d > -L$, and hence either $\log_2 3 \sum_{s < d} \eta_s > \kappa(d-1)\mu$, an event of probability $O((\log y)^{C-A})$ by Markov's inequality, which is $o((\log y)^{-e_q(C) + \varepsilon})$ under the stated condition, or $M_d > a := (1 - \kappa)(d-1)\mu - L - \hat{u}_1 \geq L((1 - \kappa)C\mu - 1) - \log_2 3$. By the Azuma–Hoeffding inequality, $\mathbb{P}(M_d > a) \leq \exp(-2a^2 / ((d-1)(\log_2 3)^2)) \leq 2^{-L(e_q^{(\kappa)}(C) - o(1))}$ with $e_q^{(\kappa)}(C) = 2((1 - \kappa)C\mu - 1)^2 / (C(\log_2 3)^2 \ln 2) \rightarrow e_q(C)$ as $\kappa \rightarrow 0$. $\square$

Lean: exact form, in `formal/Problems/Juggler/FateOneSided.lean`, by exponential moments and without the martingale, carrying out the remark after Proposition 9.3. `OneSided.badMass` is the tilted mass $\sum_{|w|=t, w \text{ } L\text{-bad}} \# [w] x^{o(w)}$ of the bad cylinders of depth t ; bad

words are prefix-closed (`OneSided.LBad_of_LBad_append`) and a cylinder splits into its two children (`OneSided.cylinder_split`), so H_q on the bad parents gives the affine recursion $\text{badMass}(t+1) \leq a_q \text{badMass}(t) + (x-1) \text{err}(2x)^t$, $a_q = 1 + (x-1)q$, for $1 \leq t < d$ (`OneSided.badMass_succ_le`, under `OneSided.OneSidedShare`, the hypothesis with its additive error err left symbolic), which unrolls to $xa_q^t N + (x-1) \text{err} t (2x)^t$ (`OneSided.badMass_le`). Lemma 8.1 and the Markov tilt put every odd failure in a bad cylinder of depth d with at least $p_C d$ odd letters (`OneSided.oddFailures_card_le_badMass`), so `OneSided.one_sided_bound` is the bound $(xa_q^{d-1} N + (x-1) \text{err}(d-1)(2x)^{d-1})/x^{p_C d}$ at every $y \geq 1$, every $d \geq \max(1, CL(y))$ and every tilt $x \geq 1$, with no ε ; at the re-centring tilt $x = p_C(1-q)/(q(1-p_C))$ the main term is $(x/a_q) N e^{-dD(p_C \| q)}$ (`OneSided.one_sided_bound_kl`, on `OneSided.tilt_pow_ratio` and `tilt_exponent_eq_kl`), the exponent $e_q^{\text{Ch}}(C)$ of Proposition 9.3. The error term carries $(2x)^{d-1}/x^{p_C d}$ where the Markov step above carries 2^d : the tilt is paid on the error too, so its absorption into $y(\log y)^{-r}$ needs $A > C(1 + (1-p_C) \log_2 x) + r$ in place of $A > C + r$. That absorption, the substitution $d = \lceil CL(y) \rceil$ and the displayed form with ε are not formalized.

So no lower bound on odd shares and no vanishing error are needed: if no $L(y)$ -bad cylinder of depth below $41 \log_2(\log 2y / \log N_0)$ sends more than 55% of its members to an odd state, every positive integer reaches 1.

The barrier word (observation; this hypothesis is doubtful). Unlike $H(C, A)$, which is asked at the single depth $d(y)$, $H_q(C, A)$ is asked at *every* depth below $d(y)$, and the shallow depths carry an obstruction of the same shape as the absorbed-cylinder one. Consider the *barrier word* $w = OE^{t-1}$: one odd letter, then even letters, with t maximal subject to badness. Its prefix walk is $u_s = \log_2 3 - s$, so w is $L(y)$ -bad exactly while $t < L(y) + \log_2 3$, and its depth is $O(L(y))$, far inside the window. On $[w]$ the map is the plain composite $G = \lfloor \cdot^{1/2} \rfloor^{t-1} \lfloor n^{3/2} \rfloor$, which is monotone, and its value sits in (N_0, N_0^2) at every scale — one more even letter would cross the floor, one fewer leaves the value below its square. So the image $G((y, 2y])$ is an interval of about $v_0 \log v_0 / \log y$ integers with v_0 bounded, and it narrows to a point: computed exactly, its width is 126, 91, 6760, 3271, 1609, 799, 3, 3, 2, 3, 1 at $y = 10^7, 10^{15}, 10^{25}, 10^{50}, 10^{100}, 10^{200}, 10^{435}, 10^{1000}, 10^{2000}, 10^{5000}, 10^{20000}$. At $y \approx 10^{20000}$, $G(y+1) = G(2y) = 4593$, an odd number: *every* member of $[OE^{13}]$ has next letter O , and an explicit 20000-digit witness is given in the dossier. Then $\#[wO] = \#[w]$, so $H_q(C, A)$ requires $(1-q)\#[w] \leq y(\log y)^{-A}$, that is $\#[w] \leq 10^{-88}y$ at $A = C = 19$, against a fair share $2^{-14} = 6.1 \cdot 10^{-5}$: a deficit of 84 orders of magnitude, growing without bound with y . What is not established is a lower bound on $\#[w]$ — a bounded-depth nested-floor parity count, not formalized and not proved here — so this is not a refutation. But $H_q(C, A)$ should not be presumed to hold, and Theorem 9.1 and its corollaries are stated conditionally for that reason. The pressure and no-momentum forms of Section 9.2 are untouched: they constrain an average over the live mass, not each cylinder.

Lean: the consequence, in `formal/Problems/Juggler/FateOneSidedCorollary.lean`, in the pattern of Corollary 8.4. `OneSided.OneSidedBound` is $H_q(C, A)$ at a scale, `OneSided.oneSidedExponent` the exponent $e_q^{\text{Ch}}(C) = CD(p_C \| q) / \ln 2$ of Proposition 9.3, and `OneSided.oddFailures_le_of_one_sided` the absorption of the exact bound above into $y(\log y)^{-e}$ for all large y , every $e < e_q^{\text{Ch}}(C)$ and $A > C(1 + \log_2 x) + 1 + e$ at the re-centring tilt x , on Gibbs' inequality $D(p_C \| q) \geq 0$ (`OneSided.klDiv_nonneg`) and $e^{-dD} \leq \Lambda^{-e_q^{\text{Ch}}(C)}$, $(2x)^{d-1} \leq \Lambda^{C(1+\log_2 x)}$ for $d = \lceil CL(y) \rceil$ (`OneSided.exp_le_rpow_scale`, `OneSided.pow_le_rpow_scale`). Then Theorem 7.2: `OneSided.implies_conjecture_of_contagion` keeps the contagion bound as a hypothesis at an exponent λ with $1 - \lambda < e$, `OneSided.one_sided_implies_conjecture` discharges it through the unconditional Theorem 5.3 at exponent $\frac{13}{40}$, so that $H_q(C, A)$ at all large scales with $e_q^{\text{Ch}}(C) > \frac{27}{40}$ gives the conjecture with nothing else assumed, and `OneSided.exact_share_implies_conjecture` is the remark above with no error term, where A disappears. The condition on A is sufficient, not the paper's $A > C + e_q(C)$: the exact form pays the tilt on the error and bounds d crudely. The numerical forms, the least C at each q , are not formalized.

9.2 The pressure form

Let $N = y/2$, let $\tau(n) = \min\{t \geq 1 : J^t(n) \leq N_0\}$, with $\min \emptyset = \infty$, be the entrance time into the floor, call n *live at depth* t if $\tau(n) > t$, and write $o_t(n)$ for the number of odd letters among $n, J(n), \dots, J^{t-1}(n)$. By Lemma 8.1, $\tau(n) > t$ implies $u_t(n) > -L(y)$. The stopping is essential: $J(1) = 1$ is odd, so an orbit that has reached 1 has an all- O tail and the unstopped odd count of every start in R grows linearly. For $\theta > 0$ put $a_\theta = \frac{1}{2}(1 + e^\theta)$.

Hypothesis $P_\theta(C)$ (live pressure). For all sufficiently large y , with $d = d(y)$,

$$\frac{1}{N} \sum_{\substack{n \text{ odd} \in (y, 2y] \\ \tau(n) > d}} e^{\theta o_d(n)} \leq a_\theta^d e^{o(d)}.$$

Theorem 9.2 (pressure form). Assume $P_\theta(C)$ with $C \geq 5$ and $\theta = \theta_C = \log(p_C/(1 - p_C))$. Then for every $\varepsilon > 0$ and all large y ,

$$\#\{n \text{ odd} \in (y, 2y] : \tau(n) > d(y)\} \leq \frac{y}{2} \left(\frac{\log 2y}{\log N_0} \right)^{-(e(C) - \varepsilon)},$$

the bound of Theorem 8.3. Hence $P_{\theta_C}(C)$ with $C \geq 19$ implies the conjecture using Theorem 1, and with $C \geq 18$ under the conditional exponent of Appendix C.

Proof. If $\tau(n) > d$ then $u_d > -L$, i.e. $o_d \geq p_C d$. Hence

$$\#\{\tau > d\} \leq \#\{\tau > d, o_d \geq p_C d\} \leq e^{-\theta p_C d} \sum_{\tau(n) > d} e^{\theta o_d(n)} \leq N e^{-\theta p_C d} a_\theta^d e^{o(d)}.$$

At $\theta = \theta_C$, $\theta p_C - \log a_\theta = D(p_C \| \frac{1}{2})$, so the count is $\leq N \exp(-dD(p_C \| \frac{1}{2})(1 - o(1))) \leq N 2^{-(e(C) - \varepsilon)L}$. $\square$

Lean: exact form, in `formal/Problems/Juggler/FatePressure.lean`, on the live weight of `LiveCountWeight` (starts in $\{1, \dots, N\}$ that stay above N_0 for d steps). `livePressure` is $\sum_n \text{live } x^{o_d(n)}$ as the generating function of the live weight; `live_count_le_pressure` is the Markov step $\#\{\text{live}, o_d \geq k\} \leq \text{pressure}/x^k$; `live_oddCount_ge` is Lemma 8.1 on live starts ($o_d \geq p_C d$ for $d \geq CL(N)$); and `live_count_le_of_pressure` says: if the pressure at the tilt $x = p_C/(1 - p_C)$ is at most $N a_\theta^d E$, the live starts number at most $N \exp(-dD(p_C \| \frac{1}{2}))E$, by the identity $a_\theta^d / x^{p_C d} = e^{-dD(p_C \| 1/2)}$ (`tilt_value`). The substitution $E = e^{o(d)}$ and $d = \lceil CL \rceil$ that gives the displayed bound is not formalized.

For $1 \leq t < d$ let $\mu_{\theta,t}$ be the probability measure on the starts live at depth t with density proportional to $e^{\theta o_t(n)}$, and let $s_\theta(t) = \mu_{\theta,t}(J^t(n) \text{ odd})$ be the *tilted odd share*: the probability that the next letter is O when odd-heavy live prefixes are up-weighted exponentially. If the live set is empty, define $s_\theta(t) = 0$; subsequent live moments are then zero.

Hypothesis $M_{\theta,q}(C)$ (no momentum). For all large y , $\sum_{t=1}^{d(y)-1} (s_\theta(t) - q)^+ = o(d(y))$.

Proposition 9.3. $M_{\theta,1/2}(C)$ implies $P_\theta(C)$. More generally, for $0 < q < 1$ and a fixed $\theta > 0$, $M_{\theta,q}(C)$ gives

$$\#\{\tau > d\} \leq N \exp\{-d[\theta p_C - \log(1 - q + qe^\theta) - o(1)]\}.$$

If $0 < q < p_C < 1$ and the hypothesis is assumed at the optimizing tilt $\theta = \theta_{C,q} := \log \frac{p_C(1-q)}{q(1-p_C)}$, this becomes $N \exp(-dD(p_C \| q)(1 - o(1)))$, with exponent $e_q^{\text{Ch}}(C) = C D(p_C \| q) / \ln 2$, at least the Azuma exponent of Theorem 9.1. Its least C at $q = 0.5, 0.55, 0.60, 0.62$ is 19, 41, 214, 1496 (with λ^{***} : 18, 38, 198, 1369).

Proof. Dropping the condition $\tau > t + 1$ in favour of $\tau > t$ only enlarges the sum, so

$$\sum_{\tau > t+1} e^{\theta o_{t+1}} \leq \sum_{\tau > t} e^{\theta o_{t+1}} = \left(\sum_{\tau > t} e^{\theta o_t} \right) (1 + (e^\theta - 1)s_\theta(t)).$$

With $a_{\theta,q} = 1 + (e^\theta - 1)q$ and $c_\theta = (e^\theta - 1)/a_{\theta,q}$, $1 + (e^\theta - 1)s \leq a_{\theta,q} \exp(c_\theta(s - q)^+)$. Telescoping from $t = 1$ gives $\sum_{\tau > d} e^{\theta o_d} \leq N e^\theta a_{\theta,q}^{d-1} \exp(c_\theta \sum_{t < d} (s_\theta(t) - q)^+)$, which is P_θ at $q = \frac{1}{2}$. For general q , the exponential Markov step gives $\theta p_C - \log a_{\theta,q}$ at the same fixed θ . Assuming $M_{\theta,q}$ at $\theta = \log \frac{p_C(1-q)}{q(1-p_C)}$ gives $D(p_C \| q)$. $\square$

Lean: `weightGen_succ_le_share` (the one-depth step), `one_add_le_exp_excess`, `weightGen_le_pressure` (the telescoped bound) and `count_le_pressure` (the exponential Markov step), in `formal/Problems/Juggler/TiltedShare.lean`, on the word-weight framework of `RateFreeDensity`; `tilt_exponent_eq_kl` is the identity at the re-centring tilt.

Lean: the consequences of Theorem 9.2 and Proposition 9.3, in `formal/Problems/Juggler/FatePressureCorollary.lean`, in the pattern of Corollary 8.4. A failure never enters the floor, so the odd failures of $(y, 2y]$ are live starts of $\{1, \dots, 2y\}$ at every depth (`Pressure.oddFailures_subset_live`); `Pressure.PressureBound` is $P_\theta(C)$ at a scale with $e^{o(d)}$ quantified as $(\log y)^\varepsilon$, and `Pressure.NoMomentumBound` is $M_{\theta,q}(C)$ on the live weight at the re-centring tilt with the $o(d)$ as δd . The exact forms above bound the failures by $2y e^{-dD} (\log y)^\varepsilon$, with $D = D(p_C \| \frac{1}{2})$ for the pressure form and $D = D(p_C \| q) - c_x \delta$, $c_x = (x - 1)/a_{x,q}$, for the no-momentum form (`Pressure.momentumExponent`); the shared absorption `Pressure.absorb` turns that into $y(\log y)^{-e}$ for every $e < CD/\ln 2 - \varepsilon$ and all large y (`Pressure.oddFailures_le_of_pressure`, `Pressure.oddFailures_le_of_noMomentum`). Then Theorem 7.2, with the contagion bound as a hypothesis (`Pressure.pressure_conj_of_contagion`, `Pressure.noMomentum_conj_of_contagion`) or discharged through the unconditional Theorem 5.3 at exponent $\frac{13}{40}$ (`Pressure.pressure_implies_conjecture`, `Pressure.noMomentum_implies_conjecture`): either hypothesis at all large scales, with its exponent above $\frac{27}{40}$, gives the conjecture with nothing else assumed. The numerical forms are not formalized.

9.3 Scope of the moment criterion

(a) Any $o(\log \log y)$ initial depths are free. A letter contributes a factor at most e^θ to the moment, so bias — however extreme — at any set of depths of size $o(d(y))$ is absorbed by the $e^{o(d)}$. In particular the depth-five split (Paper B's Conjecture 7.3) and every split to any fixed depth, or to depth $s_0(y)$ for any $s_0 = o(\log \log y)$, is irrelevant to the reduction.

(b) A tower comparison in a model population. Suppose a model cylinder O^t has size $N\beta^{t-1}$. Its moment relative to the fair reference total $N e^\theta a_\theta^{t-1}$ is $(\beta e^\theta / a_\theta)^{t-1}$. This ratio decays precisely when $\beta < (1 + e^{-\theta})/2$, about 0.836 at $\theta_{19} \approx 0.396$. Its count alone exceeds the reference survival exponent only when $\beta > 2^{-e(C)/C}$, about 0.981 at $C = 19$.

These are algebraic comparisons to a reference mass. They do not bound $s_\theta(t)$ in an actual stopped population, whose total live mass may be much smaller. A model tower below 0.836 is therefore not, by itself, a proof of the no-momentum hypothesis.

(c) One-sidedness and aggregation. $M_{\theta,q}$ bounds the accumulated positive excess of a weighted average over depths. Under-populated odd continuations can offset excess within the same depth; over-populated cylinders contribute in proportion to their tilted weight. In Walsh terms, $e^{\theta X_s} = a_\theta + b_\theta(-1)^{J^s(n)}$ with $b_\theta/a_\theta = -\tanh(\theta/2)$, so the unstopped moment is

$$\frac{1}{N} \sum_n e^{\theta o_d(n)} = a_\theta^d \left(1 + \sum_{\emptyset \neq T} (-\tanh \frac{\theta}{2})^{|T|} \frac{W_T}{N} \right), \quad W_T = \sum_n \prod_{s \in T} (-1)^{J^s(n)} :$$

Walsh sums of high order are exponentially down-weighted ($\tanh(0.198) = 0.196$ per letter), whereas $H(C, A)$ asks for a separate supremum bound on each $L(y)$ -bad cylinder at the target depth.

(d) Other sufficient statistics. The proof of Theorem 9.1 also works if the odd-share bound fails only on bad atoms of total probability $O((\log y)^{-B})$ at each depth, with $B > e_q(C)$: the expected accumulated excess is $O(d(\log y)^{-B})$, and Markov's inequality absorbs it.

Lean: this variant, on the exponential-moment proof, in `formal/Problems/Juggler/FateOneSidedAtoms.lean`. `OneSided.OneSidedShareExc` allows at each depth an exceptional set of words of total cylinder mass at most exc on which the share bound may fail; an exceptional bad atom sends at most its whole mass to its odd child, which costs $(x - a_q)\#[w]x^{o(w)}$ in the tilted mass, so the recursion of Theorem 9.1's Lean proof reads $\text{badMass}(t+1) \leq a_q \text{badMass}(t) + (x-1)(\text{err}(2x)^t + (1-q)\text{exc}x^t)$ (`OneSided.badMass_succ_le_exc`), with the exact bound `OneSided.one_sided_bound_exc` and its re-centred form `OneSided.one_sided_bound_k1_exc`. With $\text{exc} = y(\log y)^{-B}$ the absorption (`OneSided.oddFailures_le_of_exc`) needs $B > C \log_2 x + 1 + e$, against $A > C(1 + \log_2 x) + 1 + e$ for the error: the exceptional atoms are not doubled at each depth. `OneSided.exc_implies_conjecture` runs this weakest one-sided form to the conjecture with nothing else assumed when $e_q^{\text{Ch}}(C) > \frac{27}{40}$, and `OneSided.exc_conj_of_contagion` with the contagion bound as a hypothesis. The condition on B is sufficient and not the $B > e_q(C)$ of the Markov absorption above, which is not restated.

For comparison, let $D(w) = \#[wO] - \frac{1}{2}\#[w]$, $\mathcal{C}_t = \sum_{|w|=t} \#[w]^2$, and let W_T be the Walsh sums on the same full set of odd starts. Parseval gives the exact identities

$$\sum_w D(w)^2 = 2^{-t-2} \sum_{S \subseteq \{0, \dots, t-1\}} |W_{S \cup \{t\}}|^2 = \frac{1}{2}\mathcal{C}_{t+1} - \frac{1}{4}\mathcal{C}_t.$$

Lean: the second equality, $\sum_w D(w)^2 = \frac{1}{2}\mathcal{C}_{t+1} - \frac{1}{4}\mathcal{C}_t$, is `CylinderEnergy.sum_bias_sq` in `formal/Problems/Juggler/FateCylinderEnergy.lean`, for an arbitrary finite set of starts; it rests on the cylinder splitting $\#[w] = \#[wE] + \#[wO]$ (`CylinderEnergy.wordCount_split`) and is otherwise the algebra of $(b - \frac{a+b}{2})^2 = \frac{a^2+b^2}{2} - \frac{(a+b)^2}{4}$. The Parseval equality with the Walsh sums is not formalized; no Walsh transform appears in that file.

An additional quantitative bound on these sums can supply an exceptional-atom estimate. No rate-independent equivalence with the restricted bad-cylinder hypothesis is asserted. Full-population fair collision bounds also face the absorbed-cylinder obstruction of Section 8.3.

Lean: the supply, in `formal/Problems/Juggler/FateEnergyAtoms.lean`, for the energy of the $L(y)$ -bad words. `Energy.badEnergy` is $\sum_w \text{bad} D(w)^2$ over the odd starts of $(y, 2y]$, and `Energy.biasEnergy` the sum over all words, equal to $\frac{1}{2}\mathcal{C}_{t+1} - \frac{1}{4}\mathcal{C}_t$ (`Energy.biasEnergy_eq`, on `CylinderEnergy.sum_bias_sq`). A bad atom violating $\#[wO] \leq q\#[w]$ for $q > \frac{1}{2}$ has $D(w) > (q - \frac{1}{2})\#[w]$, so the squared masses of the bad violators sum to at most the bad energy over $(q - \frac{1}{2})^2$, and Cauchy-Schwarz over at most 2^t atoms gives $(\sum_{\text{bad violators}} \#[w])^2 \leq 2^t \text{badEnergy} / (q - \frac{1}{2})^2$ (`Energy.mass_violators_le`). Hence $\text{badEnergy}_t \leq (q - \frac{1}{2})^2 \text{exc}^2 / 2^t$ at every depth $1 \leq t < d$ is the one-sided hypothesis with exceptional atoms of mass exc and no error term (`Energy.oneSidedShareExc_of_energy`), and `Energy.energy_implies_conjecture` runs it to the conjecture: the bad energy bound with $\text{exc} = y(\log y)^{-B}$ at the depths below $\lceil CL(y) \rceil$ at all large scales (`Energy.EnergyBound`), $\frac{1}{2} < q < p_C$, $B > C \log_2 x + 1 + e$ and $\frac{27}{40} < e < e_q^{\text{Ch}}(C)$, and nothing else. The restriction to bad words is forced: the unrestricted energy (`Energy.badEnergy_le_biasEnergy`) is dominated by absorbed orbits, whose cylinders are fully biased. The exact enumeration of Section 11 then finds the bad energy itself at its worst-case value $\frac{1}{4} \sum_w \text{bad} \#[w]^2$ from depth 16–20 at every computable scale, because bad words that have dipped to a bounded value collapse onto single deterministic futures; the reduction stands, the hypothesis has no numerical support, and the per-cylinder forms it feeds inherit the warning.

Lean: the collapsed component, in `formal/Problems/Juggler/FateCollapse.lean`, the exact layer of the observation that the collapsed mass, taken as a whole, is nearly fair. For a finite set of starts, `Collapse.fiber` counts the starts whose t -th iterate is a given value, and the next-letter bias of the starts whose iterate lands in a window $[a, b]$ of values (`Collapse.windowBias`) is the alternating sum $-\sum_{v=a}^b (-1)^v \text{fiber}(v)$ (`Collapse.windowBias_eq_sum`), hence at most the total variation of the fiber profile over the window plus one fiber (`Collapse.abs_windowBias_le`, by summation by parts). One step of the map sums fibers over preimages, the

even numbers of $[v^2, (v+1)^2)$ plus at most one odd preimage (`Collapse.fiber_succ_eq`, on Lemma 3.1's even block in fiber form), and the even branch smooths: with the depth- t profile between m_v and M_v on the double block $[v^2, (v+2)^2)$, consecutive block sums differ by at most $(v+1)(M_v - m_v) + 2M_v$ (`Collapse.blockSum_sub_le`). Hence `Collapse.collapse_bias_le`: the bias of a collapsed window at depth $t+1$ is at most $\sum_v ((v+1)(M_v - m_v) + 2M_v)$ plus twice the mass arriving through odd preimages plus one fiber. With a smooth profile, $M_v - m_v$ of order $\bar{f}/v$ on a block of size $\bar{f}v$, the first term is $O(\bar{f})$ per value against a window mass of $\bar{f}v$ per value: the collapsed mass entering through the even branch is fair up to a relative $O(1/v)$, and the excess share of the moment criterion sits in the odd branch and in the high walkers. Runs of even letters are self-smoothing: a sandwich $m \leq \text{fiber}(t, \cdot) \leq M$ on $[v^2, (v+1)^2)$ with odd branch at most P gives $vm \leq \text{fiber}(t+1, v) \leq (v+1)M + P$ (`Collapse.fiber_succ_sandwich`), so after two even steps the bias of a window at depth $t+2$ is bounded by the depth- t sandwich with the relative oscillation not amplified and a loss of a relative $O(1/w)$ at the final scale (`Collapse.collapse_bias_two_step`); iterated, the relative oscillation at the end of an even run is the one at its start plus $O(1/w)$ at the last scale. What the even run cannot repair is the oscillation it starts with, the profile after the preceding odd run: its first even step is the share law of Section 4, and its block average is Proposition 4.4. Not here: any bound on the variation of a fiber profile beyond that propagation, and the parity of the odd-preimage mass, which is the share law at bounded scale.

(e) Bounded-depth statistics and the walk model. Fix k . The probability measure on parity words that is fair to depth k and all- O afterwards satisfies every cylinder-count statement of depth $\leq k$ exactly, and under it a positive proportion of model walks never crosses the barrier. An argument for the Tao-type bound that used only depth- $\leq k$ cylinder statistics of the Juggler map and the walk logic would prove the same bound for that measure. Thus finite-depth statistics plus the envelope walk alone are insufficient. This model example does not rule out a proof using additional arithmetic properties of the Juggler map.

9.4 Where the difficulty lies

$P_\theta(C)$ says that the exponentially tilted distribution of live parity words of length $d \asymp \log_2 \log y$ has the fair pressure $\log a_\theta$: odd-heavy live words — those with about $e^\theta/(1+e^\theta) \approx 60\%$ odd letters, the words the tilt selects — are not over-populated at an exponential rate in d . Typical such words contain odd runs of length ≥ 4 , so the nested towers of height ≥ 4 that stop Paper B at depth five sit inside almost every relevant cylinder; but no individual cylinder is required to split fairly, only the tilted average over the $\approx 2^{0.97d}$ typical cylinders at each depth. This is a statement about averaging over cylinders at unbounded depth — a mean over characters, not a supremum — of a kind for which analytic number theory sometimes has tools where pointwise bounds fail. We record the question and do not pursue an analytic approach here.

9.5 Scale-averaged pressure is sufficient

A cumulative pressure estimate suffices even when individual scales have large spikes. Fix $C > 1$, $\theta > 0$, $0 < q < 1$, and $\eta \geq 0$. Put $a = 1 - q + qe^\theta$, $r = C(\theta p_C - \log a)/\log 2$, and at $y = 2^k$ define

$$Z_k = \sum_{\substack{n \text{ odd} \in (2^k, 2^{k+1}] \\ \tau(n) > d(2^k)}} e^{\theta o_{d(2^k)}(n)}, \quad \rho_k = \frac{Z_k}{2^{k-1} a^{d(2^k)}}.$$

Choose a fixed k_0 above the verified target.

Theorem 9.4 (scale-average criterion). Suppose, for every $\varepsilon > 0$,

$$\sum_{k=k_0}^K \rho_k \ll_\varepsilon K^{1+\eta+\varepsilon}.$$

If $r - \eta > 3/8$, the written Theorem 5.19 implies universal termination. With the unconditional Lean baseline in its place, $r - \eta > 103/203$ suffices. The cumulative arithmetic bound itself remains open.

Proof. A nonterminating start remains live and has $o_d > p_C d$. Exponential Markov therefore bounds its odd reciprocal mass in the k -th block by $O(\rho_k(k+1)^{-r})$. Summing the cumulative hypothesis on dyadic groups of indices gives

$$\sum_{k \leq K} \rho_k(k+1)^{-r} = O(K^\beta) \quad \text{for every } \beta > \max(1 + \eta - r, 0).$$

The logarithmic boundary case is absorbed by the strict inequality. The even-tree argument of Theorem 7.2 bounds total failure mass by $O((\log X)^\beta \log \log X)$. Choose $\max(1 + \eta - r, 0) < \beta < 5/8$ and use Theorem 5.19 to contradict a nonempty failure class. For the baseline replace $5/8$ by $100/203$. $\square$

The baseline implication, with the actual stopped Juggler weights, is `ScaleAverage.pressure_average_conjecture` in `FateScaleAverage.lean`; its fourteen-theorem audit is separate from the historical Paper C audit. The formal corollary `ScaleAverage.pressure_average_conjecture_of_ooee` also proves the $3/8$ implication with `OOEEProductionBound` explicit. Its reusable proof accepts any supplied contagion exponent; the written OOEE proof supplies that premise here. No estimate of the actual cumulative pressure is asserted.

10. The free term and limits of the reductions

10.1 The free term is the infinite-depth live mass

Recall the free term ψ_F of (6.1). For odd n , $n \in F \iff \lfloor n^{3/2} \rfloor \in F$ (Theorem 6.1), and $n \in F \iff \tau(n) = \infty$ since $[1, N_0] \subseteq R$. Let $\mathbb{P}_x^{\log}$ be the $1/n$ -weighted measure on the odd $n \in (\sqrt{x}, x]$.

Proposition 10.1 (normalized live limit). Put

$$q_{OO}(t) = \frac{8}{t} \sum_{\substack{\sqrt{x} < n \leq x \\ \text{word}_2(n) = OO}} \frac{1}{n}, \quad x = e^t.$$

Whenever the conditioning set is nonempty,

$$\psi_F(t) = q_{OO}(t) \lim_{d \rightarrow \infty} \mathbb{P}_x^{\log}(\tau(n) > d \mid \text{word}_2(n) = OO). \quad (10.1)$$

The limit is decreasing. For large t , $q_{OO}(t) = 1 + O(e^{-\delta t})$ for some $\delta > 0$. If a hypothesis in Sections 8–9 gives a live-start bound with rate $r > 0$, then, for every $0 < \varepsilon < r$,

$$\psi_F(t) \ll_\varepsilon \left(\frac{t}{2 \log N_0} \right)^{-(r-\varepsilon)}. \quad (10.2)$$

For the one-sided and no-momentum forms, the same proofs bound live starts as well as failures. The rate is the case-dependent rate in Theorem 4.

Proof. At fixed x the conditioning set is finite, and $F = \{n : \tau(n) = \infty\}$. Continuity for decreasing events gives (10.1), including its normalization factor. The depth-two parity estimate for odd starts gives OO -count $y/4 + O(y^{1-\delta'})$ on dyadic blocks. Partial summation, or the same exponential-sum bound on initial subintervals, gives the asserted reciprocal-mass asymptotic for q_{OO} .

Cover $(\sqrt{x}, x]$ by downward dyadic blocks, allowing one partial block at the lower end. On $(y, 2y]$ the reciprocal mass of live starts is at most their count divided by y , hence

$O((\log y / \log N_0)^{-(r-\varepsilon)})$. There are $O(t)$ blocks and $\log y \geq t/2 - O(1)$. Multiplication by $8/t$, the definition of ψ_F , proves (10.2). No equality between an unweighted block ratio and its reciprocal-weighted ratio is used. $\square$

This connects the free term with live counts. The decreasing limit does not provide a rate of convergence in depth; the finite-depth estimates remain additional sufficient hypotheses.

10.2 What the upper recursion proves

Equations (6.1) and the fiber upper bound give, for large t ,

$$\varphi_F(t) \leq \frac{1}{2}\varphi_F(t/2) + \frac{1}{3}\varphi_F(3t/4) + \frac{1}{4}\psi_F(t) + O(e^{-\delta t}). \quad (10.3)$$

The coefficient is $1/3$ with the proved upper share; the ideal fiber model would replace it by $1/4$. If c denotes either coefficient, the homogeneous exponent $e_c > 0$ is the root of $\frac{1}{2}2^{e_c} + c(4/3)^{e_c} = 1$. Numerically it is 0.3391 for $c = 1/3$, and $1 - \lambda_{\text{ideal}} = 0.5073$ for $c = 1/4$. These values describe the upper comparison, not a proved exact decay rate of a nonempty failure set.

Corollary 10.2 (upper comparison). If $\psi_F(t) = O(t^{-r})$ for some $r > 0$, then $\varphi_F(t) = O(t^{-a})$ for every $0 < a < \min(r, e_c)$, using $c = 1/3$ in (10.3). The ideal model has the analogous assertion with $c = 1/4$. The upper inequality alone does not imply $F = \emptyset$.

Proof. For such a , $h = \frac{1}{2}2^a + c(4/3)^a < 1$. Choose a large initial scale T and then K so that $\varphi_F(t) \leq Kt^{-a}$ on $[T/2, T]$, and so that the forcing in (10.3) is bounded by $(1-h)Kt^{-a}$ for $t \geq T$. Induction on intervals $[T, (4/3)^j T]$ proves the bound. At the endpoint $a = e_c$ there is no slack; in particular an endpoint estimate cannot be inferred by the same argument. A positive multiple of t^{-e_c} satisfies the homogeneous comparison on a tail, so that comparison alone does not force the function to vanish. The lower contagion estimates and this upper comparison do not establish a matching asymptotic. $\square$

10.3 The depth-uniformity budget

Section 9.3(e) limits what fixed-depth statistics plus the envelope walk alone can prove. The form of Theorem 8.3 says how uniform that information must be *along the cylinder-count route*.

Proposition 10.3 (depth-uniformity budget). Suppose the available cylinder discrepancy bound is

$$\left| \#[w]_y - 2^{-(d-1)}y/2 \right| \leq K2^{\kappa d}y^{1-\delta_0}2^{-cd},$$

for the relevant bad words, with constants $K, \delta_0, c > 0$, $\kappa \geq 0$, and a scale threshold independent of d . At $d = C \log_2 \log y + O(1)$, the resulting error majorant after summing over at most 2^d words is $o(y(\log y)^{-r})$ for every fixed $r > 0$ if and only if $cC < 1$.

Proof. Divide the error majorant by $y(\log y)^{-r}$ and take logarithms. The result is

$$O(1) + ((\kappa + 1)C + r) \log \log y - \Theta((\log y)^{1-cC}).$$

It tends to $-\infty$ exactly when $cC < 1$. If $cC \geq 1$, this majorant is insufficient; the actual error may still be much smaller. Without uniform control of the prefactor and the scale threshold, a bound at every fixed depth would not justify any conclusion at growing depth. $\square$

At $C = 19$ this requires the saving exponent to lose less than a factor $2^{1/19} = 1.037$ per depth; at $C = 41$, $2^{1/41} = 1.017$. Weyl differencing loses a factor 2^c with $c \geq 1$ per differencing (Paper B's chain $\frac{1}{24} \rightarrow \frac{1}{96}$ from depth three to four is $c = 2$), and with such a loss the cylinder-count route reaches depth $\approx \log_2 \log y$ — one unit of $L(y)$, enough for the all- E word to reach the floor, not for the Chernoff tail, which needs $C \geq 5$ units. The proposition is narrow by design: it concerns methods whose only available error estimate deteriorates exponentially in the depth and are used through the per-cylinder count of Theorem 8.3. A method that organizes its errors differently, that proves a polylogarithmic relative saving uniformly in the depth (as $H(C, A)$ asks), or that addresses the pressure statement directly is not covered by it.

11. Numerical observations

The following archived numerical experiments accompany the theorems. Orbit steps and integer counts use exact arithmetic; phases, weighted statistics and fair-coin comparisons use floating-point arithmetic. Random samples use fixed seeds (`research.juggler_sequence.fate_contagion`, `research.juggler_sequence.tao_reduction`; artifacts in Appendix B), they do not establish asymptotic estimates and they enter no analytic proof. Detailed tables are kept in the repository notes [14].

Fibers, blocks, closure. Over $3375 \leq m < 10^5$ and spot ranges at 10^6 and 10^7 , the mean of G_m/H_m is 0.5000, 0.5004, 0.5027; the minimum on good fibers is 0.328 ($m = 1003635$, $\alpha_m \approx \frac{1}{3}$); no good fiber falls below $\frac{1}{3}H_m - 2$. On the blocks $m' \in [20, 60) \cup [200, 230) \cup [1000, 1010) \cup \{3000, 5000\}$ the deviation of $|U(m')|$ from its main term is at most $0.72\sqrt{|I(m')_{\text{odd}}|}$, i.e. square-root scale.

The share law and the extremes. A complete census of $\Phi(m)$ for $m \leq 2 \cdot 10^5$ (`research.juggler_sequence.oefiber_share`) finds 2481 fibers with $G_m = 0$ and 2508 with $G_m = H_m$, and $H_m \geq 1$ throughout. The mean of G_m/H_m is 0.498 on the decade $[10^2, 10^3)$ and 0.500 on every decade above it, and the empty rate falls 8.0, 3.0, 1.35, 0.90 per cent over the decades from 10^2 to $2 \cdot 10^5$, tracking $1/H_m$. The archived sampling code uses $\hat{\beta}_m = \alpha_m H_m$, whereas Lemma 4.5 uses $\beta_m = \alpha_m(H_m - 1)$. Near 10^6 , 432 fibers with $|\hat{\beta}_m| \leq 1.2$ give mean absolute model error 0.0195 against $1/H_m = 0.015$, and every extreme fiber found has $\hat{\beta}_m \in [-0.79, 0.15]$, inside the model window of Corollary 4.6(2). The extremes sit $\asymp m^{1/3}$ apart as Remark 4.8 predicts: in $m \in [10^5, 1.2 \cdot 10^5)$, where that period is 46.4, their median spacing is 48 and 244 of 306 gaps are one period. Empty fibers occur singly or in adjacent pairs and never in threes. The observed empty rate exceeds the $(25/108)/H_m$ of Corollary 4.6(3) by about 1.4, and the sample is consistent with a discretization effect: of the 297 empty fibers in $[10^5, 1.3 \cdot 10^5)$, 174 have $S(\hat{\beta}_m, \theta_m) = 0$ while 123 have $S > 0$ and miss only because the fiber samples it at 31 points. Consistently, the ratio falls 1.49, 1.44, 1.36 across the last three decades. The closure of the Lean-verified seed $[1, 260]$ under the two productions of Section 3, computed exactly up to 10^9 , has density between 0.45 and 0.49 on the complete dyadic blocks $(2^k, 2^{k+1}]$, $16 \leq k \leq 28$. The last reported block is truncated to $(2^{29}, 10^9]$; its density is 0.4529, against 0.25 for the E -only closure, and its realized recursion coefficients at 10^9 are 0.9998 for E (theory 1) and 0.3332 for OE (reference $\frac{1}{3}$). The separate decomposition sample at $x = 10^6$ has OO -type share 0 against reference share 0.2987. This finite closure is not a two-way closed fate class; its omitted production illustrates the OO term without establishing an asymptotic property of F .

Survival above the certified floor. Because every $n \leq N_0 = 3.5 \cdot 10^8$ reaches 1, the statistic “ $J^t(n) \leq N_0$ for some $t \leq d$ ” is a finite computation, and its complement is the live set of Theorem 8.3 (contained in the envelope-bad cylinders). For random odd starts in $(y, 2y]$, the fraction still above N_0 after d steps is compared with the odd-start fair-coin survival $\Pr[u_t > -L(y) \forall t \leq d \mid u_1 = \log_2 3 - 1]$:

y	$L(y)$	$d = 11$	$d = 21$	$d = 31$	$d = 36$
10^{12}	0.526	0.221/0.221	0.082/0.084	0.037/0.038	0.027/0.028
10^{15}	0.841	0.251/0.267	0.100/0.106	0.044/0.047	0.032/0.035
10^{20}	1.249	0.295/0.309	0.135/0.138	0.068/0.068	0.049/0.049
10^{30}	1.826	0.442/0.439	0.175/0.175	0.098/0.095	0.075/0.070
10^{50}	2.558	0.554/0.554	0.267/0.264	0.148/0.143	0.101/0.097

Entries are empirical/fair at the depths retained in the archived record: 40000 starts at 10^{12} , 10^{15} , 10^{20} , and 20000 at 10^{30} , 10^{50} . The sampler draws integers in $(y, 2y]$ and rounds even draws up to the next odd integer; this differs from uniform odd sampling only at the two

endpoints, with total variation $1/y$. A resource cap is conservatively recorded as no observed descent, so empirical survival is an upper bound for the sampled live fraction when a cap is reached.

The separate pressure runs use 40000 starts at each of $10^{20}, 10^{30}, 10^{50}$, with no capped orbits. At these scales, the recorded extrema of $s_\theta(t)$ over $1 \leq t \leq 40$ and $\theta \in \{0.396, 0.6\}$ are approximately 0.4728 and 0.5337. At the four recorded depths 10, 20, 30, 40, the live-moment ratio to the fair-coin comparison ranges from 0.9521 to 1.0812, using outward rounding. The archived 10^{12} pressure run has one capped orbit; its unknown later letters are omitted by that estimator. It is excluded from this pressure comparison and cannot certify an upper bound on the full live moment. These finite observations do not establish the hypotheses of Section 9 at depths tending to infinity.

Exact cylinder energies (observation). An exact enumeration of every odd start of $(y, 2y]$ to depth 30 at $y = 10^4, 10^5, 10^6, 10^7$ (`research.juggler_sequence.cylinder_energy_measure`, no sampling) records the bias energy $\sum_w D(w)^2$ over all words, over the $L(y)$ -bad words and over the live starts, against the fair-coin value $N/4$ and the worst case $\frac{1}{4} \sum_w \# [w]^2$. Over all words the worst case is reached by depth 16–22 at every scale, the absorbed-cylinder obstruction of Section 8.3 in numbers. Over the $L(y)$ -bad words it is reached by depth 16–20 as well, on cylinders of up to 11939 members at 10^7 : a bad word whose walk has dipped near $-L(y)$ has collapsed onto a bounded value above the floor, and its cylinder’s later letters are the deterministic orbit of that value. The mass of bad atoms violating $\# [wO] \leq 0.55 \# [w]$ falls with the scale in the dense window (0.47, 0.26, 0.12, 0.05 of the bad mass at depth 10) and is a constant 0.42–0.49 at depth 20. These finite observations do not test the hypotheses of Sections 8–10 at their own scales; they show that their per-cylinder forms live, at every computable scale, in a regime where cylinders are parity-constant.

The excess of the bad cylinders (observation). The same enumeration measures the quantity $H_q(C, A)$ bounds, namely $\max_w \text{bad}(\# [wO] - q \# [w])$, at every depth of the hypothesis’s own window $1 \leq t < d(y)$ (`research.juggler_sequence.bad_cylinder_excess`). At these scales the additive error is inert — $y(\log y)^{-A}$ is below 10^{-38} at $y = 10^7$ for $A \geq 32$ — so the honest comparison is with a fair coin on the same cylinder sizes, whose largest excess is $\sqrt{2 \log W \cdot n_{\max}}/2$. The largest A the measured excess permits over the window at $C = 19$ is 2.62, 2.84, 2.61, 1.59 at $y = 10^4, \dots, 10^7$, against a fair-coin value of 2.62, 2.86, 3.10, 3.34: the map is indistinguishable from a coin at the two smaller scales and measurably worse at the two larger. At a fixed shallow depth the shares move the other way, the three largest bad cylinders at depth 4 having odd shares 0.53, 0.53, 0.52 at 10^4 and 0.5001, 0.4994, 0.5008 at 10^7 , because the number of distinct iterates a cylinder sits over widens with y . What does not move with y is the top of the window: the distinct iterates per bad cylinder at $t = d(y) - 1$ are 1.14, 1.02, 1.01, 1.00, so there the cylinders are parity-constant at every scale measured, and the hypothesis reduces to whether the mass of the largest such cylinder — a fraction of order 10^{-3} here, with no visible decay — falls below $(\log y)^{-A}$. Three decades of y cannot see a $(\log y)^{-19}$; these observations neither establish nor refute the hypotheses of Sections 8–10. The backward computation of the barrier word, recorded with Theorem 9.1 above and in its dossier (`research.juggler_sequence.barrier_collapse`), reaches $y = 10^{20000}$ and is what turns that measurement into a statement about $H_q(C, A)$ at the scales where it is asserted.

12. Conclusions and open estimates

The strongest written contagion exponent is now $5/8$, from the three actual productions E, OE and OOE. Appendix E contains the new analytic proof; Section 5.9 proves its weighted assembly and physical cutoffs. Independent review and complete analytic Lean verification remain outstanding. The two-production exponent $100/203$ remains the fully machine-checked baseline.

One sufficient remaining target is

$$\#\{n \text{ odd in } (y, 2y] : \tau(n) > \lceil CL(y) \rceil\} \ll y(\log y)^{-e}, \quad e > 3/8,$$

for a fixed C and all sufficiently large y . This would imply universal termination by Theorem 7.2. The unconditional Lean baseline gives the same implication for $e > 103/203$. No such arithmetic rate is proved. Universal termination alone gives no uniform stopping-time bound here: Theorem 7.3 concerns eventual entry.

Live pressure and its scale-averaged form are sufficient routes to the missing rate. The no-momentum hypothesis supplies a sufficient condition for the pressure bound. The earlier localized Appendix C and ideal-production models retain their stated hypotheses; they do not limit the new written exponent or establish further production inequalities. No nontrivial cycle or unbounded orbit is excluded.

Appendix A. Lean names

All in `formal/Problems/Juggler/FateContagion.lean` unless noted. The root `formal/Problems/JugglerFatePaper.lean` imports exactly the thirty-seven modules named here and builds with `lake build Problems.JugglerFatePaper` without `sorry` and without `native_decide`; `formal/AxiomCheckPaperC.expected` records the axioms of every name below. Lean certifies the exact combinatorial identities and the two abstract lemmas listed here, not the analytic density estimates.

Statement	Lean
backward closure	<code>BackwardClosed, backwardClosed_iterate</code>
Lemma 2.1, fate classes closed	<code>reachesOne_backwardClosed,</code> <code>not_reachesOne_backwardClosed,</code> <code>ancestor_backwardClosed,</code> <code>escapes_backwardClosed,</code> <code>reachesOne_floorPower, escapes_floorPower</code>
Lemma 2.1, trichotomy	<code>Periodic, periodic_of_repeat,</code> <code>fate_trichotomy</code> (from <code>cycles_or_escapes</code>)
Lemma 2.1, exclusion	<code>reachesOne_bounded, reachesOne_not_escapes,</code> <code>cycle_basin_bounded,</code> <code>cycle_basin_not_escapes,</code> <code>reachesOne_not_cycle_basin,</code> <code>periodic_iterate_mod</code>
Lemma 3.1 (even block)	<code>floorPower_even_block, even_block_mem,</code> <code>even_block_card</code>
Lemma 3.2 (OE fiber)	<code>sqrt_sqrt_eq_iff, floorPower_oe_fiber,</code> <code>oe_fiber_mem, oe_fiber_disjoint</code>
Theorem 6.1 (odd generation)	<code>ForwardClosed, reachesOne_forwardClosed,</code> <code>not_reachesOne_forwardClosed,</code> <code>escapes_forwardClosed,</code> <code>mem_iff_floorPower_mem,</code> <code>exists_odd_ancestor,</code> <code>exists_odd_ancestor_ge_three,</code> <code>nonempty_iff_odd_image_mem, odd_mem_iff</code>
Lemma 8.1 (envelope descent)	<code>iterate_le_of_envelope,</code> <code>mem_of_envelope_floor,</code> <code>reachesOne_of_itinerary_envelope; power</code> <code>envelope power_bound_word</code> (Paper A layer)

Statement	Lean
Lean floor $N_0 = 260$, in Problems/Juggler/TerminationFloor257.lean	reachesOne_of_lt_two_hundred_sixty_one
Lemma 4.7 (cube fibers), in Problems/Juggler/CubeFiber.lean	cube_fiber_range, cube_fiber_sqrt_even, cube_fiber_even_image, even_cube_fiber_full, cube_fiber_sqrt_odd, cube_fiber_alternating, odd_cube_fiber_alternating
Shared counting of Lemmas 4.1, 4.1' and 4.3 (separated sequences in a window), in Problems/Juggler/FateWindowCount.lean	WindowCount.StepGe, WindowCount.StepLe, WindowCount.span_ge, WindowCount.span_le, WindowCount.mono_of_stepGe, WindowCount.window_card_le, WindowCount.window_card_le_nat
Proposition 4.4, the exact layer and the deduction from the exponential-sum bounds (the bounds are hypotheses), in Problems/Juggler/FateBlockAverage.lean	BlockAverage.blockE, BlockAverage.oddBlock, BlockAverage.mem_oddBlock, BlockAverage.mem_oeFiber_iff_cell34, BlockAverage.U, BlockAverage.oddBlock_card_eq, BlockAverage.U_card_eq, BlockAverage.psi, BlockAverage.slowSum, BlockAverage.fastSum, BlockAverage.productSum, BlockAverage.four_card_U, BlockAverage.block_average_of_bounds, BlockAverage.block_average_bound, BlockAverage.psi_succ, BlockAverage.abs_psi, BlockAverage.oddBlock_filter_eq, BlockAverage.slowSum_eq_fibers, BlockAverage.oeFiber_card_succ_diff, BlockAverage.abs_alt_sum_le, BlockAverage.slowSum_abs_le, BlockAverage.card_Ico_block, BlockAverage.oddBlock_card_le, BlockAverage.oddBlock_card_ge, BlockAverage.oddBlock_quarter_close, BlockAverage.block_average_two_bounds, BlockAverage.block_average_asymptotic, BlockAverage.block_average_bound_two

Statement	Lean
Section 5.1 without its exponential sums, and Theorem 5.3 at exponent 13/40 unconditionally, in Problems/Juggler/FateProduction.lean	Production.sqrt_floor_exp, Production.floor_exp_ge_half, Production.family_E, Production.good_fiber_logMass_ge, Production.family_OE, Production.errE, Production.errOE, Production.errAdd, Production.two_le_exp, Production.one_div_floor_exp_le, Production.sum_Ioc_le_sum_Ioc_pred, Production.halfLogMass_split, Production.production_two, Production.rate2, Production.coef2, Production.err2, Production.rate2_ge, Production.rate2_le, Production.rate2_pos, Production.coef2_ge, Production.err2_nonneg, Production.err2_le, Production.production_two_sum, Production.zeta2_pos, Production.zeta2_antitone, Production.errors_le, Production.errors_vanish, Production.contagion_elementary, Production.logMass_contagion_elementary, Production.failures_logMass_ge, Production.conjecture_of_tao_rate, Production.conjecture_of_cylinder_bound
Corollary 5.5(2), the log-mass clause at exponent 13/40, in Problems/Juggler/FateProduction.lean	Production.failures_logMass_ge
Section 5.7 and Appendix D, the production words and their prefix-freeness, in Problems/Juggler/FateProductionWords.lean	FateProductionWords.Vword, FateProductionWords.Vword_length, FateProductionWords.Vword_oddCount, FateProductionWords.Vword_eq_odd_cons, FateProductionWords.Vword_prefix_iff, FateProductionWords.Vword_six_prefixFree, FateProductionWords.Vword_five
Section 4.3, the exact layer of the share law (the expansion, the range, the integral), in Problems/Juggler/FateShareLaw.lean	ShareLaw.taylor_three_halves, ShareLaw.xval_expansion, ShareLaw.xval_expansion_fiber, ShareLaw.phi, ShareLaw.phiRange, ShareLaw.phi_ge_min, ShareLaw.phi_le_chord, ShareLaw.phi_sub_le, ShareLaw.exists_phi_sub_eq, ShareLaw.phiRange_le_half_iff, ShareLaw.extremeMeasure, ShareLaw.extremeMeasure_eq_zero, ShareLaw.extremeMeasure_piece1, ShareLaw.extremeMeasure_piece2, ShareLaw.extremeMeasure_piece3, ShareLaw.extremeMeasure_piece4, ShareLaw.integral_extremeMeasure

Statement	Lean
Section 10(d), the cylinder-splitting identity, in Problems/Juggler/FateCylinderEnergy.lean	CylinderEnergy.itinerary_succ_append, CylinderEnergy.wordCount, CylinderEnergy.wordCount_split, CylinderEnergy.energy, CylinderEnergy.bias, CylinderEnergy.energy_succ, CylinderEnergy.sum_bias_sq
Appendix D.1, the exact landing windows (D.1) and (D.2), in Problems/Juggler/FateLandingWindow.lean	LandingWindow.cell34, LandingWindow.cell34_eq_floorPower_two, LandingWindow.le_cell34_iff, LandingWindow.cell34_lt_iff, LandingWindow.windowStart, LandingWindow.windowStart_exists, LandingWindow.windowStart_le_iff, LandingWindow.lt_windowStart_iff, LandingWindow.exact_endpoints, LandingWindow.setOf_cell34_mem_Ico, LandingWindow.exact_endpoints_iterate, LandingWindow.setOf_iterate_mem_Ico
Lemma 4.1 (sweep), in Problems/Juggler/FateSweep.lean	Sweep.cell, Sweep.sweep_cell, sweep_fract_lt_half, sweep_fract_ge_half, sweep_ceil, sweep_rep_le_half, sweep_rep_gt_half
Lemma 4.1' (monotone pairing), in Problems/Juggler/FateSweepMonotone.lean	Sweep.sweep_monotone_cell, sweep_monotone_fract_lt_half, sweep_monotone_fract_ge_half, sweep_monotone_ceil, sweep_monotone_rep_le_half, sweep_monotone_rep_gt_half
Lemma 5.1 (recursion), in Problems/Juggler/FateRecursion.lean	recursion_lemma
Section 6.2, Proposition 6.3(i) and the exact layer of (6.1), in Problems/Juggler/FateFirstLetter.lean	MinimalMember, minimalMember_odd, minimalMember_image_odd, minimal_failure_odd_odd, exists_minimal_failure, first_letter_trichotomy, first_letter_pieces_disjoint, first_letter_split, shellLogMass, shellLogMass_split, floorPower_odd_lt, floorPower_odd_injective, ooPiece, sum_image_ooPiece, card_image_ooPiece
Lemma 8.2 (Chernoff count), in Problems/Juggler/FateChernoff.lean	weightGen_one, count_oddCount_ge_le, count_oddCount_ge_real_le, entropyLog, klHalf, klHalf_eq, klHalf_nonneg, tilt_value, count_oddCount_ge_le_exp, count_oddCount_ge_le_kl, LBad, pC, chernoffExponent, logb_two_three_le, half_le_pC, pC_lt_one, LBad_oddCount_ge, LBad_count_le
Theorem 8.3 (explicit form), in Problems/Juggler/FateChernoff.lean	cylinder, oddFailures, EnvelopeBad, oddFailures_subset_bad_cylinders, oddFailures_card_le, LBad_of_envelopeBad, oddFailures_card_le_chernoff, scaleRatio, scaleL, depth, cylinder_even_root_empty, oddFailures_card_le_explicit

Statement	Lean
Theorem 9.2 (pressure form), in Problems/Juggler/FatePressure.lean	livePressure, live_count_le_pressure, envelopeBad_of_liveTo, LBad_of_liveTo, live_oddCount_ge, live_count_le_of_pressure
Proposition 9.3 (pressure telescoping), in Problems/Juggler/TiltedShare.lean	oddMass, tiltedShare, weightGen_succ_le_share, one_add_le_exp_excess, weightGen_le_pressure, count_le_pressure, NoMomentum, count_le_of_noMomentum, tilt_exponent_eq_kl, MeanShare, weightGen_le_of_meanShare, MeanShareOff, initial_depths_are_free, tower_ratio_lt_one
Lemma 5.2 (seed), in Problems/Juggler/FateSeed.lean	exists_ge_three_of_backwardClosed, seed_lemma, seed_constant_pos
Theorem 7.2 (Tao-type rate, contagion as a hypothesis), in Problems/Juggler/FateTaoReduction.lean	logMass_le_oddLogMass, oddLogMass_le_of_dyadic, tao_rate_implies_empty, tao_rate_implies_conjecture
Corollary 8.4 (the conjecture from a cylinder bound), in Problems/Juggler/FateCylinderCor ollary.lean	CylinderBound, one_le_depth, chernoffExponent_nonneg, oddFailures_eventually_le, cylinder_bound_implies_conjecture
Shared numerics of Lemmas 4.2, 4.3 and 8.2 (roots by powering, Bernoulli at a point), in Problems/Juggler/FateNumerics.lean	Numerics.rpow_le_iff_pow, Numerics.le_rpow_iff_pow, Numerics.rpow_lt_iff_pow, Numerics.lt_rpow_iff_pow, Numerics.bernoulli_ge, Numerics.bernoulli_le
Lemma 4.2 (fiber parity), in Problems/Juggler/FateFiberParity.lean	FiberParity.xval, FiberParity.two_xval, FiberParity.floor_two_xval, FiberParity.cell_xval_even_iff, FiberParity.xval_step, FiberParity.xval_step_ge, FiberParity.xval_step_le, FiberParity.xval_step_mono, FiberParity.oeFiber, FiberParity.mem_oeFiber, FiberParity.oeFiber_eq_image, FiberParity.oeFiber_card, FiberParity.evenImageCount, FiberParity.fiber_ge_rpow, FiberParity.fiber_lt_rpow, FiberParity.rpow_four_thirds_succ_ge, FiberParity.rpow_four_thirds_succ_le, FiberParity.rpow_two_thirds_succ_le, FiberParity.oeFiber_card_ge, FiberParity.oeFiber_card_le, FiberParity.Am, FiberParity.alpha, FiberParity.eps, FiberParity.Good, FiberParity.eps_le, FiberParity.step_ge, FiberParity.step_le, FiberParity.fiber_parity_good

Statement	Lean
Lemma 4.3 (thin fibers), in Problems/Juggler/FateThinFibers.lean	FiberParity.span_ge_of_step, FiberParity.arc_count_le, FiberParity.Am_step_le, FiberParity.Am_step_ge, FiberParity.eps_antitone, FiberParity.bad_mem_arc, FiberParity.two_rpow_third_le, FiberParity.two_rpow_two_thirds_le, FiberParity.rpow_two_thirds_ge, FiberParity.eps_div_eps_double, FiberParity.Am_double_sub_le, FiberParity.bad_count_le, FiberParity.bad_block_logMass_le, FiberParity.bad_sum_dyadic_le, FiberParity.eps_pow_two_mul, FiberParity.two_rpow_neg_third_le, FiberParity.bad_logMass_le
Theorem 5.3 given (5.2), Theorem 7.3, Corollary 8.4 through (5.2), in Problems/Juggler/FateContagionBound.lean	productionRate, productionCoeff, productionRate_pos, productionRate_ge, productionRate_le, productionRate_le_one, productionCoeff_ge, productionCoeff_nonneg, zeta, zeta_antitone, le_rpow_div_of_pow_le, zeta_pos_49, seedConst, gA, gA_seed, logMass_ge_gA, contagion_of_production_inequality, logMass_contagion_of_production, oddFailures_eq_empty, tao_rate_iff_conjecture, conjecture_of_cylinder_bound_of_production
Proposition 4.4 (its two exponential-sum bounds, the fast sum and the product sum; the exact layer, the slow sum, the block count and the deduction of (4.1) and of the asymptotic form from those two bounds are Lean), the share law 4.5 and Corollary 4.6 (the phase expansion, the range of the quadratic phase and the integral 25/108 are Lean, the equidistribution and the measure identifications are not), the production inequality (5.2) (its E-family and OE-fiber family are Lean with explicit errors and give Theorem 5.3 at exponent 13/40 unconditionally, and the averaged OE family of Section 5.8 gives it at 100/203; the block-average family and the ladder are not Lean and are no longer needed for the exponent), Corollaries 5.4 and 5.5 in the range $100/203 < \lambda < \lambda_{\text{ideal}}$ (at 100/203 both are Lean, in Problems/Juggler/FateDyadicDensity.lean), Sections 8–10 except Lemma 8.2, the explicit form of Theorem 8.3, Corollary 8.4, the exact form of Theorem 9.2 and Proposition 9.3, Appendix C; Theorem 5.3 in the range $100/203 < \lambda < \lambda_{\text{ideal}}$, and Theorems 7.2, 7.3 and Corollary 8.4 with the contagion bound as a hypothesis outside the exponents of Section 5.8	human proofs

Statement	Lean
Theorem 9.1 (one-sided form, exact, by exponential moments), in Problems/Juggler/FateOneSided.lean	OneSided.cylinder_split, OneSided.LBad_of_LBad_append, OneSided.sum_allWords_succ, OneSided.sum_pow_oddCount_le, OneSided.badWeight, OneSided.badWeight_nonneg, OneSided.badWeight_le_card, OneSided.badMass, OneSided.OneSidedShare, OneSided.badMass_succ_le, OneSided.card_cylinder_zero_le, OneSided.badMass_one_le, OneSided.badMass_le, OneSided.oddFailures_card_le_badMass, OneSided.one_sided_bound, OneSided.klDiv, OneSided.tilt, OneSided.tilt_ge_one, OneSided.tilt_pow_ratio, OneSided.one_sided_bound_kl
Theorem 9.1's consequence (the conjecture from the one-sided hypothesis), in Problems/Juggler/FateOneSidedCorollary.lean	OneSided.OneSidedBound, OneSided.OneSidedExact, OneSided.oneSidedExponent, OneSided.OneSidedShare.mono, OneSided.oneSidedBound_of_exact, OneSided.klDiv_nonneg, OneSided.exp_le_rpow_scale, OneSided.pow_le_rpow_scale, OneSided.oddFailures_le_of_one_sided, OneSided.implies_conjecture_of_contagion, OneSided.one_sided_implies_conjecture, OneSided.exact_share_implies_conjecture
Section 9.2's consequences (the conjecture from the pressure and no-momentum hypotheses), in Problems/Juggler/FatePressureCorollary.lean	Pressure.oddFailures_subset_live, Pressure.PressureBound, Pressure.NoMomentumBound, Pressure.momentumExponent, Pressure.absorb, Pressure.oddFailures_le_of_pressure, Pressure.pressure_conj_of_contagion, Pressure.pressure_implies_conjecture, Pressure.oddFailures_le_of_noMomentum, Pressure.noMomentum_conj_of_contagion, Pressure.noMomentum_implies_conjecture
Section 10(d), Theorem 9.1 with exceptional atoms, in Problems/Juggler/FateOneSidedAtoms.lean	OneSided.OneSidedShareExc, OneSided.oneSidedShareExc_of_share, OneSided.OneSidedBoundExc, OneSided.oneSidedBoundExc_of_bound, OneSided.badMass_succ_le_exc, OneSided.badMass_le_exc, OneSided.one_sided_bound_exc, OneSided.main_term_eq, OneSided.one_sided_bound_kl_exc, OneSided.pow_le_rpow_scale_gen, OneSided.tail_le, OneSided.oddFailures_le_of_exc, OneSided.exc_conj_of_contagion, OneSided.exc_implies_conjecture

Statement	Lean
Section 10(d), the bias energy supplying the exceptional atoms, in Problems/Juggler/FateEnergyAtoms.lean	OneSided.OneSidedShareExc.mono_err, Energy.bias, Energy.energyOn, Energy.biasEnergy, Energy.badEnergy, Energy.violators, Energy.badEnergy_le_biasEnergy, Energy.card_allWords, Energy.mass_violators_le, Energy.oneSidedShareExc_of_energy, Energy.wordCount_cylinder, Energy.biasEnergy_eq, Energy.EnergyBound, Energy.oneSidedBoundExc_of_energy, Energy.energy_conj_of_contagion, Energy.energy_implies_conjecture
Section 10(d), the collapsed component, in Problems/Juggler/FateCollapse.lean	Collapse.fiber, Collapse.window, Collapse.windowBias, Collapse.card_filter_window, Collapse.card_window, Collapse.windowBias_eq_sum, Collapse.abs_alt_sum_le, Collapse.Icc_eq_Ico, Collapse.abs_windowBias_le, Collapse.lt_sq_succ_of_floorPower_eq, Collapse.pre, Collapse.iterate_succ_eq, Collapse.fiber_succ, Collapse.pre_filter_even, Collapse.blockSum, Collapse.oddPart, Collapse.oddPart_nonneg, Collapse.fiber_succ_eq, Collapse.evenCount, Collapse.evenCount_le, Collapse.le_evenCount, Collapse.blockSum_sub_le, Collapse.sum_abs_oddPart_sub_le, Collapse.collapse_bias_le, Collapse.fiber_succ_sandwich, Collapse.collapse_bias_two_step

Statement	Lean
Section 8.4, the unconditional criteria at $C = 30$ with every constant a numeral, in Problems/Juggler/FateCertified.lean	Certified.e_ge, Certified.e_le, Certified.log_le_of_pow_le, Certified.le_log_of_pow_le, Certified.logb_two_le, Certified.le_logb_two, Certified.log_two_le, Certified.le_log_two, Certified.le_logb_three, Certified.logb_three_le, Certified.le_log_lo, Certified.le_log_hi, Certified.log_lo_le, Certified.log_hi_le, Certified.logb_three_pos, Certified.pC_thirty_ge, Certified.pC_thirty_le, Certified.klHalf_ge_of, Certified.klHalf_le_of, Certified.klHalf_thirty_ge, Certified.klHalf_thirty_le, Certified.chernoffExponent_thirty_gt, Certified.chernoffExponent_thirty_lt, Certified.cylinder_bound_thirty, Certified.pressure_thirty, Certified.klDiv_half, Certified.oneSidedExponent_half, Certified.logb_tilt_thirty_le, Certified.one_sided_thirty
Lemma 5.14 (block lock), in Problems/Juggler/FateBlockLock.lean	BlockLock.gridRes, BlockLock.fract_grid, BlockLock.gridRes_mul_inj, BlockLock.grid_reindex, BlockLock.grid_half_count, BlockLock.grid_near_count, BlockLock.span_bounds, BlockLock.block_count, BlockLock.block_chain, BlockLock.block_lock
Lemma 5.15 (fiber lock), in Problems/Juggler/FateFiberLock.lean	FiberParity.evenImageCount_eq_fract, FiberParity.eps_mul_Hlen_le, FiberParity.exists_coprime_approx, FiberParity.fiber_block_lock, FiberParity.fiber_lock
The resonance count behind Theorem 5.16, in Problems/Juggler/FateResonanceCount.lean	FiberParity.Resonant, FiberParity.resonant_mem_arc, FiberParity.Am_window_le, FiberParity.shifted_arc_count_le, FiberParity.resonance_count_one, FiberParity.resonance_count_le, FiberParity.resonance_count_le'
Theorem 5.16 (poor-fiber tail) and the non-poor fiber bound of Corollary 5.17, in Problems/Juggler/FatePoorTail.lean	FiberParity.Poor, FiberParity.oeFiber_nonempty, FiberParity.resonant_mono, FiberParity.poor_resonant, FiberParity.poor_count_le, FiberParity.poor_count_le', FiberParity.poor_block_logMass_le, FiberParity.poor_sum_dyadic_le, FiberParity.poor_logMass_le, FiberParity.nonpoor_fiber_logMass_ge

Statement	Lean
Theorem 5.18 and its consequences (Section 5.8), in <code>Problems/Juggler/FatePoorProduction.lean</code>	<code>Production.family_OE_averaged,</code> <code>Production.err0Eavg,</code> <code>Production.errAddAvg,</code> <code>Production.production_two_averaged,</code> <code>Production.coef2avg,</code> <code>Production.err2avg,</code> <code>Production.coef2avg_ge,</code> <code>Production.coef2avg_nonneg,</code> <code>Production.err2avg_nonneg,</code> <code>Production.err2avg_le,</code> <code>Production.production_two_averaged_sum,</code> <code>Production.errorsAvg_le,</code> <code>Production.errorsAvg_vanish,</code> <code>Production.zeta2avg_antitone,</code> <code>Production.contagion_averaged,</code> <code>Production.zeta2avg_pos,</code> <code>Production.logMass_contagion_averaged,</code> <code>Production.failures_logMass_averaged,</code> <code>Prod</code> <code>uction.conjecture_of_tao_rate_averaged,</code> <code>Production.conjecture_of_cylinder_averaged</code>
Corollary 5.4 and Corollary 5.5 at the fate classes (Section 5.5), in <code>Problems/Juggler/FateDyadicDensity.lean</code>	<code>Density.blockCount,</code> <code>Density.block_logMass_le,</code> <code>Density.block_index,</code> <code>Density.halfLogMass_le_blocks,</code> <code>Density.exists_block_of_shell,</code> <code>Density.natDensity_of_shell_bound,</code> <code>Density.natDensity_averaged,</code> <code>Density.reachesOne_logMass_averaged,</code> <code>Density.reachesOne_natDensity_averaged,</code> <code>Density.failures_natDensity_averaged,</code> <code>Density.basin_logMass_averaged,</code> <code>Density.basin_natDensity_averaged,</code> <code>Density.escapes_logMass_averaged,</code> <code>Density.escapes_natDensity_averaged</code>

Appendix B. Constants and artifacts

Exponents. Roots of $2^{-\lambda} + \sum_i c_i e_i^\lambda = 1$:

recursion	terms (c_i, e_i)	root
elementary sweep only	$(\frac{2}{21}, \frac{3}{4})$	0.1385
block average only (λ^*)	$(\frac{1}{3}, \frac{3}{8})$	0.3774
block average + sweep (adversarial 1/7)	$(\frac{5}{21}, \frac{3}{8}), (\frac{2}{21}, \frac{3}{4})$	0.4051
block average + pairing (named intermediate)	$(\frac{1}{9}, \frac{3}{8}), (\frac{2}{9}, \frac{3}{4})$	0.4480
+ <i>OEOEE</i> (named intermediate; elementary, audited)	$\dots, (\frac{1}{27}, \frac{9}{32})$	0.4801
+ $V_3 = \textit{OEOEOEE}$ (named intermediate; elementary, audited)	$\dots, (\frac{1}{81}, \frac{27}{128})$	0.4891
+ $V_4 = \textit{OEOEOEOEE}$ (named intermediate; elementary, audited)	$\dots, (\frac{1}{243}, \frac{81}{512})$	0.4916

recursion	terms (c_i, e_i)	root
+ $V_5 = OEOEOEOEOEE$ (named intermediate; elementary, audited)	$\dots, (\frac{1}{729}, \frac{243}{2048})$	0.4924
+ $V_6 = OEOEOEOEOEOEE$ (λ^{**} , earlier finite-production route; elementary, audited)	$\dots, (\frac{1}{2187}, \frac{729}{8192})$	0.4926
pairing + $OEOEE$ (λ^{***} , Appendix C, conditional)	pairing terms and $(\frac{1}{9}, \frac{9}{32})$	0.5392
depth-two ideal-share model	$(\frac{1}{3}, \frac{3}{4})$	0.4927
two productions at the averaged coefficient, $\eta_0 \downarrow 0$ (λ_{ideal} , Theorem 5.18; Lean to 100/203)	$(\frac{1}{3} - \frac{2}{3}\eta_0, \frac{3}{4})$	0.4927
+ $OEOEE$, $OEOEE$ (closed: fibers $P^{5/32}$; Lemma 3.9 leftover $P^{89/96}$)	$\dots, (\frac{2}{27}, \frac{27}{64})$	0.5561
pairing + $OEOEE$ + $OEOEE$ (model comparison)	pairing terms and $(\frac{4}{27}, \frac{9}{32})$	0.5665
conditional infinite $V_k = (OE)^{k-1}OEE$ model	$\dots, (3^{-(k+2)}, (\frac{3}{4})^k \frac{3}{8})_{k \geq 1}$	0.4927
conditional infinite V_k model and $OEOEE$		0.5769
conditional O -runs ≤ 2 , pairing share (Section 5.7)	transfer matrix	0.6247
conditional O -runs ≤ 2 , ideal-share model	transfer matrix	0.7180
conditional O -runs ≤ 3 , pairing share	transfer matrix	0.7095
conditional O -runs ≤ 3 , ideal-share model	transfer matrix	0.8414
abstract ceiling, all O -runs at the ideal share (Proposition 5.12)	$2^{-\lambda} + \frac{1}{3}(\frac{3}{2})^\lambda = 1$	1

Theorem 5.19 additionally attains the written exponent 5/8, from coefficients 1, 33/100 and 11/100 at scales 1/2, 3/4 and 9/16. Its exact certificate is in Section 5.9; the $OEOEE$ analytic input is not yet completely formalized.

Depth constants. The historical least integer values below use the three explicitly named rate thresholds; the threshold $1 - \lambda_{\text{ideal}} = 0.5073$ of the second route gives the same integers as 0.5074 in every column; intermediate V_2 through V_5 have complementary thresholds 0.5199, 0.5109, 0.5084, and 0.5076, respectively.

criterion	λ^{**} , rate 0.5074	pairing, rate 0.5520	λ^{***} , rate 0.4608
fair Chernoff	19	20	18
one-sided Azuma, $q = 0.5, 0.55, 0.60, 0.62$	19, 41, 223, 1586	20, 44, 240, 1715	18, 39, 206, 1451
optimized pressure, same q	19, 41, 214, 1496	20, 43, 230, 1618	18, 38, 198, 1369

Depth $d(y) = \lceil 20L(y) \rceil$ (the conservative a-fortiori table) with $N_0 = 3.5 \cdot 10^8$: 25 at 10^{20} , 72 at 10^{100} , 138 at 10^{1000} , 204 at 10^{10000} .

Decimal values in the tables are rounded; all strict inequalities use the roots of the displayed defining equations.

Artifacts (repository sneakyweasel/bt1ab; SHA-256):

- data/research/juggler/fate_contagion/summary.json
SHA-256: d322b8549e2239c5fc78279c7a579332766a3287811f73ee4d225d6eb9e5cc7b
- data/research/juggler/tao_reduction/summary.json
SHA-256: 2316e5339e3670d283c1983a5b5b0c11bc92101ab327c477072b7897619a587f
- formal/Problems/Juggler/FateContagion.lean
SHA-256: 834198a6b0e8cc13c9fb91d9ad938f62ce5e783a3cc9009fb38334c913d5b7e2
- formal/Problems/Juggler/CubeFiber.lean
SHA-256: 2b0add4108651218c43fb741a207d3d6860525a2ad13ad1341ece9fc5c37f8e9
- formal/Problems/Juggler/TiltedShare.lean
SHA-256: 7dcafa45a2a0e9bca182a713b495d791cf812e4c70d6472c67bf85b0b736aef4
- formal/Problems/Juggler/FateRecursion.lean
SHA-256: e6d050acc924867abcf5c03b4a3e59b62363f40bbba452e2ced83afe474dd435
- formal/Problems/Juggler/FateFirstLetter.lean
SHA-256: a383c715010cb1e7323359a414aeaa99242550f59f7213049cc528232e37875a
- formal/Problems/Juggler/FateBlockAverage.lean
SHA-256: 5cef5c273577376b72799732865415a91c6b91f8bdb06e79bd398d9c478d0b07
- formal/Problems/Juggler/FateProduction.lean
SHA-256: 7cc24ff9f95692ead6b594aa703677910daf4936f24becce7b13da2c4105df2e
- formal/Problems/Juggler/FateProductionWords.lean
SHA-256: a712db485e4c48029ce9a9afcad1cfea1240f4bb4dda1cb7537661ae1f3fdb3e
- formal/Problems/Juggler/FateOneSided.lean
SHA-256: 7ae0aa4cab061a24ef51eff381a0f4bbd4e4569ae2578f04e4c3677b5999bddc
- formal/Problems/Juggler/FateOneSidedCorollary.lean
SHA-256: 481b4cecbabd7291d551b814f733b9cfd78dec50a936a31fd9ab6544408c714e
- formal/Problems/Juggler/FatePressureCorollary.lean
SHA-256: cdd1ca9cbb59c4bb0a5fb24702f12bb8e27c21365f9b28e515819f874e9b437e
- formal/Problems/Juggler/FateOneSidedAtoms.lean
SHA-256: db2601560b54b95a5ac5b2f5f16c0d83e3b4cf4182f1d20795c47db26c0a9ce3
- formal/Problems/Juggler/FateEnergyAtoms.lean
SHA-256: e98ccccbb25140d41ede236466f7e0f522c433a4c80fcaa330abfb1388653b1a
- formal/Problems/Juggler/FateCollapse.lean
SHA-256: 56ba2f731fdd02ea1d19a6e6ef909534947c8c30b48828c72f8009901ed06e06
- formal/Problems/Juggler/FateCertified.lean
SHA-256: ff47a23462fb591264125859b9d511bb2cec046da7a71fbad5d17b35a5581c90

- `formal/Problems/Juggler/FateShareLaw.lean`
SHA-256: `a229f4fa4cf48cb5b003c746672947eec10bf93d7287ad7eecaa3f84d5236e7c`
- `formal/Problems/Juggler/FateCylinderEnergy.lean`
SHA-256: `274cd6b960bdd1de8c1b637bd2dee86c8cbe67e883e25937812dadcb9e98d6a0b`
- `formal/Problems/Juggler/FateLandingWindow.lean`
SHA-256: `ef28985363d3b855e0a9ec4b8fba687855d34ab8cf1ff9b355a50e9377e29545`
- `formal/Problems/Juggler/FateWindowCount.lean`
SHA-256: `9d790163433899c73993a24b434ff20ae45aba3dbe47b717507dba840307b6a3`
- `formal/Problems/Juggler/FateSweep.lean`
SHA-256: `a5ede4a6d25cec71d6b6c7fb99e577df3335366175dd22f361581456669a72cc`
- `formal/Problems/Juggler/FateChernoff.lean`
SHA-256: `3b808649443805fa3dd41b89333afe886f3cbc7e36f1bdae23f6c93fe95a7032`
- `formal/Problems/Juggler/FatePressure.lean`
SHA-256: `f69ad74fcaed87b692451113cf72eeadc1182efc0ac8121e255e6b079bf2bbf2`
- `formal/Problems/Juggler/FateCylinderCorollary.lean`
SHA-256: `e2ffc98f56268a31826b86f54115e0fed949e62651d05936d545cd829761b4d2`
- `formal/Problems/Juggler/FateNumerics.lean`
SHA-256: `8340b3a8a4ff2ce35bd77b7697ca95b9d36e5d61ea4173782287d0839b381737`
- `formal/Problems/Juggler/FateFiberParity.lean`
SHA-256: `bf2b1c6058c3d00f0f9cf24f11bb7bee7a17a75375c2ddc7223e065621e64131`
- `formal/Problems/Juggler/FateThinFibers.lean`
SHA-256: `3fbcd3e9f7adf13967e7494691aa8f62779630e0446fe81e813486947478cb74`
- `formal/Problems/Juggler/FateContagionBound.lean`
SHA-256: `d8991fd05c076875a0e8fd75692a10bb31930c289a3812de07212b620645acca`
- `formal/Problems/Juggler/FateBlockLock.lean`
SHA-256: `b4fdc02e5b41b8759156f5aa1b3f208854d2e6c994a3963e24aa164df498bed7`
- `formal/Problems/Juggler/FateFiberLock.lean`
SHA-256: `f82aceb5a06642f7f55ea1b1bbee317ea487c46e97cbf52aecbd920dc9d10007`
- `formal/Problems/Juggler/FateResonanceCount.lean`
SHA-256: `f39eca082166b63d15989946e8ae54279bb9a4b779d5440d0c970212883f4aa1`
- `formal/Problems/Juggler/FatePoorTail.lean`
SHA-256: `773c579e2ba303dc49f1488020ea142b0b35ea26573707ebfd6afca7c4dd284b`
- `formal/Problems/Juggler/FatePoorProduction.lean`
SHA-256: `0fc332ef6b54bebbe4de215b03ac582a2d0cc38d82a391d23856624599940283`

- `formal/Problems/Juggler/FateDyadicDensity.lean`
SHA-256: f5065c952302de36981e147f5ae95804eaa16d34a38459025a65e10b5a24e4fe
- `formal/Problems/JugglerFatePaper.lean`
SHA-256: ce57dde1ab5f47f98277f4b870641f864727a4136b617627773e531e2b6753c2
- `formal/AxiomCheckPaperC.expected`
SHA-256: b8569df4c76a94d8665940b04dba86739d773a796d8683e7801ede5c654ddc24
- `src/research/juggler_sequence/fate_contagion.py`
SHA-256: 27b94f873d4cc9fc4f424443db058c8b3f7c6217e9bdbde7cc17c984829e9c46
- `src/research/juggler_sequence/tao_reduction.py`
SHA-256: 7d4cc3e0dbf197232c084e0aa6fe89cff7deb77bdab2d52523f5bcfc58a9e14f
- `docs/theory/figures/render_paper_c_figures.py`
SHA-256: 955e5771d36ba103a5192cb28432244436a2ced78edf482d532d1fe6fe9afbd0

The archived records retain the parameter names and values of their original runs (including the older pairing-only threshold). Current exponents and depth constants are checked separately in `data/research/juggler/paper_c_publication_review/validation.json`. The numerical experiments of Section 11 use fixed random seeds. Their code is run by `python -m research.juggler_sequence.fate_contagion` and `python -m research.juggler_sequence.tao_reduction`. The current code reports current rate constants; archived parameter labels are preserved in the supplied JSON. The figures are regenerated by `python docs/theory/figures/render_paper_c_figures.py`.

Appendix C. An earlier conditional strengthening

This appendix depends on one analytic hypothesis beyond the main text. No statement of Sections 1–12 uses it; it improves the constants of Theorems 1, 3 and 4 and nothing else.

C.1 The hypothesis

For odd n put $m = \lfloor n^{3/2} \rfloor$, $v = \lfloor m^{3/2} \rfloor$, $\psi(y) = (-1)^{\lfloor y \rfloor}$, and

$$\psi_1 = \psi(n^{3/2}), \quad \psi_2 = \psi(m^{3/2}), \quad \psi_3 = \psi(v^{1/2}),$$

the signs of the parities of m , of $\lfloor m^{3/2} \rfloor$ and of $\lfloor v^{1/2} \rfloor$ — the second, third and fourth letters of the itinerary of n along the branch *OOE*.

Hypothesis L (localized triple parity discrepancy with a slow twist). For every $\varepsilon > 0$ there is P_0 such that for all $P \geq P_0$, every interval $I \subseteq (P, 2P]$ of length $Y \geq P^{1/2}$, every $(a, b, c) \in \{0, 1\}^3 \setminus \{(0, 0, 0)\}$, and every integer ℓ with $|\ell| \leq P^{1/24}$,

$$\left| \sum_{n \in I, n \text{ odd}} \psi_1^a \psi_2^b \psi_3^c e\left(\frac{\ell}{2} n^{9/16}\right) \right| \leq Y P^{-1/24+\varepsilon}.$$

Status. Hypothesis L is Theorem 4.12 of the working draft [12] (Section 3.5 there), whose Lemma 4.10 removes the twist by partial summation after the Weyl differencing and whose proof runs the seven-step argument of its Theorems 4.4 and 4.7 (exact linearization of the nested floors, differencing with $H = P^{1/12}$, a cell decomposition, second-derivative tests per cell) over a sub-dyadic interval, exhibiting the three absolute terms — two partial end cells

$5.2P^{3/8}$, the pure passenger $3.8P^{7/16}$, the majorant end cells $17P^{1/4}$ — that do not scale with the number of summands. Because [12] is an unrefereed working draft, we keep the statement as a hypothesis here and rely on nothing but the statement; what *is* proved in this appendix is everything that follows from it. (The derivation below coincides with Corollary 4.13 of [12].)

C.2 The *OOEEE* production on even blocks

For $m' \geq 2$ let $I(m') = [m'^{32/9}, (m' + 1)^{32/9})$, an interval of length $Y = \frac{32}{9}m'^{23/9}(1 + O(1/m'))$ at scale $P = m'^{32/9}$, so that $Y \asymp P^{23/32} \geq P^{1/2}$. Let

$$\mathcal{O}(m') = \{n \text{ odd} \in I(m') : \text{word}_5(n) = \text{OOEEE}, J^5(n) = m'\}.$$

Lemma C.1 (transparent nesting at the fifth letter). For odd $n \geq 3$, with m, v as above, $0 \leq n^{9/16} - v^{1/4} \leq n^{-15/16}$. Consequently $\lfloor v^{1/4} \rfloor = \lfloor n^{9/16} \rfloor$ unless $\{n^{9/16}\} < n^{-15/16}$.

Proof. $v \leq m^{3/2} \leq n^{9/4}$ gives $v^{1/4} \leq n^{9/16}$. For the other direction, $m \geq n^{3/2} - 1$ and $v \geq m^{3/2} - 1$ give $v \geq (n^{3/2} - 1)^{3/2} - 1 \geq n^{9/4} - \frac{3}{2}n^{3/4} - 1$, and then, by the mean value theorem for $u \mapsto u^{1/4}$ on $[v, n^{9/4}]$, $n^{9/16} - v^{1/4} \leq (n^{9/4} - v)/(4v^{3/4}) \leq (\frac{3}{2}n^{3/4} + 1)/(4v^{3/4})$. Since $v \geq \frac{1}{2}n^{9/4}$ for $n \geq 3$, $v^{3/4} \geq 2^{-3/4}n^{27/16}$, and the bound is at most $2^{3/4}(\frac{3}{2}n^{3/4} + 1)/(4n^{27/16}) \leq n^{-15/16}$ for $n \geq 3$. The floors of two numbers in $[n^{9/16} - n^{-15/16}, n^{9/16}]$ differ only if an integer lies in that interval, i.e. if $\{n^{9/16}\} < n^{-15/16}$. $\square$

Lemma C.2 (the exceptional set is small). Let $\mathcal{E}(I) = \{n \text{ odd} \in I : \{n^{9/16}\} < n^{-15/16}\}$. For $I \subseteq (P, 2P]$ of length $Y \geq P^{1/2}$, $|\mathcal{E}(I)| \ll YP^{-1/16}$.

Proof. $\mathcal{E}(I) \subseteq \{n \text{ odd} \in I : \{n^{9/16}\} \in [0, P^{-15/16})\}$. By the Erdős–Turán inequality [15, Theorem 2.5], for any $H \geq 1$,

$$|\mathcal{E}(I)| \leq \frac{Y}{2}P^{-15/16} + C\left(\frac{Y}{H} + \sum_{h=1}^H \frac{1}{h} \left| \sum_{n \in I \text{ odd}} e(hn^{9/16}) \right| \right).$$

Write $n = 2k + 1$ and $F_h(k) = h(2k + 1)^{9/16}$; then $F'_h(k) = \frac{9}{8}h(2k + 1)^{-7/16}$ is monotone with $0 < F'_h \leq \frac{9}{8}hP^{-7/16} \leq \frac{1}{2}$ for $h \leq H := \frac{4}{9}P^{7/16}$, so the Kusmin–Landau inequality [13, Theorem 2.1] gives $|\sum_k e(F_h(k))| \ll 1/\min F'_h \ll P^{7/16}/h$. Hence $|\mathcal{E}(I)| \ll YP^{-15/16} + YP^{-7/16} + P^{7/16}$, and $P^{7/16} \leq YP^{-1/16}$ for $Y \geq P^{1/2}$. $\square$

Lemma C.3 (a Fourier expansion of the fifth letter). Let $J = \lfloor P^{1/24} \rfloor$. There are coefficients b_q , $0 < |q| \leq J$, with $|b_q| \ll 1/|q|$, and a nonnegative trigonometric polynomial Δ_J of degree J with all coefficients at most $1/(J + 1)$, such that for every real y , $|\psi(y) - \sum_{0 < |q| \leq J} b_q e(qy/2)| \leq 2\Delta_J(y/2)$. Moreover, for $I \subseteq (P, 2P]$ of length $Y \geq P^{1/2}$,

$$\sum_{n \in I \text{ odd}} \Delta_J(\tfrac{1}{2}n^{9/16}) \ll YP^{-1/24} + P^{7/16} \log P, \quad \left| \sum_{n \in I \text{ odd}} e(\tfrac{q}{2}n^{9/16}) \right| \ll \frac{P^{7/16}}{|q|} \quad (0 < |q| \leq J).$$

Proof. $\psi(y) = 2 \cdot \mathbf{1}[\{y/2\} < \frac{1}{2}] - 1$, and Vaaler's theorem [10] approximates the indicator of $[0, \frac{1}{2})$ modulo 1 by a trigonometric polynomial of degree J with coefficients $\ll 1/|q|$ and constant term $\frac{1}{2}$, with error at most a majorant Δ_J of the stated form; the constant terms cancel in $2 \cdot \mathbf{1} - 1$. For the two sums, write $n = 2k + 1$: the phase $\frac{q}{2}(2k + 1)^{9/16}$ has derivative $\frac{9q}{16}(2k + 1)^{-7/16}$, monotone, of absolute value between $\frac{9}{16}|q|(2P)^{-7/16}$ and $\frac{9}{16}P^{1/24-7/16} < \frac{1}{2}$, so Kusmin–Landau gives $\ll P^{7/16}/|q|$. The Δ_J sum is its constant term, $\leq Y/(J + 1)$, plus $(J + 1)^{-1} \sum_{0 < |r| \leq J} |\sum_n e(\frac{r}{2}n^{9/16})| \ll P^{7/16} \log J/J$. $\square$

Proposition C.4 (conditional on Hypothesis L). For $m' \geq 2$,

$$|\mathcal{O}(m')| = \frac{1}{16} \# \{n \text{ odd} \in I(m')\} + O_\varepsilon(|I(m')| m'^{-4/27+\varepsilon}),$$

every $n \in \mathcal{O}(m')$ has $J^4(n) \in E(m')$, and for a backward-closed A and $m' \in A$, $\mathcal{O}(m') \subseteq A$ with log-mass at least $\frac{1}{9m'}(1 - O(m'^{-4/27+\varepsilon}))$.

Proof. Along $OOEEE$ the states $J^0(n), \dots, J^4(n)$ are n (odd), m (odd), v (even), $\lfloor v^{1/2} \rfloor$ (even), $\lfloor \sqrt{\lfloor v^{1/2} \rfloor} \rfloor = \lfloor v^{1/4} \rfloor$ (even; the inner identity is the exact nesting of square roots, Lemma 3.2), and $J^5(n) = \lfloor \sqrt{J^4(n)} \rfloor$. If $J^5(n) = m'$ then $J^4(n) \in [m'^2, (m'+1)^2)$ is even, i.e. $J^4(n) \in E(m')$, so $n \in A$ by Lemma 3.1 and backward closure. For the count, let $I = I(m')$, $P = m'^{32/9}$, $Y = |I|$. For $n \in I \setminus \mathcal{E}(I)$, Lemma C.1 gives $\lfloor v^{1/4} \rfloor = \lfloor n^{9/16} \rfloor$, and $n \in I$ means $n^{9/16} \in [m'^2, (m'+1)^2)$; hence for such n with $\text{word}_5(n) = OOEEE$, $J^4(n) = \lfloor n^{9/16} \rfloor \in [m'^2, (m'+1)^2)$ and $J^5(n) = m'$ automatically. Therefore

$$|\mathcal{O}(m')| - \#\{n \text{ odd} \in I : \text{word}_5(n) = OOEEE\} \leq |\mathcal{E}(I)| \ll YP^{-1/16}$$

by Lemma C.2. Next, for odd n the indicator of $OOEEE$ is $\frac{1}{16}(1 - \psi_1)(1 + \psi_2)(1 + \psi_3)(1 + \psi_4)$ with $\psi_4 = \psi(v^{1/4})$: the factor $1 - \psi_1$ enforces m odd, so that $J^2(n) = v$ and ψ_2 is its parity sign; $1 + \psi_2$ enforces v even, so that $J^3(n) = \lfloor v^{1/2} \rfloor$ and ψ_3 is its sign; $1 + \psi_3$ enforces $J^3(n)$ even, so that $J^4(n) = \lfloor v^{1/4} \rfloor$ and ψ_4 is its sign. Expanding the product gives the main term $\frac{1}{16}\#\{n \text{ odd} \in I\}$ and fifteen sign sums $\pm \frac{1}{16} \sum_n \psi_1^a \psi_2^b \psi_3^c \psi_4^d$, $(a, b, c, d) \neq 0$.

Sums with $d = 0$. These are Hypothesis L with $\ell = 0$: each is $\leq YP^{-1/24+\varepsilon}$.

Sums with $d = 1$. Replace $\psi_4 = \psi(v^{1/4})$ by $\psi(n^{9/16})$: by Lemma C.1 the two agree off $\mathcal{E}(I)$, so the change is $\leq 2|\mathcal{E}(I)| \ll YP^{-1/16}$. Then expand $\psi(n^{9/16})$ by Lemma C.3:

$$\sum_n \psi_1^a \psi_2^b \psi_3^c \psi(n^{9/16}) = \sum_{0 < |q| \leq J} b_q \sum_n \psi_1^a \psi_2^b \psi_3^c e\left(\frac{q}{2}n^{9/16}\right) + O\left(\sum_n \Delta_J\left(\frac{1}{2}n^{9/16}\right)\right),$$

using $|\psi_i| = 1$ for the error. If $(a, b, c) \neq 0$, each inner sum is Hypothesis L with $\ell = q$, $|q| \leq J \leq P^{1/24}$, and $\sum_q |b_q| \ll \log P$, so the total is $\ll YP^{-1/24+\varepsilon} \log P$. If $(a, b, c) = 0$ the inner sums are the pure sums of Lemma C.3, $\ll P^{7/16}/|q|$, and their total against $|b_q| \ll 1/|q|$ is $\ll P^{7/16}$. The majorant term is $\ll YP^{-1/24} + P^{7/16} \log P$ by Lemma C.3.

Collecting, with $P^{7/16} \log P \leq YP^{-1/16} \log P$ for $Y \geq P^{1/2}$, every error is $\ll YP^{-1/24+\varepsilon}$, and $P^{-1/24} = m'^{-4/27}$. For the log-mass, each $n \in \mathcal{O}(m')$ has $1/n \geq (m'+1)^{-32/9}$, and $\frac{1}{16} \cdot \frac{1}{2} \cdot \frac{32}{9} m'^{23/9} = \frac{1}{9} m'^{23/9}$. $\square$

C.3 The improved exponent

Theorem C.5 (conditional on Hypothesis L). Theorem 5.3 holds for every $\lambda < \lambda^{***} \approx 0.5392$, the root of

$$2^{-\lambda} + \frac{1}{9}\left(\frac{3}{8}\right)^\lambda + \frac{2}{9}\left(\frac{3}{4}\right)^\lambda + \frac{1}{9}\left(\frac{9}{32}\right)^\lambda = 1.$$

Proof. Add to the three families of Section 5.1 a fourth: for $m' \in A \cap (x^{9/64}, x^{9/32} - 1]$, the set $\mathcal{O}(m')$; then $I(m') \subseteq (\sqrt{x}, x]$ because $m'^{32/9} > \sqrt{x}$ and $(m'+1)^{32/9} \leq x$. Its members are odd with odd $\lfloor n^{3/2} \rfloor$, hence disjoint from the even members and from the OE -type members of the first three families, and distinct m' give disjoint $I(m')$. By Proposition C.4 the family has log-mass at least $\frac{1}{9}(1 - \varepsilon_7(x))(g_A(9t/32) - 2x^{-9/32})$ with $\varepsilon_7 \rightarrow 0$, so (5.2) gains the term $(\frac{1}{9} - \varepsilon_7)g_A(9t/32)$. Lemma 5.1 applies with the additional pair $(e_4, c_4) = (\frac{9}{32}, \frac{1}{9})$ and the new root. $\square$

Downstream, every constant of Sections 7–10 improves: the rate threshold of Theorems 7.2–7.3 becomes $e > 1 - \lambda^{***} \approx 0.4608$; the least depth constant of Corollary 8.4 and Theorem 9.2 is $C = 18$ ($e(18) \approx 0.480$); in the one-sided form $C(0.5) = 18$, $C(0.55) = 39$. Pairing plus $OOEEE$ sits at 0.5392, above the depth-two ceiling 0.4927, because $OOEEE$ is an extra production. A further production from the words $OOOEE$ and $OOEOE$ on even blocks would give $\lambda = 0.5561$ and $C = 18$; those fibers have length $P^{5/32}$, below the threshold of the localized triple, and Lemma 3.9's trivial bound on the kernel proof is the whole interval, so the analogue of Hypothesis L for the kernel estimate of [12] is not available (fate-contagion note §7.4). Hypothesis L concerns

consecutive odd starts in an interval and says nothing about the odd images S_{odd} ; the free term ψ_F is untouched by this appendix.

Appendix D. The finite nested productions

This appendix gives the analytic input used in (5.10). Only the fixed words $V_k = (OE)^{k-1}OEE$, $1 \leq k \leq 6$, are used. Constants may depend on k and on an arbitrarily small $\varepsilon > 0$; no numerical constant or estimate uniform in an unbounded depth is asserted. The classical estimates are those used in Proposition 4.4.

D.1 Exact endpoints and smooth weights

Put $F(u) = \lfloor u^{3/4} \rfloor$ and $\Phi(a) = \lceil a^{4/3} \rceil$. For integers $a < b$,

$$\{n : a \leq F(n) < b\} = [\Phi(a), \Phi(b)) \cap \mathbb{Z}. \quad (\text{D.1})$$

If n has word V_k , repeated use of Lemma 3.2 gives $J^{2i}(n) = F^i(n)$ for $0 \leq i \leq k$, followed by the last even step. With $w = F^{k-1}(n)$, its final state is $J^{2k+1}(n) = \lfloor w^{3/8} \rfloor$. Thus the exact landing window over $m \geq 2$ is

$$I_k(m) = [\Phi^{k-1}(a), \Phi^{k-1}(b)) \cap \mathbb{Z}, \quad a = \lceil m^{8/3} \rceil, \quad b = \lceil (m+1)^{8/3} \rceil. \quad (\text{D.2})$$

It contains precisely the desired preimages after imposing the parity conditions of V_k . Let $s = (3/4)^{k-1}$, $\rho_k = 3s/8$, and $P = m^{1/\rho_k}$. The mean value theorem, applied successively to $u^{4/3}$, gives endpoint error $O_k(P^{1-s})$ relative to the smooth window $[m^{1/\rho_k}, (m+1)^{1/\rho_k})$. Consequently

$$|I_k(m)| = \rho_k^{-1} m^{1/\rho_k - 1} (1 + o(1)), \quad \frac{|I_k(m) \triangle I_k^{\text{sm}}(m)|}{|I_k^{\text{sm}}(m)|} = O_k(P^{-5s/8}). \quad (\text{D.3})$$

Here and below interval length and integer count differ by at most a bounded endpoint error.

Lean: (D.1) is `LandingWindow.exact_endpoints` and `LandingWindow.setOf_cell34_mem_Ico`, and the nested window (D.2) is `LandingWindow.exact_endpoints_iterate`, in `formal/Problems/Juggler/FateLandingWindow.lean`. There F is `cell34` and $\Phi(a) = \lceil a^{4/3} \rceil$ is `windowStart`, the least n with $a^4 \leq n^3$, which is what the ceiling means in exact arithmetic; the two are a Galois connection, and that is the whole proof. `cell34_eq_floorPower_two` is $J^2 = F$ on an OE step. The smooth comparison (D.3), the endpoint error and the multiplicities are not formalized.

We also need the multiplicity of an inner-layer integer after earlier layers have been summed out. For a fixed $i \geq 1$, put $p_i = (4/3)^i$. Induction in (D.1) gives

$$\Phi^i(u) = u^{p_i} + O_i(u^{p_i-4/3}).$$

For $i = 1$ the error is $O(1)$; each further inverse step multiplies the previous error by $O(u^{p_i/3})$, and its own rounding error is smaller. Taking the difference at $u+1$ and u , and then counting odd integers, shows that the number of odd n with $F^i(n) = u$ is

$$\omega_i(u) = \frac{1}{2} p_i u^{p_i-1} + O_i(u^{p_i-4/3} + 1) = \frac{1}{2} p_i u^{p_i-1} (1 + O_i(u^{-1/3})). \quad (\text{D.4})$$

The last form uses $p_i \geq 4/3$. This weight has a smooth monotone main term and a pointwise power-saving error. On any layer interval of scale W , partial summation transfers bounds for all initial partial sums to this main weight; the pointwise error costs $O_i(W^{-1/3})$ relative to the total mass. At most two partial end fibers occur at each grouping step. Their relative size is at most $O_k(P^{-5s/8})$, or a stronger power, since the shortest relative layer interval has length $W^{5/8}$.

D.2 Averaged cancellation inside the short fibers

Let v range over consecutive integers in an interval at scale W , of length $L \gg W^{5/8}$, contained in a fixed dyadic range. Write $B_v = [v^{4/3}, (v+1)^{4/3}) \cap \mathbb{Z}$. The following bound holds on either parity class of u , and on the full integer lattice:

$$\sum_v \left| \sum_{u \in B_v} \psi(u^{3/2}) \right| \ll LW^{1/3} W^{-1/6} (\log W)^2. \quad (\text{D.5})$$

The same right side bounds the sum over any initial subinterval of the displayed v -range.

To see this, use Vaaler at $Q = \lfloor W^{1/6} \rfloor$. A nonzero Fourier mode on a parity class has phase $q(2r+b)^{3/2}/2$. Its derivative at the beginning of a v -fiber is $\alpha_q(v) = \frac{3}{2}qv^{2/3} + O(qW^{-2/3})$, and changes by $O(qW^{-1/3})$ across that fiber. As a function of v , the leading frequency increases in steps comparable to $\delta = qW^{-1/3}$. The full lattice has the same estimates with a different fixed leading constant.

Fibers whose derivative range meets an integer, or lies within a constant multiple of δ of one, are counted trivially. In each unit frequency interval there are $O(1)$ such fibers; there are $O(L\delta + 1)$ unit intervals. They contribute $O(qL + W^{1/3})$, since a fiber has $O(W^{1/3})$ points. For the remaining fibers, Kusmin–Landau bounds the mode by the reciprocal distance of its derivative range to the nearest integer. Grouping those distances in intervals of length comparable to δ gives a harmonic sum. Including the two incomplete unit intervals at the ends, the total for this mode is

$$\ll qL + L \log W + W^{1/3}(1 + q^{-1} \log W) \ll qL + L \log W. \quad (\text{D.6})$$

The last step uses $L \gg W^{5/8}$. On an initial subinterval use its length in the first bound, and the full L to dominate the endpoint terms; thus the bound needed for partial summation is uniform. Summing (D.6) against coefficients $O(1/q)$ gives $O(LQ + L(\log W)^2)$. The error majorant contributes $O(LW^{1/3}/Q + LQ + L \log W)$, by its $O(1/Q)$ coefficients. These prove (D.5). Bounded coefficients depending only on v may be inserted after the inner absolute value.

In terms of the outer variable scale $Z = W^{4/3}$, the relative saving in (D.5) is $Z^{-1/8}$, up to logarithms. This averaged estimate does not assert cancellation on every individual fiber, which Lemma 4.7 shows would be false.

D.3 The final one-variable estimate

On an interval K of scale W and length $L \asymp W^{5/8}$, each nonconstant product of the three signs

$$(-1)^u, \quad \psi(u^{3/2}), \quad \psi(u^{3/4})$$

has sum $O_\varepsilon(LW^{-1/6+\varepsilon})$. The same majorant holds for every initial subinterval of K .

If the fast sign occurs, expand the fast and slow waves at $Q = \lfloor W^{1/6} \rfloor$. A nonzero fast mode has phase $(qu^{3/2} + ru^{3/4} + bu)/2$, with $b \in \{0, 1\}$, $q \neq 0$, and $|q|, |r| \leq Q$. Its second derivative has constant sign and magnitude comparable to $|q|W^{-1/2}$. The second-derivative test bounds the sum by

$$\ll L|q|^{1/2}W^{-1/4} + |q|^{-1/2}W^{1/4}.$$

The coefficient sums, the two majorants and their bounded product give

$$\ll (LW^{-1/4}Q^{1/2} + W^{1/4} + L/Q)(\log W)^2 \ll LW^{-1/6}(\log W)^2.$$

For the slow majorant, the derivative is $\asymp |r|W^{-1/4}$, below $1/2$, so Kusmin–Landau gives the needed $O(W^{1/4}/|r|)$ mode bound.

If only the slow wave occurs, pairing consecutive level sets of $\lfloor u^{3/4} \rfloor$ gives $O(LW^{-1/4} + W^{1/4})$. With the extra sign $(-1)^u$, each level set instead has alternating sum $O(1)$, giving the same bound. The lone sign $(-1)^u$ sums to $O(1)$. All arguments retain these majorants on initial subintervals, including their partial end cells. These estimates also apply with the smooth weights (D.4).

D.4 Assembly for the six words

For $w_i = F^i(n)$, the indicator of V_k on odd starts is

$$2^{-2k}(1 + \psi(n^{3/2})) \prod_{i=1}^{k-1} (1 - (-1)^{w_i})(1 + \psi(w_i^{3/2}))(1 + \psi(w_{k-1}^{3/4})). \quad (\text{D.7})$$

For $k = 1$ the product is empty. Every factor records the corresponding next parity after the preceding restrictions, so (D.7) is an exact indicator identity; the iterates w_i are never replaced by a single floor of a power of n .

Expand (D.7). Its constant term gives 2^{-2k} times the number of odd integers in $I_k(m)$. For each other term, select the earliest layer with a remaining sign. Earlier layers sum to the weight (D.4), including its negligible remainder. If that layer has a fast sign and is not the final layer, group by the next-layer integer and apply (D.5). An additional sign $(-1)^u$ is handled by splitting into its two parity classes. All later-layer factors are constant on the grouped fiber and have magnitude at most one. If that layer has only $(-1)^u$, sum alternately within each next-layer fiber; partial summation gives an error at most a constant times the smooth weight per fiber, saving a factor $Z^{-1/4}$ at layer scale Z . If no earlier sign remains, apply Section D.3 on the final layer.

The scale of layer i is $Z_i = P^{(3/4)^i}$, and its interval length has order $Z_i^{1-\rho_k/(3/4)^i}$. The final layer has relative length $5/8$; every preceding layer has a longer relative interval. Thus all applications above meet their length condition. The weakest fast-layer saving is $Z_{k-2}^{-1/8} = P^{-s/6}$, and the final-layer estimate saves $Z_{k-1}^{-1/6} = P^{-s/6}$. For $k = 1$ only Section D.3 is needed. The weight and endpoint errors in D.1 are smaller. There are finitely many terms for each $k \leq 6$, so

$$\#\{n \in I_k(m) : \text{word}_{2k+1}(n) = V_k\} = 2^{-2k} \#\{n \in I_k(m) : n \text{ odd}\} + O_{k,\varepsilon}(|I_k(m)|P^{-s/6+\varepsilon}). \quad (\text{D.8})$$

The location and length of the fiber, together with $s/(6\rho_k) = 4/9$, turn this into the two-sided logarithmic mass estimate

$$\sum_{\substack{n \in I_k(m) \\ \text{word}_{2k+1}(n) = V_k}} \frac{1}{n} = \frac{3^{-k}}{m} \left(1 + O_{k,\varepsilon}(m^{-4/9+\varepsilon})\right). \quad (\text{D.9})$$

For shell sums, discard the bounded number of terminal source integers whose exact fiber meets a shell boundary. The upper endpoint perturbation in (D.3) is smaller than a source step at that endpoint, and the lower endpoint is treated identically. Their reciprocal source mass is exponentially small in the shell logarithm. Summing (D.9) therefore gives the claimed coefficient 3^{-k} at source scale $\rho_k t$, with vanishing errors. Finally the prefix-free words give disjoint source sets, and the subtraction in (5.9) proves (5.10). This completes the analytic input to Theorem 5.3. $\square$

Appendix E. Actual OOE poor-fiber tail

This appendix supplies the written analytic input to Theorem 5.19. It derives the short-interval estimate needed for the actual nested floors, with all four parity guards. It does not invoke Paper B's global estimate as an unproved localization.

E.1 Statement and exact objects

For positive integers n put

$$O(n) = \lfloor n^{3/2} \rfloor, \quad Q(n) = \lfloor \sqrt{n} \rfloor, \quad a(n) = n + (n \bmod 2), \quad w(n) = \log \frac{a(n) + 1}{a(n) - 1}.$$

Let F_m contain exactly those n for which n and $O(n)$ are odd, $O^2(n)$ and $Q(O^2(n))$ are even, and $Q^2(O^2(n)) = m$. Thus F_m is the actual OOE predecessor fibre, with all four parity guards. Write

$$W_m = \sum_{n \in F_m} w(n), \quad R_m = W_m / w(m).$$

Theorem E.1 (poor-fibre tail). For every fixed $0 < \eta < 1/9$,

$$\sum_{m \geq U: R_m < 1/9 - \eta} \frac{1}{m} \ll_{\eta} U^{-7/9} \quad (U \geq 1). \quad (\text{E.1})$$

The implied constant and the onset of the estimates may depend on η . No numerical onset is asserted. The proof also applies to the set where $|R_m - 1/9| > \eta$. It does not claim uniform convergence at every target. The exact empty fibres in the branch dossier are consistent with (E.1).

Set $B(a) = \text{ceil}(a^{2/3})$ and $A_m = B(B(m^4))$. The formal, unguarded preimage of m under $Q^2 \circ O^2$ is exactly

$$I_m = [A_m, A_{m+1}) \cap \mathbb{N}. \quad (\text{E.2})$$

Indeed $O(n) \geq a$ iff $n^3 \geq a^2$, and $Q^2(v) = m$ iff $m^4 \leq v < (m+1)^4$. The ceilings give

$$m^{16/9} \leq A_m \leq m^{16/9} + 2.$$

Consequently, with $P = m^{16/9}$ and H_m the number of odd integers in I_m ,

$$|I_m| = \frac{16}{9} m^{7/9} + O(1), \quad H_m = \frac{8}{9} m^{7/9} + O(1), \quad I_m \subseteq [P, 2P] \quad (\text{E.3})$$

for all sufficiently large m . Interval length here means the difference of its real endpoints; integer counts differ by at most a constant. On odd n in I_m , $w(n) = 2/P$ times $1 + O(1/m)$, uniformly. Also $w(m) = 2/m$ times $1 + O(1/m)$. Writing $C_m = |F_m|$ gives

$$R_m = \frac{8}{9} \frac{C_m}{H_m} + O(m^{-7/9}). \quad (\text{E.4})$$

The larger error $m^{-7/9}$, rather than m^{-1} , retains the integer rounding in H_m . All constants in (E.2)–(E.4) are absolute.

E.2 Classical analytic inputs

Write $e(t) = \exp(2 \cdot \pi \cdot i \cdot t)$. We use the following classical forms on arbitrary real intervals, retaining an absolute endpoint term when necessary:

- If f is monotone and stays at distance at least λ from every integer, the Kusmin–Landau bound is $O(1 + \lambda^{-1})$.
- If $\lambda \leq |f''| \leq C \lambda$, then the second-derivative bound on an interval of length L is $O_C(1 + L \sqrt{\lambda} + \lambda^{-1/2})$.
- For N consecutive terms of modulus at most one and integer $1 \leq D \leq N$, van der Corput differencing bounds the square of their sum by $O(N^2/D + (N/D) \sum_{1 \leq h < D} |T_h|)$, with overlap correlations T_h .

- The one- and three-dimensional Erdős–Turán–Koksma inequalities bound unnormalized interval or box discrepancy by the count divided by K , plus the finite Fourier sums weighted by the product of $1/\max(1, |k_j|)$, for frequencies at most K .

These are the same tools as in Paper B, Sections 3–4. For accessible statements and a proof of the differencing inequality, see Jammes, Section 2. The second-derivative bound is also stated as Theorem E.10 in Bergelson–Richter. The discrepancy reference is Kuipers–Niederreiter, *Uniform Distribution of Sequences*, Chapter 2, already indexed as `kuipers-niederreiter-1974-uniform-distribution`. No qualitative Hardy-field or global-count localization is used.

The finite differencing bullet is now kernel-checked, with constants 2 and 4 and exact overlap length $N-h$, in `WeylDifferencing.lean`. Its exponential-sum specialization works directly on the odd lattice. The proof mapping distinguishes this classical input from its phase-specific application. The first-derivative estimate and the quantitative second-derivative test are now also kernel-checked. The latter has constants 4 and 8 on the unit lattice, and 8 and 4 on the odd lattice. Uniform curvature of the actual carry-cell phases and their unweighted $O(P^{3/8})$ sums are now kernel-checked in `OOEECurvature.lean`, including the growing shift range and explicit floor conditions. The proof mapping retains the extra endpoint required by the derivative test. The mixed-mode proof now supplies the carry partition, weighted sums, Fourier errors, original phase comparison and differencing. Pure slow modes and three-coordinate discrepancy remain separate Lean obligations.

E.3 The short mixed estimate

For real $x \geq 3$ define $X(x) = x^{3/2}$, $Z(x) = x^{9/8}$; for integer n put $M(n) = \text{floor}(X(n))$, $Y(n) = M(n)^{3/2}$.

Lemma E.2 (fixed mixed modes). Fix constants $K, L_0 \geq 1$. For every integer triple (i, j, k) with $\max(|i|, |j|, |k|) \leq K$ and $(i, j) \neq (0, 0)$, and every interval I in $[P, 2P]$ of length at most $L_0 P^{7/16}$,

$$\sum_{n \in I, n \text{ odd}} e\left(\frac{i}{2}X(n) + \frac{j}{2}Y(n) + \frac{k}{2}Z(n)\right) \ll_{K, L_0} P^{13/32}. \quad (\text{E.5})$$

K is fixed independently of P . This lemma expressly excludes the pure slow modes $(0, 0, k)$. No estimate uniform in growing K is asserted.

E.3.1 The carry approximation on a short interval

Let $b(t) = t - 1/2$. For integer $R \geq 2$ set

$$b_R(t) = - \sum_{1 \leq |r| \leq R} \frac{e(rt)}{2\pi i r}, \quad E_R(t) = \min\{1, (R||t||)^{-1}\},$$

with $E_R = 1$ at integers. The ordinary truncated Fourier series satisfies $b = b_R + O(E_R)$, including at integers; Paper B Lemma 4.3 proves this pointwise assertion. We re-estimate its total error at the current length.

For any interval J in $[P, 3P]$ of length at most $L_0 P^{7/16}$, the second-derivative test on the odd lattice gives

$$\left| \sum_{n \in J, n \text{ odd}} e(rX(n)) \right| \ll_{L_0} P^{3/16} |r|^{1/2} + P^{1/4} |r|^{-1/2} + 1. \quad (\text{E.6})$$

Here $x = 2t + 1$ only changes fixed curvature constants. Apply the one-dimensional discrepancy inequality with $R = \text{floor}(P^{1/8})$. Its error is at most

$$O\left(P^{7/16}/R + P^{3/16}R^{1/2} + P^{1/4} + \log(2R)\right) = O(P^{5/16}).$$

Splitting distances to the nearest integer into dyadic ranges starting at $1/R$ therefore gives

$$\sum_{n \in J, n \text{ odd}} E_R(X(n)) \ll_{L_0} P^{5/16} \log P. \quad (\text{E.7})$$

For clarity: at distance at most z there are $O(z P^{7/16} + P^{5/16})$ points. Each dyadic range contributes $O(P^{7/16}/R)$ from its length, while the discrepancy contribution forms a geometric sum. This also counts exact integer values of X . The same bound holds for $X(n+2h)$ by translation of the odd interval. No dyadic exceptional set has been multiplied by a short-interval length ratio.

E.3.2 Differencing, with a bounded number of carry cells

Suppose $j \neq 0$. Conjugating if necessary, take $u = j/2 > 0$; then $1/2 \leq u \leq K/2$. Use odd-lattice shifts $2h$ with $1 \leq h \leq D = \text{floor}(P^{1/16})$. The overlap interval still has length $O_{L_0}(P^{7/16})$. Write

$$\theta = \{X(n)\}, \quad \delta(x) = X(x+2h) - X(x), \quad g(n) = M(n+2h) - M(n).$$

The one-signed linearization, proved by factorization in Paper B Lemma 7.1, is

$$Y(n) = \frac{3}{2}M(n)n^{3/4} - \frac{1}{2}n^{9/4} + E(n), \quad 0 \leq E(n) \ll n^{-3/4}.$$

It gives the exact identity

$$\Delta_h Y(n) = A_h(n) + \frac{3}{2}g(n)(n+2h)^{3/4} - \frac{3}{2}\theta\Delta_h(n^{3/4}) + \Delta_h E(n), \quad (\text{E.8})$$

where $\Delta_h f(x) = f(x+2h) - f(x)$ and

$$A_h(x) = \frac{3}{2}x^{3/2}\Delta_h(x^{3/4}) - \frac{1}{2}\Delta_h(x^{9/4}).$$

Removing the last two terms of (E.8) from the exponential costs at most

$$O_K\left(hP^{7/16-1/4} + P^{7/16-3/4}\right) = O_K(P^{1/4}). \quad (\text{E.9})$$

The function delta is increasing and $\delta' = O(hP^{-1/2})$. Its floor therefore cuts the overlap interval into

$$O_{L_0}(1 + hP^{7/16-1/2}) = O_{L_0}(1) \quad (\text{E.10})$$

cells, including the partial endpoint cells. On each cell freeze $G = \text{floor}(\delta)$ and let $z = \delta - G$. Then $0 \leq z < 1$ is monotone, G is of order $h \sqrt{P}$, and

$$g = G + \kappa, \quad \kappa = z + b(X(n)) - b(X(n+2h)) \in \{0, 1\}. \quad (\text{E.11})$$

Define the smooth cell phases

$$F_{G,\epsilon}(x) = uA_h(x) + \frac{3u}{2}(G + \epsilon)(x+2h)^{3/4} + \frac{i}{2}\delta(x) + \frac{k}{2}\Delta_h Z(x), \quad \epsilon = 0, 1.$$

The exponential remaining after (E.9) equals, exactly,

$$(1-z)e(F_{G,0}) + ze(F_{G,1}) + (b(X(n)) - b(X(n+2h)))(e(F_{G,1}) - e(F_{G,0})). \quad (\text{E.12})$$

This follows by substituting (E.11) into $(1 - \kappa)e(F_{G,0}) + \kappa e(F_{G,1})$. There is no independence assumption.

E.3.3 Every local error term

The identity $A_h(x) = x^{9/4} a(2h/x)$, with $a(t) = (3/2)((1+t)^{3/4} - 1) - (1/2)((1+t)^{9/4} - 1)$, has $a(0) = a'(0) = 0$. Writing $a(t) = t^2 a_2(t)$, whose first two derivatives are bounded near zero, shows $A_h'' = O(h^2 P^{-7/4})$. Thus on each fixed cell

$$F_{G,\epsilon}''(x) = -\frac{9}{32}u(G+\epsilon)(x+2h)^{-5/4} + O_K(h^2 P^{-7/4} + hP^{-3/2} + hP^{-15/8}). \quad (\text{E.13})$$

Its curvature is comparable to $u h P^{-3/4}$, with a fixed sign and bounded ratio, uniformly in $h \leq P^{1/16}$. The second-derivative estimate and partial summation for z and $1-z$ bound the first two terms of (E.12) by

$$O_{K,L_0} \left(P^{7/16} (uh)^{1/2} P^{-3/8} + (uh)^{-1/2} P^{3/8} + 1 \right) = O_{K,L_0}(P^{3/8}). \quad (\text{E.14})$$

The endpoint term $P^{3/8}$ is retained. There are only $O_{L_0}(1)$ cells.

Replace the two b functions in (E.12) by b_R . By (E.7) and $|e(F_1) - e(F_0)| \leq 2$, their total remainder is $O(P^{5/16} \log P)$, even after the partition. Each nonzero Fourier mode now has phase $F_{G,\epsilon} + rX(x)$ or $F_{G,\epsilon} + rX(x+2h)$. Its curvature is comparable to $|r|P^{-1/2}$: the competing curvature divided by that quantity is $O_K(h P^{-1/4}/|r|) = O_K(P^{-3/16})$. Hence the modes, summed with their $O(1/|r|)$ coefficients over $1 \leq |r| \leq R = P^{1/8}$, cost

$$O_{K,L_0} \left(P^{3/16} R^{1/2} + P^{1/4} + \log(2R) \right) = O_{K,L_0}(P^{1/4}). \quad (\text{E.15})$$

Combining (E.9), (E.14), (E.7), and (E.15), every overlap correlation obeys $|T_h| = O_{K,L_0}(P^{3/8})$. In particular, the carry error is smaller than this bound; $\log P$ is absorbed by the gap $3/8 - 5/16 = 1/16$.

If the odd count N in I is at least D , differencing gives

$$|S|^2 \ll_{K,L_0} P^{7/8-1/16} + P^{7/16+3/8} = O_{K,L_0}(P^{13/16}). \quad (\text{E.16})$$

If $N < D$ the trivial estimate is already stronger. This proves (E.5) for $j \neq 0$. If $j = 0$ and $i \neq 0$, the undifferenced smooth phase has curvature comparable to $|i|P^{-1/2}$, since the k term has curvature $O_K(P^{-7/8})$. Its sum is $O_{K,L_0}(P^{1/4})$, which also proves (E.5). This completes Lemma E.2.

E.4 The pure slow modes and the actual last guard

Let $V(n) = \text{floor}(Y(n))$. Uniformly for n in $[P, 2P]$,

$$|\sqrt{V(n)} - n^{9/8}| \ll P^{-3/8}. \quad (\text{E.17})$$

Indeed replacing $\text{sqrt}(\text{floor}(Y))$ by $\text{sqrt}(Y) = M^{3/4}$ costs $O(P^{-9/8})$; replacing $M = X - \text{theta}$ by X in the $3/4$ power costs $O(P^{-3/8})$. The mean value theorem applies on positive intervals bounded below by constant multiples of the relevant powers. Thus replacing Z by $\text{sqrt}(V)$ in a fixed-frequency exponential sum costs $O_K(P^{1/16})$, negligible in (E.5). We use (E.17) for exponential sums, not to assert equality of floor parities.

Put $\alpha_m = (9/8)m^{2/9}$. In the odd coordinate $n = 2t + 1$, the derivative of $(k/2)n^{9/8}$ is $(9k/8)n^{1/8}$. On I_m it differs from $k \alpha_m$ by at most

$$C_0 |k| m^{-7/9} \quad (\text{E.18})$$

for an absolute C_0 . This follows from (E.3) and the derivative bound for $x^{1/8}$, since $P^{-7/8} P^{7/16} = m^{-7/9}$. The derivative is monotone.

For fixed integers $K \geq 1$ and real $C > 2 C_0 K$, suppose

$$\|k\alpha_m\| > Cm^{-7/9} \quad (1 \leq k \leq K). \quad (\text{E.19})$$

For sufficiently large m the derivative of each pure slow phase stays in one interval between consecutive integers and at distance at least $(C/2)m^{-7/9}$ from them. Kusmin–Landau gives

$$\left| \sum_{n \in I_m, n \text{ odd}} e\left(\frac{k}{2}Z(n)\right) \right| \ll m^{7/9}/C + 1 \ll H_m/C + 1. \quad (\text{E.20})$$

Negative k follow by conjugation. Equation (E.17) gives the same bound with an additional $O_K(P^{1/16})$ for the actual $\sqrt{V}$ phase.

E.5 Joint parity, resonance inclusion, and the tail

Apply three-dimensional box discrepancy to the actual points

$$\left(\{X(n)/2\}, \{Y(n)/2\}, \{\sqrt{V(n)}/2\} \right), \quad n \in I_m, n \text{ odd}.$$

The box $[1/2, 1)$ times $[0, 1/2)$ times $[0, 1/2)$ specifies exactly the three remaining OOE guards. Its volume is $1/8$. Half-open boundaries encode integer-floor parity exactly, including square and cube coincidences.

For fixed K and C as above and m satisfying (E.19), Lemma E.2 controls every mode with $(i, j) \neq (0, 0)$, and (E.20) controls the pure slow modes. Therefore

$$|C_m/H_m - 1/8| \leq A/K + A \log(2K)/C + O_{K, L_0}(P^{-1/32}) + O_K(P^{-3/8}) + O_K(H_m^{-1}), \quad (\text{E.21})$$

where A is absolute; fixed logarithmic frequency weights are absorbed in the indicated constants. For any $\epsilon > 0$, first choose K so the first term is less than $\epsilon/3$; then choose $C > 2 C_0 K$ so the second is less than $\epsilon/3$; finally choose m large so the remaining terms total less than $\epsilon/3$. This order keeps K and C independent of m .

Together with (E.4), this proves: for every $\eta > 0$ there are fixed K_η, C_η and m_η such that

$$m \geq m_\eta, |R_m - 1/9| > \eta \implies \exists 1 \leq q \leq K_\eta : \|q\alpha_m\| \leq C_\eta m^{-7/9}. \quad (\text{E.22})$$

For example, take $\epsilon = \eta/4$ and enlarge m_η until the error in (E.4) is at most $\eta/4$. Strict poor inequalities $R_m < 1/9 - \eta$ are contained in this absolute-deviation set.

For fixed q , the derivative of $q\alpha_m$ is $(q/4)m^{-7/9}$. On $[M, 2M)$, each window within $C_\eta M^{-7/9}$ of an integer has preimage length at most $8 C_\eta 2^{7/9}/q$. There are $O(q M^{2/9} + C_\eta + 1)$ possible integer levels. Counting integers in these bounded intervals, then summing over $q \leq K_\eta$, gives

$$\#\{M \leq m < 2M : m \text{ satisfies the right side of (E.22)}\} \ll_\eta M^{2/9}. \quad (\text{E.23})$$

The same bound for deficient targets follows beyond m_η ; increasing its constant absorbs the finite initial set. Division by M and summation over dyadic blocks starting at U yields (E.1). This proves Theorem E.1.

E.6 What the coefficient now supplies

For any subset A of the positive integers, any finite target cutoff X , and fixed $0 < \eta < 1/9$, Theorem E.1 and $w(m) \leq 4/m$ give

$$\sum_{m \in A, m \leq X} W_m \geq (1/9 - \eta) \sum_{m \in A, m \leq X} w(m) - C_\eta. \quad (\text{E.24})$$

The constant is independent of A and X . Separate the poor targets and use their bounded total w -mass; all W_m are nonnegative. Fibres for different targets are disjoint. If A is backward closed under the actual Juggler map, every source in these fibres also belongs to A .

The physical source cutoffs and three-production recursion are proved in Section 5.9. This analytic input is a written proof pending independent review. The main mixed-mode estimate has a separate Lean proof; the complete poor-target and production conclusions are not yet certified by that formalization.

Acknowledgments and use of AI

Large language models assisted the development of this work, including the prose, proposed proof arguments, the Lean formalizations, and the computations. AI assistance is not independent mathematical validation. The formalization covers only the results identified in Section 1.4 and Appendix A. The author is responsible for the statements, proofs, code, and the decision to make this version public.

Availability and version

This is version 1.2.0 of Paper C, of 22 September 2026. It is a preprint, has not been refereed, and is not deposited. It adds the written OOE poor-fiber proof, contagion at $5/8$, the sufficient rate threshold $3/8$, and an updated proof-status map. The fully machine-checked baseline remains $100/203$. It also retains the corrected distinction between finite and unbounded stopping-word moments in Section 5.7. The current Zenodo version is 1.1.0 (doi:10.5281/zenodo.22865705) of 21 September 2026, which added Section 5.8 with its Lean layer, the corresponding rows of Section 1.4 and Appendices A and B, the remarks of Sections 5.7, 6.3 and 7.1, and the exponent λ_{ideal} in the statements; version 1.0.0 (doi:10.5281/zenodo.22678165) is of 9 September 2026. The concept DOI 10.5281/zenodo.22678164 resolves to the latest version, and the record is at <https://zenodo.org/records/22865705>. The author's ORCID is 0009-0004-1939-3382. A later revision goes up through the record's new-version operation, which keeps the concept DOI, rather than as a new deposit. The changes between versions are listed in the build guide, `PAPER_C_BUILD.md`, in the repository. The accompanying source package contains the Markdown and LaTeX sources, figures, the local Lean dependency closure with its pinned toolchain, numerical audit code, archived experiment records, and a manifest of file hashes. The repository is `sneakyweasel/btlab`. The original development notes remain historical records; where their statements differ, the corrected proofs and scopes in this version of Paper C govern this preprint.

References

1. C. A. Pickover, *Computers and the Imagination: Visual Adventures Beyond the Edge*, St. Martin's Press, New York, 1991, ch. 40.
2. OEIS Foundation Inc., "Juggler sequence: if n even then $\lfloor \sqrt{n} \rfloor$ else $\lfloor n^{3/2} \rfloor$," Sequence A094683 in *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org/A094683> (accessed 9 September 2026).
3. J. C. Lagarias, "The $3x + 1$ problem and its generalizations," *Amer. Math. Monthly* 92 (1985), 3–23. doi:10.1080/00029890.1985.11971528.
4. R. Terras, "A stopping time problem on the positive integers," *Acta Arith.* 30 (1976), 241–252. doi:10.4064/aa-30-3-241-252.
5. C. J. Everett, "Iteration of the number-theoretic function $f(2n) = n$, $f(2n + 1) = 3n + 2$," *Adv. Math.* 25 (1977), 42–45. doi:10.1016/0001-8708(77)90087-1.
6. T. Tao, "Almost all orbits of the Collatz map attain almost bounded values," *Forum Math. Pi* 10 (2022), e12. doi:10.1017/fmp.2022.8.

7. I. Krasikov and J. C. Lagarias, “Bounds for the $3x + 1$ problem using difference inequalities,” *Acta Arith.* 109 (2003), 237–258. doi:10.4064/aa109-3-4.
8. J. C. Lagarias (ed.), *The Ultimate Challenge: The $3x+1$ Problem*, American Mathematical Society, Providence, RI, 2010. doi:10.1090/mbk/078.
9. E. W. Weisstein, “Juggler Sequence,” *MathWorld — A Wolfram Web Resource*, <https://mathworld.wolfram.com/JugglerSequence.html>.
10. J. D. Vaaler, “Some extremal functions in Fourier analysis,” *Bull. Amer. Math. Soc. (N.S.)* 12 (1985), 183–216. doi:10.1090/S0273-0979-1985-15349-2.
11. P. Cochin, “Lower Bounds for Cycle Lengths in the Juggler Map” (Paper A), Zenodo version 1.0.2, 20 September 2026, doi:10.5281/zenodo.22865237; concept DOI 10.5281/zenodo.22676452 for all versions; revision of 21 September 2026 at docs/theory/juggler_finite_dynamics_note.md in the repository <https://github.com/sneakyweasel/btlab/>.
12. P. Cochin, “Five-Step Descent Certificates for the Juggler Map: Parity Statistics of Nested Floor Powers” (Paper B), Zenodo version 1.0.0, 21 September 2026, doi:10.5281/zenodo.22864934; concept DOI 10.5281/zenodo.22864933 for all versions; revision of 20 September 2026 at docs/theory/juggler_parity_discrepancy_note.md, same repository.
13. S. W. Graham and G. Kolesnik, *Van der Corput’s Method of Exponential Sums*, London Mathematical Society Lecture Note Series 126, Cambridge University Press, Cambridge, 1991. doi:10.1017/CBO9780511661976.
14. P. Cochin, research notes on fate contagion, the Tao-type reduction and the poor-fiber tail for the Juggler map, 2026; docs/theory/juggler_fate_contagion_note.md, docs/theory/juggler_tao_reduction_note.md and docs/theory/juggler_oe_poor_fiber_tail_note.md, same repository.
15. L. Kuipers and H. Niederreiter, *Uniform Distribution of Sequences*, Wiley-Interscience, New York, 1974.