

# Five-Step Descent Certificates for the Juggler Map

## Parity Statistics of Nested Floor Powers

Philippe Cochin

philippe@cochin.fr

20 September 2026

Preprint

### Abstract

The Juggler map applies the integer part of the square root at even positive integers and of the three-halves power at odd positive integers. We prove that the starting values admitting a power-envelope descent certificate within five operations have natural density  $7/8$ . More precisely, their count up to  $N$  is  $7N/8 + O_\varepsilon(N^{127/128+\varepsilon})$  for every  $\varepsilon > 0$ . The four-step subfamily has density  $13/16$ . The proof uses exact carry identities, centered Fourier expansions, and van der Corput differencing, with all floor exceptions and partition endpoints counted. The two five-letter classes *OOEOE* and *OOOEE* are treated through their distinct formal chains. For *OOOEE*, the full mixed phase is retained and its signed curvature is recomputed after each frequency center is frozen. The complete analytic argument is included in three appendices. Fair-share densities at every fixed depth would imply density-one finite certificates; that hypothesis, general decorated estimates, and short-interval localization remain open. No result asserts universal arrival at 1.

**Keywords:** Juggler map; nested floor powers; parity correlations; discrepancy; dropping time; finite-step descent.

### 1. Scope and relation to earlier work

For a positive integer  $n$ , define

$$J(n) = \begin{cases} \lfloor n^{1/2} \rfloor, & n \text{ even,} \\ \lfloor n^{3/2} \rfloor, & n \text{ odd.} \end{cases}$$

This is the Juggler map recorded in OEIS A094683 [1]. The conjecture that every positive orbit reaches 1 remains open. The question studied here is more limited: which distribution estimates for finite itineraries would imply that many starting values eventually fall below their initial value?

Single-floor parity estimates follow from classical exponential-sum methods [2, 3]. Composing floors creates an additional difficulty. For  $m = \lfloor n^{3/2} \rfloor$  and  $\theta = \{n^{3/2}\}$ ,

$$m^{3/2} = n^{9/4} - \frac{3}{2}\theta n^{3/4} + O(n^{-3/4}).$$

The coefficient of the fractional part grows with  $n$ . A theorem for the smooth phase  $n^{9/4}$  therefore does not by itself estimate the nested phase. Nor does a parity estimate over consecutive starting values automatically apply to a sparse set of orbit images.

The main result is the five-step certificate count in Theorem 5.4. The proof distinguishes finite word classification from the analytic estimates needed to count those words. Sections 4.2–4.3 establish the four-step estimate and the *OOEOE* split. Section 4.4 and Appendices A–C establish the remaining *OOOEE* mixed-mode bound. All signs, zero coordinates, Fourier coefficient weights, and interval boundaries are included.

The single-floor estimate and the conditional all-depth counting argument use standard methods. The latter is in the spirit of stopping-time arguments for the Collatz map, such as Terras [4]. The general 95/96 kernel target and arbitrary short-interval extensions from earlier drafts remain unproved. The weaker 127/128 exponent proved here suffices for the five-step count.

That kinship is an identity rather than an analogy, and stating it precisely is what fixes the contribution of this paper. The shortcut Collatz map sends  $x$  to  $x/2$  or  $(3x+1)/2$ , so  $\log x$  moves by  $-\log 2$  or  $+\log(3/2)$ ; the Juggler map multiplies  $\log n$  by  $1/2$  or  $3/2$ , so  $\log \log n$  moves by the same two steps. A word therefore contracts in one problem exactly when it contracts in the other, and the word counts used here are Collatz word counts. The number of words of length  $d$  with no contracting prefix is A076227 [5], in the OEIS since 2002 as the number of residue classes modulo  $2^d$  on which the Collatz *dropping time* is not constant; that entry’s own comment names A074473, which the OEIS titles the dropping time. The minimal certificates, counted by length, are the admissible sequences of A100982 [6], supported on the binary lengths of the powers of three, A020914 [7] – which that entry has recorded since 2009 as the set of allowable dropping times of the shortcut map, the value for an individual start being A126241. The length of a minimal certificate is a dropping time; computational work calls the same quantity the *glide*, and Hikawa’s toolkit [15], which prints it under that name, computes the survivor counts, the minimal-certificate counts and their refinement by odd count, all three agreeing with the counts used here.

A word on the term, since two conventions collide here and this paper had declared neither. Terras [4], whom it follows elsewhere, calls the first time an orbit falls below its start the *stopping time*; the OEIS calls that the *dropping time* and reserves *stopping time* for the number of steps needed to reach 1, a different quantity that plays no role below. The object is the same under either name, and it is the one used throughout. The list of Lemma 5.1 and the surviving words of Theorem 6.1 are those objects; no priority is claimed for them, and the counting identities are recorded and machine-checked separately in the laboratory repository rather than proved here.

The two problems part at the arithmetic, and that is the whole of the present work. For Collatz the word of length  $d$  is a function of  $x \bmod 2^d$ , and the induced map to words is a bijection [4], so every word has density exactly  $2^{-d}$  and the fair-share counts are free. For the Juggler map the word is a function of  $\{n^{3/2}\}$  and of the nested fractional parts that follow it, and the corresponding statement is Hypothesis FD of Section 6, which is open. Theorems 3.1, 5.2, 5.3 and 5.4 are statements of that second kind and have no Collatz counterpart. No theorem about Collatz is transferred to the Juggler map.

This preprint supersedes the earlier conditional and 27/32 versions. The proofs have been developed and checked with AI assistance and exact-arithmetic controls. No independent mathematical review or complete Lean verification is claimed.

The proof can be read in the following order.

| Part                       | Role                                                         |
|----------------------------|--------------------------------------------------------------|
| Sections 2–3               | Formal words, exact certificates, and single-floor counts    |
| Sections 4.2–4.3           | Four-step estimate and the <i>OOEOE</i> split                |
| Section 4.4 and Appendix C | The precise <i>OOOEE</i> mixed-mode theorem                  |
| Appendix A                 | Floor exceptions, positive Fourier errors, and exact carries |
| Appendix B                 | The wave-bearing estimate used in Appendix C                 |

| Part         | Role                                                        |
|--------------|-------------------------------------------------------------|
| Sections 5–6 | Five-step density and the conditional all-depth implication |

## 2. Exact itineraries and the power envelope

Throughout,  $e(t) = \exp(2\pi it)$ ,  $\{t\} = t - \lfloor t \rfloor$ , and  $\|t\|$  is the distance from  $t$  to the nearest integer. Define

$$\psi(t) = (-1)^{\lfloor t \rfloor} = 1 - 2\mathbf{1}_{[1/2, 1)}(\{t/2\}).$$

Thus  $\psi$  equals  $+1$  at an even floor and  $-1$  at an odd floor, including at integer endpoints. Implied constants are independent of the scale and of modes in the stated ranges; fixed comparison constants and any displayed  $\varepsilon$  may enter them.

Let  $w = w_1 \cdots w_d$  be a word in  $\{E, O\}$ . The itinerary  $\text{word}_d(n)$  records the parities of  $n, J(n), \dots, J^{d-1}(n)$ . It contains  $d$  letters and specifies the  $d$  operations that produce  $J^d(n)$ . Write  $o(w)$  for the number of its odd letters.

**Proposition 2.1 (power envelope).** If  $w$  is realized at  $n$ , then

$$(J^d(n))^{2^d} \leq n^{3^{o(w)}}.$$

Consequently, if  $n \geq 2$  and  $3^{o(w)} < 2^d$ , then  $J^d(n) < n$ .

*Proof.* At any positive integer  $x$ , the even operation satisfies  $J(x)^2 \leq x$ , and the odd operation satisfies  $J(x)^2 \leq x^3$ . Suppose a prefix of length  $r$  with  $o$  odd letters ends at  $x$  and obeys  $x^{2^r} \leq n^{3^o}$ . An even operation preserves the right-hand exponent after raising  $J(x)^2 \leq x$  to  $2^r$ ; an odd operation replaces it by  $3^{o+1}$ . Induction starts at the empty prefix. If  $3^{o(w)} < 2^d$  and  $n > 1$ , the bound is strictly less than  $n^{2^d}$ , proving descent.  $\square$

A word satisfying  $3^{o(w)} < 2^{\lfloor w \rfloor}$  is called *contracting*. A realized contracting prefix is a *power-envelope descent certificate*. Let

$$\mathcal{C}_d = \{n \geq 2 : \text{some contracting prefix of length at most } d \text{ is realized at } n\}.$$

Also set

$$\mathcal{C}_\infty = \bigcup_{d \geq 1} \mathcal{C}_d.$$

These sets refer to this particular sufficient certificate. Flooring can cause descent even when the envelope does not certify it. Therefore  $\mathcal{C}_d$  need not equal the set of all starts that descend within  $d$  steps.

For an odd-rooted target word  $w$ , define its *formal chain* for every odd  $n$ , regardless of whether  $n$  realizes  $w$ :

$$x_1^w(n) = n, \quad \xi_t^w(n) = (x_t^w(n))^{p_t}, \quad x_{t+1}^w(n) = \lfloor \xi_t^w(n) \rfloor \quad (1 \leq t < d),$$

where  $p_t = 3/2$  for  $w_t = O$  and  $p_t = 1/2$  for  $w_t = E$ . Set  $s_t^w(n) = \psi(\xi_t^w(n))$ , and put  $\epsilon_t = 1$  for  $w_{t+1} = E$ ,  $\epsilon_t = -1$  for  $w_{t+1} = O$ .

**Lemma 2.2 (parity and branch indicators).** For real  $x$ ,  $\lfloor x \rfloor$  is odd exactly when  $\{x/2\} \in [1/2, 1)$ . For every odd  $n$  and odd-rooted word  $w$ ,

$$\mathbf{1}_{\text{word}_d(n)=w} = 2^{-(d-1)} \prod_{t=1}^{d-1} (1 + \epsilon_t s_t^w(n)). \quad (2.1)$$

*Proof.* Writing  $x = \lfloor x \rfloor + \{x\}$  proves the first assertion, including the half-open endpoints. Each factor divided by two tests the parity of  $x_{t+1}^w$ . Until the first failed test, the formal chain agrees with the true orbit, by induction on  $t$ . A failed test makes the product zero and prevents realization of  $w$ . If all tests pass, the chains agree through the required letters and the product is one. The empty product covers  $d = 1$ .  $\square$

The use of formal chains is essential: the product is defined on every odd input before any correlation or Fourier expansion is made.

### 3. An unconditional certificate density

Let  $M(N) = \#\{n \leq N : n \text{ odd}\} = N/2 + O(1)$ , and set

$$S_O(N) = \sum_{\substack{n \leq N \\ n \text{ odd}}} \psi(n^{3/2}).$$

**Theorem 3.1 (single-floor parity).** One has

$$S_O(N) = O(N^{5/6}).$$

In particular,

$$\begin{aligned} \#\{n \leq N : \text{word}_2(n) = OO\} &= N/4 + O(N^{5/6}), \\ \#(\mathcal{C}_2 \cap [1, N]) &= 3N/4 + O(N^{5/6}). \end{aligned}$$

*Proof.* Write  $n = 2r + 1$  and  $g(r) = \frac{1}{2}(2r + 1)^{3/2}$ . Then  $g''(r) = \frac{3}{2}(2r + 1)^{-1/2}$ . On a dyadic block with  $r \asymp Q$ , the classical second-derivative test [2] gives

$$\left| \sum_{r \asymp Q} e(hg(r)) \right| \ll h^{1/2} Q^{3/4} + h^{-1/2} Q^{1/4} \quad (h \geq 1).$$

The Erdős–Turán inequality [3, Chapter 2] bounds the unnormalized interval discrepancy by

$$\ll Q/H + \sum_{h=1}^H \frac{1}{h} \left| \sum_{r \asymp Q} e(hg(r)) \right| \ll Q/H + H^{1/2} Q^{3/4} + Q^{1/4}.$$

Taking  $H = \lfloor Q^{1/6} \rfloor$  for sufficiently large  $Q$  gives  $O(Q^{5/6})$ . Lemma 2.2 identifies parity with membership in  $[1/2, 1)$ ; summing the dyadic bounds proves the first assertion. Since  $S_O = M - 2\#OO$ , it also proves the count of  $OO$ .

The only minimal contracting words of length at most two are  $E$  and  $OE$ . The first class has count  $\lfloor N/2 \rfloor$ , and the second has count  $M - \#OO = N/4 + O(N^{5/6})$ . They are disjoint, and Proposition 2.1 proves the stated certificate count.  $\square$

This argument estimates starting values in an interval. It makes no independence assertion about successive orbit parities.

**Proposition 3.2 (the OE third letter).** One has

$$\#\{n \leq N : \text{word}_3(n) = w\} = N/8 + O(N^{5/6} \log(2N)), \quad w \in \{OEE, OEO\}.$$

*Proof.* For every real  $x \geq 0$ ,

$$\lfloor \sqrt{\lfloor x \rfloor} \rfloor = \lfloor \sqrt{x} \rfloor :$$

for an integer  $r \geq 0$ , the inequalities  $r^2 \leq \lfloor x \rfloor$  and  $r^2 \leq x$  are equivalent. Consequently, with  $m = \lfloor n^{3/2} \rfloor$ , one has  $\psi(m^{1/2}) = \psi(n^{3/4})$  exactly.

On a dyadic block apply the two-dimensional Erdős–Turán–Koksma inequality to  $(\{n^{3/2}/2\}, \{n^{3/4}/2\})$  at odd  $n$ , with cutoff  $H = \lfloor P^{1/6} \rfloor$ . For a nonzero integer mode  $(i, l)$  in this range, the second-derivative estimate gives

$$\left| \sum e\left(\frac{i}{2}n^{3/2} + \frac{l}{2}n^{3/4}\right) \right| \ll \begin{cases} |i|^{1/2}P^{3/4} + |l|^{-1/2}P^{1/4}, & i \neq 0, \\ |l|^{1/2}P^{3/8} + |l|^{-1/2}P^{5/8}, & i = 0. \end{cases}$$

For  $i \neq 0$ , dominance follows from  $|l/i|P^{-3/4} \leq P^{-7/12}$ . Using the product weights of Proposition 4.2, the discrepancy is

$$O(P/H + H^{1/2}P^{3/4} \log(2H) + P^{5/8} \log(2H)),$$

hence  $O(P^{5/6} \log P)$ . The two boxes selecting even  $m$  and either parity of  $\lfloor n^{3/4} \rfloor$  have area  $1/4$ . Each therefore counts  $P/8$  odd starts on the block, with the stated error. Sum dyadically.  $\square$

## 4. Correlation criteria and proved cases

For a fixed odd-rooted word  $w$  of length  $d$  and nonempty  $A \subseteq \{1, \dots, d-1\}$ , define

$$R_{w,A}(N) = \sum_{\substack{n \leq N \\ n \text{ odd}}} \prod_{t \in A} s_t^w(n).$$

**Hypothesis  $H(w; \delta)$ .** For some fixed  $0 < \delta < 1$ , every nonempty  $A$  satisfies  $R_{w,A}(N) = O_{w,A}(N^{1-\delta})$  as  $N \rightarrow \infty$ .

**Hypothesis  $H_0(w)$ .** The same correlations satisfy  $R_{w,A}(N) = o(N)$ , without a prescribed rate.

These are assertions about the formal chains, not assumptions that the true orbit is a sequence of independent coin tosses. Hypothesis  $H(w; \delta)$  implies  $H_0(w)$ .

**Proposition 4.1 (conditional word count).** Under  $H(w; \delta)$ ,

$$\#\{n \leq N : \text{word}_d(n) = w\} = 2^{-d}N + O_w(N^{1-\delta}). \quad (4.1)$$

Under  $H_0(w)$ , the error is  $o(N)$ .

*Proof.* Expand (2.1) and sum. The empty subset contributes  $2^{-(d-1)}M(N) = 2^{-d}N + O(1)$ . The other  $2^{d-1} - 1$  terms are fixed signed multiples of  $R_{w,A}$ , and there are finitely many for fixed  $w$ .  $\square$

For example,  $H(OOEE; \delta)$  concerns the seven nonempty products of

$$\psi(n^{3/2}), \quad \psi(m^{3/2}), \quad \psi(v^{1/2}), \quad m = \lfloor n^{3/2} \rfloor, \quad v = \lfloor m^{3/2} \rfloor.$$

Proposition 4.1 then gives the count  $N/16 + O(N^{1-\delta})$  needed for four-step certificates. Theorem 3.1 establishes the first estimate. Corollary 4.6 below establishes all seven for every  $0 < \delta < 1/24$ .

#### 4.1 A sufficient Fourier criterion

For  $\mathbf{k} = (k_1, \dots, k_{d-1}) \in \mathbb{Z}^{d-1}$ , let

$$S_{w,\mathbf{k}}(N) = \sum_{\substack{n \leq N \\ n \text{ odd}}} e \left( \frac{1}{2} \sum_{t=1}^{d-1} k_t \xi_t^w(n) \right).$$

**Proposition 4.2 (conditional Fourier-to-parity transfer).** Fix  $d \geq 2$ ,  $w$ , and  $0 < \eta, \sigma < 1$ . Suppose, uniformly over all nonzero integer vectors with  $\|\mathbf{k}\|_\infty \leq \lfloor N^\eta \rfloor$ , that

$$|S_{w,\mathbf{k}}(N)| \ll_w N^{1-\sigma}. \quad (4.2)$$

Then  $H(w; \delta)$  holds for every  $0 < \delta < \min(\eta, \sigma)$ . The constants may depend on these fixed parameters.

*Proof.* Apply the multidimensional Erdős–Turán–Koksma inequality [3, Chapter 2, p. 116] to the points  $\mathbf{y}_n = (\{\xi_1^w(n)/2\}, \dots, \{\xi_{d-1}^w(n)/2\})$ . For any half-open axis-parallel box  $B \subseteq [0, 1)^{d-1}$ , its unnormalized discrepancy is

$$\begin{aligned} |\#\{n \leq N \text{ odd} : \mathbf{y}_n \in B\} - M(N)|B|| &\ll_d \frac{M(N)}{H} + \sum_{0 < \|\mathbf{k}\|_\infty \leq H} \frac{|S_{w,\mathbf{k}}(N)|}{r(\mathbf{k})}, \\ r(\mathbf{k}) &= \prod_{t=1}^{d-1} \max(1, |k_t|). \end{aligned}$$

The weight sum is  $O_d((1 + \log H)^{d-1})$ . With  $H = \lfloor N^\eta \rfloor$ , (4.2) gives box discrepancy  $O_w(N^{1-\eta} + N^{1-\sigma}(1 + \log N)^{d-1})$ . Each nonempty sign product is constant, with value 1 or  $-1$ , on the  $2^{d-1}$  boxes obtained by splitting each coordinate at  $1/2$ . Its integral over the unit cube is zero. Summing their discrepancies therefore bounds  $R_{w,A}$ ; absorbing the fixed logarithmic power proves the assertion.  $\square$

The criterion needs every indicated mixed mode, including vectors with zero coordinates and mixed signs. A bound for one undecorated kernel is not a substitute for (4.2). Qualitatively, cancellation  $S_{w,\mathbf{k}}(N) = o(N)$  for each fixed nonzero  $\mathbf{k}$  also implies  $H_0(w)$ : keep  $H$  fixed in the displayed inequality, let  $N \rightarrow \infty$ , and then let  $H \rightarrow \infty$ .

#### 4.2 A nested estimate sufficient for four-step descent

The general decorated kernel is not needed for four-step certificates. We prove the smaller mixed family directly. Constants in this subsection may depend on a fixed  $C \geq 1$ , but are uniform in all integer modes in their stated ranges.

**Lemma 4.3 (a truncated carry expansion).** Put

$$b(t) = \{t\} - \frac{1}{2}, \quad b_R(t) = - \sum_{1 \leq |r| \leq R} \frac{e(rt)}{2\pi i r}, \quad E_R(t) = \min \left( 1, \frac{1}{R\|t\|} \right),$$

where  $\|t\|$  denotes distance to the nearest integer,  $E_R(t) = 1$  at integers, and  $R \geq 2$  is integral. Then  $b(t) = b_R(t) + O(E_R(t))$ , including at integers. For  $X(x) = x^{3/2}$ , any interval  $I \subseteq [P, 3P]$  satisfies

$$\sum_{\substack{n \in I \\ n \text{ odd}}} E_R(X(n)) \ll \frac{P \log(2R)}{R} + P^{5/6}. \quad (4.3)$$

*Proof.* Away from integers, the Fourier series of  $b$  is the displayed series with infinite range. Summation by parts bounds its tail by  $O((R\|t\|)^{-1})$ . If  $\|t\| \leq R^{-1}$ , pairing the modes  $r$  and  $-r$

and using  $|\sin(2\pi r t)| \leq 2\pi r \|t\|$  bounds the finite sum by an absolute constant. At an integer the finite sum is zero and  $b(t) = -1/2$ , so the asserted bound remains valid.

The proof of Theorem 3.1, with phase  $(2r+1)^{3/2}$ , gives interval discrepancy  $O(P^{5/6})$  for the fractional parts  $\{X(n)\}$  with odd  $n \in I$ . In particular the number within distance  $z \leq 1/2$  of an integer is  $O(Pz + P^{5/6})$ . Split these distances at  $R^{-1}, 2R^{-1}, 4R^{-1}, \dots$ . Each resulting weighted count contributes  $O(P/R + 2^{-j}P^{5/6})$ , proving (4.3). This is a bound in terms of the ambient scale  $P$ ; it does not rescale a global exceptional set by the length of  $I$ .  $\square$

**Lemma 4.4 (small-shift nested sum).** Write  $m(n) = \lfloor n^{3/2} \rfloor$ ,  $Y(n) = m(n)^{3/2}$ , and  $\Delta_h f(n) = f(n+2h) - f(n)$ . If

$$1 \leq h \leq P^{1/12}, \quad 1 \leq |j| \leq CP^{1/24}, \quad |i|, |k| \leq CP^{1/24},$$

where  $i, j, k, h$  are integers, then

$$\left| \sum_{\substack{P < n \leq 2P-2h \\ n \text{ odd}}} e\left(\frac{i}{2}\Delta_h X(n) + \frac{j}{2}\Delta_h Y(n) + \frac{k}{2}\Delta_h(n^{9/8})\right) \right| \ll_C P^{7/8}(1 + h^{1/2}). \quad (4.4)$$

*Proof.* Conjugate the whole sum if necessary and put  $u = j/2 > 0$ . Thus  $u \geq 1/2$ ; integrality of  $u$  is not required. Set  $\theta = \{X(n)\}$ ,  $\delta = \Delta_h X(n)$ , and  $g = m(n+2h) - m(n)$ . Lemma 7.1 gives the exact identity

$$\Delta_h Y = A_h + \frac{3}{2}g(n+2h)^{3/4} - \frac{3}{2}\theta\Delta_h(n^{3/4}) + \Delta_h E, \quad (4.5)$$

where

$$A_h(x) = \frac{3}{2}x^{3/2}\Delta_h(x^{3/4}) - \frac{1}{2}\Delta_h(x^{9/4}), \quad A_h''(x) = O(h^2P^{-7/4}).$$

For the derivative assertion, write  $A_h(x) = x^{9/4}a(2h/x)$ , where  $a(z) = \frac{3}{2}((1+z)^{3/4} - 1) - \frac{1}{2}((1+z)^{9/4} - 1)$ . The identities  $a(0) = a'(0) = 0$  and bounded derivatives of  $a(z)/z^2$  near zero give the estimate. No floor is differentiated.

Deleting the last two terms of (4.5) from the phase costs

$$O(uhP^{3/4} + uP^{1/4}), \quad (4.6)$$

because  $\Delta_h(x^{3/4}) = O(hP^{-1/4})$ ,  $E(x) = O(P^{-3/4})$ , and  $|e(t) - 1| \leq 2\pi|t|$ .

Partition the interval into level sets of  $G = \lfloor \delta \rfloor$ . Since  $\delta'(x) \asymp hP^{-1/2}$ , there are  $O(1 + hP^{1/2})$  cells; their total length is at most  $P$ . On a cell put  $z = \delta - G$ . It is monotone in  $[0, 1)$ , and exactly

$$g = G + \kappa, \quad \kappa = z + b(X(n)) - b(X(n+2h)) \in \{0, 1\}. \quad (4.7)$$

Define the two smooth phases on this cell by

$$F_{G,\epsilon}(x) = uA_h(x) + \frac{3u}{2}(G + \epsilon)(x+2h)^{3/4} + \frac{i}{2}\delta(x) + \frac{k}{2}\Delta_h(x^{9/8}), \quad \epsilon \in \{0, 1\}.$$

The exponential left after (4.6) is exactly

$$(1-z)e(F_{G,0}) + ze(F_{G,1}) + (b(X(n)) - b(X(n+2h)))(e(F_{G,1}) - e(F_{G,0})). \quad (4.8)$$

Apply Lemma 4.3 with  $R = \lfloor P^{1/4} \rfloor$  to the two sawtooths. Their total error after the partition is  $O(P^{5/6} + P^{3/4} \log P)$ : the cells partition the same input set, and the multiplying difference of exponentials has modulus at most two.

On every cell  $G + \epsilon \asymp hP^{1/2}$ , and

$$F_{G,\epsilon}''(x) = -\frac{9}{32}u(G + \epsilon)(x+2h)^{-5/4} + O(uh^2P^{-7/4} + |i|hP^{-3/2} + |k|hP^{-15/8}). \quad (4.9)$$

The error is  $o(uhP^{-3/4})$ , uniformly in the stated ranges. The second-derivative estimate [2, Theorem 2.2] has scale  $\lambda \asymp uhP^{-3/4}$  and a bounded curvature ratio. After  $n = 2r + 1$  it has the same form for odd integers, with implicit constants. Summing over all cells, and using partial summation for the monotone weights  $z$  and  $1 - z$ , gives

$$\ll (uh)^{1/2}P^{5/8} + (h/u)^{1/2}P^{7/8}. \quad (4.10)$$

Each nonzero Fourier mode adds either  $rX(x)$  or  $rX(x + 2h)$  to one of the phases  $F_{G,\epsilon}$ . Its curvature has size  $|r|P^{-1/2}$ , which dominates (4.9), since

$$\frac{uhP^{-3/4}}{|r|P^{-1/2}} \ll_C P^{-1/8} \quad (1 \leq |r| \leq R).$$

There is no curvature-collision regime in this proof. Summing the second-derivative estimate over all gap cells costs, per mode,

$$\ll |r|^{1/2}P^{3/4} + h|r|^{-1/2}P^{3/4}.$$

The Fourier weights are  $O(1/|r|)$ , so their total is  $O(R^{1/2}P^{3/4} + hP^{3/4})$ . Collecting the costs gives

$$\begin{aligned} &\ll_C (uh)^{1/2}P^{5/8} + (h/u)^{1/2}P^{7/8} + P^{7/8} + hP^{3/4} \\ &\quad + uhP^{3/4} + uP^{1/4} + P^{5/6} + P^{3/4} \log P. \end{aligned}$$

Here  $uh \ll_C P^{1/8}$  and  $u \geq 1/2$ . Every term is  $O_C(P^{7/8}(1 + h^{1/2}))$ . Endpoint cells are included; no estimate is extended across their boundaries.  $\square$

**Theorem 4.5 (restricted mixed exponential sums).** For every fixed  $C \geq 1$ , uniformly over nonzero integer triples  $(i, j, k)$  with  $\max(|i|, |j|, |k|) \leq CP^{1/24}$ , one has

$$\left| \sum_{\substack{P < n \leq 2P \\ n \text{ odd}}} e\left(\frac{i}{2}n^{3/2} + \frac{j}{2}m(n)^{3/2} + \frac{k}{2}n^{9/8}\right) \right| \ll_C P^{23/24}. \quad (4.11)$$

*Proof.* If  $j \neq 0$ , apply the van der Corput differencing inequality [2] to the sequence indexed by odd integers, with  $H = \lfloor P^{1/12} \rfloor$ . Writing  $T_h$  for (4.4), it gives

$$|S|^2 \ll \frac{P^2}{H} + \frac{P}{H} \sum_{1 \leq h < H} |T_h| \ll_C \frac{P^2}{H} + P^{15/8}H^{1/2} \ll_C P^{23/12}.$$

If  $j = 0$ , the phase is smooth. For  $i \neq 0$ , its curvature is  $\asymp |i|P^{-1/2}$ , because  $|k/i|P^{-3/8} \ll_C P^{-1/3}$ . The bound is  $O_C(P^{37/48} + P^{1/4})$ . For  $i = 0$ , one has  $k \neq 0$ , curvature  $\asymp |k|P^{-7/8}$ , and bound  $O_C(P^{7/12} + P^{7/16})$ . Both are smaller than (4.11).  $\square$

**Corollary 4.6 (three formal signs).** Put  $v = \lfloor m^{3/2} \rfloor$ . Every sign class of

$$(\psi(n^{3/2}), \psi(m^{3/2}), \psi(v^{1/2}))$$

on odd  $n \leq N$  has count

$$\frac{N}{16} + O(N^{23/24}(\log(2N))^3). \quad (4.12)$$

In particular,  $H(OOEE; \delta)$  holds for every  $0 < \delta < 1/24$ .

*Proof.* On a dyadic block,

$$|v^{1/2} - n^{9/8}| \leq |v^{1/2} - m^{3/4}| + |m^{3/4} - X^{3/4}| \ll P^{-9/8} + P^{-3/8}.$$

Replacing the third phase coordinate in (4.11) by  $v^{1/2}$  costs  $O_C(|k|P^{5/8}) = O_C(P^{2/3})$ . Apply the three-dimensional Erdős–Turán–Koksma inequality, as in Proposition 4.2, to the points

$$(\{X/2\}, \{Y/2\}, \{v^{1/2}/2\}),$$

with cutoff  $\lfloor P^{1/24} \rfloor$  on this block. Its box discrepancy is  $O(P^{23/24}(\log(2P))^3)$ . The eight half-cube boxes encode the three signs exactly. Summing over dyadic blocks proves (4.12) and bounds all seven nonempty sign products by the same error. The cutoff is chosen separately on each block; no uniformity from a large block is assumed on smaller ones.  $\square$

This establishes the hypothesis used for four-step certificates. It does not estimate the expanding third-level coordinate  $v^{3/2}$  or every formal chain required at length five. The next subsection handles the *OOEOE* chain.

### 4.3 The fifth-letter split of OOEO

Write

$$X = n^{3/2}, \quad m = \lfloor X \rfloor, \quad Y = m^{3/2}, \quad v = \lfloor Y \rfloor, \quad U = v^{1/2}, \quad w = \lfloor U \rfloor, \quad W = w^{3/2}.$$

The four formal coordinates for *OOEOE* are  $X, Y, U, W$ . The new floor is handled by centering its Fourier expansion at the integer part of a smooth coefficient. This introduces many intervals; the next two lemmas keep their boundary cost explicit.

**Lemma 4.7 (a centered Fourier expansion).** For  $0 \leq \beta \leq 1$ , put

$$a_r(\beta) = \int_0^1 e(-(\beta + r)t) dt.$$

For integral  $T \geq 2$ , uniformly in real  $t$  and  $\beta$ ,

$$e(-\beta\{t\}) = \sum_{|r| \leq T} a_r(\beta) e(rt) + O(E_T(t)), \quad |a_r(\beta)| + |a'_r(\beta)| \ll (1 + |r|)^{-1}. \quad (4.13)$$

Here  $E_T$  is as in Lemma 4.3. If  $B = N + \beta$ ,  $N \in \mathbb{Z}$ , then  $e(-B\{t\}) = e(-Nt)e(-\beta\{t\})$  exactly. On an interval where  $N = \lfloor B(x) \rfloor$  is fixed and  $\beta(x)$  is monotone, the sum over  $|r| \leq T$  of the supremum norms and total variations of  $a_r(\beta(x))$  is  $O(\log(2T))$ .

*Proof.* The integral defines the coefficients also at the removable singularities of

$$a_r(\beta) = \frac{1 - e(-\beta)}{2\pi i(r + \beta)}.$$

Integration by parts gives the asserted bounds for  $|r| \geq 2$ , including for the derivative with respect to  $\beta$ ; the remaining coefficients and their derivatives are bounded by their integrals. For  $|r| \geq 2$ , the coefficient is  $(1 - e(-\beta))/(2\pi ir) + O(r^{-2})$ , uniformly in  $\beta$ . The tail of the first series is bounded by the argument of Lemma 4.3, and the absolutely convergent remainder has tail  $O(T^{-1})$ . Near an integer the paired  $r, -r$  terms are bounded as in that lemma. At the integer itself both sides are bounded and  $E_T = 1$ . This proves (4.13) at every point. Finally the variation of  $\beta$  on one interval is at most one, and summing the coefficient bounds gives the logarithm.  $\square$

**Bounded-residual extension.** The same expansion holds for every real  $\beta$  with  $|\beta| \leq B_0$ , where  $B_0$  is fixed, with constants depending only on  $B_0$ . Moreover, for a real function  $\beta(x)$  on an interval  $I$ , the condition  $\|\beta\|_{\infty, I} + \text{TV}_I(\beta) \leq B_0$  implies

$$\sum_{|r| \leq T} (\|a_r(\beta)\|_{\infty, I} + \text{TV}_I(a_r \circ \beta)) \ll_{B_0} \log(2T).$$

Monotonicity is not required for this extension.

*Proof.* For  $|r| > 2B_0 + 2$ , the integral formula gives  $a_r(\beta) = (1 - e(-\beta))/(2\pi ir) + O_{B_0}(r^{-2})$ , and  $|a_r| + |a'_r| \ll_{B_0} (1 + |r|)^{-1}$ . The remaining finitely many coefficients and derivatives are bounded directly by their integrals, including at  $r + \beta = 0$ . The same paired-tail argument proves the pointwise remainder; bounded cutoffs can be absorbed in the constant since  $E_T(t) \geq \min(1, 2/T)$ . Finally  $\text{TV}_I(a_r \circ \beta) \leq \sup_{|v| \leq B_0} |a'_r(v)| \text{TV}_I(\beta)$ . Sum the harmonic bounds. Complex conjugation gives the expansion with the opposite sign in the exponent.  $\square$

**Lemma 4.8 (mixed sums over the frequency intervals).** Set

$$J_0 = P^{1/48}, \quad H = \lfloor P^{1/12} \rfloor.$$

Let  $1 \leq |k| \leq J_0$ ,  $1/2 \leq u \leq J_0/2$ , and  $|i| \leq J_0$ . Partition a subinterval of  $[P, 2P]$  into  $D$  intervals  $I$ , with

$$D \ll |k|P^{9/16}, \quad |I| \ll L := P^{7/16}/|k|.$$

For a fixed constant  $C$ , define

$$M_I = \sup_{\substack{|R| \leq C|k|P^{9/16} \\ I' \subseteq I}} \left| \sum_{\substack{n \in I' \\ n \text{ odd}}} e\left(\frac{i}{2}X(n) + uY(n) + \frac{k}{2}n^{27/16} + Rn^{9/8}\right) \right|.$$

The second supremum ranges over subintervals. Then

$$\sum_I M_I \ll_C P^{23/24}. \quad (4.14)$$

The constants in the partition assumptions are fixed. In particular, this is a bound after summing all the frequency intervals, not a rescaling of a full-block discrepancy estimate.

*Proof.* Fix a shift  $1 \leq h < H$ . The exact carry argument of Lemma 4.4 applies on every  $I$ . The additional smooth part has differenced second derivative

$$O(|i|hP^{-3/2} + |k|hP^{-21/16} + |R|hP^{-15/8}) = O_C(J_0hP^{-3/2} + |k|hP^{-21/16}).$$

This is  $o(uhP^{-3/4})$ . Thus the zero-mode curvature and the dominance of every nonzero carry mode are unchanged.

Intersecting the frequency intervals with the gap cells produces  $O(D + hP^{1/2})$  cells in total: on an interval of length  $\ell$ , the gap function has variation  $O(hP^{-1/2}\ell)$ . Retaining these boundaries, the zero carry modes cost

$$\ll (uh)^{1/2}P^{5/8} + (h/u)^{1/2}P^{7/8} + D(uh)^{-1/2}P^{3/8}.$$

With carry cutoff  $R_0 = \lfloor P^{1/4} \rfloor$ , the nonzero modes cost

$$\ll R_0^{1/2}P^{3/4} + hP^{3/4} + DP^{1/4}.$$

These follow by applying the second-derivative estimate on every intersection, exactly as in (4.10). The weights  $z, 1 - z$  in the carry identity have uniformly bounded variation on each intersection.

The discarded floor terms cost  $O(uhP^{3/4} + uP^{1/4})$ . The truncated carry error costs  $O(P^{5/6} + P^{3/4} \log P)$  over the whole partition. For each local sum, its two endpoints  $n, n + 2h$  lie in the same  $I$ , so this error is bounded by a fixed multiple of  $\sum_{n \in I} E_{R_0}(X(n))$ . The disjoint intervals therefore charge (4.3) only once. The same bounds hold after taking the suprema over  $R$  and subintervals, because their local bounds use only  $|I|$ , its gap-cell count, and this nonnegative error sum.

If  $B_h$  denotes the sum over  $I$  of those supremum bounds for the differenced sums, the collected costs imply

$$B_h \ll_C P^{7/8}(1 + h^{1/2}) + |k|P^{15/16}h^{-1/2}. \quad (4.15)$$

Indeed  $uh \leq P^{5/48}/2$ ,  $u \geq 1/2$ , and  $DP^{1/4} \ll |k|P^{13/16} \leq P^{5/6}$ .

Apply van der Corput's inequality to each local sum, padding by zero inside an interval of length  $O(L)$  when taking a subinterval. Since  $H \ll L$ , its squared bound is

$$O\left(\frac{L}{H} \left(\#I + \sum_{h < H} |T_{I,h}|\right)\right),$$

where  $\#I$  counts odd integers and endpoint constants are harmless. Cauchy-Schwarz over the  $D$  intervals, together with  $DL \ll P$ , gives

$$\begin{aligned} \left(\sum_I M_I\right)^2 &\ll_C \frac{P^2}{H} + \frac{P}{H} \sum_{h < H} B_h \\ &\ll_C \frac{P^2}{H} + P^{15/8}H^{1/2} + |k|P^{31/16}H^{-1/2} \ll_C P^{23/12}. \end{aligned}$$

The three largest exponents are respectively  $2 - 1/12$ ,  $15/8 + 1/24$ , and  $1/48 + 31/16 - 1/24$ , all equal to  $23/12$ . This proves (4.14), including partial endpoint intervals.  $\square$

**Theorem 4.9 (four formal coordinates).** Uniformly over nonzero integer quadruples with  $\max(|i|, |j|, |\ell|, |k|) \leq P^{1/48}$ ,

$$\left| \sum_{\substack{P < n \leq 2P \\ n \text{ odd}}} e\left(\frac{i}{2}X + \frac{j}{2}Y + \frac{\ell}{2}U + \frac{k}{2}W\right) \right| \ll P^{23/24} \log(2P). \quad (4.16)$$

*Proof.* We first bound the error in (4.13) along  $U(n)$ . The comparison  $U = n^{9/8} + O(P^{-3/8})$  from Corollary 4.6 gives, for positive integer  $q$ ,

$$\left| \sum e(qU) \right| \ll q^{1/2}P^{9/16} + q^{-1/2}P^{7/16} + qP^{5/8}.$$

Erdős-Turán with cutoff  $\lfloor P^{1/8} \rfloor$  gives interval discrepancy  $O(P^{7/8})$  for  $\{U(n)\}$ . The distance-strip argument of Lemma 4.3 therefore gives

$$\sum E_T(U(n)) \ll P \log(2T)/T + P^{7/8}. \quad (4.17)$$

All sums here are over the same odd-input block.

When  $k = 0$ , Theorem 4.5 and the comparison for  $U$  prove (4.16). Suppose  $k \neq 0$ , and put  $\theta = \{X\}$ ,  $\xi = \{U\}$ . Taylor's theorem, with its second derivative bounded on the unit floor intervals, gives

$$\begin{aligned} U &= n^{9/8} - \frac{3}{4}\theta n^{-3/8} + O(P^{-9/8}), \\ W &= n^{27/16} - \frac{9}{8}\theta n^{3/16} - \frac{3}{2}n^{9/16}\xi + O(P^{-9/16}). \end{aligned} \quad (4.18)$$

For clarity,  $U = m^{3/4} + O(P^{-9/8})$ . Also  $W = v^{3/4} - (3/2)v^{1/4}\xi + O(P^{-9/16})$ ,  $v^{3/4} = m^{9/8} + O(P^{-9/16})$ , and  $v^{1/4} = n^{9/16} + O(P^{-15/16})$ ; expanding  $m = X - \theta$  proves both displayed formulas.

Set

$$B(x) = \frac{3k}{4}x^{9/16}, \quad C(x) = \frac{9k}{16}x^{3/16}, \quad T = \lfloor P^{1/8} \rfloor.$$

Partition by  $N = \lfloor B(x) \rfloor$ , and put  $\beta = B - N$ . Since  $|B'| \asymp |k|P^{-7/16}$ , this partition has  $D \ll |k|P^{9/16}$  intervals, each of length  $O(P^{7/16}/|k|)$ . The two endpoint intervals can be shorter. The total variation of  $\beta$  on any one interval is at most one.

Use (4.18), then expand  $e(-B\{U\})$  by Lemma 4.7. Formula (4.17) bounds its total truncation error by  $O(P^{7/8} \log P)$ ; the initial Taylor error is  $O(|k|P^{7/16})$ . On a fixed interval and for  $|r| \leq T$ , set

$$R = r - N + \ell/2.$$

This is a fixed frequency on that interval. The corresponding phase is

$$\frac{i}{2}X + \frac{j}{2}Y + \frac{k}{2}n^{27/16} + RU - C\theta.$$

The first formula in (4.18) replaces it by

$$\frac{i}{2}X + \frac{j}{2}Y + \frac{k}{2}n^{27/16} + Rn^{9/8} - (C + \frac{3}{4}Rn^{-3/8})\theta + O(|R|P^{-9/8}).$$

The essential cancellation is the exact identity

$$C + \frac{3}{4}Rn^{-3/8} = \frac{3}{4}n^{-3/8}(B + R) = \frac{3}{4}n^{-3/8}(r + \beta + \ell/2). \quad (4.19)$$

Its absolute value is  $O((T + P^{1/48})P^{-3/8})$ , so the remaining  $\theta$ -term can be deleted. Since  $|R| \ll |k|P^{9/16}$ , all these replacements, summed with the coefficient masses, cost

$$O((|k|P^{7/16} + P^{5/8}(T + P^{1/48})) \log P) \ll P^{3/4} \log P.$$

Thus it remains to estimate the smooth-frequency phases in Lemma 4.8. If  $j \neq 0$ , conjugation if needed makes  $u = |j|/2 \geq 1/2$ . Partial summation for the coefficients  $a_r(\beta)$ , whose summed supremum norms and variations are  $O(\log P)$  on each interval, and (4.14) give  $O(P^{23/24} \log P)$ .

If  $j = 0$ , differentiate with  $R$  fixed. The remaining phase  $\Phi = iX/2 + kn^{27/16}/2 + Rn^{9/8}$  satisfies

$$\begin{aligned} \Phi''(x) &= \frac{3i}{8}x^{-1/2} + \frac{297k}{512}x^{-5/16} + \frac{9R}{64}x^{-7/8} \\ &= \frac{243k}{512}x^{-5/16} + O(|i|P^{-1/2} + (T + |\ell| + 1)P^{-7/8}). \end{aligned}$$

The last equality substitutes  $R = -B(x) + (r + \beta(x) + \ell/2)$  only after differentiation. The error is  $o(|k|P^{-5/16})$ . Summing the second-derivative bounds over every frequency interval costs

$$O(|k|^{1/2}P^{27/32} + D|k|^{-1/2}P^{5/32}) \ll |k|^{1/2}(P^{27/32} + P^{23/32}).$$

The logarithmic coefficient mass still leaves this below (4.16). All zero coordinates and both signs of every mode have now been covered.  $\square$

**Corollary 4.10 (the OOEEO split).** Each of the sixteen formal sign classes of  $(\psi(X), \psi(Y), \psi(U), \psi(W))$  over odd  $n \leq N$  has count

$$N/32 + O(N^{47/48}). \quad (4.20)$$

In particular, for each  $a \in \{OOEOE, OOEOO\}$ ,

$$\#\{n \leq N : \text{word}_5(n) = a\} = N/32 + O(N^{47/48}).$$

The hypothesis  $H(OOEEOE; 1/48)$  holds.

*Proof.* Apply four-dimensional Erdős–Turán–Koksma on each dyadic block with cutoff  $\lfloor P^{1/48} \rfloor$ . Theorem 4.9 bounds its box discrepancy by

$$O(P^{47/48} + P^{23/24}(\log(2P))^5) = O(P^{47/48}).$$

The sixteen half-cubes have volume  $1/16$ ; the number of odd inputs is  $P/2 + O(1)$ . Summing dyadically proves (4.20). All fifteen nonempty sign products are bounded by summing these discrepancies. The signs  $(-1, +1, -1, \pm 1)$  select the two actual words by Lemma 2.2. This does not count the formal chain through the expanding third-level coordinate  $v^{3/2}$ .  $\square$

#### 4.4 The OOOEE mixed-mode estimate

For this subsection the formal coordinates are

$$X = n^{3/2}, \quad m = \lfloor X \rfloor, \quad Y = m^{3/2}, \quad v = \lfloor Y \rfloor, \quad Z = v^{3/2}, \quad w = \lfloor Z \rfloor, \quad U = \sqrt{w}.$$

They differ from the chain in Section 4.3.

**Theorem 4.11 (OOOEE mixed modes).** For every fixed  $C \geq 1$  and every  $\varepsilon > 0$ , uniformly over nonzero integer quadruples with  $\max(|i|, |j|, |k|, |l|) \leq CP^{1/24}$ ,

$$\left| \sum_{\substack{P < n \leq 2P \\ n \text{ odd}}} e\left(\frac{1}{2}(iX + jY + kZ + lU)\right) \right| \ll_{C, \varepsilon} P^{127/128 + \varepsilon}. \quad (4.21)$$

*Proof.* Appendix C provides the full argument using Lemmas A.1–A.4 and Theorem B.1. The cases with  $k = 0$  are treated directly. For  $k \neq 0$ , two differencings use half-shift cutoffs  $P^{1/48}$  and  $P^{1/24}$ . The complete second correlation has exponent  $31/32$ , including all positive errors. The two outer inequalities give exponents  $63/64$  and  $127/128$ ; their diagonal terms are smaller.  $\square$

**Corollary 4.12 (the OOOE split).** Every formal parity-sign class of  $(X, Y, Z, U)$  on odd  $n \leq N$  has count

$$N/32 + O_\varepsilon(N^{127/128 + \varepsilon}). \quad (4.22)$$

In particular *OOOEE* and *OOOEO* each have that count, and  $H(\text{OOOEE}; \delta)$  holds for every  $0 < \delta < 1/128$ .

*Proof.* On each dyadic block apply Erdős–Turán–Koksma at cutoff  $\lfloor P^{1/24} \rfloor$  to the four coordinates divided by two, as in Proposition 4.2. The sixteen half-cubes all have volume  $1/16$ , and there are  $P/2 + O(1)$  odd inputs. Absorb the fixed logarithmic weights in (4.21) into epsilon, then sum dyadically with each block’s own cutoff. The signs  $(-1, -1, +1, \pm 1)$  select the stated actual words. Summing the sixteen discrepancies also controls every nonempty sign product. The end of Appendix C gives the endpoint conventions and full dyadic argument.  $\square$

### 5. Finite-depth descent densities

**Lemma 5.1 (minimal certificates through length five).** The contracting words of length at most five with no proper contracting prefix are exactly

$$E, \quad OE, \quad OOE, \quad OOOE, \quad OOEEO. \quad (5.1)$$

*Proof.* An  $E$ -rooted word already contracts at its first letter, and an  $OE$ -rooted word contracts at its second. Every remaining word begins with  $OO$ . At length three, even two odd letters give  $3^2 > 2^3$ , so no new certificate appears. At length four, contraction requires at most two odd letters, giving only *OOEE*. After excluding that prefix, the possible fourth-level prefixes are *OOEO*, *OOOE*, and *OOOO*. At length five, contraction requires at most three odd letters because  $3^3 < 2^5 < 3^4$ . The first two prefixes must therefore end with  $E$ , and the third cannot contract. This proves (5.1).  $\square$

These five words are the minimal certificates of length at most five counted by A100982 [6]. That sequence is supported on the binary lengths of the powers of three [7], which is why length three carries no certificate; the enumeration above verifies its first terms rather than establishing them.

The list, and the statement for every length — a minimal certificate of length  $L$  exists exactly when a power of three lies in  $[2^{L-1}, 2^L)$ , its odd count is then forced by  $L$ , and its last letter is  $E$  — are machine-checked in `formal/Problems/Juggler/PaperBCertificates.lean`

and `PaperBCertificateLengths.lean`. Length six adds no minimal certificate and length seven adds exactly three, `OOEOOEE`, `OOEOOEE` and `OOEOOEE`; that is a count of words, not a density statement, since each would need its own mixed-mode estimate. Both facts are checked in `PaperBFiveStepDensity.lean`.

**Theorem 5.2 (unconditional four-step density).** One has

$$\#(\mathcal{C}_4 \cap [1, N]) = \frac{13N}{16} + O(N^{23/24}(\log(2N))^3).$$

*Proof.* Lemma 5.1 gives the disjoint prefix classes  $E$ ,  $OE$ , and  $OOEE$ . Theorem 3.1 supplies the first two counts. The  $OOEE$  class is the sign class  $(-1, +1, +1)$  in Corollary 4.6, so has count  $N/16 + O(N^{23/24}(\log(2N))^3)$ . Their densities add to  $1/2 + 1/4 + 1/16 = 13/16$ .  $\square$

**Theorem 5.3 (a five-step subfamily of density  $27/32$ ).** Let

$$\mathcal{D}_5 = \mathcal{C}_4 \cup \{n \geq 2 : \text{word}_5(n) = \text{OOEOE}\}.$$

Then  $\mathcal{D}_5 \subseteq \mathcal{C}_5$  and

$$\#(\mathcal{D}_5 \cap [1, N]) = 27N/32 + O(N^{47/48}).$$

In particular the lower natural density of  $\mathcal{C}_5$  is at least  $27/32$ .

*Proof.* The added prefix is disjoint from  $E$ ,  $OE$ ,  $OOEE$ , and  $3^3 < 2^5$  certifies its contraction. Add Corollary 4.10 to Theorem 5.2, absorbing the smaller four-step error. The main terms sum to  $13/16 + 1/32 = 27/32$ . This does not assert that  $\mathcal{D}_5$  exhausts  $\mathcal{C}_5$ .  $\square$

**Theorem 5.4 (full five-step certificate density).** For every  $\varepsilon > 0$ ,

$$\#(\mathcal{C}_5 \cap [1, N]) = \frac{7N}{8} + O_\varepsilon(N^{127/128+\varepsilon}).$$

In particular  $\mathcal{C}_5$  has natural density  $7/8$ . This counts the power-envelope certificate class, not every start that may happen to descend within five operations.

*Proof.* By Lemma 5.1,  $\mathcal{C}_5$  is the disjoint union of  $\mathcal{D}_5$  and the  $OOOEE$  class. Add Theorem 5.3 and Corollary 4.12. Their main terms are  $27N/32 + N/32 = 7N/8$ ; the former error  $O(N^{47/48})$  is smaller.  $\square$

The partition behind Theorem 5.4 — at depth five the cylinders of the five words of Lemma 5.1 have sizes 16, 8, 2, 1, 1, are pairwise disjoint and exhaust the certified words, so that  $1/2 + 1/4 + 1/16 + 1/32 + 1/32 = 7/8$  is cylinder arithmetic rather than an identity between rational literals — and the assembly of the count as a sum of five class counts are machine-checked in `PaperBFiveStepDensity.lean`; the analytic estimates for those counts are not.

The exact fractions and their analytic requirements can be read together:

| Certificate family                      | Minimal prefixes added | Density | Status        |
|-----------------------------------------|------------------------|---------|---------------|
| $\mathcal{C}_1$                         | $E$                    | $1/2$   | Unconditional |
| $\mathcal{C}_2$                         | $OE$                   | $3/4$   | Theorem 3.1   |
| $\mathcal{C}_4$                         | $OOEE$                 | $13/16$ | Theorem 5.2   |
| $\mathcal{D}_5 \subseteq \mathcal{C}_5$ | $OOEOE$                | $27/32$ | Theorem 5.3   |
| $\mathcal{C}_5$                         | $OOOEE$                | $7/8$   | Theorem 5.4   |

The formal  $OOOEE$  chain in Section 4.4 uses the expanding third-level coordinate  $v^{3/2}$ ; the  $OOEOE$  chain in Section 4.3 uses  $v^{1/2}$  at that level. The proofs therefore concern distinct families. Neither proof asserts fair-share counts for every actual itinerary at every depth.

## 6. The fixed-depth reduction to density-one certificates

**Hypothesis FD.** For every fixed  $d \geq 1$  and every odd-rooted word  $w$  of length  $d$ ,

$$\#\{n \leq N : \text{word}_d(n) = w\} = 2^{-d}N + o_w(N).$$

No common error rate in  $d$  is assumed. The family  $H_0(w)$  over all such words implies FD.

**Theorem 6.1 (conditional density-one certificate theorem).** Under FD,  $\mathcal{C}_\infty$  has natural density one.

*Proof.* Put  $p = \log 2 / \log 3 > 1/2$ . A word with no contracting prefix through length  $d$  must in particular have at least  $pd$  odd letters at length  $d$ . All such words begin with  $O$ . For a uniformly chosen odd-rooted word of length  $d$ , the number of odd letters is distributed as  $1 + B_{d-1}$ , where  $B_{d-1}$  is a binomial random variable with parameters  $(d-1, 1/2)$ . This is an auxiliary count on words, not a stochastic model asserted for the Juggler orbit.

Choose  $q = q_d := p - (1-p)/(d-1)$ . This is the largest threshold the step admits: it makes  $pd - 1 \geq q(d-1)$  an equality, so every admissible  $q$  is at most  $q_d$ . The one remaining constraint,  $q_d > 1/2$ , holds at every  $d \geq 4$  and fails at  $d = 3$ , because  $q_4 = (4p-1)/3 > 1/2$  reduces to  $p > 5/8$ , which is  $2^8 > 3^5$ , and  $q_3 = (3p-1)/2 \leq 1/2$  reduces to  $p \leq 2/3$ , which is  $2^3 < 3^2$ . No “sufficiently large  $d$ ” clause is needed. The sharp threshold  $q = p$  is itself unavailable, and only because the leading  $O$  is spent before the binomial begins: the step at  $q = p$  reduces to  $p \geq 1$ . For  $t > 0$ , the exponential Markov inequality and the binomial generating function give

$$\Pr(B_{d-1} \geq q(d-1)) \leq \left( \frac{1+e^t}{2e^{qt}} \right)^{d-1}.$$

Both steps are machine-checked in `formal/Problems/Juggler/PaperBMarkov.lean`, stated over binomial coefficients rather than over a probability space because the count here is over words: `sum_choose_mul_pow` is the generating function and `chernoff_density` the Markov bound. The optimisation below is an identity there rather than a differentiation — `tilt_gives_theta` shows the right-hand side at  $t = \log(q/(1-q))$  is exactly  $\vartheta(q)^{d-1}$ . The logarithm of the expression in parentheses is zero at  $t = 0$  and has derivative  $1/2 - q < 0$  there. Hence some fixed  $t > 0$  makes it a number  $\vartheta < 1$ ; optimising  $t$  gives  $\vartheta(q) = q^{-q}(1-q)^{q-1}/2 = e^{-\text{KL}(q\|1/2)}$ . Write  $\vartheta_d := \vartheta(q_d)$ . Since  $\vartheta$  decreases on  $(1/2, 1)$  and  $q_d$  increases in  $d$ , the  $\vartheta_d$  decrease, so  $\vartheta_d \leq \vartheta_4 < 1$  for every  $d \geq 4$ . The strict inequality  $\vartheta(q) < 1$  for  $q \neq 1/2$  — the step that makes this bound decay rather than merely not grow — is machine-checked in `formal/Problems/Juggler/PaperBChernoff.lean` as `theta_lt_one`, resting on strict Gibbs (`klDiv_pos`); the fate layer’s `klDiv_nonneg` gives only  $\vartheta \leq 1$ , which is vacuous at every depth.

For fixed  $d$ , FD and a finite sum over surviving words show that the natural density of  $\mathbb{N} \setminus \mathcal{C}_d$  is their number divided by  $2^d$ . That inference is finite additivity of density and nothing more, which is why FD is needed at each fixed  $d$  separately and no uniformity in  $d$  is claimed; it is machine-checked as `density_of_finite_union` in `formal/Problems/Juggler/PaperBDensity.lean`, together with the  $d \rightarrow \infty$  passage below (`exceptional_density_zero`). Neither formalises FD itself, which is open, so Theorem 6.1 remains conditional exactly as stated. It is at most  $\frac{1}{2}\vartheta_d^{d-1}$  for every  $d \geq 4$ . Since  $\mathcal{C}_d \subseteq \mathcal{C}_\infty$ ,

$$\overline{\text{dens}}(\mathbb{N} \setminus \mathcal{C}_\infty) \leq \frac{1}{2}\vartheta_d^{d-1}.$$

Letting  $d \rightarrow \infty$  proves the assertion. In this argument  $N \rightarrow \infty$  is taken first, with  $d$  fixed.  $\square$

The conclusion concerns descent below the initial value. It does not prove that a descended value reaches 1, or that subsequent orbit samples avoid an exceptional set. Even a density-zero exceptional set can contain a nontrivial cycle. No interchange of the two limits, and no assertion of finite stopping time for every start, follows from Theorem 6.1.

**Remark 6.2 (the threshold, and what it leaves).** The threshold  $q_d$  is chosen to be sharp rather than convenient, and the difference is not a constant factor. An earlier version of

this proof took  $q = (p + 1/2)/2$ , which is admissible at every depth but halves the distance to  $1/2$  when the  $-1$  costs only  $O(1/d)$ . The exponents are

$$-\log \vartheta((p + \tfrac{1}{2})/2) = 0.0085959587, \quad -\log \vartheta(p) = 0.0346881852,$$

so the convenient threshold reports 24.8% of the exponent its own inequality supplies.

The sharp value is not an accident of the binomial. A word has no contracting prefix only if  $o \log 3 \geq d \log 2$  at the endpoint, so the event is a large deviation of the same walk whose step is  $\log(3/2)$  or  $-\log 2$  with equal probability; its Cramér rate  $\varrho = \min_t \mathbb{E}[e^{tX}] = 0.9659065532$  and  $\vartheta(p)$  are the same number, and the inequality above reaches it. The words this bound counts are those of A076227 [5], so its decay is a statement about Collatz residue classes as well. There the fair-share densities that Hypothesis FD assumes are a theorem of Terras [4]; that asymmetry, and not the combinatorics, is what the analytic sections of this paper are for.

At  $q_d$  the bound has a closed form. Writing  $\text{KL}(q) = -\log \vartheta(q)$ , so  $\text{KL}'(q) = \log(q/(1-q))$ ,

$$\tfrac{1}{2}\vartheta(q_d)^{d-1}/\varrho^d \longrightarrow \tfrac{1}{2}e^{\text{KL}(p)}e^{(1-p)\text{KL}'(p)} = p^p(1-p)^{1-p}\left(\frac{p}{1-p}\right)^{1-p} = p,$$

every factor of  $1-p$  cancelling, so the bound of Theorem 6.1 reads

$$\overline{\text{dens}}(\mathbb{N} \setminus \mathcal{C}_d) \leq p \varrho^d (1 + O(1/d)),$$

with both constants explicit and no unspecified  $t$ .

What remains between this and the truth is a polynomial together with an oscillation, and the oscillation is not a defect of the measurement. Writing  $\beta = p$  for the threshold, the exact count satisfies

$$N_d/2^d = \psi(\{d\beta\}) \varrho^d d^{-3/2} (1 + o(1)),$$

with  $\psi$  a non-constant function on the circle: over twenty consecutive depths the ratio keeps a spread of 1.07 at  $d \approx 1000, 2000, 4000$  alike, and binning by  $\{d\beta\}$  collapses it, the within-bin scatter being one tenth of the across-bin range. The reason is arithmetic rather than probabilistic: at each fixed  $d$  the endpoint  $o \log 3 - d \log 2$  lies on a lattice of spacing  $\log 3$  whose offset  $-d \log 2 \bmod \log 3$  equidistributes, so the barrier at 0 is approached from a different distance at every depth, namely  $1 - \{d\beta\}$ .

Neither the exponential rate nor the  $d^{-3/2}$  is claimed here, and the rate is forty years old. It is Theorem D of Lagarias [14]:  $1 - F(k) \leq 2^{-\eta k}$  with  $\eta = 1 - H(\lambda^{-1}) = 0.0500444728$ ,  $H$  the binary entropy, the remark after it supplying the matching lower bound so that the estimate is two-sided on the exponential scale. That is exactly  $\vartheta(p)$  of this section, and not by numerical coincidence but by an identity:  $2^{-\eta} = 2^{H(p)-1} = p^{-p}(1-p)^{p-1}/2 = \vartheta(p)$ , the two agreeing to every digit computed. Hikawa [9, Section 6] proves the same rate independently in the weight basis,  $\log_2 W(d) = \gamma d + O(\log d)$  with  $\gamma = \lambda H(1/\lambda)$  and  $c = \lambda(1 - H(1/\lambda)) = \lambda D(1/\lambda \| \tfrac{1}{2})$  bits per unit weight; since lengths are  $\lambda$  times weights his constants are Lagarias's  $\eta$  rescaled,  $\gamma = H(p)/p$  and  $c = \eta/p$ , and  $c/\lambda = 0.0500444728$  recovers  $\eta$ . He remarks that the rate appears to be new for these sequences; that remark is incorrect, and it is repeated here only to say so. What is his is the weight-basis form and Conjecture 7.1. The  $d^{-3/2}$  power is Hikawa's Conjecture 7.1 [9, Section 7], motivated there by the residual  $\log_2 W(d) - \gamma d$  tracking a ballot-type correction  $-\tfrac{3}{2} \log_2 d$  to within 0.3 bits over  $100 \leq d \leq 10^4$ . No priority over either is asserted. Of the parity-vector line, [11] was read in full and states no asymptotic; of [10] only the abstract was available to this revision, and of [9] the author's version note on the record, from which Sections 6 and 7 are cited; the September preprint [17], a third paper of the line, is listed because it exists, nothing of its text having been available. Nor does [12] bear on the asymptotic: it refines the same count by a local cut statistic and represents the words as partitions inside a Beatty–Fermers shape: exact finite structure, with no growth constant and no oscillation. That Beatty structure is a predecessor for any reading of the admissible lengths A020914 through their

Beatty complement A054414. What this section offers is the oscillating prefactor  $\psi$  : that the ratio depends on  $d$  only through  $\{d\beta\}$ , that it is non-constant, and the barrier-step mechanism below that explains why. Novelty is claimed for that, and it survives the nearest prior result. Winkler [13, Section 7] derives growth constants and exact normalized oscillations for A100982 and its siblings A174795 and A174796. A100982 is the minimal-certificate count of this paper,  $2N_{d-1} - N_d$  — in the notation of the recursion below,  $b_{d-1}M_{d-1}$ , the survivors on the barrier that lose their even extension — and summing that identity writes  $N_d/2^d$  as a tail of his sequence, so his results do bear on  $\psi$ . They do not determine it. His Corollary 12 normalises  $a_3(r)$  by its lower cycle bound and proves that the ratio has  $\liminf$  equal to 1 and  $\limsup$  equal to  $\alpha/(\alpha - 1)$  with  $\alpha = \log_2 3$ , the lower value attained exactly on the record minima of  $\{r\alpha\}$  and the upper approached along the record maxima. That is an exact statement about the extremes of two sparse sets of orders, not an asymptotic for a general  $r$ , and it carries no error term at all. The envelope has width 2.7095112914 and transfers termwise to the tail, while the entire oscillation of  $\psi$  spans a factor 1.06 : too coarse by a factor near thirty to resolve the shape. Both corollaries were verified against the counts of this paper.

Consequently  $(N_d/2^d)/(p\varrho^d) \cdot d^{3/2}$  does not converge. Its values 16.4, 17.0, 16.6 at  $d = 640, 1280, 2560$  are samples of  $\psi/p$  and not approaches to a limit; the mean of  $\psi$  is about 10.9, which is the constant such measurements had previously been read as estimating. The convenient threshold admits no reading of this kind at all: its overshoot is  $1.2 \cdot 10^6$  at  $d = 320$  and  $6.9 \cdot 10^{17}$  at  $d = 1280$ , growing without bound.

The shape of  $\psi$  is known and its existence is not, and the structure behind it is exact down to one missing limit.

*The count recursion.* Write  $N_d$  for the surviving words of length  $d$ ,  $M_d$  for those of them with  $o_d = \lceil d\beta \rceil$  exactly — sitting on the barrier — and  $b_d = \lceil (d+1)\beta \rceil - \lceil d\beta \rceil$ . Every survivor has two extensions; if the barrier does not rise both survive, and if it rises then a survivor strictly above the barrier still keeps both while one exactly on it loses the even extension and only that one. Hence

$$N_{d+1} = 2N_d - b_d M_d, \quad P_{d+1} = P_d \left(1 - \frac{1}{2} b_d R_d\right),$$

with  $P_d = N_d/2^d$  and  $R_d = M_d/2^d/P_d$  the share of survivors on the barrier. This is exact and elementary; it is checked against the integer counts and machine-checked in `formal/Problems/Juggler/PaperBBarrierStep.lean` and, as an identity of integer counts, in `PaperBCertificateRecursion.lean`.

This recursion is not new. On A076227 [5] it is Zarubin's formula of 11 August 2019, which reads  $a(L_k) = 2a(L_k - 1) - A_{100982}(k)$  at the lengths  $L_k$  where the barrier rises and  $a(L) = 2a(L - 1)$  where it does not — the display above, in the entry's indexing — and Winkler's formula of 12 September 2017 gives the same counts as a Pascal-type sum [8]. The entry also carries, from 21 July 2026, Kazunobu Hikawa's statement of the surviving-word object itself: the binary words  $v(1) \dots v(n)$  with  $2^m < 3^{v(1)+\dots+v(m)}$  for every  $m$  [9]. Nothing in this subsection is claimed as new. It is restated because the argument needs it in this form, and it is machine-checked so that the form is certified rather than assumed.

The increments are tabulated too: T. D. Noe's frequency comment on A186008 [16], February 2011, lists  $1/2, 1/4, 1/16, 1/16, 3/128, 7/256, 3/256, \dots$ , which are  $b_{d-1}M_{d-1}/2^d$  here, the certificate density gained at each carrying length, and A186009 [16] is the sequence of nonzero minimal-certificate counts. The telescoped form of the recursion, writing  $N_d/2^d$  as the mass of the minimal certificates beyond depth  $d$  plus the survivors at the cutoff, and the doubling  $N_{d+1} = 2N_d$  exactly at the lengths of A054414 are machine-checked in `PaperBCertificateRecursion.lean`; that exchanging two adjacent barrier letters costs exactly the mass on the barrier, the elementary fact under the discontinuities of  $\psi$ , is machine-checked in `PaperBJumpTransposition.lean`.

*The right coordinate is  $R$ .* Conditioned on survival, the law of  $m_d = o_d - \lceil d\beta \rceil$  depends on the depth only through  $\{d\beta\}$ , and far more cleanly than  $\psi$  does, because a conditional law carries no  $\varrho^d d^{-3/2}$  to divide out. Advancing the phase by  $\beta$  shifts the barrier word one place

and so applies one more update,

$$\Pi_{\varphi+\beta} = T_{b(\varphi)}\Pi_{\varphi}/(1 - \tfrac{1}{2}b(\varphi)\Pi_{\varphi}(0)),$$

where  $T_0\pi(m) = \frac{1}{2}(\pi(m) + \pi(m-1))$  and  $T_1\pi(m) = \frac{1}{2}(\pi(m) + \pi(m+1))$ . Because  $T_0$  preserves mass and sends  $\pi(0)$  to  $\pi(0)/2$ , this gives  $R(\varphi + \beta) = R(\varphi)/2$  whenever  $\varphi < 1 - \beta$ , confirmed to  $10^{-12}$ . Averaging the logarithm of the recursion over the circle recovers the rate,  $\log \varrho = \int_0^1 \log(1 - \frac{1}{2}b(\varphi)R(\varphi)) d\varphi$ , to  $3 \cdot 10^{-5}$  — a constraint rather than a restatement, since  $\varrho$  is known in closed form.

*Why a polynomial appears at all.* A geometric tail  $r^m$  keeps its shape under both updates, reproducing with factor  $[\frac{1}{2}(1 + 1/r)]^{1-\beta} [\frac{1}{2}(1 + r)]^{\beta}$ . That expression is stationary exactly at  $r = (1 - \beta)/\beta = e^{-\lambda^*}$ , and there it equals  $-\text{KL}(\beta \| \frac{1}{2}) = \log \varrho$  identically. So the characteristic equation has a **double** root:  $\varrho$  is the *minimum* of that expression, a second variational formula in the tail variable dual to the Cramér one in the tilt variable. A simple root would give a pure exponential; a double root is the critical case and forces the polynomial factor.

*Where  $\psi$  comes from.* Combining the recursion with  $P_d = \psi(\{d\beta\})\varrho^d d^{-3/2}$  makes  $\log \psi$  a coboundary over the rotation,

$$\log \psi(\varphi + \beta) - \log \psi(\varphi) = f(\varphi), \quad f = \log(1 - \tfrac{1}{2}bR) - \log \varrho,$$

whose mean vanishes by the identity above, so  $\hat{\psi}(n) = \hat{f}(n)/(e^{2\pi i n \beta} - 1)$  — reproduced to a few percent across fifteen modes.  $R$  is a step function whose jumps sit on  $\{k\beta\}$ , so  $f$  jumps there and at  $1 - \beta$ , and a coboundary spreads those over the same orbit:  $\psi$ 's jumps are not an accident of the barrier arithmetic but the shape of a coboundary. The Ostrowski structure in  $\hat{\psi}$  is inherited from  $\hat{f}$ , not manufactured by the divisor —  $\hat{f}$  is five times smaller at the resonances than away from them, which is what keeps  $\psi$  bounded. It is in fact of bounded variation, and its jumps are the survivor sequence itself. The jump of  $\psi$  at  $\{n\beta\}$  is

$$a_n = \frac{a_1 N_n}{(2\varrho)^{n-1}} \quad (n \geq 1),$$

reproduced to  $2.3 \cdot 10^{-14}$  over  $n \leq 3000$  by a transfer-operator propagation that uses no measured data; with the normalisation of  $\psi$  above,  $a_n = 2\varrho a_1 \psi(\{n\beta\}) n^{-3/2}(1 + o(1))$ , so the amplitudes decay with the same power as the counts they come from, and the steepening power law that local differencing had suggested was the bias of the differencing. The one constant is closed form,

$$a_1 = \frac{1}{2\varrho\sqrt{2\pi\beta(1-\beta)}} = 0.427956804.$$

The reason is a Fourier identity. Let  $A_n$  be the probability that the walk of Remark 6.2 is nonnegative at step  $n$ , divided by  $\varrho^n$ . Then  $\sqrt{n} A_n = \Phi(\{n\beta\})(1 + O(1/n))$  with the ladder profile  $\Phi(x) = r^{1-x}/((1-r)\sqrt{2\pi\beta(1-\beta)})$ ,  $r = (1 - \beta)/\beta$ , whose mean  $\kappa = 1.541814521$  is the non-lattice local-limit constant, and  $\hat{\psi}(k) = \hat{\Phi}(k)G(e^{-2\pi i k \beta})$  for every  $k$ , where  $G(w) = \sum_d N_d w^d / (2\varrho)^d$ . So the jump of  $\psi$  at the origin,  $2\varrho a_1 = 1/\sqrt{2\pi\beta(1-\beta)}$ , is the jump of  $\Phi$  there; mode by mode, with  $\psi$  measured at depth  $10^6$ , the ratio  $|\hat{\psi}(k)|/|\hat{\Phi}(k)G(e^{-2\pi i k \beta})|$  is  $1.0015 \pm 0.0020$  over  $k \leq 256$  with no trend in  $k$ , and explicitly  $\hat{\psi}(k) = -(2\varrho a_1 / 2\pi i k) G(e^{-2\pi i k \beta})$  for  $k \neq 0$ . Summing the jumps gives  $2\varrho a_1 (G(1) - 1)$ , and the mean of  $\psi$  is  $\kappa G(1) = 10.8927$  with  $G(1) = 7.0649$  a convergent series, so the summability of the jump spectrum and the convergence of  $G(1)$  are one statement. These are verified computations, recorded with their tests in the repository; the existence of  $\psi$  as a limit is not among them.

*What is open, and what is excluded.* One link is unproved: that the iteration converges,  $R_d \rightarrow R(\{d\beta\})$ . Everything above is exact or elementary given it. Three natural attacks are excluded, the third by argument rather than by measurement. Birkhoff contraction on the

product of positive operators would give geometric memory loss, and the loss is polynomial. A small-divisor obstruction would show as an excess at  $\beta$ 's convergent denominators, and the convergence is uniform across them. A change of weight would restore a spectral gap, and none does: at weight  $r^m$  the log essential radius is the same expression whose stationary point is  $r^*$ , so it meets  $\log \varrho$  there and lies strictly above it at every other  $r$ . Equivalently a Foster–Lyapunov drift  $V(m) = \theta^m$  is pinned at exactly  $\varrho$  when  $\theta = 1/r^*$  and is useless otherwise. That one is proved rather than measured.

The literature reaches the driven case but stops short of this one. Oçafraïn treats absorbing boundaries that move periodically, and two of those results bear on the construction above. For a boundary that moves at all there is *no* quasi-stationary distribution in the strict sense, so the phase-indexed family  $R(\varphi)$  is forced rather than chosen; and for a finite state space with a periodic boundary a quasi-ergodic distribution exists by an extension of Darroch–Seneta, which covers the truncated rational barrier. The general moving-boundary theorems assume a conditional Doeblin condition that is equivalent, in the time-homogeneous case, to *exponential uniform* convergence to quasi-stationarity — which this process fails precisely by being critical. So the missing combination is driven *and* critical, and it is now the hypothesis of a named theorem rather than an absence in the record.

The driving itself is not the obstruction. For a rational barrier  $p/q$  the period map is autonomous, and the constant in the convergence  $\text{TV} \sim c/d$  is flat in  $q$ : across six convergents of  $\beta$  from  $q = 19$  to  $q = 24727$  the exponent stays  $d^{-1}$  and  $c$  stays 20.1, the last two agreeing to six digits. Every barrier in the family is a bounded perturbation of the same line,  $|\lceil ts \rceil - ts| < 1$  uniformly in  $t$  and  $s$ , which is why  $q \rightarrow \infty$  costs nothing. What remains is the criticality itself.

None of this changes Theorem 6.1, whose conclusion follows from any  $\vartheta < 1$ . The algebra of the threshold — that  $q_d$  saturates the step, that no admissible threshold exceeds it, that  $q = p$  is blocked only by  $p < 1$ , and that  $d \geq 4$  is the first usable depth — is machine-checked in `formal/Problems/Juggler/PaperBThreshold.lean`, and the tilt, its closed form and the arithmetic of the barrier gap in `formal/Problems/Juggler/PaperBTilt.lean`.

## 7. Exact floor defects and separate analytic questions

For  $n \geq 3$ , write

$$X = n^{3/2}, \quad m = \lfloor X \rfloor, \quad \theta = X - m, \quad Y = m^{3/2}, \quad v = \lfloor Y \rfloor, \quad \theta_2 = Y - v.$$

**Lemma 7.1 (one-signed linearization).** One has

$$m^{3/2} = \frac{3}{2}mn^{3/4} - \frac{1}{2}n^{9/4} + E(n), \quad 0 \leq E(n) \leq \frac{3}{8}(n^{3/2} - 1)^{-1/2} \leq \frac{1}{2}n^{-3/4}.$$

*Proof.* Let  $a = \sqrt{m}$ ,  $b = \sqrt{X}$ . Direct factorization gives

$$E = a^3 - \frac{3}{2}a^2b + \frac{1}{2}b^3 = \frac{1}{2}(a - b)^2(2a + b) \geq 0.$$

Moreover  $\theta = b^2 - a^2$ , and  $Ea \leq \frac{3}{8}\theta^2$  follows from  $4a(2a + b) \leq 3(a + b)^2$ , equivalent to  $(5a + 3b)(a - b) \leq 0$ . Thus  $E \leq \frac{3}{8}m^{-1/2}$ , which is at most  $\frac{3}{8}(X - 1)^{-1/2}$ . Finally  $(X - 1)^{-1/2} \leq \frac{4}{3}X^{-1/2}$  for  $n \geq 3$ .  $\square$

**Lemma 7.2 (gap and carry).** For an integer  $h \geq 1$ , set  $\Delta_h X(n) = X(n+2h) - X(n)$ . Then

$$m(n+2h) - m(n) = \lfloor \Delta_h X(n) \rfloor + \kappa_h(n), \quad \kappa_h(n) = \mathbf{1}_{\{X(n)\} + \{\Delta_h X(n)\} \geq 1} \in \{0, 1\}.$$

*Proof.* Expand  $X(n) + \Delta_h X(n)$  into integer and fractional parts, and take its floor. The sum of the two fractional parts lies in  $[0, 2)$ .  $\square$

For  $1 \leq h \leq P/4$  and  $n \in (P, 2P]$ , differentiation gives  $(\Delta_h X)'(n) \asymp hP^{-1/2}$ . The level sets of  $\lfloor \Delta_h X \rfloor$  consequently form  $O(1 + hP^{1/2})$  intervals. Full intervals have length  $\asymp P^{1/2}/h$ ; the two endpoint intervals can be shorter. A coefficient involving this floor is frozen only on these intervals. A derivative estimate proved there must be summed over the partition, or replaced by a separate global argument with controlled errors.

**Lemma 7.3 (next-level defect).** One has

$$v^{3/2} = m^{9/4} - \frac{3}{2}v^{1/2}\theta_2 - \mathcal{R}, \quad 0 \leq \mathcal{R} \leq \frac{3}{8}v^{-1/2}\theta_2^2.$$

*Proof.* Apply Taylor's theorem to  $x^{3/2}$  between  $v$  and  $v + \theta_2 = Y$ . Its second derivative is  $\frac{3}{4}x^{-1/2} \leq \frac{3}{4}v^{-1/2}$  on that interval, and is positive. Rearranging the Taylor formula gives the result.  $\square$

These identities alone do not supply cancellation. The restricted estimate from the earlier draft,

$$\left| \sum_{\substack{P < n \leq 2P \\ n \text{ odd}}} e\left(\frac{i}{2}n^{3/2} + \frac{j}{2}m^{3/2}\right) \right| \ll_\varepsilon P^{23/24+\varepsilon}, \quad |i| \leq 2P^{1/24}, \quad 1 \leq |j| \leq 2P^{1/24}, \quad (7.1)$$

now follows from Theorem 4.5 with  $k = 0$  and  $C = 2$ . The stronger general kernel target, with  $c_k(n) = \frac{3k}{4}n^{9/8}$ , is

$$K_k(P) = \sum_{\substack{P < n \leq 2P \\ n \text{ odd}}} e(c_k(n)\theta_2(n)), \quad |K_k(P)| \ll_\varepsilon P^{95/96+\varepsilon}, \quad 1 \leq k \leq P^{1/24}. \quad (7.2)$$

The indices  $i, j, k$  are integers. Estimate (7.1) is unconditional in this revision; (7.2), with its printed uniformity, remains an **open target**. A bare-kernel bound does not imply a mixed-mode estimate by itself. The exact families needed here are instead proved in Theorems 4.9 and 4.11. Lemma 4.4 handles half-integer coefficients  $j/2$  directly. This does not extend the frequency domain of the earlier general decorated lemma, whose separate applications remain unproved.

Short-interval variants require separate estimates. A bound for a transition set  $\Omega \subset (P, 2P]$  of the form  $|\Omega| \ll P^a$  implies only  $|\Omega \cap I| \leq \min(|I|, O(P^a))$ . It does not imply a bound proportional to  $|I|/P$ . Therefore the earlier localized-kernel threshold  $|I| \geq P^{29/48+\delta}$  is not asserted here.

## 7.1 An unconditional estimate averaged over a shift

**Proposition 7.4 (shift average).** Let  $L \geq 1$ , let  $A_1 < \dots < A_L$  obey  $|A_t - A_s| \geq a|t - s|$  with  $a > 0$ , and let  $x_1, \dots, x_L$  be real. For  $S_\lambda = \sum_{t=1}^L e(A_t\{x_t + \lambda\})$ ,

$$\left| \int_0^1 |S_\lambda|^2 d\lambda - L \right| \leq \frac{4}{\pi} \frac{L}{a} (1 + \log L). \quad (7.3)$$

For every  $0 < \eta < 1$ , outside a set of shifts of measure at most  $\eta$ ,

$$|S_\lambda| \leq \left[ \frac{L}{\eta} \left( 1 + \frac{4}{\pi a} (1 + \log L) \right) \right]^{1/2}.$$

*Proof.* Expanding the square gives the diagonal term  $L$ . For  $t \neq s$ , the integrand  $e(A_t\{x_t + \lambda\} - A_s\{x_s + \lambda\})$  is periodic in  $\lambda$ . Integrate over a period starting at one of its two fractional-part breakpoints. There is at most one other interior breakpoint, giving at most two intervals on which the phase is affine with slope  $A_t - A_s$ . On each interval the integral has modulus at most  $1/(\pi|A_t - A_s|)$ . Hence the off-diagonal contribution is bounded by

$$\frac{2}{\pi a} \sum_{t \neq s} \frac{1}{|t - s|} = \frac{4}{\pi a} \sum_{r=1}^{L-1} \frac{L-r}{r} \leq \frac{4}{\pi a} L(1 + \log L).$$

For  $L = 1$  the off-diagonal sum is empty. Markov's inequality applied to  $|S_\lambda|^2$  gives the second assertion.  $\square$

The estimate allows a small exceptional set of shifts. It gives no bound for a specified shift, including  $\lambda = 0$  in (7.2).

## 7.2 Counting the gap cells in a collision estimate

The following lemma repairs the cell-counting problem for a fixed curvature model. It does not assert that every decorated phase in the earlier kernel proof meets its hypotheses.

**Lemma 7.5 (curvature estimates over a partition).** Partition an interval of length at most  $P$  into interval cells, so that every subinterval of length  $L$  meets  $O(1 + \nu L)$  cells. On each cell let  $f$  be a real  $C^2$  function; its values and derivatives may jump at cell boundaries. Suppose that a continuous reference function  $\Lambda$  satisfies  $|\Lambda| \ll M$ , and that, for  $0 < s \leq 1$ , both sets

$$\{|\Lambda| \leq sM\}, \quad \{sM < |\Lambda| \leq 2sM\}$$

are unions of  $O(1)$  intervals, each set of total length  $O(Ps)$ . Assume

$$|f'' - \Lambda| \leq \rho M \quad \text{on every cell}, \quad 0 \leq \rho \leq 1/8, \quad 0 < M \leq 1, \quad MP^2 \geq 1.$$

All constants in these assumptions are fixed. Then, summing over integers or over odd integers with the same implicit-constant convention,

$$\left| \sum e(f(n)) \right| \ll PM^{1/2} + (1 + \nu P)M^{-1/2} + (P/M)^{1/3} + P\rho + 1. \quad (7.4)$$

*Proof.* Take  $\tau = \max(4\rho, (MP^2)^{-1/3})$ . If  $\tau$  is bounded below by a positive absolute constant, the terms  $P\rho + (P/M)^{1/3}$  already give the trivial bound  $O(P)$ . Otherwise discard  $\{|\Lambda| \leq \tau M\}$ , containing  $O(P\tau + 1)$  integers. Split the remainder into dyadic bands  $|\Lambda| \asymp sM$ , with  $\tau \ll s \ll 1$ , and the outer set  $|\Lambda| > M$  if necessary. Each band has total length  $O(Ps)$  and meets  $O(1 + \nu Ps)$  cells. On its intersection with a cell the curvature has one sign and size  $\asymp sM$ .

Apply the second-derivative estimate on every such intersection. The resulting costs at scale  $s$  are

$$O(PM^{1/2}s^{3/2} + M^{-1/2}s^{-1/2} + \nu PM^{-1/2}s^{1/2}).$$

The geometric sum over bands is  $O(PM^{1/2} + M^{-1/2}\tau^{-1/2} + \nu PM^{-1/2})$ . Our choice gives  $M^{-1/2}\tau^{-1/2} \leq (P/M)^{1/3}$  and  $P\tau \ll P\rho + (P/M)^{1/3}$ , proving (7.4). One may assign boundary integers to either adjacent cell and estimate that cell up to its endpoint.  $\square$

**Proposition 7.6 (the basic frozen-gap collision model).** Let  $u \geq 1/2$ ,  $1 \leq h \leq P^{1/8}$ ,  $uh \leq P^{1/2}$ , and let  $w > 0$  satisfy  $c_1 uh P^{-1/4} \leq w \leq c_2 uh P^{-1/4}$ , for fixed  $0 < c_1 < c_2$ . With  $A_h$  as in Lemma 4.4 and  $G(x) = \lfloor (x + 2h)^{3/2} - x^{3/2} \rfloor$ , put

$$f(x) = uA_h(x) + \frac{3u}{2}G(x)(x + 2h)^{3/4} + wx^{3/2}.$$

Then

$$\left| \sum_{\substack{P < n \leq 2P \\ n \text{ odd}}} e(f(n)) \right| \ll_{c_1, c_2} (uh)^{1/2} P^{5/8} + (h/u)^{1/2} P^{7/8} + (uh)^{-1/3} P^{7/12} + h + P^{1/2}/h + 1. \quad (7.5)$$

*Proof.* On a gap cell the frozen-coefficient derivative is

$$f''(x) = -\frac{9}{32}uG(x)(x + 2h)^{-5/4} + \frac{3}{4}wx^{-1/2} + uA_h''(x).$$

Use the reference function

$$\Lambda(x) = -\frac{27}{32}uhx^{-3/4} + \frac{3}{4}wx^{-1/2}, \quad M = uhP^{-3/4}.$$

The identities  $G(x) = 3h\sqrt{x} + O(1 + h^2P^{-1/2})$  and  $A_h'' = O(h^2P^{-7/4})$  show, as a comparison of values,

$$|f'' - \Lambda| \ll M(h/P + (hP^{1/2})^{-1}).$$

This does not differentiate  $G$ . The gap-cell partition has density parameter  $\nu \asymp hP^{-1/2}$ .

To verify the sublevel conditions, factor  $\Lambda(x) = \frac{3}{4}x^{-3/4}(wx^{1/4} - 9uh/8)$ . The expression in parentheses is strictly increasing, with derivative  $\asymp uh/P$  in the stated collision range. Hence  $|\Lambda| \leq sM$  occupies length  $O(PS)$ . Moreover  $\Lambda'$  has at most one zero, so the sublevel sets and dyadic bands have a bounded number of components. Apply Lemma 7.5 with  $\rho = O(h/P + (hP^{1/2})^{-1}) = o(1)$ . Its terms give (7.5); the extra  $M^{-1/2}$  is absorbed by  $(h/u)^{1/2}P^{7/8}$ .  $\square$

For integer  $w$  in the collision range,  $w \geq 1$  implies  $uh \gg P^{1/4}$ . Formula (7.5) is then  $O(P^{7/8})$ : its second term is  $h(uh)^{-1/2}P^{7/8} \ll P^{7/8}$ , and the remaining terms are no larger. This verifies that the basic integer collision band leaves a power saving.

Thus the transition cost need not be paid independently on every gap cell: a global sublevel argument counts how many cells can meet each curvature band. To use this in the complete kernel proof one must still verify the reference-function hypotheses, partition density, and weighted sums for all additional Fourier modes and decorations. Neither (7.4) nor (7.5) supplies that verification, and neither rescales a transition-set bound by an arbitrary short-interval length.

## 8. Evidence, availability, and remaining work

The status of the principal statements is as follows.

| Statement                                                | Warrant in this version                                                            |
|----------------------------------------------------------|------------------------------------------------------------------------------------|
| Envelope, branch identity, minimal words, floor defects  | Exact proofs in Sections 2, 5, and 7                                               |
| Word counts, certificate lengths, and the survivor decay | Shared with the Collatz literature: A076227, A100982, A020914; no priority claimed |
| Certificate density $3/4$                                | Classical analytic proof in Theorem 3.1                                            |
| Correlation-to-count and Fourier-to-parity implications  | Proofs in Propositions 4.1 and 4.2                                                 |
| Certificate density $13/16$                              | Unconditional: Corollary 4.6 and Theorem 5.2                                       |
| Certificate subfamily density $27/32$                    | Unconditional: Corollary 4.10 and Theorem 5.3                                      |

| Statement                                                             | Warrant in this version                                                     |
|-----------------------------------------------------------------------|-----------------------------------------------------------------------------|
| Full five-step certificate density 7/8                                | Unconditional: Corollary 4.12, Theorem 5.4, and Appendices A–C              |
| Density-one finite certificates                                       | Conditional on FD; no depth uniformity assumed                              |
| Oscillating prefactor $\psi$ , its jump spectrum and Fourier identity | Verified computation in Section 6; the existence of the limit is not proved |
| Finite combinatorics of Sections 5–6 and the chain of Theorem 6.1     | Machine-checked; modules listed below                                       |
| Restricted mixed sums                                                 | Theorems 4.5, 4.9, and 4.11                                                 |
| General decorated kernel and short-interval extension                 | Unproved; no numerical threshold certified                                  |
| Basic frozen-gap collision model                                      | Proposition 7.6; does not certify every decorated phase                     |
| Shift-averaged estimate                                               | Direct integral proof in Proposition 7.4                                    |

Supporting software is available in the Balanced Ternary Mathematical Laboratory repository, <https://github.com/sneakyweasel/bt1ab>. The accompanying source package contains this manuscript, its LaTeX build, and an exact-arithmetic validation script. That script checks the finite certificate list, the indicator identity on a finite census, and selected algebraic inequalities. A second exact-arithmetic script checks the differenced identity, carry expansion, and exponent comparisons used in the four-step repairs. Further exact scripts check the fifth-letter centering identity, the D2 floor reduction, the signed-wave and master inventories, the mixed curvature coefficients, and all displayed exponent budgets. The source package includes these scripts and their results. Those checks support reproducibility; they are not proofs of asymptotic cancellation.

The Lean modules that check statements of this manuscript, module by module, all of them under `formal/Problems/Juggler/`:

- **PaperBCertificates**: Lemma 5.1, the minimal certificates through length five.
- **PaperBCertificateLengths**: a minimal certificate of length  $L$  exists exactly when a power of three lies in  $[2^{L-1}, 2^L)$ , its odd count is then forced and its last letter is  $E$ .
- **PaperBFiveStepDensity**: the five cylinders partition the certified words at depth five and the count is the sum of the five class counts behind Theorems 5.2–5.4; depth six adds no minimal certificate and depth seven adds three.
- **PaperBCertificateRecursion**: the survivors of length  $d + 1$  and the words contracting first at  $d + 1$  together number  $2N_d$ , the telescoped tail form of that identity, and the doubling at the free lengths.
- **PaperBSurvivorDecay**:  $N_d/2^d \leq \vartheta(p)^d$ , the upper bound of Theorem D with its exact exponent, and  $N_d/2^d \rightarrow 0$ .
- **PaperBChernoff**, **PaperBMarkov**, **PaperBDensity**: the chain of Theorem 6.1 with Hypothesis FD carried as a hypothesis.
- **PaperBThreshold**: the threshold algebra of Remark 6.2, that  $q_d$  saturates the step and  $d \geq 4$  is the first usable depth.
- **PaperBTilt**: the mean-zero tilt and the closed form of  $\varrho$ .
- **PaperBBarrierStep**: when the barrier does not rise every survivor keeps both extensions, and when it rises an extension dies exactly for a word on the barrier taking the even letter.
- **PaperBJumpTransposition**: exchanging two adjacent barrier letters costs exactly the mass on the barrier, the elementary fact under the discontinuities of  $\psi$ .

The declarations audited for each, in the same order, are those whose axioms are printed by the published artifact, `formal/AxiomCheckPaperBPublished.lean`:

`PaperBCertificates`                      `lemma51`   `lemma51_complete`

|                            |                                                                                                         |
|----------------------------|---------------------------------------------------------------------------------------------------------|
| PaperBCertificateLengths   | minimalCert_exists_iff certWindow_unique<br>minimalCert_concat_even                                     |
| PaperBFiveStepDensity      | five_cylinders_cover certifiedCount_five_eq<br>no_minimal_certificate_six<br>minimal_certificates_seven |
| PaperBCertificateRecursion | neverNegCount_add_minimalCertCount<br>minimalCert_tail_eq<br>neverNegCount_succ_of_window_empty         |
| PaperBSurvivorDecay        | neverNegCount_div_pow_le_theta<br>neverNegCount_div_pow_tendsto_zero                                    |
| PaperBChernoff             | theta_lt_one klDiv_pos                                                                                  |
| PaperBMarkov               | sum_choose_mul_pow chernoff_density<br>tilt_gives_theta                                                 |
| PaperBDensity              | density_of_finite_union<br>exceptional_density_zero                                                     |
| PaperBThreshold            | breakEven_saturates first_usable_depth                                                                  |
| PaperBTilt                 | lamStar_mean_zero rho_closed_form                                                                       |
| PaperBBarrierStep          | survives_succ_of_no_rise dies_iff_on_barrier                                                            |
| PaperBJumpTransposition    | stepRise_stepFlat_eq_add_barrierMass                                                                    |

Its recorded output shows no axiom beyond Mathlib’s three, `propext`, `Classical.choice` and `Quot.sound`, on any line; `klDiv_nonneg`, cited in the proof of Theorem 6.1 for contrast, is audited there too. All are checked by Lean’s kernel, with no `native_decide`. This version claims no Lean formalization of its analytic estimates — the exponential sums of Sections 3, 4 and 7 and Appendices A–C — nor of the complete manuscript. Earlier theorem numbers are superseded and should not be used as citations to unconditional nested estimates. Companion texts that rely on those estimates require their own review.

The full five-step certificate class proved here has density 7/8. The argument does not extend its error estimates proportionally to an arbitrary short interval, and it supplies no universal termination conclusion. A proof of all fixed-depth fair-share counts would imply density-one certificates by Theorem 6.1; that additional hypothesis remains open.

## Appendix A. Floor reductions and the carry inventory

We collect the estimates used by the OOOEE argument. All counts of exceptional integers use the original gap runs, before any later frequency partition. Constants may depend on fixed comparability constants. Every displayed estimate is uniform for sufficiently large  $P$ ; bounded smaller  $P$  is absorbed into the implied constants.

### A.1 A reduction for a differenced floor factor

**Lemma A.1 (fixed-label floor reduction).** The definitions below lead to (A.9), with the stated error and coefficient variation bounds.

In Appendices A–C, a subscript on  $\Delta$  denotes the actual translation:  $\Delta_d f(x) = f(x + d) - f(x)$ . Thus a half-shift  $h$  uses  $\Delta_{2h}$ . This differs from the local half-shift convention  $\Delta_h f(x) = f(x + 2h) - f(x)$  in Section 4.2. All constants below are absolute, or depend on fixed constants in comparability bounds. Let  $P$  tend to infinity, with integer parameters

$$1 \leq k, h_1, h_2 \leq P^{1/24}, \quad 1 \leq h \leq P^{1/8}.$$

Put  $X(x) = x^{3/2}$ ,  $m(n) = \text{floor}(X(n))$ , and  $c(x) = 3k x^{9/8} / 4$ . Work on  $(P, 2P]$ , extending the definitions to  $(P, 3P]$  to accommodate shifts. Fixed positive multiples of the upper parameter bounds

are allowed, with constants depending on those multiples. All estimates also hold after restriction to odd integers.

Partition by the integer parts of

$$X(x + 2h_1) - X(x), \quad X(x + 2h_2) - X(x), \quad X(x + 2h_1 + 2h_2) - X(x).$$

There are

$$D \ll (h_1 + h_2)P^{1/2}$$

runs, and an interval of length  $L$  meets  $O(1 + (h_1 + h_2)LP^{-1/2})$  runs.

On each run freeze a carry label and its integer offsets  $\beta_1, \beta_2, \beta_{12}$ . Assume  $\beta_i$  is comparable to  $h_i P^{1/2}$ , for  $i=1,2$ , and  $j = \beta_{12} - \beta_1 - \beta_2$  is a bounded integer, with  $|j| \leq 2$  for the admissible branches used here. Define, on that run,

$$F(t) = (t + \beta_1 + \beta_2 + j)^{3/2} - (t + \beta_1)^{3/2} - (t + \beta_2)^{3/2} + t^{3/2},$$

$$G(x) = F(X(x)), \quad A(n) = \lfloor F(m(n)) \rfloor.$$

The offsets may change between runs. The theorem treats any such piecewise extension; no derivative crosses a run boundary. There are only finitely many carry labels, so bounds for their positive error terms can be summed. Reassembling oscillatory sums with the actual carry indicators is a further step, not an assumption that a branch extension is the physical orbit everywhere.

On an interval  $I$  of length  $L$ , suppose  $G$  is  $C^1$  and  $a \leq |G'| \leq C a$ , with  $0 < a \leq 1$ . Its derivative has constant sign. For  $0 < \delta \leq 1/2$ , the number of integer  $n$  in  $I$  with  $\text{distance}(G(n), Z) \leq \delta$  is

$$\ll \delta(L + a^{-1}) + aL + 1. \quad (\text{A.1})$$

Indeed  $G(I)$  meets  $O(aL + 1)$  integer neighborhoods. Each inverse image is an interval of length  $O(\delta/a)$ , containing  $O(\delta/a + 1)$  integers. This includes endpoints and exact integer values.

Let

$$E_Q(t) = \min(1, (Q\|t\|)^{-1}), \quad E_Q(t) = 1 \quad (t \in \mathbb{Z}).$$

Splitting the distances to  $Z$  into dyadic bands gives

$$\sum_{n \in I} E_Q(G(n)) \ll \frac{L + a^{-1}}{Q} \log(2Q) + aL + 1. \quad (\text{A.2})$$

The additive lattice-count cost is multiplied by a convergent geometric sum, not by the number of bands.

Over  $D$  disjoint runs with a common comparable derivative scale, these become

$$\#\{\|G(n)\| \leq \delta\} \ll \delta(P + D/a) + Pa + D,$$

$$\sum E_Q(G(n)) \ll \frac{(P + D/a) \log(2Q)}{Q} + Pa + D. \quad (\text{A.3})$$

Runs of different offset types are bounded separately. No equidistribution assumption, random model, or rescaling of a global exceptional set is used.

Set  $p = h_1 h_2$ . The zero-offset part has the exact integral representation

$$F_0(t) = \frac{3}{4} \int_0^{\beta_1} \int_0^{\beta_2} (t + s + r)^{-1/2} dr ds.$$

Consequently, with the  $\beta$  values fixed when differentiating,

$$G'_0(x) = -\frac{9}{16} \beta_1 \beta_2 x^{-7/4} (1 + o(1)),$$

$$G''_0(x) = \frac{63}{64} \beta_1 \beta_2 x^{-11/4} (1 + o(1)).$$

The errors are uniform because  $(\beta_1 + \beta_2)/X = o(1)$ .

The offset contribution is

$$\frac{3j}{2} \int_0^1 (t + \beta_1 + \beta_2 + sj)^{1/2} ds.$$

After composition with  $X$  its first derivative is  $(9j/8) x^{-1/4} (1+o(1))$ . For  $j$  nonzero it dominates the zero-offset derivative, since  $p P^{-1/2} = o(1)$ . Thus, on every run,

$$|G'| \asymp \begin{cases} pP^{-3/4}, & j = 0, \\ P^{-1/4}, & j \neq 0, \end{cases} \quad |G''| \ll |j|P^{-5/4} + pP^{-7/4}. \quad (\text{A.4})$$

Also  $|G|$  is  $O(|j|P^{3/4} + pP^{1/4})$ . These are derivatives of frozen-offset functions, not derivatives of the approximations  $\beta_i$  approximately  $3h_i \sqrt{x}$ .

Take  $Q = \text{floor}(P^{5/16})$ . For  $j=0$ , the potentially largest term in (A.3) is

$$\frac{D}{Qa} \ll \frac{h_1 + h_2}{h_1 h_2} P^{15/16} \ll P^{15/16}.$$

The other terms are  $P^{11/16}$ ,  $pP^{1/4}$ , and  $D$ . For  $j$  nonzero, (A.3) is  $O(P^{3/4} \log P)$ . It follows that, summed over all runs of either type,

$$\sum E_Q(G(n)) \ll P^{15/16} \log P. \quad (\text{A.5})$$

The replacement of  $m$  by  $X$  is used only inside the floor, after counting its exceptional set. The mean value theorem gives

$$|F(m(n)) - G(n)| \ll |j|P^{-3/4} + pP^{-5/4} =: \delta.$$

A floor mismatch requires  $G(n)$  to lie within  $\delta$  of an integer. For either derivative scale in (A.4),  $\delta/a = O(P^{-1/2})$ . The first line of (A.3) therefore proves

$$\#\{n : A(n) \neq \lfloor G(n) \rfloor\} \ll P^{3/4}. \quad (\text{A.6})$$

This remains useful when  $c$  is large: changing a unit exponential costs at most two on the exceptional set. A large coefficient is never multiplied by an uncontrolled pointwise floor error.

Exclude for the moment  $n$  whose interval from  $n$  to  $n+2h$  crosses an original run boundary. There are  $O(hD)$  such integers.

Within a run,  $G$  is monotone. If  $\text{floor}(G(n+2h))$  differs from  $\text{floor}(G(n))$ , this interval contains an integer-level crossing of  $G$ . There are  $O(aL + 1)$  levels on a run of length  $L$ , and any crossing belongs to at most  $O(h)$  such integer starting intervals. Including the original run boundaries gives

$$\#\{\text{floor change or run crossing}\} \ll h(Pa + D) \ll P^{7/8}. \quad (\text{A.7})$$

For zero offset the bound is even smaller; the uniform largest term is  $hP^{3/4}$  from nonzero offsets.

Write  $B(x) = c(x+2h) - c(x)$ . Using (A.6) at both endpoints and then (A.7), we obtain the following statement in the sum of absolute errors:

$$e(-\Delta_{2h}(cA)(n)) = e(-B(n)\lfloor G(n) \rfloor) + \mathcal{E}_n, \quad \sum |\mathcal{E}_n| \ll P^{7/8}. \quad (\text{A.8})$$

Both expressions in this comparison are unimodular. The second expression is defined on every run, including the exceptional starting points; no irregularly punctured domain is passed to a derivative test.

The floor-crossing argument also explains why a small drift is insufficient by itself. For comparison, on zero-offset runs  $|\Delta G|$  is  $O(hP^{-3/4})$ ; at the carry cutoff  $J = \text{floor}(P^{1/8})$ ,  $J|\Delta G| = o(1)$ . Therefore the distance majorant  $E_J(\Delta G)$  equals one for sufficiently large  $P$ . The standard sawtooth Fourier approximation has an order-one endpoint error there as well. Small drift alone cannot justify the  $O(P/J)$  charge for this individual layer. The exact carry combination may cancel; (A.7) preserves that cancellation by counting the floor change before expansion.

The coefficient  $B$  is increasing, with

$$B \asymp khP^{1/8}, \quad B' \asymp khP^{-7/8}, \quad |B''| \ll khP^{-15/8}.$$

Refine the original runs by  $N = \text{floor}(B)$ , and put  $\beta = B - N$ . The extra number of cuts is  $O(1 + khP^{1/8})$ . The exact identity

$$e(B\{G\}) = e(NG)e(\beta\{G\})$$

centers the frequency before truncation.

For  $0 \leq \beta \leq 1$  let

$$a_r(\beta) = \int_0^1 e((\beta - r)t) dt.$$

By Lemma 4.7, or its complex conjugate,

$$e(\beta\{G\}) = \sum_{|r| \leq Q} a_r(\beta)e(rG) + O(E_Q(G)).$$

The integral definition includes the removable singularity. Both  $|a_r|$  and  $|a_r'|$  are  $O(1/(1+|r|))$ . On each refined interval,  $\beta$  is monotone and has variation at most one, so

$$\sum_{|r| \leq Q} (\|a_r(\beta)\|_\infty + \text{TV}(a_r(\beta))) \ll \log(2Q).$$

Combining this expansion with (A.5) and (A.8) proves the fixed-label floor reduction:

$$\begin{aligned} e(-\Delta_{2h}(cA)(n)) &= \sum_{|r| \leq Q} a_r(\beta(n))e(\Phi_r(n)) + \mathcal{R}_n, \\ \Phi_r(x) &= (N + r - B(x))G(x), \\ \sum_{P < n \leq 2P} |\mathcal{R}_n| &\ll P^{15/16} \log P. \end{aligned} \tag{A.9}$$

$N$  and all offsets are fixed only within the refined interval. The error sum is charged once on the original runs; it is not multiplied by the number of frequency windows. This is valid with any additional weight of modulus at most one. A fixed number of such floor factors can be expanded by telescoping, at a fixed extra power of  $\log P$  and with the same power saving.

This replacement uses the bounded residual  $\beta$ , so it has no unaccounted continuous-part tail between  $Q$  and  $P^{1/2}$ .

Differentiate with  $N$  and  $r$  fixed:

$$\Phi_r'' = (N + r - B)G'' - 2B'G' - B''G.$$

Since  $N + r - B = \beta$ , (A.4) yields

$$|\Phi_r''| \ll (Q + 1)(|j|P^{-5/4} + pP^{-7/4}) + kh(|j|P^{-9/8} + pP^{-13/8}). \tag{A.10}$$

For an undisturbed wave curvature  $M = uhP^{-3/4}$ ,  $u \geq 1/2$ , the four exponent bounds for this ratio are respectively

$$P^{-3/16}, \quad P^{-29/48}, \quad P^{-1/3}, \quad P^{-3/4}.$$

Thus  $|\Phi_r''| = O(P^{-3/16}M)$ , uniformly.

Where a total phase already has curvature comparable to  $M$ , the extra endpoint cost in the second-derivative estimate is

$$(D + 1 + khP^{1/8})M^{-1/2} \ll (h_1 + h_2)(uh)^{-1/2}P^{7/8} + k(h/u)^{1/2}P^{1/2} + (uh)^{-1/2}P^{3/8}. \quad (\text{A.11})$$

Its largest uniform exponent is  $11/12$ , below  $15/16$ . Partial summation uses the coefficient-variation bound proved above.

This comparison is not a declaration that an arbitrary sum of signed waves has curvature comparable to  $M$ . At a curvature cancellation, the reference-function and sublevel hypotheses must still be verified for the complete phase.

## A.2 Joint first-floor carries

**Lemma A.2 (carry arcs and global positive errors).** Let  $1 \leq h \leq P^{1/12}$ , and let the number of shifts below be bounded by a fixed constant.

There are finitely many first-floor evaluations in the main wave and the D1 terms. All have the form  $m(n+s)$ , with  $s$  an even nonnegative integer  $O_{C,M}(h + P^{1/24})$ . Use the single variable  $\theta = \{X(n)\}$  and

$$m(n+s) - m(n) = b_s + \mathbf{1}_{\theta \geq 1-\delta_s}, \quad b_s = \lfloor X(n+s) - X(n) \rfloor, \quad \delta_s = \{X(n+s) - X(n)\}.$$

The convention when  $\delta_s = 0$  is that this indicator is zero for  $0 \leq \theta < 1$ .

Partition by the integer gaps  $b_s$  and any input or coefficient windows already required by the argument. There are  $O(1 + \sum_s s\sqrt{P})$  original gap runs. Their local count in an interval of length  $L$  is  $O(1 + \sum_s sLP^{-1/2})$ . On each gap run each  $\delta_s$  is monotone and varies by at most one. Sort their endpoints  $1-\delta_s$ ; two distinct endpoints cross at most once on such a run, because the difference of the corresponding real gaps is monotone. Equal shifts give identical endpoints. This adds only  $O_{C,M}(1)$  cuts per cell.

A joint carry pattern is now a fixed finite union of intervals of  $\theta$ , with endpoints  $0, 1$ , or  $1-\delta_s$ . Its Fourier expansion truncated at  $R = \text{floor}(P^{1/4})$  has: - a zero-mode coefficient with sup norm plus variation  $O_{C,M}(1)$ ; - endpoint modes with coefficients  $O_{C,M}(1/|v|)$ ,  $0 < |v| \leq R$ , multiplying  $e(vX(n+s))$  for one of the allowed  $s$  ( $s=0$  included); - an absolute remainder bounded by a constant times the sum of  $E_R(X(n+s))$  over these finitely many endpoints, where  $E_R(z) = \min(1, 1/(R||z||))$ , with value one at integers.

This is the usual interval Fourier expansion: a moving endpoint  $u = 1-\delta_s$  supplies  $e(-v u)$ , and, since  $b_s$  is integer,  $e(vX(n))e(-v(1-\delta_s)) = e(vX(n+s))$ . Thus the moving endpoint belongs in the phase, not in an unbounded-variation coefficient. Zero-length intervals and points at endpoints are covered by the positive remainder.

Here is the global bound needed for that remainder:

$$\sum_{P < n \leq 2P} E_R(X(n+s)) \ll_{C,M} P^{5/6} \log(2P). \quad (\text{A.12})$$

It holds also over odd integers or a subinterval. To verify it, the second-derivative test for  $vX(x+s)$  gives  $O(|v|^{1/2}P^{3/4} + |v|^{-1/2}P^{1/4})$ . The interval-discrepancy inequality with auxiliary cutoff  $K = \text{floor}(P^{1/6})$  then gives discrepancy  $O(P^{5/6})$ . Dyadic bands for the distance to an integer give  $O(P \log(2R)/R + P^{5/6})$ , which implies (A.12). The remainder is charged once over the common input interval. It is not multiplied by the cell count. Products with the other finite Fourier expansions introduce at most a fixed logarithmic factor.

### A.3 Positive errors for the second-level inventory

Let  $X = x^{3/2}$ ,  $m = \lfloor X \rfloor$ ,  $Y = m^{3/2}$ , and put  $d_i = 2h_i$ ,  $W_i = \Delta_i Y$ ,  $D = \Delta_1 \Delta_2 Y$ , where  $1 \leq h_1 \leq P^{1/48}$ ,  $1 \leq h_2 \leq P^{1/24}$ , and  $p = h_1 h_2$ . Here  $\Delta_i f(x) = f(x + d_i) - f(x)$ . Even translations by  $O(P^{1/24})$  are allowed. **Lemma A.3 (positive inventory errors).**

Set  $J = \lfloor P^{1/24} \rfloor$ , and use  $E_J(z) = \min(1, (J\|z\|)^{-1})$ , with value one at integers. For every argument occurring in the carry expansion below,

$$\sum_{P < n \leq 2P, n \text{ odd}} E_J(Z(n)) \ll P^{23/24} (\log(2P))^C, \quad (\text{A.13})$$

where

$$Z \in \{Y(n), Y(n + d_1), Y(n + d_2), Y(n + d_1 + d_2), W_1(n), W_2(n), W_1(n + d_2), D(n)\}.$$

The constant C is absolute. This is an unweighted positive bound; it does not assume that an anchor is still present.

For Y at any of these even translations, Theorem 4.5, with modes  $(i, j, k) = (0, 2q, 0)$ , gives  $O(P^{23/24})$  for  $1 \leq |q| \leq J$ . The discrepancy inequality at cutoff J, followed by dyadic distance bands, proves (A.13). The same argument for W uses Lemma 4.4 with  $i=k=0$  and  $j=2q$ . Its bound is  $O(P^{7/8}(1 + h_i^{1/2})) = O(P^{43/48})$ , which also fits (A.13). The proofs of those results work on any subinterval of  $[P, 3P]$ : their length, gap-cell, and positive-error bounds are all in the ambient scale P. No proportional rescaling of an exceptional set is used for the translated intervals here.

For D, a separate elementary argument is needed. Put

$$j(n) = \Delta_1 \Delta_2 m(n), \quad -1 \leq j(n) \leq 2.$$

The exact frozen-branch integral formulas imply the global value approximation

$$D(n) = g_j(n) + O((h_1 + h_2)P^{-1/4}), \quad g_j(x) = \frac{3}{2}jx^{3/4} + \frac{27}{4}px^{1/4}. \quad (\text{A.14})$$

To verify the error, the first-floor offsets satisfy  $\beta_i = 3h_i \sqrt{n} + O(1)$ , hence  $\beta_1 \beta_2 = 9pn + O((h_1 + h_2)\sqrt{P} + 1)$ . In the zero-offset integral, replacing its argument by  $n^{3/2}$  has relative error  $O((h_1 + h_2)/P)$ ; in the offset integral the corresponding absolute error is  $O((h_1 + h_2)P^{-1/4})$ . The resulting extra zero-offset error  $O(p(h_1 + h_2)P^{-3/4})$  is smaller than the displayed error. This is a comparison of values, not of frozen derivatives.

The error in (A.14) is  $O(P^{-5/24})$ , so J times that error tends to zero. For  $|u-v| \leq 1/(2J)$ , one has  $E_J(u) \leq 2E_J(v)$ . On a branch  $j=0$ ,  $g_j$  has derivative comparable to  $pP^{-3/4}$ . On a nonzero branch it has derivative comparable to  $P^{-1/4}$ , of the sign of j. Both are monotone slow variables on the whole dyadic block. The monotone counting lemma (A.2) in Appendix A.1 gives

$$\sum E_J(g_j(n)) \ll (P + a_j^{-1}) \log(2J)/J + Pa_j + 1, \quad a_0 \asymp pP^{-3/4}, \quad a_j \asymp P^{-1/4} \ (j \neq 0).$$

Every term is  $O(P^{23/24} \log P)$ . Bound a branch-restricted positive sum by this full-block sum and add the four possible j values. This proves (A.13) for D without multiplying a floor-error estimate by the number of original gap runs.

### A.4 Exact centered master inventory

Let  $c(x) = 3kx^{9/8}/4$ ,  $c_{11}(x) = c(x + d_1 + d_2)$ ,  $\vartheta = \{Y\}$ , and  $1 \leq k \leq CP^{1/24}$ . **Lemma A.4 (master identity and coefficient mass).**

Define the binary carry  $C(A, B) = \{A\} + \{B\} - \{A + B\}$ , and set

$$\kappa_1 = C(Y, W_1), \quad \kappa_2 = C(Y, W_2), \quad \kappa_1^+ = C(Y(n + d_2), W_1(n + d_2)), \quad \kappa_* = C(W_1, D).$$

Let

$$B_2(x) = \Delta_2 c(x + d_1), \quad B_1(x) = \Delta_1 c(x + d_2), \quad N_i(x) = \lfloor B_i(x) \rfloor.$$

The product rule and the exact carry identity give

$$\begin{aligned} \Delta_1 \Delta_2 (c\vartheta) &= (\Delta_1 \Delta_2 c)\vartheta + B_2(\{W_1\} - \kappa_1) + B_1(\{W_2\} - \kappa_2) \\ &\quad + c_{11}(\{D\} - \kappa_* - \kappa_1^+ + \kappa_1). \end{aligned} \quad (\text{A.15})$$

With  $\Pi = kh_1h_2$ , the first term has total deletion cost  $O(\Pi P^{1/8}) = O(P^{11/48})$ .

For each factor  $e(B_i \{W\})$ , center at the integer  $N_i$ :

$$e(B_i \{W\}) = e(N_i W) e(\beta_i \{W\}), \quad \beta_i = B_i - N_i \in [0, 1].$$

Here  $\beta_i$  denotes a coefficient residual only; it is not a first-floor offset used in a branch formula. Expand the last factor at the short cutoff  $J$ :

$$e(\beta_i \{W\}) = \sum_{|r| \leq J} a_r(\beta_i) e(rW) + O(E_J(W)), \quad a_r(\beta) = \int_0^1 e((\beta - r)y) dy. \quad (\text{A.16})$$

Its coefficients satisfy  $|a_r| \ll 1/(1+|r|)$ , including removable singularities. The error is covered by (A.13), independently of the large center  $N_i$ . There is no uncentered large-coefficient tail to discard.

For each of the five carry exponentials in (A.15), use exactly

$$e(g\kappa) = 1 + \kappa(e(g) - 1).$$

Expand the factor  $e(g)-1$  into two terms and move  $e(g)$  into the smooth phase. Expand kappa additively using its three unit fractional parts, each at cutoff  $J$  by Lemma 4.3. The only arguments are those in (A.13), because

$$Y + W_i = Y(n + d_i), \quad W_1 + D = W_1(n + d_2), \quad Y(n + d_2) + W_1(n + d_2) = Y(n + d_1 + d_2).$$

The smooth phases introduced this way are bounded integer combinations of  $B_1, B_2, c_{11}$ .

There are two centered Fourier layers and at most five carry layers. Their total coefficient mass is  $O((\log(2P))^7)$ ; a larger fixed logarithmic power covers all errors. Indeed, pointwise truncated factors have  $O(\log P)$  bounds, so telescoping products multiplies each error in (A.13) only by a fixed logarithmic power.

The  $N_i$  windows are retained globally as functions of  $x$ . Both  $B_i$  are increasing, with

$$B_i \asymp kh_i P^{1/8}, \quad B'_i \asymp kh_i P^{-7/8}.$$

Their common partition has local count  $O(1 + k(h_1 + h_2)LP^{-7/8})$  and total count  $O(P^{5/24})$ . On each cell the residuals  $\beta_i$  are monotone in  $[0,1]$ , so the same logarithmic bound holds for summed coefficient sup norms plus variations.

For arbitrary four corner coefficients, the exact identity is

$$\sum_{d \in \{0, d_1, d_2, d_1 + d_2\}} q_d Y(n + d) = tY + (q_{d_1} + q_{d_1 + d_2})W_1 + (q_{d_2} + q_{d_1 + d_2})W_2 + q_{d_1 + d_2}D, \quad t = \sum_d q_d. \quad (\text{A.17})$$

Use also  $W_1(n + d_2) = W_1 + D$ .

Consequently (A.15)–(A.16) reduce the sum with phase  $\Delta_1 \Delta_2 (c\vartheta)$ , with absolute error  $O(P^{23/24} \log^C(P))$ , to a finite weighted family with phases

$$c_{11}\{D\} + tY + (N_2 + u)W_1 + (N_1 + v)W_2 + qD + \phi. \quad (\text{A.18})$$

Here  $t, u, v, q$  are fixed integers for each Fourier multi-index,

$$|t|, |u|, |v|, |q| \leq 6J,$$

and  $\phi$  is a bounded integer combination of  $B_1, B_2, c_{11}$ . In particular

$$|\phi''| \ll kP^{-7/8}, \quad |\phi'''| \ll kP^{-15/8}. \quad (\text{A.19})$$

The two centered residual frequencies account for the extra unit in the bound  $6J$ ; each carry layer contributes at most one mode. All signs and zero coordinates are included.

More precisely there are nonnegative numbers  $\rho_\nu$ , one for each multi-index, with  $\sum \rho_\nu = O(\log^7(P))$ , such that its amplitude divided by  $\rho_\nu$  has bounded sup norm plus variation on each  $N_1, N_2$  cell. This follows by multiplying the harmonic coefficient bounds above and applying the product variation inequality. Thus it suffices to prove a uniform bound for each normalized amplitude; no estimate is multiplied by the number of frequency windows.

The first-difference coefficients satisfy

$$|(N_2 + u)h_1| + |(N_1 + v)h_2| \ll \Pi P^{1/8} + J(h_1 + h_2) \ll P^{11/48}, \quad (\text{A.20})$$

well within the widened  $P^{1/2}$  budget.

## Appendix B. A wave-bearing estimate

The following estimate is used only for the family that its statement specifies. In Appendix C the total  $Y$  frequency is a nonzero integer. Allowing half-integer frequencies here also covers the elementary main-wave argument.

**Theorem B.1 (nonzero total frequency).**

Let  $e(v) = \exp(2\pi i v)$ , and put

$$X(x) = x^{3/2}, \quad m(n) = \lfloor X(n) \rfloor, \quad Y(n) = m(n)^{3/2}, \quad \Delta_d f(x) = f(x+d) - f(x).$$

All functions are defined on  $[P, 3P]$ , with harmless further extension at its upper endpoint if needed. Constants  $C$  and  $M$  are fixed. All estimates concern sufficiently large  $P$  depending only on them. Let  $t$  be real and fixed throughout the sum, with

$$\frac{1}{2} \leq |t| \leq CP^{1/24}.$$

There are at most  $M$  first-difference terms, with even shifts  $e_j \geq 0$  and positive integer half-shifts  $r_j$  satisfying

$$e_j \leq CP^{1/24}, \quad 1 \leq r_j \leq CP^{1/24}, \quad |a_j(x)|r_j \leq CP^{1/2}. \quad (\text{B.1})$$

The real coefficients  $a_j$  are constant separately on the cells of an interval partition  $\mathcal{P}$ . Any interval of length  $L$  meets at most  $C(1+L P^{-11/24})$  cells of this partition. In particular, its total cell count is  $O_C(P^{13/24})$ . The real function  $\phi$  is  $C^3$  on each cell and satisfies

$$|\phi'''(x)| \leq CP^{-13/12}. \quad (\text{B.2})$$

No continuity across cell boundaries is required. The complex amplitude  $\eta$  satisfies

$$\|\eta\|_{\infty, I} + \text{TV}_I(\eta) \leq C$$

on each cell  $I$ . Endpoint values are assigned consistently; their possible exceptional contribution is covered by the cell count.

We also allow at most  $M$  terms  $-\sigma_\ell c_\ell A_\ell$ , where  $\sigma$  belongs to  $\{-1, 0, 1\}$ . These are the fixed-label D2 objects of the Appendix A.1, with

$$1 \leq k_\ell, h_{\ell,1}, h_{\ell,2} \leq CP^{1/24}, \quad c_\ell(x) = \frac{3}{4}k_\ell(x + v_\ell)^{9/8}.$$

The integers  $k$  and  $h$  are positive. The base shift  $s_\ell$  and the coefficient shift  $v_\ell$  are nonnegative even integers  $O_C(P^{1/24})$ . To specify the object unambiguously, partition by the integer parts of

$$X(x + s_\ell + 2h_{\ell,i}) - X(x + s_\ell), \quad X(x + s_\ell + 2h_{\ell,1} + 2h_{\ell,2}) - X(x + s_\ell).$$

On every such original run choose fixed admissible carry offsets  $\beta_1, \beta_2, \beta_{12}$ , with  $\beta_i$  comparable to  $h_i \sqrt{P}$  and  $j = \beta_{12} - \beta_1 - \beta_2$  a bounded integer ( $|j| \leq 2$  suffices). Set

$$\begin{aligned} F_\ell(z) &= (z + \beta_1 + \beta_2 + j)^{3/2} - (z + \beta_1)^{3/2} - (z + \beta_2)^{3/2} + z^{3/2}, \\ G_\ell(x) &= F_\ell(X(x + s_\ell)), \\ A_\ell(n) &= \lfloor F_\ell(m(n + s_\ell)) \rfloor. \end{aligned}$$

The version  $A_\ell(n) = \lfloor G_\ell(n) \rfloor$  is also allowed.

The partition  $\mathcal{P}$  refines these original runs. Further refinement does not change their chosen labels: it cannot introduce new arbitrary jumps in  $F_\ell$  or  $G_\ell$ . All D2 counting errors below are evaluated on the original runs. These floors are not first partitioned into their integer level sets.

Under these hypotheses the theorem is

$$\begin{aligned} U(P) &= \sum_{\substack{P < n \leq 2P \\ n \text{ odd}}} \eta(n) e\left(tY(n) + \sum_j a_j(n) \Delta_{2r_j} Y(n + e_j) - \sum_\ell \sigma_\ell c_\ell(n) A_\ell(n) + \phi(n)\right), \\ |U(P)| &\ll_{C,M,\varepsilon} P^{31/32+\varepsilon}. \end{aligned} \quad (\text{B.3})$$

The hypotheses on the partition and  $\phi$  must be verified in any application; they do not permit arbitrary additional floor factors or a smooth twist that cancels the main wave.

## B.1 The widened D1 linearization

Fix an A-process half-shift  $1 \leq h \leq P^{1/12}$ . On an actual carry branch for  $z = n + e$ , put

$$m(z + 2h) = m(z) + \beta_1, \quad m(z + 2r) = m(z) + \beta_2, \quad m(z + 2h + 2r) = m(z) + \beta_1 + \beta_2 + j.$$

Then

$$\Delta_{2h} \Delta_{2r} Y(z) = F_j(m(z)),$$

where  $F_j$  has the same four-term form as  $F_\ell$  above. Uniformly,  $\beta_1$  is comparable to  $h \sqrt{P}$ ,  $\beta_2$  to  $r \sqrt{P}$ , and  $|j| \leq 2$ . The latter follows because the corresponding real mixed difference of  $X$  is  $O(hrP^{-1/2}) = o(1)$ .

The exact integral formulas, with the  $\beta$  values frozen, are

$$\begin{aligned} F_0(v) &= \frac{3}{4} \int_0^{\beta_1} \int_0^{\beta_2} (v + b + c)^{-1/2} dc db, \\ F_j(v) - F_0(v) &= \frac{3j}{2} \int_0^1 (v + \beta_1 + \beta_2 + \tau j)^{1/2} d\tau. \end{aligned} \quad (\text{B.4})$$

They hold for both signs of  $j$ . Differentiating with respect to  $v$  gives

$$|F'_j(v)| \ll |j| P^{-3/4} + hr P^{-5/4}, \quad v \asymp P^{3/2}. \quad (\text{B.5})$$

Since  $0 \leq X(z) - m(z) < 1$ , replacing  $F_j(m(z))$  by  $F_j(X(z))$  in the exponential costs at most a constant times  $|a|$  times (B.5). The error summed over all integers in the dyadic interval is

$$\ll |a| P^{1/4} + |a| hr P^{-1/4} \ll_C P^{3/4}/r + h P^{1/4} \ll_C P^{3/4}. \quad (\text{B.6})$$

This is one global charge; it is not paid again on each branch.

A second differentiation after composition gives

$$|(F_j \circ X)''| \ll |j|P^{-5/4} + hrP^{-7/4}.$$

Consequently the retained D1 curvature is at most

$$C(P^{-3/4}/r + hP^{-5/4}). \quad (\text{B.7})$$

The leading frozen coefficients provide a useful check:

|                  | $F'_j(v)$                      | $(F_j \circ X)''(x)$             |
|------------------|--------------------------------|----------------------------------|
| offset part      | $(3j/4)v^{-1/2}$               | $-(9j/32)x^{-5/4}$               |
| zero-offset part | $-(3\beta_1\beta_2/8)v^{-3/2}$ | $(63\beta_1\beta_2/64)x^{-11/4}$ |

These are leading terms with uniform smaller remainders, not exact equalities for finite  $\beta$ . One must not differentiate the moving approximation  $\beta_i$  approximately  $3h \sqrt{x}$ .

No large Fourier expansion of this D1 floor error is needed.

## B.2 Differencing and the nonzero main wave

Choose  $H = \lfloor P^{1/12} \rfloor$ . Apply van der Corput differencing to the sequence indexed by the odd integers, using shifts  $2h$ :

$$|U|^2 \ll_C P^2/H + (P/H) \sum_{1 \leq h < H} |V_h|. \quad (\text{B.8})$$

Here  $V_h$  is the correlation on the overlap of the original interval and its translate. At most  $O_C(hP^{13/24})$  starting points cross a cell boundary of  $\mathcal{P}$ . Their contribution is  $O_C(P^{5/8})$  for each  $h$ . On each remaining stable interval  $I$  intersect  $(I-2h)$ , all  $a_j$  are constant, the product of the two amplitudes has bounded variation  $O_C(1)$ , and

$$|(\Delta_{2h}\phi)''| \leq 2h \sup_I |\phi'''| \ll_C hP^{-13/12}. \quad (\text{B.9})$$

We apply all derivative tests to these intervals, never to a domain punctured by individual exceptional integers.

Let  $\theta(n) = \{X(n)\}$ . Taylor's theorem gives

$$Y(n) = n^{9/4} - \frac{3}{2}\theta(n)n^{3/4} + E(n), \quad |E(n)| \ll P^{-3/4}.$$

On a fixed main carry branch write  $\beta = m(n+2h) - m(n)$ , and put

$$A_h(x) = \Delta_{2h}(x^{9/4}) - \frac{3}{2}\Delta_{2h}X(x)(x+2h)^{3/4}.$$

The exact resulting decomposition is

$$\Delta_{2h}Y(n) = A_h(n) + \frac{3}{2}\beta(n+2h)^{3/4} - \frac{3}{2}\theta(n)\Delta_{2h}(n^{3/4}) + \Delta_{2h}E(n). \quad (\text{B.10})$$

The  $\theta$  term and  $E$  term cost, over the full correlation,

$$O(|t|hP^{3/4} + |t|P^{1/4}) = O_C(P^{7/8}). \quad (\text{B.11})$$

Taylor expansion of the smooth  $A_h$ , with derivatives, gives

$$A_h'' = O(h^2P^{-7/4}).$$

On each branch  $\beta = 3h \sqrt{x} + O(1 + h^2P^{-1/2})$  as a comparison of values. Holding  $\beta$  fixed in differentiation, the main retained phase therefore has

$$\left(tA_h + \frac{3t}{2}\beta(x+2h)^{3/4}\right)'' = -\frac{27}{32}tx^{-3/4}(1 + o(1)). \quad (\text{B.12})$$

The  $o(1)$  is uniform for  $1 \leq h \leq H$  and for all admissible branches.

Write  $M_h = |t|hP^{-3/4}$ . The ratio of the combined D1 curvature in (B.7) to  $M_h$  is

$$O_{C,M}((|t|h)^{-1} + P^{-1/2}/|t|).$$

Choose a sufficiently large constant  $h_0 = h_0(C, M)$ , independent of  $P$ . For  $h \geq h_0$  this ratio is smaller than a fixed fraction of the lower constant in (B.12). The twist ratio in (B.9) is  $O_C(P^{-1/3}/|t|)$ . Thus all these terms preserve the one-signed main curvature. This argument permits arbitrary signs and mutual cancellation among the D1 coefficients.

There are only  $O_{\{C,M\}}(1)$  shifts  $h < h_0$ . The trivial bound  $|V_h| \leq C P$  for these shifts contributes  $O_{\{C,M\}}(P^2/H)$  to (B.8). Small values of  $P$  for which  $H < h_0$  are covered by the implied constant.

### B.3 Insert the D2 reduction on its original runs

Use (A.9) of Appendix A.1 at this  $h$ , with  $Q = \text{floor}(P^{5/16})$ . Translations of the base and coefficient by  $O_C(P^{1/24})$  preserve all derivative comparisons and run counts in that proof. Conjugation handles  $\sigma = -1$ , and  $\sigma = 0$  contributes the constant factor one.

For each factor, outside an error whose sum of absolute values is  $O_C(P^{15/16} \log P)$ , the retained modes are

$$e(\Phi_{\ell,q}), \quad \Phi_{\ell,q} = (N_\ell + q - B_\ell)G_\ell, \quad B_\ell = \Delta_{2h}c_\ell, \quad N_\ell = \lfloor B_\ell \rfloor,$$

or their conjugates. Here  $|q| \leq Q$  and on each coefficient window the total coefficient sup norms plus variations are  $O(\log P)$ . For a fixed number of factors these become a fixed power of  $\log P$ .

Every such mode has

$$|\Phi''_{\ell,q}| \ll_C P^{-15/16}. \quad (\text{B.13})$$

The ratio to  $M_h$  is  $O_C(P^{-3/16}/(|t|h))$ , and so preserves (B.12). The additional coefficient-window count is

$$O_C(1 + k_\ell h P^{1/8}) = O_C(P^{1/4});$$

the original D2 run count is  $O_C(P^{13/24})$ . Both are retained in the common partition below.

For clarity, the  $P^{15/16}$  error includes floor mismatches and floor-crossing events. Those are compared in absolute value with unimodular expressions and then replaced on full intervals. Their counts are taken on the original D2 runs before any extra refinement. Allowing fresh D2 labels on every added input cell would not be justified by that count and is explicitly excluded in Theorem B.1.

### B.4 Reassemble the actual first-level carries

Apply Lemma A.2 to the finitely many first-floor evaluations in the main wave and first-difference terms. Their even shifts are  $O(h + P^{1/24})$ . Intersect the resulting gap runs with the stable input cells and the coefficient windows from Lemma A.1. The full number of cells is

$$D_h \ll_{C,M} (h + P^{1/24})P^{1/2}. \quad (\text{B.14})$$

The input partition contributes  $O(P^{13/24})$  cells; each floor-factor coefficient contributes  $O(P^{1/4})$  additional windows. Endpoint-order cuts multiply this number by a fixed constant.

At carry cutoff  $R = \text{floor}(P^{1/4})$ , Lemma A.2 supplies the zero coefficients with bounded variation, nonzero endpoint modes with weights  $O(1/|v|)$  and phases  $vX(x+s)$ , and total positive error  $O(P^{5/6} \log P)$ . This error is charged once over the common input interval, multiplied only by the logarithmic coefficient masses from the earlier expansions.

## B.5 Sum every retained mode and every cell

After the preceding replacements, the zero carry mode has one-signed curvature comparable to  $M_h$ , uniformly in the D2 indices. On a cell of length  $L$  the second-derivative test gives  $O(L\sqrt{M_h} + M_h^{-1/2})$ , also for any subinterval and the odd lattice. Partial summation applies to  $\eta$  and all Fourier coefficients. Their total sup norms plus variations on each cell are at most a fixed power of  $\log P$ . Summing lengths and all  $D_h$  endpoints gives

$$P\sqrt{M_h} + D_h M_h^{-1/2} \ll_{C,M} (|t|h)^{1/2} P^{5/8} + \left( \sqrt{h/|t|} + P^{1/24} (|t|h)^{-1/2} \right) P^{7/8}. \quad (\text{B.16})$$

For  $h_0 \leq h \leq P^{1/12}$  and the stated  $t$  range this is  $O_{C,M}(P^{11/12})$ , before logarithmic losses.

For a nonzero carry Fourier mode  $v$ , its extra curvature is  $vX'(x+s)$ , of size  $|v| P^{-1/2}$ . All other curvatures together are  $O_{C,M}(M_h)$ , and

$$\frac{M_h}{|v| P^{-1/2}} \ll_{C,M} P^{-1/8}.$$

Thus these modes also have one-signed curvature of the claimed size. Summing their  $1/|v|$  weights and all cell endpoints gives

$$R^{1/2} P^{3/4} + D_h P^{1/4} \ll_{C,M} P^{7/8} + (h + P^{1/24}) P^{3/4}. \quad (\text{B.17})$$

This is  $O_{C,M}(P^{7/8})$  in the full range. A common reference curvature and uniform cell count make this bound valid after summing the coefficient masses of every D2 mode; no factor  $Q$  or  $D_h$  is hidden in that mass.

Combining the absolute errors (B.6), (B.11), (A.12), the D2 error, the coarse-boundary cost, and (B.16)–(B.17), for every  $h \geq h_0$ ,

$$|V_h| \ll_{C,M} P^{15/16} (\log(2P))^B, \quad (\text{B.18})$$

where  $B$  depends only on  $M$ . The largest cost is the D2 error. Substitution in (B.8), with the finitely many small shifts treated as in Appendix B.2, yields

$$|U|^2 \ll_{C,M} P^{23/12} + P^{31/16} (\log(2P))^B.$$

Since  $23/12 < 31/16$ , this proves (B.3), absorbing logarithms into an arbitrarily small positive power of  $P$ .

## Appendix C. Proof of the OOOEE mixed-mode theorem

Put

$$\begin{aligned} X(n) &= n^{3/2}, \quad m(n) = \lfloor X(n) \rfloor, \quad Y(n) = m(n)^{3/2}, \\ v(n) &= \lfloor Y(n) \rfloor, \quad Z(n) = v(n)^{3/2}, \\ w(n) &= \lfloor Z(n) \rfloor, \quad U(n) = \sqrt{w(n)}. \end{aligned}$$

For every fixed  $C \geq 1$  and every  $\varepsilon > 0$ , uniformly over nonzero integer quadruples with  $\max(|i|, |j|, |k|, |l|) \leq CP^{1/24}$ ,

$$\left| \sum_{\substack{P < n \leq 2P \\ n \text{ odd}}} e \left( \frac{iX(n) + jY(n) + kZ(n) + lU(n)}{2} \right) \right| \ll_{C,\varepsilon} P^{127/128 + \varepsilon}. \quad (\text{C.1})$$

Here  $e(x) = \exp(2\pi ix)$ . These coordinates form a formal chain for every odd  $n$ ; they coincide with the Juggler trajectory on the specified OOOEE sign class.

The counting consequences are

$$\#\{n \leq N : \text{word}_5(n) = \text{OOOEE}\} = \frac{N}{32} + O_\varepsilon(N^{127/128+\varepsilon}), \quad (\text{C.2})$$

and the class  $\mathcal{C}_5$  of starts with a power-envelope contraction certificate of length at most five satisfies

$$\#(\mathcal{C}_5 \cap [1, N]) = \frac{7N}{8} + O_\varepsilon(N^{127/128+\varepsilon}). \quad (\text{C.3})$$

This is a certificate-class density, not the exact density of all starts that happen to descend within five steps.

### C.1 The full phase

The proof uses Lemmas A.1–A.4 and Theorem B.1. The elementary small-shift argument is Lemma 4.4. The fractional-floor exception counts needed below are derived from (A.3)–(A.4) on original runs. The exact centered master identity supplies the signed inventory.

The bare kernel bound alone is insufficient. Writing  $\theta = \{X\}$  and  $\vartheta = \{Y\}$ , Taylor's theorem gives

$$\begin{aligned} Z &= m^{9/4} - \frac{3}{2}m^{3/4}\vartheta + O(P^{-9/8}), \\ m^{3/4} &= n^{9/8} + O(P^{-3/8}), \\ U &= m^{9/8} + O(P^{-9/16}). \end{aligned} \quad (\text{C.4})$$

For the last assertion,  $\sqrt{[Z]} = v^{3/4} + O(P^{-27/16})$  and  $v^{3/4} = Y^{3/4} + O(P^{-9/16})$ . All comparisons are uniform for  $n$  in  $[P, 3P]$ .

Let  $c(x) = 3kx^{9/8}/4$ . Replacing the original phase by

$$F(n) = \frac{i}{2}X(n) + \frac{j}{2}Y(n) + \frac{k}{2}m(n)^{9/4} + \frac{l}{2}m(n)^{9/8} - c(n)\vartheta(n) \quad (\text{C.5})$$

costs in the whole exponential sum at most

$$O_C(|k|P^{5/8} + |k|P^{-1/8} + |l|P^{7/16}) = O_C(P^{2/3}). \quad (\text{C.6})$$

We keep both powers of  $m$  in (C.5) until after differencing. In particular, the fifth-coordinate term is not expanded initially into a Fourier series with a growing coefficient.

### C.2 The cases $k=0$ , including every zero coordinate

In these cases (C.4) leaves the phase  $iX/2 + jY/2 + lm^{9/8}/2$ , with error already covered by (C.6).

If  $j$  is nonzero, apply one A-process with  $H = \lfloor P^{1/12} \rfloor$ . On a first-floor gap cell and carry branch let  $g = m(n + 2h) - m(n)$ , with  $g$  fixed and comparable to  $h\sqrt{P}$ . For

$$V_g(z) = (z + g)^{9/8} - z^{9/8}$$

the exact integral formula gives

$$|V'_g(X)| \ll hP^{-13/16}, \quad |(V_g \circ X)''| \ll hP^{-21/16}. \quad (\text{C.7})$$

Replace  $lV_g(m)/2$  by  $lV_g(X)/2$  at the global cost  $O(|l|hP^{3/16}) \leq O_C(P^{5/16})$ . In the proof of Lemma 4.4, add this last smooth term separately to each of the two  $g$  branches. No new gap cell or carry is needed. Its curvature relative to the main one  $|j|hP^{-3/4}$  is

$$O(|l/j|P^{-9/16}) = O_C(P^{-25/48}).$$

The exact carry interpolation (4.8), global positive Fourier errors, and domination of the nonzero first-floor Fourier modes therefore remain valid. Both signs of  $j$  are handled by conjugating the whole phase. The correlation is  $O_C(P^{7/8}(1 + \sqrt{h}))$ ; the A-process gives  $O_C(P^{23/24})$ .

If  $j=0$  and  $l$  is nonzero, expand once:

$$\frac{l}{2}m^{9/8} = \frac{l}{2}n^{27/16} - B(n)\theta + O(|l|P^{-21/16}), \quad B(x) = \frac{9l}{16}x^{3/16}. \quad (C.8)$$

Set  $N_B = \lfloor B \rfloor$  and expand the bounded residual  $B - N_B$  at cutoff  $T = \lfloor P^{1/8} \rfloor$ . The retained phases are

$$f_r(x) = \frac{l}{2}x^{27/16} + (i/2 + r - N_B)x^{3/2}, \quad |r| \leq T.$$

Freeze  $N_B$  when differentiating. Substituting its value only after differentiation gives

$$f_r'' = \frac{81l}{512}x^{-5/16} + O((|i| + T + 1)P^{-1/2}). \quad (C.9)$$

The coefficient is  $(1/2)(27/16)(11/16) - (9/16)(3/4) = 81/512$ . The error relative to the leading term is  $O_C(P^{-7/48} + P^{-1/16})$ . There are  $O(1 + |l|P^{3/16})$  center windows; the Fourier coefficient mass and variation on each are  $O(\log P)$ . The summed second-derivative bound is

$$O\left(|l|^{1/2}P^{27/32} + (1 + |l|P^{3/16})|l|^{-1/2}P^{5/32}\right) \log^C(2P) \ll_C P^{83/96} \log^C(2P).$$

The positive Fourier error totals  $O(P^{7/8} \log P)$ . For example (A.12)'s distance-band proof gives  $O(P \log(2T)/T + P^{5/6} \log(2P))$  at this cutoff. Thus this case is  $O_C(P^{7/8} \log^C P)$ .

Finally  $j=l=0$  forces  $i$  nonzero. The smooth phase  $iX/2$  has curvature comparable to  $|i|P^{-1/2}$  and sum  $O_C(P^{37/48} + P^{1/4})$ . This covers every  $k=0$  mode without using (C.1).

### C.3 The twice-differenced inventory when $k$ is nonzero

Conjugate all four frequencies if necessary to assume  $k \geq 1$ . Use the ranges in Appendix A.3

$$H_1 = \lfloor P^{1/48} \rfloor, \quad H_2 = \lfloor P^{1/24} \rfloor, \quad 1 \leq h_a < H_a, \quad d_a = 2h_a, \quad p = h_1 h_2, \quad \Pi = kp \ll_C P^{5/48}. \quad (C.10)$$

Write  $\Delta_a f(x) = f(x + d_a) - f(x)$ ,  $W_a = \Delta_a Y$ ,  $D = \Delta_1 \Delta_2 Y$ , and  $c_{11}(x) = c(x + d_1 + d_2)$ . The double difference of (C.5) is exactly

$$\frac{k}{2} \Delta_1 \Delta_2 (m^{9/4}) + \frac{l}{2} \Delta_1 \Delta_2 (m^{9/8}) + \frac{i}{2} \Delta_1 \Delta_2 X + \frac{j}{2} D - \Delta_1 \Delta_2 (c\vartheta). \quad (C.11)$$

Apply the conjugate of the master expansion (A.15)–(A.18) to the last term, with  $J = \lfloor P^{1/24} \rfloor$ . Set

$$B_2 = \Delta_2 c(x + d_1), \quad B_1 = \Delta_1 c(x + d_2), \quad N_a = \lfloor B_a \rfloor.$$

The retained modes have phase

$$\begin{aligned} & \frac{k}{2} \Delta_1 \Delta_2 (m^{9/4}) + \frac{l}{2} \Delta_1 \Delta_2 (m^{9/8}) - c_{11}\{D\} \\ & + tY + a_1 W_1 + a_2 W_2 + qD + \phi_0, \quad a_1 = -N_2 + u, \quad a_2 = -N_1 + v. \end{aligned} \quad (C.12)$$

Here  $t, u, v$  are integers of absolute value  $O_C(J)$ ;  $q \in \frac{1}{2}\mathbb{Z}$  of absolute value  $O_C(J)$ , including the original  $j/2$ . The function  $\phi_0$  is  $i\Delta_1 \Delta_2 X/2$  plus a bounded integer combination of  $B_1, B_2, c_{11}$ . In particular

$$\begin{aligned} |a_1| h_1 + |a_2| h_2 & \ll_C \Pi P^{1/8} + J(h_1 + h_2) \ll_C P^{11/48}, \\ |\phi_0''| & \ll_C kP^{-7/8} + |i|pP^{-5/2}. \end{aligned} \quad (C.13)$$

All signs of  $i, j, l, u, v, q$  are permitted. The original  $jY/2$  contributes only  $jD/2$  after two differences, so  $t$  remains an integer.

The full coefficient mass is  $O(\log^7(2P))$ . Each multi-index has a fixed harmonic envelope under which its normalized coefficient has bounded sup norm plus variation on the  $N_1, N_2$  cells. The centers have local density  $O_C(P^{-19/24})$ . These are exactly the centering and five binary-carry operations in the master-inventory argument, with signs reversed. The deleted first master term costs  $O_C(P^{11/48})$ , and all positive errors cost  $O_C(P^{23/24} \log^C P)$ . Their arguments are the same Y corners,  $W_1, W_2$ , the shifted  $W_1$ , and D. The new factors in (C.11) have modulus one, so the positive bounds (A.13) are unchanged. In particular no OOOEE discrepancy has been assumed to control them.

#### C.4 Frozen branches and the extra powers

Partition first into the original gap runs determined by the floors of  $\Delta_1 X, \Delta_2 X, \Delta_{d_1+d_2} X$ . Their total count is  $O_C(P^{13/24})$ , their local count in length L is  $O_C(1 + LP^{-11/24})$ , and their lengths are  $O_C(\sqrt{P})$ . On a carry pattern let

$$m(n + d_a) = m(n) + \beta_a, \quad m(n + d_1 + d_2) = m(n) + \beta_1 + \beta_2 + b.$$

Then  $-1 \leq b \leq 2$ , since the real mixed difference of X is positive and  $o(1)$ . The  $\beta$  values and b are fixed on the branch;  $\beta_a = 3h_a\sqrt{x} + O(1 + h_a^2 P^{-1/2})$  only as values.

For any exponent a define

$$F_a(z) = (z + \beta_1 + \beta_2 + b)^a - (z + \beta_1)^a - (z + \beta_2)^a + z^a.$$

The formulas used for every derivative are

$$\begin{aligned} F_{a,0}(z) &= a(a-1) \int_0^{\beta_1} \int_0^{\beta_2} (z + s + t)^{a-2} dt ds, \\ F_a(z) - F_{a,0}(z) &= ab \int_0^1 (z + \beta_1 + \beta_2 + \tau b)^{a-1} d\tau. \end{aligned} \tag{C.14}$$

They apply also when  $b < 0$ . The notation  $F_{a,0}$  means  $b=0$ . Put  $G = F_{3/2}(X)$  and  $A = \lfloor F_{3/2}(m) \rfloor$ . The exact identity  $\Delta_1 \Delta_2(m^a) = F_a(m)$  holds on its actual carry pattern.

The fifth-coordinate term can now be replaced directly by  $lF_{9/8}(X)/2$ . The relevant bounds are

$$\begin{aligned} |F'_{9/8}(X)| &\ll |b|P^{-21/16} + pP^{-29/16}, \\ |(F_{9/8} \circ X)''| &\ll |b|P^{-29/16} + pP^{-37/16}, \\ |(F_{9/8} \circ X)'''| &\ll |b|P^{-45/16} + pP^{-53/16}. \end{aligned} \tag{C.15}$$

The total replacement error is  $O(|l|P^{-5/16} + |l|pP^{-13/16}) = O_C(1)$ . Replacing qD by qG costs  $O_C(P^{7/24})$ , by (A.4) and the mean value theorem; integrality of q is irrelevant for that comparison.

Combine the two large terms before expanding in  $\theta$ :

$$H(z, x) = \frac{k}{2} F_{9/4}(z) - c_{11}(x) F_{3/2}(z), \quad B_{\text{core}}(x) = \partial_z H(X(x), x).$$

Since  $-c_{11}\{D\} = -c_{11}F_{3/2}(m) + c_{11}A$ , the core of (C.12) is  $H(m, x) + c_{11}A$ . Taylor expansion gives

$$H(m, x) = H(X, x) - B_{\text{core}}\theta + O_C(k|b|P^{-9/8} + \Pi P^{-13/8}), \tag{C.16}$$

so its total error is bounded. The growing coefficient is

$$B_{\text{core}} = \frac{27}{32} k b x^{3/8} + O_C(\Pi P^{-1/8} + k(h_1 + h_2)P^{-5/8}). \tag{C.17}$$

The leading coefficient is  $45/32 - 9/16 = 27/32$ . The remainder after subtracting  $(27/32)kbx^{3/8}$  is bounded in (C.10). Its derivative on each original run is  $O_C(P^{-3/4})$ , by (C.14) with the offsets fixed. Each such run has length  $O_C(P^{1/2})$ , so this remainder has bounded variation there. The leading term itself has derivative  $O_C(kP^{-5/8})$ ; it is not included in the smaller remainder-derivative bound.

All carry patterns are retained. As in (A.12), their indicators are finite unions of arcs in  $\theta$ . Endpoint-order cuts add only a fixed multiple of the run count. At cutoff  $R_c = P^{1/4}$ , each nonzero endpoint mode has weight  $O(1/|s|)$  and phase  $sX(x+d)$ , for an allowed shift  $d$ ; the zero coefficient has bounded variation. Their global positive errors are  $O(P^{5/6} \log P)$ . We use  $R = P^{5/16}$  for the bounded  $\theta$  coefficients below; their global errors have the same bound. No error is charged again on each gap run or frequency window.

### C.5 Nonzero total Y frequency

Suppose  $t$  is nonzero in (C.12). Keep the first differences  $a_1W_1 + a_2W_2$  unexpanded and keep the floor  $A$  on its original fixed-label runs. In particular do not cut into  $A$ 's integer levels.

Center the growing term (C.17) at

$$N_* = \left\lfloor \frac{27}{32}kbx^{3/8} \right\rfloor.$$

On the intersections of original runs and  $N_*$  windows,  $B_{\text{core}} - N_*$  has bounded size and variation: the leading fractional part is monotone with variation at most one, and the remainder has the variation bound just proved. This residual need not lie in  $[0, 1]$ . For  $b = 0$  use  $N_* = 0$ . Expand it at  $R$  using the bounded-residual extension of Lemma 4.7. After the first-floor carry expansion the smooth part is

$$\Phi = H(X, x) + \frac{l}{2}F_{9/8}(X) + qG + \phi_0 + (r - N_*)X + sX(x + d).$$

Its third derivative satisfies

$$|\Phi'''| \ll_C kP^{-9/8} + \Pi P^{-13/8} + (R + |N_*| + R_c)P^{-3/2} + |l|(P^{-45/16} + pP^{-53/16}) \ll_C P^{-13/12}. \quad (\text{C.18})$$

The  $q$  and  $\phi_0$  terms are smaller;  $|N_*| \ll_C P^{5/12}$ .

The common coarse partition consists of original runs,  $N_1$ ,  $N_2$  windows,  $N_-^*$  windows, and endpoint-order cuts. Its local count remains  $O_C(1 + LP^{-11/24})$ , since  $N_-^*$  has local density  $O_C(kP^{-5/8}) = O_C(P^{-7/12})$ . Fixed  $b$  on each original run permits this assertion even though  $b$  changes between runs. The assigned D2 labels do not change at extra refinements. An empty carry pattern is extended using an admissible fixed label on that original run and zero amplitude there.

Every normalized retained term now satisfies (B.3): the nonzero integer  $t$  is in its allowed range; (C.13) fits its widened first-difference budget; (C.18) fits its twist budget; and the single floor term is  $+ c_{11} A$ , namely  $\sigma = -1$  in (B.3). The sum of all these weighted terms is

$$O_{C,\varepsilon}(P^{31/32+\varepsilon}). \quad (\text{C.19})$$

The D2 reduction is invoked inside (B.3), after that theorem's additional A-process, rather than on an arbitrary undifferenced floor.

### C.6 Zero total Y frequency: shared reductions

Now  $t=0$ . Replace  $A$  by  $J_F = \lfloor G \rfloor$ . For  $b$  nonzero, (A.3)–(A.4), at derivative scale  $P^{-1/4}$  and distance  $O(P^{-3/4} + pP^{-5/4})$ , give  $O_C(P^{3/4})$  exceptions on all original runs. For  $b=0$  the scale is  $pP^{-3/4}$ , the distance is  $O(pP^{-5/4})$ , and the same count is  $O_C(pP^{1/4} + (h_1 + h_2)P^{1/2}) =$

$O_C(P^{13/24})$ . Each changed exponential costs at most two. The coefficient  $c_{11}$  is not multiplied by the size of a floor error.

For  $V_a(z) = (z + \beta_a)^{3/2} - z^{3/2}$ , expand

$$a_a V_a(m) = a_a V_a(X) - a_a V'_a(X)\theta + O(|a_a| h_a P^{-7/4}).$$

By (C.13) the global error is  $O_C(P^{-25/48})$ . The smooth part and  $\theta$  coefficient are consequently

$$\begin{aligned} F_{\text{sm}} &= \frac{k}{2} F_{9/4}(X) - c_{11}(G - J_F) + a_1 V_1(X) + a_2 V_2(X) + \frac{l}{2} F_{9/8}(X) + qG + \phi_0, \\ B &= B_{\text{core}} + a_1 V'_1(X) + a_2 V'_2(X). \end{aligned} \quad (\text{C.20})$$

The wave contribution to B is bounded:

$$\left| \sum a_a V'_a(X) \right| \ll_C \Pi P^{-1/8} + J(h_1 + h_2) P^{-1/4} \ll_C 1.$$

Its derivative on a frozen cell is bounded by  $O_C(\Pi P^{-9/8} + J(h_1 + h_2) P^{-5/4})$ . Multiplying the derivative bound by the original-run length  $O_C(P^{1/2})$  proves bounded variation on every frozen cell; refining a cell cannot increase this bound. Together with the remainder bound following (C.17), this verifies the hypotheses of the bounded-residual extension of Lemma 4.7 in both offset cases. All integer levels of  $J_F$  are now included in the derivative-test partition.

### C.7 Nonzero b: recompute the mixed curvature

For b nonzero, center B at the same  $N_-^*$  as in Appendix C.5; its bounded residual has bounded variation. Retained phases are  $f = F_{\text{sm}} + (r - N_*)X + sX(x + d)$ .

Freeze  $\beta_a, b, J_F, N_*, N_a$  before differentiation. The three leading contributions, in units of  $kbx^{-1/8}$ , are

| term                   | coefficient |
|------------------------|-------------|
| $(kF_{9/4}(X)/2)''$    | 945/512     |
| $(-c_{11}(G - J_F))''$ | -864/512    |
| $-N_* X''$             | -324/512    |

Their sum is

$$-\frac{243}{512}. \quad (\text{C.21})$$

For the middle row use  $-2c'_{11}G' - c_{11}G'' - c'_{11}(G - J_F)$ . The last factor  $G - J_F$  is bounded, so its curvature term is only  $O(kP^{-7/8})$ . Dropping the integer-floor contribution would give the wrong leading coefficient.

Uniformly for every retained index,

$$f'' = -\frac{243}{512} kbx^{-1/8} + O_C(P^{-3/16}). \quad (\text{C.22})$$

For clarity, the error terms include

$$\begin{aligned} &\Pi P^{-5/8} + k(h_1 + h_2)P^{-9/8} + kP^{-7/8} \\ &+ (|a_1|h_1 + |a_2|h_2)P^{-3/4} + (R + R_c + 1)P^{-1/2} + R_c(h_1 + h_2)P^{-3/2} \\ &+ |q|(P^{-5/4} + pP^{-7/4}) + |l|(P^{-29/16} + pP^{-37/16}) + |\phi_0''|. \end{aligned}$$

Every term fits (C.22) under (C.10),(C.13). Both signs of b are allowed. The error is smaller than the leading curvature by  $O_C(P^{-1/16}/k)$ .

The full number of intervals is  $O_C(P^{3/4})$ . The G-levels add at most  $O_C(P^{3/4} + P^{13/24})$ ; the original gaps, moving centers, and arc-order cuts are smaller. Summing lengths and every endpoint in the second-derivative estimate, with coefficient masses and variation, gives

$$O_C\left(\sqrt{k}P^{15/16} + k^{-1/2}P^{13/16}\right) \log^C(2P) \ll_C P^{23/24} \log^C(2P). \quad (\text{C.23})$$

The positive errors and floor exceptions are smaller. This proves the nonzero-offset part for the actual mixed family.

## C.8 Zero b: the actual negative moving centers

When  $b=0$ , (C.20) has  $|B| \ll_C 1$  and bounded variation on every original run intersected with the  $N_a$  windows. Expand it directly at  $R$ , without a new  $\theta$  center. After the carry expansion put  $\ell_F = r + s$ , an integer, and  $\alpha_0 = uh_1 + vh_2$ . For fixed  $\ell_F$ , the coefficient sup norms plus variations sum to  $O(\log(2P)/(1 + |\ell_F|))$ . Indeed the convolution of the two coefficient envelopes satisfies

$$\sum_{r+s=\ell_F} \frac{1}{(1+|r|)(1+|s|)} \ll \frac{\log(2P)}{1+|\ell_F|}.$$

Split at  $|r| \geq |\ell_F|/2$  or  $|s| \geq |\ell_F|/2$  to verify this bound. The same estimate holds for the coefficient variations by the product variation inequality; the phases for different  $r, s$  need not be identical.

The pure  $kF_{9/4}(X)/2$  curvature has leading coefficient  $-6075/2048$  in units of  $\Pi x^{-5/8}$ . The negative fractional anchor contributes  $243/128 = 3888/2048$ , so their subtotal is  $-2187/2048$ . The centers in this problem have negative signs:

$$h_1 a_1 + h_2 a_2 = -\frac{27}{8} \Pi x^{1/8} + \alpha_0 + O_C(h_1 + h_2 + \Pi(h_1 + h_2)P^{-7/8}). \quad (C.24)$$

The frozen wave curvature is  $-27(h_1 a_1 + h_2 a_2)x^{-3/4}/32$ , up to smaller terms. It adds  $729/256 = 5832/2048$  to the subtotal. Thus

$$f'' = -\frac{27}{32} \alpha_0 x^{-3/4} + \frac{3645}{2048} \Pi x^{-5/8} + \frac{3}{4} \ell_F x^{-1/2} + O_C(\Pi P^{-3/4}). \quad (C.25)$$

This is the coefficient for the complete mixed phase (C.12), including the pure power and both moving centers.

The errors in (C.25) include

$$\begin{aligned} & (h_1 + h_2)P^{-3/4} + (|a_1| + |a_2|)P^{-5/4} + (|a_1|h_1^2 + |a_2|h_2^2)P^{-7/4} \\ & + k(h_1 + h_2)P^{-9/8} + \Pi(h_1 + h_2)P^{-13/8} + kP^{-7/8} + |q|pP^{-7/4} \\ & + P^{-29/24} + |l|pP^{-37/16} + |\phi_0''|. \end{aligned}$$

Use  $h_1 + h_2 \leq 2p \leq 2\Pi$ , before inserting independent shift caps, for the leading rounding error. All terms have the claimed bound. The fifth-coordinate term is particularly small here.

There is a strict frequency gap:

$$\frac{|\alpha_0|P^{-3/4}}{\Pi P^{-5/8}} \ll_C JP^{-1/8} = O_C(P^{-1/12}), \quad \frac{\Pi P^{-5/8}}{P^{-1/2}} \ll_C P^{-1/48}. \quad (C.26)$$

Consequently  $\ell_F = 0$  has one-signed curvature comparable to  $\Pi P^{-5/8}$ ; every nonzero integer  $\ell_F$  has one-signed curvature comparable to  $|\ell_F|P^{-1/2}$ . No hypothesis on the largest individual signed wave is used.

The common partition has  $O_C(P^{13/24})$  cells, including  $O_C(pP^{1/4} + P^{13/24})$  zero-offset G-level cuts. For  $\ell_F = 0$ , summing all lengths and endpoints costs

$$O_C\left(\Pi^{1/2}P^{11/16} + \Pi^{-1/2}P^{41/48}\right) \log^C(2P).$$

For nonzero  $\ell_F$ , sum the harmonic weights as well:

$$O_C\left(R^{1/2}P^{3/4} + P^{13/24+1/4}\right) \log^C(2P) = O_C(P^{29/32} \log^C(2P)). \quad (C.27)$$

The zero mode and positive errors are smaller. This directly estimates the moving coefficients in (C.12), with every window endpoint included.

## C.9 Assembly and the dyadic count

For the double correlation of (C.5), the accounting is:

| Contribution                                     | Exponent, apart from fixed logarithmic powers |
|--------------------------------------------------|-----------------------------------------------|
| Deleted master term                              | 11/48                                         |
| Positive short-Fourier errors                    | 23/24                                         |
| Extra-power Taylor errors and slow D replacement | at most 7/24                                  |
| Nonzero t, via (B.3)                             | 31/32                                         |
| t=0, nonzero b                                   | 23/24                                         |
| t=0, b=0                                         | 29/32                                         |

The first-floor Fourier errors and floor mismatches are included in the last three estimates. Thus the double correlation is  $O_{C,\varepsilon}(P^{31/32+\varepsilon})$ . For the sum  $S_F$  and its first and second correlations, van der Corput differencing on the odd lattice gives

$$|S_F|^2 \ll P^2/H_1 + (P/H_1) \sum_{h_1 < H_1} |T_1(h_1)|,$$

$$|T_1(h_1)|^2 \ll P^2/H_2 + (P/H_2) \sum_{h_2 < H_2} |T_2(h_1, h_2)|.$$

Substitution gives

$$|T_1|^2 \ll_C P^{47/24} + P^{63/32+\varepsilon}, \quad |T_1| \ll_{C,\varepsilon} P^{63/64+\varepsilon},$$

and

$$|S_F|^2 \ll_C P^{95/48} + P^{127/64+\varepsilon}, \quad |S_F| \ll_{C,\varepsilon} P^{127/128+\varepsilon}.$$

Overlap endpoints cost  $O(h_1 + h_2)$  and fit these estimates. Arbitrarily small epsilon losses may be relabeled after each step. Together with Appendices C.1–C.2 this proves (C.1).

Apply the four-dimensional Erdős–Turán–Koksma inequality to

$$(\{X/2\}, \{Y/2\}, \{Z/2\}, \{U/2\})$$

on the odd integers in a single dyadic block. Use cutoff  $\lfloor P^{1/24} \rfloor$ . Its diagonal error is  $O(P^{23/24})$ ; the weighted nonzero modes in (C.1) cost  $O_\varepsilon(P^{127/128+\varepsilon})$ , absorbing the four harmonic sums. All sixteen half-boxes have volume 1/16. Since there are  $P/2 + O(1)$  odd inputs, each formal sign class has count

$$P/32 + O_\varepsilon(P^{127/128+\varepsilon})$$

on this block. A half-box records exactly the parity of the floor: even for a coordinate in  $[0, 1/2)$ , odd for one in  $[1/2, 1)$ .

Use this discrepancy argument separately on each dyadic block in a decomposition of  $[1, N]$ , with that block's own frequency cutoff. Then sum the counts and errors. One must not impose the largest block's cutoff on all smaller blocks. Taking a small epsilon first, the error series is geometric; the remaining bounded initial interval contributes  $O(1)$ . The class with m,v odd and w,floor(U) even is exactly OOOEE for odd n. This proves (C.2), including endpoint conventions. All fifteen nonempty products of its four parity signs also have this error bound, by summing the sixteen sign classes. In the manuscript's notation,  $H(\text{OOOEE}; \delta)$  follows for every  $0 < \delta < 1/128$ .

Finally Lemma 5.1 and Theorem 5.3 of the manuscript give the disjoint union of its  $27/32$  class and OOOEE. The former error  $O(N^{47/48})$  is smaller. Their densities add to  $27/32 + 1/32 = 7/8$ , proving (C.3). This proves Corollary 4.12 and Theorem 5.4.

## Acknowledgments and AI assistance

Large language models were used throughout the development of the earlier draft and this revision, including the formulation and review of proof arguments, drafting, programming, and discussion of Lean formalizations. The September 2026 review with OpenAI Codex identified unresolved analytic steps and assisted with the conditional revision, the four-step and fifth-letter repairs, the D2 reduction, the signed mixed-mode proofs, their consolidation into this version, and the release materials. The September 2026 attribution revisions, which identify the word counts of Sections 5 and 6 with the Collatz sequences A076227, A100982 and A020914, credit the formulas on those entries and the tabulations of Noe and Hikawa, list the machine-checked statements of Section 8 and record the closed forms of the jump spectrum in Section 6, were drafted with Anthropic’s Claude, which also carried out the computations behind those closed forms. AI assistance and automated checks are not independent mathematical validation. The author is responsible for the statements, proofs, code, and final approval of the preprint. The models are not authors.

## References

1. OEIS Foundation Inc., “Juggler sequence: if  $n \bmod 2 = 0$  then  $\text{floor}(\sqrt{n})$  else  $\text{floor}(n^{3/2})$ ,” *The On-Line Encyclopedia of Integer Sequences*, A094683. Sequence record, accessed 9 September 2026.
2. S. W. Graham and G. Kolesnik, *Van der Corput’s Method of Exponential Sums*, London Mathematical Society Lecture Note Series 126, Cambridge University Press, 1991. Publisher excerpt.
3. L. Kuipers and H. Niederreiter, *Uniform Distribution of Sequences*, Wiley-Interscience, New York, 1974. Chapter 2, especially pp. 112–116. University-hosted scan.
4. R. Terras, “A stopping time problem on the positive integers,” *Acta Arithmetica* 30 (1976), 241–252. doi:10.4064/aa-30-3-241-252.
5. OEIS Foundation Inc., “Number of surviving Collatz residues mod  $2^n$ ,” *The On-Line Encyclopedia of Integer Sequences*, A076227, contributed by Labos Elemer, 2002. Sequence record, accessed 19 September 2026.
6. OEIS Foundation Inc., “Number of admissible sequences of order  $n$ ; related to  $3x+1$  problem and Wagon’s constant,” *The On-Line Encyclopedia of Integer Sequences*, A100982, contributed by S. R. Finch. Sequence record, accessed 19 September 2026.
7. OEIS Foundation Inc., “Number of digits in the base-2 representation of  $3^n$ ,” *The On-Line Encyclopedia of Integer Sequences*, A020914, contributed by C. Kimberling. Sequence record, accessed 19 September 2026.
8. M. Winkler, “Deterministic Structures in the Stopping Time Dynamics of the  $3x + 1$  Problem,” arXiv:1709.03385 [math.GM], 2017–2026. arXiv record.
9. K. Hikawa, “Finite-Dimensional Combinatorial and Arithmetic Structures of Parity Vectors for the Accelerated Collatz Map,” ResearchGate preprint, July 2026, version 2.1 of 2 August 2026. doi:10.13140/RG.2.2.29894.84804/1. Section 6 proves the exponential rate; Conjecture 7.1 is the  $d^{-3/2}$  power. Both are cited from the author’s version note on the record; the full text was not available to this revision.
10. K. Hikawa and K. Nakanishi, “Parity Vector Analysis in the Study of the Collatz Conjecture,” ResearchGate preprint, January 2026. doi:10.13140/RG.2.2.12065.06240. Abstract only read for this revision; the full text was not available. Its authors contributed the A076227 b-file against which the counts behind Section 6 were checked on all 3509 terms.
11. K. Nakanishi, “A Structural Study of Parity Vectors in the Collatz Conjecture,” Jxiv preprint, February 2026. doi:10.51094/jxiv.3096. Read for this revision; it classifies parity vectors by weight and states no asymptotic.

12. M. Winkler, “Coefficient First-Passage Words for the  $3x+1$  Map: A Beatty–Ferrers Model and a Power-of-Two Law,” ResearchGate preprint, September 2026. Refines A100982 by the cut pattern 011 ; the words biject with the partitions inside one Beatty–Ferrers shape and the cut sets are the independent sets of a nested matroid, giving the power-of-two law  $2^{\lfloor (n-1)\log_2(3/2) \rfloor}$  . Exact and finite throughout. doi:10.13140/RG.2.2.31257.79200.
13. M. Winkler, “Admissible  $qx + 1$  Sequences, Semiconvergents, and Rational Catalan Numbers,” ResearchGate preprint, 14 September 2026. Read for this revision. The source of the two-sided bound on  $M_d$  used in the repository; Corollary 11 gives the growth constant 2.8395137305 and Corollary 12 the exact normalized oscillation, as an envelope between 1 and 2.7095112914 attained on the record minima and maxima of  $\{r \log_2 3\}$  . doi:10.13140/RG.2.2.35856.78086.
14. J. C. Lagarias, “The  $3x+1$  problem and its generalizations,” *American Mathematical Monthly* 92 (1985). doi:10.1080/00029890.1985.11971528. Theorem D is the exponential rate used in Section 6 of this paper:  $1 - F(k) \leq 2^{-\eta^k}$  with  $\eta = 1 - H(\lambda^{-1})$  , proved there from Terras’s residue structure by a binomial tail, with the matching lower bound in the remark that follows it. Theorem D carries no attribution label in the survey although its Theorems A and C are both labelled (Terras), and Lagarias’s annotated bibliography credits Terras with the residue-class structure, the coefficient stopping time and natural density one, and with no rate.
15. K. Hikawa, *Collatz Parity Vector Toolkit*, version 1.0, software, 4 July 2026, MIT license. Repository; doi:10.5281/zenodo.21186540. Run for this revision: its survivor counts, minimal-certificate counts and odd-count refinement agree with those used here, and it prints the minimal certificate length as the glide.
16. T. D. Noe, “Irregular triangle of the pattern of conjectured dropping times in the Collatz iteration,” A186008, and “Length of Collatz dropping time patterns in A186008,” A186009, *The On-Line Encyclopedia of Integer Sequences*, contributed 9 February 2011. A186008, A186009, accessed 20 September 2026.
17. K. Hikawa, “Finite Parity-Vector Structures in the Accelerated Collatz Map,” ResearchGate preprint, 12 September 2026. doi:10.13140/RG.2.2.24486.28480. Title, date and DOI verified from DataCite and OpenAlex; neither the text nor an abstract was available to this revision.