

Bayesian Approaches to Statistics and Modeling Case Study • Part III

Mark Kurzeja

The Data

- Last time: we found that a normal regression model failed to capture skew and did not take into account the implicit groupings in the data
- Should we include the high school variable? Should we cluster based on IQ, age, or both? What belief are we expressing if we do?

	kid_score	mom_hs	mom_iq	mom_age
1	65	1	121	27
2	98	1	89	25
3	85	1	115	27
4	83	1	99	25
5	115	1	93	27
6	98	0	108	18

Belief about the World

Collect Data

Bayesian Update

The Data

- **Clustering Variables:**
 - We could cluster on whether a mother has a certain range of IQs
 - We could cluster based on the mother's age
 - Whether the mother went to high school or not is another great clustering variable
- For simplicity, we are going to cluster based on whether a mother went to high school and their IQ

	kid_score	mom_hs	mom_iq	mom_age
1	65	1	121	27
2	98	1	89	25
3	85	1	115	27
4	83	1	99	25
5	115	1	93	27
6	98	0	108	18

- **We will use**
 - Low IQ = $IQ < 85$
 - Medium IQ = $85 \leq IQ \leq 115$
 - High IQ = $115 < IQ$

Re Belief about the World

h

Collect Data

Bayesian Update

The Model

- We are going to allow all three parameters to vary according to whether a mother attended HS and their IQ group
 - For example, we will have a different β_0 , β_1 , and β_2 for mothers who are high IQ and attended HS, for those mothers that did not attend HS but also have a high IQ, etc.
- This model now has six sets of β'_0 s, β'_1 s and β'_2 s
- We are also going to say that each of the β'_0 s come from a common distribution. We will also do the same with the β'_1 s and the β'_2 s

Belief about the World

Collect Data

Bayesian Update

The Data

- Modeling in a Bayesian way forces us to state a lot of our assumptions outright
- Stating all assumptions allows someone to challenge them
- We will use a hierarchical model with varying intercepts to capture this belief

Key Point: Always ask **why** a modeling decision is being made

The belief is that the relationship between the IQ of the mother and the child varies based on the mother's education and the mother's IQ group



The Data

- Is this belief “right”? *Of course not*
- All models are subjective, Bayesian or not
- It is important to remember this as we increase the complexity of the model – these are modeling choices based on updated beliefs and not ground truth

Key Point: *All modeling is subjective*

Belief about the World

Collect Data

Bayesian Update

The Model

Multilevel Regression Form

Response Variable

Slope on IQ for a given high school status and IQ group

$$ChildIQ = \beta_{0,HS,IQG} + \beta_{1,HS,IQG} \times (momIQ) + \beta_{2,HS,IQG} \times (momAge)$$




Intercept term for a given high school status and IQ group

Slope on age for a given high school status and IQ group

Belief about the World

Collect Data


Bayesian Update



Belief about the World

Collect Data


Bayesian Update



Belief about the World

Collect Data


Bayesian Update



Belief about the World

Collect Data

Bayesian Update



The Model

- Hierarchical models often have a lot of moving parts (as we can see in the graphic)
- They can model complicated relationships in a more intuitive way than some other methods of modeling
- Downfalls: fitting one of these models in a Bayesian framework can be computationally difficult

Belief about the World

Collect Data

Bayesian Update

The Model - Updates

- Include six new intercepts (β'_0s) , β'_1s , β'_2s , and $\sigma'_i s$ – each constrained by a hierarchical relationship
- Change the distribution of the errors from a normal to a skewed normal to account for the left-skew that we observed in the posterior predictive fits

Belief about the World

Collect Data

Bayesian Update

The Model

- Just like before, when we run the model we will get posterior distributions on each of the parameters
- This time, we can compare each subgroup's intercept and slope to see if they have any relationships
- Why not incorporate a non-linear term (like IQ^2) instead?
 - Expressing belief that different IQ groups come from different distributions and not a non-linear extension of a more global process


Belief about the World

Collect Data

Bayesian Update

The Model – The Posteriors

Posterior Distributions of the Intercept




Belief about the World

Collect Data

Bayesian Update

The Model – The Posteriors

Posterior Distributions of the IQ Coefficient




Belief about the World

Collect Data

Bayesian Update

The Model – The Posteriors

Posterior Distributions of the Age Coefficient



Belief about the World


Collect Data

Bayesian Update

The Model – The Posteriors

Posterior Distribution of Skew Parameter

The skew parameter is predominatly negative indicating that the distribution of errors is left-skewed



Belief about the World


Collect Data

Bayesian Update

The Model – The Posteriors

Posterior Predictive Intervals for First Four Observations

The blue line is the observed value for Childs IQ. The red lines are the 95% predictive interval



Belief about the World


Collect Data

Bayesian Update

The Model – The Posteriors

Normalized Posterior Predictive Intervals

If a dot is in $[-1, 1]$ then the posterior predictive interval contained the true child's IQ




Belief about the World

Collect Data

Bayesian Update

The Model – The Posteriors

Histogram of Width of Predictive Intervals



Belief about the World


Collect Data

Bayesian Update

The Model – The Posteriors

Histogram of Errors

Computed as Mean Prediction - Actual IQ



Belief about the World


Collect Data

Bayesian Update

The Model – The Posteriors

Histogram of Errors - Multilevel Model

Computed as Mean Prediction - Actual IQ



Belief about the World


Collect Data

Bayesian Update

The Model – The Posteriors

Posterior Predictive Check

The true density, of Child IQs, is in blue. The generated posteriors are in grey



Belief about the World


Collect Data

Bayesian Update

The Model – The Posteriors

Histogram of Errors

Computed as Mean Prediction - Actual IQ



Belief about the World

Collect Data

Bayesian Update


The Model - Observations

- The multi-level model, for this data, isn't a massive improvement in prediction ability but is an improvement in inference – we can now tease out the individual group effects
- It provides insight into the differences between the slopes of the parameters
 - Increased variance in parameter estimates as a result
- Next steps in modeling:
 - Account for the over-inflated left-tail using a mixture distribution or another technique



Parting Notes

- Bayesian models are computationally intensive but can model very complex relationships, perform regularization, incorporate previous analysis, and express uncertainty intuitively within the same model
- Methods are becoming more feasible due to advances in sampling techniques and computational power



Parting Notes

- Modeling is an iterative process that is filled with assumptions and is subjective
- Capturing the variance of an estimate is sometimes more important than the estimate itself
- Large posterior intervals \rightarrow Large uncertainty in estimate given the current model and data!

