Aarhus University

Computer Science

Numerical Linear Algebra

Handin 9

Author
Søren M. Damsgaard

Student number
202309814

April 25, 2026

Zulu Bravo PsyOp Striker 9

Given a system of flowing thingies 1 with the following information:

1. A contains 300 L of mixture and 84 g of salt.
2. B contains 100 L of muxture and 36 g of salt.
3. a flows with 30 $L/\min$ of water
4. b flows with 15 $L/\min$ of mixture



Figure 1: Containers connected by pipes: Flowchart of the system.

The Equalizer

Determining the flow of c and d so as the amount of liquid in both containers are constant.

1. A has an input a and b which totals to 45 $L/\min$, so c needs to flow with 45 $L/\min$ to cancel out the input from a and b.
2. B has an output b and a newly determined input c, so the following should determine the flow of the remaining pipe $c - b = 30\ L/\min = d$

Two Peens In A System

Explain why the salt amounts $y_0(t)$ in A and $y_1(t)$ in B satisfy the system

$$y_0'(t) = -0.15y_0(t) + 0.15y_1(t)$$
$$y_1'(t) = 0.15y_0(t) - 0.45y_1(t)$$

The change in salt corresponds to the input and output, percentages, of mixture in each container.
for A: c outputs 45 $L/\min$ which is 15% of A hence $-0.15y_0(t)$, and b inputs 15 $L/\min$ which is 15% of B hence $+0.15y_1(t)$.
for B: c inputs so it's $+0.15y_0(t)$ and b outputs 45 $L/\min$ which is 45% of B hence $-0.45y_1(t)$.

This give us a system that can be written as:

$$y'(t) = Ay = \begin{bmatrix} -0.15 & 0.15 \\ 0.15 & -0.45 \end{bmatrix} \begin{bmatrix} y_0(t) \\ y_1(t) \end{bmatrix}$$

with the initial value:

$$y(0) = \begin{bmatrix} 84 \\ 36 \end{bmatrix}$$

To Eigen or Not To Eigen

The Author calls upon god to protect me against the horros and impending doom of using a numpy function that isn't explicitely shown in the Bibel of Numerical Linear Algebra by Jesus AKA Andrew Swan.

```python
import numpy as np

A = np.array([[-0.15, 0.15],
              [0.15, -0.45]])
# Please don't end me... \(;_;)/
eigvalues, eigvectors = np.linalg.eig(A)

print('Eig Values:\n', eigvalues)
print('Eig Vectors:\n', eigvectors)

# Output (not part of the actual code, just for reference):
Eig Values:
 [-0.08786797 -0.51213203]
Eig Vectors:
 [[ 0.92387953 -0.38268343]
 [ 0.38268343  0.92387953]]
```

Reducing the Eig thingies to just two significant values.

$$\text{Eigenvalues} \qquad \lambda_0 = -0.09 \qquad \lambda_1 = -0.51 \tag{1}$$

$$\text{Eigenvectors} \qquad v_0 = \begin{bmatrix} 0.92 \\ 0.38 \end{bmatrix} \qquad v_1 = \begin{bmatrix} -0.38 \\ 0.92 \end{bmatrix} \tag{2}$$

Solutions

Finding a solution for $y_0'(t)$ and $y_1'(t)$ using the eigen thingies, proposition 22.3 [1] says a solution is given using the eigenvalues and eigenvectors as such:

$$y(t) = c_0 v_0 e^{\lambda_0 t} + c_1 v_1 e^{\lambda_1 t}$$

Inserting the values from (1) and (2).

$$y(t) = c_0 \begin{bmatrix} 0.92 \\ 0.38 \end{bmatrix} e^{-0.09t} + c_1 \begin{bmatrix} -0.38 \\ 0.92 \end{bmatrix} e^{-0.51t}$$

To isolate and determine c_0 and c_1, the initial state for $y(0)$ can be used.

$$y(0) = \begin{bmatrix} 84 \\ 36 \end{bmatrix} = c_0 \begin{bmatrix} 0.92 \\ 0.38 \end{bmatrix} + c_1 \begin{bmatrix} -0.38 \\ 0.92 \end{bmatrix} = \begin{bmatrix} c_0 \cdot 0.92 + c_1 \cdot -0.38 \\ c_0 \cdot 0.38 + c_1 \cdot -0.92 \end{bmatrix}$$

This gives us the following system:

$$\begin{aligned} 0.92c_0 - 0.38c_1 &= 84 \\ 0.38c_0 + 0.92c_1 &= 36 \end{aligned}$$

Using the god-given gift of numpy:

```
B = np.array([[0.92, -0.38],
              [0.38, 0.92]])

b = np.array([84, 36])

c0, c1 = np.linalg.solve(A, b)
```

Appendix

Only code used is explicitely shown when applied.

References

[1] Andrew Swan. *Numerisk Lineær Algebra*. Institut for Matematik, Aarhus Universitet, 2025.