Socket Preservation with Ivory Graft and Free Gingival Graft (Tooth 11)

Clinical Case Report

Graft Material Used: 100% Ivory Dentin Graft
Application: Socket preservation after extraction of tooth 11, covered with Free Gingival Graft (FGG) from the tuberosity.
Performed by: Dr. Avi Kuperschlag



Skip to content

1 Pre-operative condition

A frontal clinical image showing the anterior maxilla prior to treatment. Tooth 11 is present but non-restorable, with aesthetic and functional concerns. The surrounding gingival architecture appears healthy, making the case suitable for extraction and socket preservation in preparation for implant placement.



2 Tooth extraction

Tooth 11 was atraumatically extracted. Examination of the socket confirmed intact bony walls and adequate vascularity, making the site suitable for grafting.



Skip to content

3 Socket grafting

Ivory Dentin Graft was placed into the socket to preserve alveolar ridge dimensions and promote natural bone regeneration.



4 Free gingival graft (FGG) from the tuberosity

A soft tissue graft was harvested from the tuberosity area and used to cover the grafted socket, providing stability and soft tissue closure.



Skip to content

5 Socket sealed and sutured

The site was sealed with the FGG and secured with sutures. A membrane could also be used as an alternative to soft tissue grafts for socket coverage.



6 Reopening 4 months post-op

Re-entry at 4 months revealed mature, regenerated bone with excellent integration of the graft, supporting ideal conditions for implant placement.



Skip to content

7 Implant Placement

A dental implant was placed with high primary stability, allowing immediate loading. The regenerated ridge fully supported the implant body.




8 Temporary crown



Skip to content


9 Pre-op CBCT


CBCT showing the compromised condition of the alveolar ridge and root of tooth 11 prior to extraction and grafting.




10 Post-op CBCT – 4 Months

CBCT 4 months post operation - The excellent integration of Ivory graft allowed the implant to be placed with high primary stability allowing immediate loading.





Pre-op CBCT



CBCT · 4 months

Skip to content

Case Outcome:

The combination of Ivory Dentin Graft and a free gingival graft from the tuberosity provided an effective socket preservation approach. At 4 months post-op, the regenerated site supported immediate implant loading with high primary stability — demonstrating a predictable, patient-friendly approach to anterior implant rehabilitation.