

Advanced Methods for Supervised Interval Variable Selection

1. Partial Least Squares Regression

2. LAR/LASSO

1

1

Objectives

- Describe LAR / LASSO.
- Explain how to use LAR / LASSO in SAS.
- Discuss advantages and disadvantages with this variable selection method.

2

2

LAR / LASSO

- Target used or not?
 - Used
- Original or constructed variables as output?
 - Original variables

3

3

Use of the LARS

- LARS can be used for two main tasks:
 - Variable selection
 - Model-fitting and prediction

4

4

LARS Used for Variable Selection

- The coefficients for the potential input variables are continuously grown from zero to the final coefficient estimate.
- The input variables with 0 as the coefficient would be rejected.

5

5

Available Algorithms

- The following algorithms are available:
 - Least Angle Regression (LAR)
 - Least Absolute Shrinkage and Selection Operator (LASSO)
 - Adaptive LASSO
 - None -- Ordinary least squares regression
 - Others – forward, backward, stepwise, and elasticnet

6

6

Least Angle Regression: Basics

- Efron, B., Hastie, T., Johnstone, I. and Tibshirani, R. (2004)
- *Least angle regression*
 - As in forward selection, a sequence of regression models is produced.
 - In each step, one parameter is added to the model.
 - The complete model (all parameters entered into the model) corresponds to the full least squares solution.
 - Complexity of the model can be optimized on validation data.

7

7

Least Angle Regression: Algorithm

- First: Zero coefficients and zero predicted response.
- Find the input variable with highest correlation with the target (least angle with...).
- A step is taken in the direction of this input variable. This creates a residual vector now considered the response.
- Determination of step length:
 - Some other input variable has the same correlation with the residual vector as the first variable does with the response.

8

8

Least Angle Regression: Algorithm


- The predicted response moves in the equiangular direction of these two input variables.
 - Movement until a third input variable has the same correlation with the residual as the two input variables now already in the model.
- A new step direction is determined:
 - Equiangular between the three input variables in the model.
- The predicted response moves again.
- Process continues until all input variables are in the model.

9

9

Least Angle Regression Geometrically

- Two covariates (predictors) x_1 and x_2 and the space $L(x_1, x_2)$ that is spanned by them
- $\mu = E(Y|X)$




10

10

Least Angle Regression Geometrically

- Three covariates (predictors) x_1 , x_2 and x_3 and the space $L(x_1, x_2, x_3)$ that is spanned by them





11

11

Coefficient Traces

For 50 steps,

Forward Selection →



12

Least Angle Regression: Mathematics

Preliminaries:

- $\sum_{i=1}^n y_i = 0$, $\sum_{i=1}^n x_{ij} = 0$, and $\sum_{i=1}^n x_{ij}^2 = 1$, for $j = 1, \dots, p$
- $\mu = E(Y|X)$, $\hat{\mu}$ is the estimated μ
- Current correlations: $c(\hat{\mu}) = X^T(Y - \hat{\mu})$
- The absolute correlations are related to the angles of the current residuals with X_j 's
- Cosine of the angle between two vectors vs. Correlation between two variables

$$\cos(a, b) = \frac{a \cdot b}{\|a\| \cdot \|b\|}$$

$$\rho_{X,Y} = \frac{\sum_{i=1}^n x_i y_i}{\sqrt{\sum_{i=1}^n x_i^2} \sqrt{\sum_{i=1}^n y_i^2}}$$

13

Least Angle Regression: Mathematics

The Equiangular Vectors:

- Let \mathcal{A} be the set of indices corresponding to covariates in the current model
- Let $X_{\mathcal{A}} = [\dots s_j X_j \dots]_{j \in \mathcal{A}}$, where $s_j = \pm 1$
- $U_{\mathcal{A}} = X_{\mathcal{A}} w_{\mathcal{A}}$: the unit vector making equal angles ($< 90^\circ$) with the columns of $X_{\mathcal{A}}$
- $X_{\mathcal{A}}^T U_{\mathcal{A}} = A_{\mathcal{A}} \mathbf{1}_{\mathcal{A}}$ and $\|U_{\mathcal{A}}\| = 1$, where $A_{\mathcal{A}}$ is a constant.
- Then $w_{\mathcal{A}} = A_{\mathcal{A}} G_{\mathcal{A}}^{-1} \mathbf{1}_{\mathcal{A}}$ and $A_{\mathcal{A}} = (\mathbf{1}_{\mathcal{A}}^T G_{\mathcal{A}}^{-1} \mathbf{1}_{\mathcal{A}})^{-1/2}$ where $G_{\mathcal{A}} = X_{\mathcal{A}}^T X_{\mathcal{A}}$

14

14

Least Angle Regression: Mathematics

Current Correlations

- ▶ $\hat{c} = X^T(Y - \hat{\mu}_{\mathcal{A}})$
- ▶ Let $\hat{C} = \max_j \{|\hat{c}_j|\}$.
- ▶ $\mathcal{A} = \{j : |\hat{c}_j| = \hat{C}\}$
- ▶ Let $a = X^T U_{\mathcal{A}}$.
- ▶ Consider $\mu(\gamma) = \hat{\mu}_{\mathcal{A}} + \gamma U_{\mathcal{A}}$. Then $c_j(\gamma) = X_j^T(Y - \mu(\gamma)) = \hat{c}_j - \gamma a_j$.
- ▶ For $j \in \mathcal{A}$, $|c_j(\gamma)| = \hat{C} - \gamma A_{\mathcal{A}}$.

15

15

Least Absolute Shrinkage and Selection Operator (LASSO)

- A constrained form of ordinary least squares is used:
 - The sum of the absolute values of the regression coefficients must be smaller than a certain value.
- The LASSO coefficients $\beta=(\beta_1, \beta_2, \beta_3, \dots, \beta_p)$ are the solution to

$$\begin{aligned} &\text{Minimize } \|y - X\beta\|^2 \\ &\text{subject to } \sum_{j=1}^p |\beta_j| \leq t \end{aligned}$$

- A quadratic programming algorithm is used to compute the coefficients

16

16

LASSO vs. Ridge Regression

- Ridge regression also shrinks the regression coefficients by imposing a penalty on their size.
- The Ridge coefficients $\beta=(\beta_1, \beta_2, \beta_3, \dots, \beta_p)$ are the solution to

$$\begin{aligned} &\text{Minimize } \|y - X\beta\|^2 \\ &\text{subject to } \sum_{j=1}^p \beta_j^2 \leq t \end{aligned}$$

- Which is equivalent to

17

17

Ridge Regression Shrinkage (Mathematics)

- The SVD of X has the form:

$$\mathbf{X} = \mathbf{U}\mathbf{D}\mathbf{V}^T$$

- Dimensions:

- U is N-by-p orthogonal matrix, with columns spanning the column space of X
- V is p-by-p orthogonal matrix, with columns spanning the row space of X
- D is p-by-p diagonal matrix, with diagonal entries $d_1 \geq d_2 \geq \dots \geq d_p \geq 0$ called the singular values of X

18

18

Ridge Regression Shrinkage (Mathematics)

- Using the SVD of \mathbf{X} we can re-write the OLS solutions

$$\begin{aligned}\mathbf{X}\hat{\beta}^{\text{ls}} &= \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\mathbf{y} \\ &= \mathbf{U}\mathbf{U}^T\mathbf{y},\end{aligned}$$

and the ridge solutions:

$$\begin{aligned}\mathbf{X}\hat{\beta}^{\text{ridge}} &= \mathbf{X}(\mathbf{X}^T\mathbf{X} + \lambda\mathbf{I})^{-1}\mathbf{X}^T\mathbf{y} \\ &= \mathbf{U}\mathbf{D}(\mathbf{D}^2 + \lambda\mathbf{I})^{-1}\mathbf{D}\mathbf{U}^T\mathbf{y} \quad \text{the shrinkage} \\ &= \sum_{j=1}^p \mathbf{u}_j \frac{d_j^2}{d_j^2 + \lambda} \mathbf{u}_j^T \mathbf{y},\end{aligned}$$

19

19


LASSO vs. Ridge Regression

- The change to the penalty function is subtle, but has a dramatic impact on the resulting estimator.
- Like ridge regression, penalizing the absolute values of the coefficients introduces shrinkage towards zero.
- However, unlike ridge regression, some of the coefficients are shrunk all the way to zero; such solutions, with multiple values that are identically zero, are said to be **sparse**.
- The penalty thereby performs a sort of continuous variable selection
- The resulting estimator was thus named the lasso, for "Least Absolute Shrinkage and Selection Operator"

20

20

Ridge Regression Shrinkage (Example)




$$\begin{aligned}\text{df}(\lambda) &= \text{tr}[\mathbf{X}(\mathbf{X}^T\mathbf{X} + \lambda\mathbf{I})^{-1}\mathbf{X}^T] \\ &= \text{tr}(\mathbf{H}_\lambda) \\ &= \sum_{j=1}^p \frac{d_j^2}{d_j^2 + \lambda}.\end{aligned}$$

Reference: T. Hastie, R. Tibshirani, and J. Friedman,
The Elements of Statistical Learning.

21

21

LASSO Shrinkage (Example)




$$s = \frac{t}{\sum_{j=1}^p |\hat{\beta}_j|}$$

Reference: T. Hastie, R. Tibshirani, and J. Friedman,
The Elements of Statistical Learning.

22

22

Ridge Shrinkage vs. LASSO Shrinkage (Example)



23

23

Adaptive LASSO

- This is a modification of the LASSO algorithm.
- The basics are the same but weights are applied to the parameters in the LASSO constraint.
- A weight vector is defined as

$$w = \frac{1}{|\hat{\beta}|^\gamma}, \text{ with } \gamma \geq 0$$

- $\hat{\beta}$ is an estimate of the parameters.

24

24

Adaptive LASSO

- The adaptive LASSO coefficients $\beta = (\beta_1, \beta_2, \beta_3, \dots, \beta_p)$ are the solution to

$$\text{Minimize } \|y - X\beta\|^2$$

$$\text{subject to } \sum_{j=1}^p |w_j \beta_j| \leq t$$

25

25

Model Selection

- Two issues are important in the model selection:
 - How many steps should be processed?

Path Stopping Criterion	Maximum Steps
Maximum Steps	200


- From which of the steps should the model be selected?

Model Selection Criterion	SBC
---------------------------	-----

26

26

How Many Steps Should Be Processed?




27

27

From Which of the Steps Should the Model Be Selected?

- How can we select the best model from the sequence of constructed models?



28

28

From Which of the Steps Should the Model Be Selected?

- The available model selection criteria include
 - SBC – Schwartz Bayesian information criterion
 - BIC – Bayesian information criterion
 - AIC – Akaike information criterion
 - AICC -- Corrected Akaike's information criterion
 - CP – Cp statistic, explained in Regression class
 - Validation – explained in DM I class
 - Cross Validation – explained in PLS slides
 - ADJRSQ, CVEX, PRESS

29

29

Performance Assessment: Loss Function

- Typical choices for quantitative response Y :

$$L(Y, \hat{f}(X)) = \begin{cases} (Y - \hat{f}(X))^2 & \text{(squared error)} \\ |Y - \hat{f}(X)| & \text{(absolute error)} \end{cases}$$

- Typical choices for categorical response G :

$$L(G, \hat{G}(X)) = I(G \neq \hat{G}(X)) \quad (0-1 \text{ loss})$$

$$L(G, \hat{p}(X)) = -2 \sum_{k=1}^K I(G = k) \log \hat{p}_k \\ = -2 \log \hat{p}_G(X) \quad (\text{log-likelihood})$$

30

30

In-sample and Extra-sample Error

- In-sample error** is the average prediction error, conditioned on the training sample \mathbf{x} . It is obtained when new responses are observed for the training set features.

$$Err_{in} = \frac{1}{N} \sum_{i=1}^N Err(x_i) = \frac{1}{N} \sum_{i=1}^N E_y E_{Y^{New}} L(Y_i^{New}, \hat{f}(x_i)).$$

- Extra-sample error** is the average prediction error when both features and responses are new (no conditioning on the training set).

31

31

In-sample Error

- For squared error, 0-1, and other loss function, it can be shown generally that

$$Err_{in} = E_y(\overline{err}) + \frac{2}{N} \sum_{i=1}^N Cov(\hat{y}_i, y_i).$$

- Can be simplified as $Err_{in} = E_y(\overline{err}) + 2 \cdot \frac{d}{N} \sigma_\varepsilon^2$ for the model $Y = f(X) + \varepsilon$ by a linear fit with d inputs.

32

32

Model Selection Criteria

- C_p statistic (when d free parameters are fitted under squared error loss):

$$C_p = \overline{err} + 2 \cdot \frac{d}{N} \hat{\sigma}_\varepsilon^2$$

where $\overline{err} = \frac{1}{N} \sum_{i=1}^N L(y_i, \hat{f}(x_i))$ is the training error rate.

- C_p estimates the in-sample error.

33

33

Model Selection Criteria

- AIC (Akaike information criterion), a more generally applicable estimate of the in-sample error when a log-likelihood loss function is used (when $N \rightarrow \infty$):

$$\begin{aligned} -2E[\log \Pr_{\hat{\theta}}(Y)] &= -\frac{2}{N} E\left(\sum_{i=1}^N \log \Pr_{\hat{\theta}}(y_i)\right) + 2 \frac{d}{N} \\ &= -\frac{2}{N} \cdot \log \text{lik} + 2 \frac{d}{N} = \frac{AIC}{N} \end{aligned}$$

- General form: $AIC = -2 \cdot (\log \text{lik}) + 2 \cdot d$.
- Choose the model giving smallest AIC over the set of models considered

34

34

Model Selection Criteria

- BIC (Bayesian information criterion) is motivated from Bayesian point of view.
- BIC tends to penalize complex models more heavily, giving preference to simpler models in selection
- Its general form is:

$$BIC = -2 \cdot (\log \text{lik}) + (\log N) \cdot d.$$

35

35

Bayesian Model Selection

- Suppose we have candidate models $M_m, m=1, \dots, M$
- with corresponding model parameters θ_m .
- Prior distribution: $\Pr(\theta_m | M_m), m=1, \dots, M$.
- Posterior probability:

$$\Pr(M_m | Z) \propto \Pr(M_m) \cdot \Pr(Z | M_m).$$
- Compare two models via posterior odds:

$$\frac{\Pr(M_m | Z)}{\Pr(M_l | Z)} = \frac{\Pr(M_m)}{\Pr(M_l)} \cdot \frac{\Pr(Z | M_m)}{\Pr(Z | M_l)}$$
- The second factor on the RHS is called the Bayes factor and describes the contribution of the data towards posterior odds.

36

36

Bayesian Approach Continued

- Unless strong evidence to the contrary, we typically assume that prior over models is uniform (non-informative prior).
- Using Laplace approximation, one can establish a simple (but approximate) relationship between posterior model probability and the BIC.
- Lower BIC implies higher posterior probability of the model. Use of BIC as model selection criterion is thus justified.

37

37

AIC or BIC?

- BIC is asymptotically consistent as a selection criterion. That means, given a family of models including the true model, the probability that BIC will select the correct one approaches one as the sample size becomes large.
- AIC does not have the above property. Instead, it tends to choose more complex models as $N \rightarrow \infty$.
- For small or moderate samples, BIC often chooses models that are too simple, because of its heavy penalty on complexity.

38

38

LARS vs. LASSO




FIGURE 3.15. Left panel shows the LARS coefficient profiles on the simulated data, as a function of the L_1 arc length. The right panel shows the Lasso profile. They are identical until the dark-blue coefficient crosses zero at an arc length of

Reference: T. Hastie, R. Tibshirani, and J. Friedman,
The Elements of Statistical Learning.

39

39

LAR / LASSO: Pros

- Variable selection and model fitting integrated in the same node
- Computationally efficient
- Original variables as output

40

40

LAR / LASSO: Cons

- Difficult to know at which step to stop the modeling process.
- Mathematically challenging to understand.
- Degrees of freedom used in the LASSO algorithm are not mathematically justified, but works in practice.

41

41

LAR / LASSO: Degrees of Freedom

- Df of an fitted vector \hat{y} as

$$df(\hat{y}) = \frac{1}{\sigma^2} \sum_{i=1}^N \text{Cov}(\hat{y}_i, y_i).$$

- $\text{Cov}(\hat{y}, y)$ refers to the sampling covariance
- After the k th step of the LAR procedure, the effective degrees of freedom of the fit vector is exactly k .
- For the lasso, at any stage $df(\hat{y})$ approximately equals the number of predictors in the model.
- A detailed study of the degrees of freedom for the lasso may be found in Zou et al. (2007).

42

42



LAR / LASSO for Variable Selection


- This demonstration illustrates how to use the LARS to perform variable selection.
- Variations of the LARS node/procedure are shown.

43

43

A Comparison of the Selection and Shrinkage

- Consider a simulated example with two correlated inputs X_1 and X_2 , with correlation ρ . We assume that the true regression coefficients are $\beta_1 = 4$ and $\beta_2 = 2$.



- A full study: Frank and Friedman (1993). These authors conclude that for minimizing the prediction error, ridge regression is generally preferable to variable subset selection, PC regression, and PLS

44

44

Questions?



45

45
