

ABSOLUTE CALIBRATION FOR WIDE-FIELD SURVEYS

GAUTHAM NARAYAN
STSCI



CALSPEC+

NOAO: Abi Saha, Tom Matheson, Sean Points, Jenna Claver

UA: Tim Axelrod, Ed Olszewski, Jay Holberg, Ivan Hubeny

STScI: Annalisa Calamida, Ralph Bohlin, Susana Deustua, Elena Sabbi, Armin Rest, John MacKenty

Harvard: Christopher Stubbs

Cambridge: Kaisey Mandel



THE UNIVERSITY
OF ARIZONA

WHAT THE FLUX

GAUTHAM NARAYAN
STSCI



CALSPEC+

NOAO: Abi Saha, Tom Matheson, Sean Points, Jenna Claver


UA: Tim Axelrod, Ed Olszewski, Jay Holberg, Ivan Hubeny

STScI: Annalisa Calamida, Ralph Bohlin, Susana Deustua, Elena Sabbi, Armin Rest, John MacKenty

Harvard: Christopher Stubbs


Cambridge: Kaisey Mandel






The statistical errors tell us that we
should have a robust sub-10%
measurement of the equation of state of
dark energy with our current sample size

“What could possibly go wrong?”




WE DON'T.


If you are sick of hearing that $w = -1$ but we're systematics dominated, then you should care about calibration.



The typical calibration chain for wide-field surveys



Most standards are extremely complex and hard to model




Most standards are extremely complex and hard to model




Definitely
not a
blackbody.

Or a tree.

I am not really confident, but I think it's a map of the tree.






Establishing new standards from the ground is hard, particularly in the NIR


(where supernovae are better standard candles, dust is less of an issue, and where JWST and WFIRST are going to look)

**OUR MOST PRECISELY CALIBRATED PHOTOMETRIC SYSTEM ISN'T
ON THE GROUND**





It's possible to simplify this mess.



Much cleaner!

EXTENDING HST'S CALIBRATION TO THE ENTIRE SKY

REBUILDING CALSPEC

REBUILDING CALSPEC

- Transfer HST's high precision calibration to ground based facilities

REBUILDING CALSPEC

- Transfer HST's high precision calibration to ground based facilities
- Obtain ground based spectroscopy, above atmosphere HST photometry of DA white dwarfs. GO 12967 (18 orbits), 13711 (60 orbits), 15113 (54 orbits, ongoing)


REBUILDING CALSPEC

- Transfer HST's high precision calibration to ground based facilities
- Obtain ground based spectroscopy, above atmosphere HST photometry of DA white dwarfs. GO 12967 (18 orbits), 13711 (60 orbits), 15113 (54 orbits, ongoing)
- Why DA WDs? Simple pure Hydrogen atmospheres - most accurate models we have

REBUILDING CALSPEC

- Transfer HST's high precision calibration to ground based facilities
- Obtain ground based spectroscopy, above atmosphere HST photometry of DA white dwarfs. GO 12967 (18 orbits), 13711 (60 orbits), 15113 (54 orbits, ongoing)
- Why DA WDs? Simple pure Hydrogen atmospheres - most accurate models we have
- Faint - $V \sim 16.5\text{--}19.5$ - can observe them directly with LSST

- ★ CALSPEC Standards
- ☆ Secondary Standards
- ★ Cycle 20+22 Targets
- ★ Cycle 25 Targets



Spectra for ~35 DA WDs (MMT, Gemini, SOAR)

HST photometry for 23 northern + equatorial candidates

Always at least three standards from any site - some close to proposed LSST deep drilling footprint

Proof-of -concept with first four objects we had data for in Cycle 20 (Narayan+, 2016)


“What could possibly go wrong? Part Deux”

Spectroscopy constrains T_{eff} , $\log(g)$

Pick targets with
 $22000 \text{K} < T_{\text{eff}} < 75000 \text{K}$
 $7 < \log(g) < 8.5$



Faint targets are distant
- interstellar extinction


Multiband HST
photometry from
drizzled images to
establish absolute flux,
constrain reddening




Last generation of white dwarf atmosphere codes = de-reddened and flattened spectra, extracted Balmer lines, separately fit for reddening.

Ignored covariance, between parameters, correlated residuals, was very susceptible to bias with lower S/N data











We took our calibration, checked against GD153 & we found issues!
 UVIS1/2 have very different QE.
 MAST data products from late 2016 on now reflect this.



"This is good, but that trend we corrected for was kinda shocking! Ralph/Susanna - how well do we know these WFC3 zero points actually?"





LESSONS FROM CYCLE 20

LESSONS FROM CYCLE 20

- WFC3 flux calibration is a lot less solid than we'd have liked

LESSONS FROM CYCLE 20

- WFC3 flux calibration is a lot less solid than we'd have liked
- Drizzling, cosmic rays AND cosmic-ray rejection, and faint background galaxies in the IR can wreak havoc with your photometry

LESSONS FROM CYCLE 20

- WFC3 flux calibration is a lot less solid than we'd have liked
- Drizzling, cosmic rays AND cosmic-ray rejection, and faint background galaxies in the IR can wreak havoc with your photometry
- Really need bluer bands to constrain Av

LESSONS FROM CYCLE 20

- WFC3 flux calibration is a lot less solid than we'd have liked
- Drizzling, cosmic rays AND cosmic-ray rejection, and faint background galaxies in the IR can wreak havoc with your photometry
- Really need bluer bands to constrain Av
- Fitting procedure has gremlins

LESSONS FROM CYCLE 20

- WFC3 flux calibration is a lot less solid than we'd have liked
Monitor CALSPEC primary standard too.
- Drizzling, cosmic rays AND cosmic-ray rejection, and faint background galaxies in the IR can wreak havoc with your photometry
- Really need bluer bands to constrain Av
- Fitting procedure has gremlins

LESSONS FROM CYCLE 20

- WFC3 flux calibration is a lot less solid than we'd have liked
Monitor CALSPEC primary standard too.
- Drizzling, cosmic rays AND cosmic-ray rejection, and faint background galaxies in the IR can wreak havoc with your photometry
Write your own HST pipeline.
- Really need bluer bands to constrain Av
- Fitting procedure has gremlins

LESSONS FROM CYCLE 20

- WFC3 flux calibration is a lot less solid than we'd have liked
Monitor CALSPEC primary standard too.
- Drizzling, cosmic rays AND cosmic-ray rejection, and faint background galaxies in the IR can wreak havoc with your photometry
Write your own HST pipeline.
- Really need bluer bands to constrain Av
Get F275W measurements.
- Fitting procedure has gremlins

LESSONS FROM CYCLE 20

- WFC3 flux calibration is a lot less solid than we'd have liked
Monitor CALSPEC primary standard too.
- Drizzling, cosmic rays AND cosmic-ray rejection, and faint background galaxies in the IR can wreak havoc with your photometry
Write your own HST pipeline.
- Really need bluer bands to constrain Av
Get F275W measurements.
- Fitting procedure has gremlins
New model + require consistency with a different approach.

LESSONS FROM CYCLE 20


- WFC3 flux calibration is a lot less solid than we'd have liked
Monitor CALSPEC primary standard too.
- Drizzling, cosmic rays AND cosmic-ray rejection, and faint background galaxies in the IR can wreak havoc with your photometry
Write your own HST pipeline.
- Really need bluer bands to constrain Av
Get F275W measurements.
- Fitting procedure has gremlins
New model + require consistency with a different approach.

1% photometry is hard. Below 1% is very hard.


"So if we just fix those things, we're good right?"

NEW PHOTOMETRY PIPELINE

- Construct flc images from MAST data products to get rid of CTE bleed trails
- Turn off CR rejection since it can change PSFs (F625W, F775W)
- Photometry in sequence of apertures, careful aperture corrections for every image
- Bayesian hierarchical model to solve for mean magnitudes, C20 to C22 zeropoint offsets, zero points, WFC3 dispersion*




NEW MODELS



- Approach 1: Fit spectra + apparent magnitudes (i.e. tied to CALSPEC standards)
<http://wdmodel.rtfd.io>
- Approach 2: Fit spectra + instrumental magnitudes of CALSPEC standards + our DA WDs to put everything on a common system
<https://github.com/taxelrod/WDFitPhot>
- Colors, parameters should agree with both methods


SDSSJ-081508



Model HST photometry + spectroscopy (observations AND their noise properties) at the same time

Gaussian Processes for the noise, MPI to distribute computation (we've run it on 32 nodes at once)

Easily extensible if you want to model binary WD spectra or add other parameters



Gemini GMOS

$$T_E = 32979.712^{+387.953}_{-375.679} \text{ K}$$

$$\log(g) = 7.292^{+0.088}_{-0.091} \text{ dex}$$

$$A_V = 0.040^{+0.007}_{-0.007} \text{ mag}$$

$$d_l = 445.621^{+3.477}_{-3.432}$$

$$\text{FWHM} = 4.210^{+1.701}_{-1.795} \text{ \AA}$$

$$f_\sigma = 0.670^{+0.149}_{-0.112}$$

$$\mu = 59.994^{+0.016}_{-0.015} \text{ mag}$$

MMT Blue Channel

$$T_E = 32138.078^{+568.803}_{-513.897} \text{ K}$$

$$\log(g) = 7.369^{+0.171}_{-0.154} \text{ dex}$$

$$A_V = 0.024^{+0.012}_{-0.012} \text{ mag}$$

$$d_l = 579.706^{+7.467}_{-7.293}$$


$$\text{FWHM} = 6.500^{+2.640}_{-2.727} \text{ \AA}$$

$$f_\sigma = 0.686^{+0.142}_{-0.107}$$

$$\mu = 59.958^{+0.025}_{-0.024} \text{ mag}$$

“What could possibly go wrong? Round Three”

ILAPHv3 f99 nodisp Photometric Residuals by Passband



Done with analysis of Cycle 20 + 22 data


We can characterize zeropoint differences relative to CALSPEC (minuscule), overall dispersion, and the RMS

The RMS is less than 0.01 mag in every single band (covering NUV to NIR) - better than 0.005 mag in the optical

WHO ORDERED THAT?

F160W?????

ILAPHv3 f99 nodisp F160W residuals vs F625W



WHAT'S GOING ON IN F160W?

WHAT'S GOING ON IN F160W?

- Not dust - no trend with Av - just brightness

WHAT'S GOING ON IN F160W?

- Not dust - no trend with Av - just brightness
 - We tried several different dust laws + fitting the dust law

WHAT'S GOING ON IN F160W?

- Not dust - no trend with Av - just brightness
 - We tried several different dust laws + fitting the dust law
 - Reddening estimates agree well with dust maps

WHAT'S GOING ON IN F160W?

- Not dust - no trend with A_V - just brightness
 - We tried several different dust laws + fitting the dust law
 - Reddening estimates agree well with dust maps
- Not the model - no trend with T_{eff} - just brightness

WHAT'S GOING ON IN F160W?

- Not dust - no trend with A_V - just brightness
 - We tried several different dust laws + fitting the dust law
 - Reddening estimates agree well with dust maps
- Not the model - no trend with T_{eff} - just brightness
 - We tried a different model grid too (both could still be wrong)

WHAT'S GOING ON IN F160W?

- Not dust - no trend with A_v - just brightness
 - We tried several different dust laws + fitting the dust law
 - Reddening estimates agree well with dust maps
- Not the model - no trend with T_{eff} - just brightness
 - We tried a different model grid too (both could still be wrong)
- Not different PSF shape for bright vs faint objects

WHAT'S GOING ON IN F160W?

- Not dust - no trend with A_V - just brightness
 - We tried several different dust laws + fitting the dust law
 - Reddening estimates agree well with dust maps
- Not the model - no trend with T_{eff} - just brightness
 - We tried a different model grid too (both could still be wrong)
- Not different PSF shape for bright vs faint objects
 - 7 different reductions of the data. Differences remain for very wide range of aperture & annulus

WHAT'S GOING ON IN F160W?

- Not dust - no trend with A_V - just brightness
 - We tried several different dust laws + fitting the dust law
 - Reddening estimates agree well with dust maps
- Not the model - no trend with T_{eff} - just brightness
 - We tried a different model grid too (both could still be wrong)
- Not different PSF shape for bright vs faint objects
 - 7 different reductions of the data. Differences remain for very wide range of aperture & annulus
- Not passband model - that'd affect all photometry

WHAT'S GOING ON IN F160W?

- Not dust - no trend with A_V - just brightness
 - We tried several different dust laws + fitting the dust law
 - Reddening estimates agree well with dust maps
- Not the model - no trend with T_{eff} - just brightness
 - We tried a different model grid too (both could still be wrong)
- Not different PSF shape for bright vs faint objects
 - 7 different reductions of the data. Differences remain for very wide range of aperture & annulus
- Not passband model - that'd affect all photometry
- Possibilities left:


WHAT'S GOING ON IN F160W?

- Not dust - no trend with A_V - just brightness
 - We tried several different dust laws + fitting the dust law
 - Reddening estimates agree well with dust maps
- Not the model - no trend with T_{eff} - just brightness
 - We tried a different model grid too (both could still be wrong)
- Not different PSF shape for bright vs faint objects
 - 7 different reductions of the data. Differences remain for very wide range of aperture & annulus
- Not passband model - that'd affect all photometry
- Possibilities left:
 - Fainter objects are longer exposures, and there is unmodeled thermal background - hard to rule out with existing data, but will with C25 data

WHAT'S GOING ON IN F160W?

- Not dust - no trend with A_V - just brightness
 - We tried several different dust laws + fitting the dust law
 - Reddening estimates agree well with dust maps
- Not the model - no trend with T_{eff} - just brightness
 - We tried a different model grid too (both could still be wrong)
- Not different PSF shape for bright vs faint objects
 - 7 different reductions of the data. Differences remain for very wide range of aperture & annulus
- Not passband model - that'd affect all photometry
- Possibilities left:
 - Fainter objects are longer exposures, and there is unmodeled thermal background - hard to rule out with existing data, but will with C25 data
 - WFC3/IR exhibits some count rate non-linearity

WDModel Conservative Fit Results




Plot created by Douglas Tucker showing the ratio of the fit values for Teff from the 2 sources vs. the Teff from Pier-Emmanuel's fits. Color-coded to show the absolute deviation of the WDmodel fit value for $E(B-V)$

Courtesy, Douglas Tucker, Deborah Gulleidge, DES

WDModel Conservative Fit Results


Grid only goes down to 16,000K



Plot created by Douglas Tucker showing the ratio of the fit values for Teff from the 2 sources vs. the Teff from Pier-Emmanuel's fits. Color-coded to show the absolute deviation of the WDmodel fit value for E(B-V)

Courtesy, Douglas Tucker, Deborah Gulleidge, DES


EXCELLENT AGREEMENT WITH DUST MAP




Also in good agreement with estimates from Green et al. 3D dust map from PS1

With full sample, should be able to fit for dust law shape independently from entire sample (assumes all objects have the same dust law).

MULTI-YEAR LCO MONITORING TO DETECT VARIABILITY



MULTI-YEAR LCO MONITORING TO DETECT VARIABILITY



CALSPEC+ STATUS

CALSPEC+ STATUS

- Dealing with non-linearity in F160W? Hic sunt dracones...

CALSPEC+ STATUS

- Dealing with non-linearity in F160W? Hic sunt dracones...
- Irrelevant for LSST - if you exclude F160W, the RMS is less than 4 mmag from 2750 Å — 1 micron

CALSPEC+ STATUS

- Dealing with non-linearity in F160W? Hic sunt dracones...
- Irrelevant for LSST - if you exclude F160W, the RMS is less than 4 mmag from 2750 Å — 1 micron
- Excellent agreement between both fitting approaches, dust map, different model grids

CALSPEC+ STATUS

- Dealing with non-linearity in F160W? Hic sunt dracones...
- Irrelevant for LSST - if you exclude F160W, the RMS is less than 4 mmag from 2750 Å — 1 micron
- Excellent agreement between both fitting approaches, dust map, different model grids
- TODO: direct comparison to GAIA distances - SOON!

CALSPEC+ STATUS

- Dealing with non-linearity in F160W? Hic sunt dracones...
- Irrelevant for LSST - if you exclude F160W, the RMS is less than 4 mmag from 2750 Å — 1 micron
- Excellent agreement between both fitting approaches, dust map, different model grids
- TODO: direct comparison to GAIA distances - SOON!
- Extends CALSPEC across the sky - Data (Calamida+ in prep) and analysis (Narayan+ in prep)

FOR JWST

FOR JWST

- F stars + solar analogs in addition to white dwarfs (not enough flux for MIRI)

FOR JWST

- F stars + solar analogs in addition to white dwarfs (not enough flux for MIRI)
- Targets selected for a “slim” Cycle 1 calibration program (K. Gordon, G. Reike...)

FOR JWST




- F stars + solar analogs in addition to white dwarfs (not enough flux for MIRI)
- Targets selected for a “slim” Cycle 1 calibration program (K. Gordon, G. Reike...)
- Detailed modeling of stellar atmospheres + dust at beyond 5 micron is an open problem

Modeling in the NIR has large differences

Three stellar SEDs,
each 8790 K

Black = Pickles SED
Red = 1732526*
Green = 1802271*

* (STIS + NICMOS)




FOR JWST

- F stars + solar analogs in addition to white dwarfs (not enough flux for MIRI)
- Targets selected for a “slim” Cycle 1 calibration program (K. Gordon, G. Reike...)
- Detailed modeling of stellar atmospheres + dust at beyond 5 micron is an open problem

FOR JWST

- F stars + solar analogs in addition to white dwarfs (not enough flux for MIRI)
- Targets selected for a “slim” Cycle 1 calibration program (K. Gordon, G. Reike...)
- Detailed modeling of stellar atmospheres + dust at beyond 5 micron is an open problem
- ALL NEW SYSTEMATICS GUARANTEED!



For now, do this with our white dwarfs. The End.

