

Near-infrared SN Ia as standard candles

Arturo Avelino

CfA, Harvard

Andrew Friedman, Kaisey Mandel, Robert P. Kirshner and Peter Challis

Pittsburgh, April 2018

The problem

Optical samples of SN Ia for cosmology have reached their limit to constrain the dark energy (DE) because of the systematic uncertainties.


- More optical data *doesn't* mean better DE constraints.
- Optical light is dimmed and reddened by dust




A solution: NIR observations!

- Near infrared (**NIR**) light is much **less sensitive to dust** than the optical wavelengths.

Optical




Near infrared






Low-z NIR sample

Compiled by **Andrew Friedman**
(UCSD):

- CfA, CSP, Literature
- 190 SN Ia with optical + NIR (YJHK) light curves






Determine distances from the NIR data

Template method


Gaussian-Process method


Gaussian-Process fit


sn2005el





Gaussian-Process fit






























Distance modulus


$$\Delta m_s(\hat{t}) \equiv m_s(\hat{t}) - \mathcal{M}(\hat{t}) - m_{0,s} \quad (11)$$

where $m_s(\hat{t})$ and $\mathcal{M}(\hat{t})$ are the apparent magnitude and the magnitude of the normalized template at phase \hat{t} , respectively. We can express this difference for all the $N_{LC,s}$ phases in a given LC as the vector,

$$\Delta \mathbf{m}_s \equiv \begin{pmatrix} \Delta m_s(\hat{t}_1) \\ \Delta m_s(\hat{t}_2) \\ \vdots \\ \Delta m_s(\hat{t}_{N_{LC,s}}) \end{pmatrix}. \quad (12)$$

Then, to determine $m_{0,s}$ we minimize the negative of the log likelihood function $L(m_{0,s})$ defined as

$$-2 \ln L(m_{0,s}) = \Delta \mathbf{m}_s^\top \cdot C^{-1} \cdot \Delta \mathbf{m}_s \quad (13)$$



Distance modulus

$$\Delta m_s(\hat{t}) \equiv m_s(\hat{t}) - \mathcal{M}(\hat{t}) - m_{0,s} \quad (11)$$

where $m_s(\hat{t})$ and $\mathcal{M}(\hat{t})$ are the apparent magnitude and the magnitude of the normalized template at phase \hat{t} , respectively. We can express this difference for all the $N_{LC,s}$ phases in a given LC as the vector,

$$\Delta \mathbf{m}_s \equiv \begin{pmatrix} \Delta m_s(\hat{t}_1) \\ \Delta m_s(\hat{t}_2) \\ \vdots \\ \Delta m_s(\hat{t}_{N_{LC,s}}) \end{pmatrix}. \quad (12)$$

Then, to determine $m_{0,s}$ we minimize the negative of the log likelihood function $L(m_{0,s})$ defined as

$$-2 \ln L(m_{0,s}) = \Delta \mathbf{m}_s^\top \cdot C^{-1} \cdot \Delta \mathbf{m}_s \quad (13)$$

where C is the $N_{LC,s}$ -dimensional covariance matrix where the (\hat{t}_i, \hat{t}_j) component is given by:

$$C_{ij} \equiv \text{Cov} (\Delta m_s(\hat{t}_i), \Delta m_s(\hat{t}_j)) \quad (14)$$

$$= \sigma_{\mathcal{M}}(\hat{t}_i) \sigma_{\mathcal{M}}(\hat{t}_j) \exp \left[-\frac{(\hat{t}_i - \hat{t}_j)^2}{2l^2} \right] + \hat{\sigma}_{m,s}^2(\hat{t}_i) \delta_{ij} \quad (15)$$

where $\sigma_{\mathcal{M}}(\hat{t})$ is the population standard deviation of the sample distribution of magnitudes at time \hat{t} , determined from Eq. (B2) during the training process used to construct the mean LC template, with the hyperparameter l computed via Eq. (A6), while $\hat{\sigma}_{m,s}^2(\hat{t}_i)$ is the photometric error of the datum $m_s(\hat{t}_i)$.

Distance modulus

$$\Delta m_s(\hat{t}) \equiv m_s(\hat{t}) - \mathcal{M}(\hat{t}) - m_{0,s} \quad (11)$$

where $m_s(\hat{t})$ and $\mathcal{M}(\hat{t})$ are the apparent magnitude and the magnitude of the normalized template at phase \hat{t} , respectively. We can express this difference for all the $N_{LC,s}$ phases in a given LC as the vector,

$$\Delta \mathbf{m}_s \equiv \begin{pmatrix} \Delta m_s(\hat{t}_1) \\ \Delta m_s(\hat{t}_2) \\ \vdots \\ \Delta m_s(\hat{t}_{N_{LC,s}}) \end{pmatrix}. \quad (12)$$

Then, to determine $m_{0,s}$ we minimize the negative of the log likelihood function $L(m_{0,s})$ defined as

$$-2 \ln L(m_{0,s}) = \Delta \mathbf{m}_s^\top \cdot C^{-1} \cdot \Delta \mathbf{m}_s \quad (13)$$

where C is the $N_{LC,s}$ -dimensional covariance matrix where the (\hat{t}_i, \hat{t}_j) component is given by:

$$C_{ij} \equiv \text{Cov} (\Delta m_s(\hat{t}_i), \Delta m_s(\hat{t}_j)) = \sigma_{\mathcal{M}}(\hat{t}_i) \sigma_{\mathcal{M}}(\hat{t}_j) \exp \left[-\frac{(\hat{t}_i - \hat{t}_j)^2}{2l^2} \right] +$$

From Eq. (13), we can calculate an analytic expression for the maximum likelihood estimator (MLE) of the apparent magnitude at B -band maximum light, $\hat{m}_{0,s}$, given by:

$$\hat{m}_{0,s} = \left[\sum_{i,j}^{N_{LC,s}} (C^{-1})_{ij} \right]^{-1} \times \sum_i^{N_{LC,s}} \left[(m_s(\hat{t}_i) - \mathcal{M}(\hat{t}_i)) \sum_j^{N_{LC,s}} (C^{-1})_{ij} \right], \quad (16)$$

with the MLE of the uncertainty of $\hat{m}_{0,s}$ given as

$$\sigma_{0,s} = \left[\sum_{i,j}^{N_{LC,s}} (C^{-1})_{ij} \right]^{-1/2}. \quad (17)$$

Distance modulus

$$\Delta m_s(\hat{t}) \equiv m_s(\hat{t}) - \mathcal{M}(\hat{t}) - m_{0,s} \quad (11)$$

where $m_s(\hat{t})$ and $\mathcal{M}(\hat{t})$ are the apparent magnitude and the magnitude of the normalized template at phase \hat{t} , respectively. We can express this difference for all the $N_{LC,s}$ phases in a given LC as the vector,

$$\Delta \mathbf{m}_s \equiv \begin{pmatrix} \Delta m_s(\hat{t}_1) \\ \Delta m_s(\hat{t}_2) \\ \vdots \\ \Delta m_s(\hat{t}_{N_{LC,s}}) \end{pmatrix}. \quad (12)$$

Then, to determine $m_{0,s}$ we minimize the negative of the log likelihood function $L(m_{0,s})$ defined as

$$-2 \ln L(m_{0,s}) = \Delta \mathbf{m}_s^\top \cdot C^{-1} \cdot \Delta \mathbf{m}_s \quad (13)$$

$$\mu_s = \hat{m}_{0,s} - \langle M_0 \rangle \quad (19)$$

with uncertainty given as

$$\sigma_{\mu,s} = \sqrt{\sigma_{0,s}^2 + \sigma_{int}^2} \quad (20)$$

where C is the $N_{LC,s}$ -dimensional covariance matrix where the (\hat{t}_i, \hat{t}_j) component is given by:

$$C_{ij} \equiv \text{Cov} (\Delta m_s(\hat{t}_i), \Delta m_s(\hat{t}_j)) = \sigma_{\mathcal{M}}(\hat{t}_i) \sigma_{\mathcal{M}}(\hat{t}_j) \exp \left[-\frac{(\hat{t}_i - \hat{t}_j)^2}{2l^2} \right] +$$

From Eq. (13), we can calculate an analytic expression for the maximum likelihood estimator (MLE) of the apparent magnitude at B -band maximum light, $\hat{m}_{0,s}$, given by:

$$\hat{m}_{0,s} = \left[\sum_{i,j}^{N_{LC,s}} (C^{-1})_{ij} \right]^{-1} \times \sum_i^{N_{LC,s}} \left[(m_s(\hat{t}_i) - \mathcal{M}(\hat{t}_i)) \sum_j^{N_{LC,s}} (C^{-1})_{ij} \right], \quad (16)$$

with the MLE of the uncertainty of $\hat{m}_{0,s}$ given as

$$\sigma_{0,s} = \left[\sum_{i,j}^{N_{LC,s}} (C^{-1})_{ij} \right]^{-1/2}. \quad (17)$$

Distance modulus

$$\Delta m_s(\hat{t}) \equiv m_s(\hat{t}) - \mathcal{M}(\hat{t}) - m_{0,s} \quad (11)$$

where $m_s(\hat{t})$ and $\mathcal{M}(\hat{t})$ are the apparent magnitude and the magnitude of the normalized template at phase \hat{t} , respectively. We can express this difference for all the $N_{LC,s}$ phases in a given LC as the vector,

$$\Delta \mathbf{m}_s \equiv \begin{pmatrix} \Delta m_s(\hat{t}_1) \\ \Delta m_s(\hat{t}_2) \\ \vdots \\ \Delta m_s(\hat{t}_{N_{LC,s}}) \end{pmatrix}. \quad (12)$$

Then, to determine $m_{0,s}$ we minimize the negative of the log likelihood function $L(m_{0,s})$ defined as

$$-2 \ln L(m_{0,s}) = \Delta \mathbf{m}_s^\top \cdot C^{-1} \cdot \Delta \mathbf{m}_s \quad (13)$$

$$\mu_s = \hat{m}_{0,s} - \langle M_0 \rangle \quad (19)$$

with uncertainty given as

$$\sigma_{\mu,s} = \sqrt{\sigma_{0,s}^2 + \sigma_{int}^2} \quad (20)$$

where C is the $N_{LC,s}$ -dimensional covariance matrix where the (\hat{t}_i, \hat{t}_j) component is given by:

$$C_{ij} \equiv \text{Cov} (\Delta m_s(\hat{t}_i), \Delta m_s(\hat{t}_j)) = \sigma_{\mathcal{M}}(\hat{t}_i) \sigma_{\mathcal{M}}(\hat{t}_j) \exp \left[-\frac{(\hat{t}_i - \hat{t}_j)^2}{2l^2} \right] +$$

From Eq. (13), we can calculate an analytic expression for the maximum likelihood estimator (MLE) of the apparent magnitude at B -band maximum light, $\hat{m}_{0,s}$, given by:

$$\hat{m}_{0,s} = \left[\sum_{i,j}^{N_{LC,s}} (C^{-1})_{ij} \right]^{-1} \times \sum_i^{N_{LC,s}} \left[(m_s(\hat{t}_i) - \mathcal{M}(\hat{t}_i)) \sum_j^{N_{LC,s}} (C^{-1})_{ij} \right], \quad (16)$$

with the MLE of the uncertainty of $\hat{m}_{0,s}$ given as

$$\sigma_{0,s} = \left[\sum_{i,j}^{N_{LC,s}} (C^{-1})_{ij} \right]^{-1/2}. \quad (17)$$

Intrinsic dispersion


Scatter in the Hubble residuals after accounting for peculiar-velocity and photometric uncertainties.

Intrinsic dispersion σ_{int} :

$$-2 \ln \mathcal{L}(\sigma_{\text{int}}^2) = \sum_s^{N_{\text{SN}}} \left[\ln \left(\sigma_{0,s}^2 + \sigma_{\text{int}}^2 + \sigma_{\mu_{\text{pec}},s}^2 \right) + \frac{\delta \mu_s^2}{\sigma_{0,s}^2 + \sigma_{\text{int}}^2 + \sigma_{\mu_{\text{pec}},s}^2} \right]$$

Blondin, Mandel, Kirshner, 2011



Hubble diagrams from Template method





Gaussian-Process method

Arturo Avelino, "Near-infrared SN Ia as standard candles"


Gaussian-Process Method



Gaussian-Process Method



Hubble diagrams from Gaussian-Process method



Combining multiple NIR bands

Arturo Avelino, "Near-infrared SN Ia as standard candles"

Distance modulus

4.3. Distance modulus from the combined NIR bands


From the distance moduli $(\mu_s^Y, \mu_s^J, \mu_s^H, \mu_s^K)$ for a given supernova s determined from each NIR band following either of the two methods described above, we determine the “total” distance modulus $\hat{\mu}_s$ in each method. First we define the vector of residuals

$$\delta\boldsymbol{\mu}_s \equiv \begin{pmatrix} \mu_s^Y - \hat{\mu}_s \\ \mu_s^J - \hat{\mu}_s \\ \mu_s^H - \hat{\mu}_s \\ \mu_s^K - \hat{\mu}_s \end{pmatrix}. \quad (25)$$


where μ_s^Z is given by either Eq. (19) or (23). Then, to determine $\hat{\mu}_s$ we minimize the negative of the likelihood function $L(\hat{\mu}_s)$ defined as

$$-2 \ln L(\hat{\mu}_s) = \delta\boldsymbol{\mu}_s^\top \cdot C_\mu^{-1} \cdot \delta\boldsymbol{\mu}_s \quad (26)$$

Hubble diagrams from Template method



Hubble diagrams from Gaussian-Process method



How good or bad are these results?




Arturo Avelino, "Near-infrared SN Ia as standard candles"

Optical Hubble diagram


Arturo Avelino, "Near-infrared SN Ia as standard candles"

Fitting the optical light curves only


SALT2



SNooPy



Optical-only Hubble diagrams



Intrinsic dispersion and wRMS summary

Band	Method	σ_{int}	wRMS (mag)
Y	Template	0.095 ± 0.018	0.129
Y	GP	0.091 ± 0.020	0.125
J	Template	0.118 ± 0.015	0.156
J	GP	0.099 ± 0.017	0.137
H	Template	0.061 ± 0.015	0.113
H	GP	0.057 ± 0.022	0.117
K_s	Template	0.138 ± 0.028	0.180
K_s	GP	0.096 ± 0.056	0.170
any $YJHK_s$	Template	0.089 ± 0.012	0.123
any $YJHK_s$	GP	0.081 ± 0.015	0.118
YJH	Template	0.055 ± 0.015	0.097
YJH	GP	0.061 ± 0.025	0.105
JHK_s	Template	0.089 ± 0.026	0.134
JHK_s	GP	0.098 ± 0.047	0.149
Optical	SALT2	0.154 ± 0.023	0.216
Optical	SNooPy	0.140 ± 0.020	0.146

RAISIN = SN IA in the IR


Tracing cosmic expansion with SN Ia in the Near Infrared

RAISIN-1

- 23 SN Ia, redshift ~ 0.3

RAISIN-2

- 24 SN Ia, redshift ~ 0.5



Take away

- NIR SN Ia are very good standard candles compared with optical observations.
- Very promising for cosmology when combining optical+NIR observations:
RAISIN program, WFIRST.

