

Computer Problem Solution

- a) Using the training data in **TrainingSamplesDCT8.mat**, what are reasonable estimates for the prior probabilities?

Solution:

Two priors probabilities, $P_Y(cheetah)$ and $P_Y(grass)$, could be estimated based on the number of vectors in the training set. The estimation of $P_Y(cheetah)$ and $P_Y(grass)$ are:

$$P_Y(cheetah) = N_{FG}/(N_{FG} + N_{BG}) \quad (1)$$

$$P_Y(grass) = N_{BG}/(N_{FG} + N_{BG}) \quad (2)$$

where

N_{BG} is the number of vectors in matrix **TrainsampleDCT_BG**

N_{FG} is the number of vectors in matrix **TrainsampleDCT_FG**

- b) Using the training data in **TrainingSamplesDCT8.mat**, compute and plot the index histograms $P_{X|Y}(x|cheetah)$ and $P_{X|Y}(x|grass)$.

Solution:

The index histograms is the following picture:




Figure 1: index histograms

The value of $P_{X|Y}(x|cheetah)$ and $P_{X|Y}(x|grass)$ is showed as following histograms:




Figure 2: index histograms

1.

$$\forall n(P(n) \rightarrow Q(n)), \quad (3)$$

where

$P(n)$ is “ n is an odd integer” and

$Q(n)$ is “ n^2 is odd.”

2. Assume $P(n)$ is true.

3. By definition, an odd integer is $n = 2k + 1$, where k is some integer.

4.

$$\begin{aligned} n^2 &= (2k + 1)^2 \\ &= 4k^2 + 4k + 1 \\ &= 2(2k^2 + 2k) + 1 \end{aligned}$$

5. $\therefore n^2$ is an odd integer. \square

c) Let $A = \{1, 2, 3\}$ and $B = \{1, 2, 3, \{1, 2, 3\}\}$:

Then, $A \in B$ and $A \subseteq B$.

d) Let $A = \{1, 3, 5\}$, $B = \{1, 2, 3, \}$, and universe $U = \{1, 2, 3, 4, 5\}$:

$$A \cup B = \{1, 2, 3, 5\},$$

$$A \cap B = \{1, 3\},$$

$$A - B = \{5\},$$

$$\bar{A} = \{2, 4\},$$

$$A - A = \emptyset.$$