


Positivity in real Schubert calculus

Slides available at snkarp.github.io



F. Sottile, "Frontiers of reality in Schubert calculus"



M. Griffon, CC BY 3.0 Deed

Steven N. Karp (University of Notre Dame)

arXiv:2309.04645 (joint with Kevin Purbhoo)

arXiv:2405.20229 (joint with Evgeny Mukhin and Vitaly Tarasov)

October 17, 2024


University of California, Los Angeles

Schubert calculus (1886)

- Divisor Schubert problem: given subspaces $W_1, \dots, W_{d(m-d)} \subseteq \mathbb{C}^m$ of dimension $m - d$, find all

d -subspaces $V \subseteq \mathbb{C}^m$ such that $V \cap W_i \neq \{0\}$ for all i .

- e.g. $d = 2$, $m = 4$ (projectivized). Given 4 lines $W_i \subseteq \mathbb{CP}^3$, find all lines $V \subseteq \mathbb{CP}^3$ intersecting all 4. Generically, there are 2 solutions.



We can see the 2 solutions explicitly when two pairs of the lines intersect.

- If the W_i 's are generic, the number of solutions V is f^\square , the number of *standard Young tableaux* of rectangular shape $d \times (m - d)$.
- Fulton (1984): "The question of how many solutions of real equations can be real is still very much open, particularly for enumerative problems."

The Grassmannian $\text{Gr}_{d,m}(\mathbb{C})$

- The *Grassmannian* $\text{Gr}_{d,m}(\mathbb{C})$ is the set of d -dimensional subspaces of \mathbb{C}^m .

$$V := \begin{matrix} (1, 0, -4, -3) \\ \nearrow \\ \vec{0} \\ \searrow \\ (0, 1, 3, 2) \end{matrix} = \begin{bmatrix} 1 & 0 & -4 & -3 \\ 0 & 1 & 3 & 2 \end{bmatrix} \in \text{Gr}_{2,4}(\mathbb{C}) = \begin{bmatrix} 1 & 1 & -1 & -1 \\ 0 & 2 & 6 & 4 \end{bmatrix}$$

$$\Delta_{1,2} = 1, \quad \Delta_{1,3} = 3, \quad \Delta_{1,4} = 2, \quad \Delta_{2,3} = 4, \quad \Delta_{2,4} = 3, \quad \Delta_{3,4} = 1$$

$$\text{Plücker relation: } \Delta_{1,3}\Delta_{2,4} = \Delta_{1,2}\Delta_{3,4} + \Delta_{1,4}\Delta_{2,3}$$

- Given $V \in \text{Gr}_{d,m}(\mathbb{C})$ as a $d \times m$ matrix, for d -subsets I of $\{1, \dots, m\}$ let $\Delta_I(V)$ be the $d \times d$ minor of V in columns I . The *Plücker coordinates* $\Delta_I(V)$ are well-defined up to a common scalar.
- $\text{Gr}_{d,m}(\mathbb{C})$ is a projective variety of dimension $d(m-d)$.


Shapiro–Shapiro conjecture

- Do there exist Schubert problems with all real solutions?

Shapiro–Shapiro conjecture (1993)

Let $W_1, \dots, W_{d(m-d)} \in \text{Gr}_{m-d,m}(\mathbb{R})$ osculate the moment curve $\gamma(t) := (\frac{t^{m-1}}{(m-1)!}, \frac{t^{m-2}}{(m-2)!}, \dots, t, 1)$ at real points. Then there exist f **real** $V \in \text{Gr}_{d,m}(\mathbb{R})$ such that $V \cap W_i \neq \{0\}$ for all i .

- e.g.



F. Sottile, "Frontiers of reality in Schubert calculus"

- This Schubert problem arises in the study of linear series in algebraic geometry, differential equations, and pole placement problems in control theory.
- Bürgisser, Lerario (2020): a uniformly random Schubert problem over \mathbb{R} has $\approx \sqrt{f}$ real solutions.

Shapiro–Shapiro conjecture and secant conjecture

- Sottile (1999) tested the conjecture and proved it asymptotically.
- Eremenko, Gabrielov (2002): cases $d \leq 2$, $m - d \leq 2$.
- Mukhin, Tarasov, Varchenko (2009): full conjecture via the *Bethe ansatz*.
- Levinson, Purbhoo (2021): topological proof of the full conjecture.

Secant conjecture, divisor form (Sottile (2003))

Let $W_1, \dots, W_{d(m-d)} \in \text{Gr}_{m-d,m}(\mathbb{R})$ *be secant to the moment curve $\gamma(t)$ along non-overlapping real intervals*. Then there exist $f \square$

real $V \in \text{Gr}_{d,m}(\mathbb{R})$ such that $V \cap W_i \neq \{0\}$ for all i .

- The Shapiro–Shapiro conjecture is a limiting case of this conjecture.
- Eremenko, Gabrielov, Shapiro, Vainshtein (2006): case $m - d \leq 2$.

Theorem (Karp, Purbhoo (2023))

The divisor form of the secant conjecture is true.

Total positivity


- *Totally positive matrices* (matrices whose minors are all positive) have been studied since the 1930's. Gantmakher and Krein (1937) showed that square totally positive matrices have positive eigenvalues.

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 4 \\ 1 & 3 & 9 \end{bmatrix} \quad \begin{aligned} \lambda_1 &= 10.6031 \dots \\ \lambda_2 &= 1.2454 \dots \\ \lambda_3 &= 0.1514 \dots \end{aligned}$$

- Lusztig (1994) introduced total positivity for algebraic groups G and flag varieties G/P . An element $V \in \text{Gr}_{d,m}(\mathbb{C})$ is *totally nonnegative* if its Plücker coordinates are all nonnegative.

$$V := \begin{bmatrix} 1 & 0 & -4 & -3 \\ 0 & 1 & 3 & 2 \end{bmatrix} \in \text{Gr}_{2,4}^{\geq 0}$$

$$\Delta_{1,2} = 1, \quad \Delta_{1,3} = 3, \quad \Delta_{1,4} = 2, \quad \Delta_{2,3} = 4, \quad \Delta_{2,4} = 3, \quad \Delta_{3,4} = 1$$



- Postnikov (2006) parametrized $\text{Gr}_{d,m}^{\geq 0}$ using *plabic graphs*.
- $\text{Gr}_{d,m}^{\geq 0}$ is related to cluster algebras, electrical networks, the KP hierarchy, scattering amplitudes, curve singularities, the Ising model, knot theory, ...

Positive Shapiro–Shapiro conjecture

Positivity conjecture (Mukhin, Tarasov (2017); Karp (2021))

Let $W_1, \dots, W_{d(m-d)} \in \operatorname{Gr}_{m-d,m}(\mathbb{R})$ osculate the moment curve $\gamma(t)$ at real points $t_1, \dots, t_{d(m-d)} \geq 0$. Then there exist $f \square$

totally nonnegative $V \in \operatorname{Gr}_{d,m}^{\geq 0}$ such that $V \cap W_i \neq \{0\}$ for all i .

- e.g. $d = 2, m = 4$. If $t_3, t_4 \rightarrow \infty$, then the 2 solutions $V \in \operatorname{Gr}_{2,4}(\mathbb{C})$ are

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & t_1 t_2 & t_1 + t_2 & 2 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} \frac{t_1+t_2}{2} & 1 & 0 & 0 \\ -t_1 t_2 & 0 & 2 & 0 \end{bmatrix}.$$

- Karp (2023): the positivity conjecture is equivalent to a conjecture of Eremenko (2015), which implies the divisor form of the secant conjecture.

Theorem (Karp, Purbhoo (2023))

The positivity conjecture is true.

- To prove it, we explicitly solve for the $\Delta_I(V)$'s over $\mathbb{C}[\mathfrak{S}_{d(m-d)}]$.

Universal Plücker coordinates

- Shapiro–Shapiro problem: given $W_1, \dots, W_{d(m-d)} \in \operatorname{Gr}_{m-d,m}(\mathbb{C})$ which osculate the moment curve $\gamma(t)$ at $t_1, \dots, t_{d(m-d)} \in \mathbb{C}$, find all

$V \in \operatorname{Gr}_{d,m}(\mathbb{C})$ such that $V \cap W_i \neq \{0\}$ for all i .

Theorem (Karp, Purbhoo (2023))

There exist linear operators $\beta_I = \beta_I(t_1, \dots, t_{d(m-d)})$ indexed by d -subsets $I \subseteq \{1, \dots, m\}$ with the following properties.

- (i) The β_I 's commute and satisfy the Plücker relations.*
- (ii) There is a bijection between the common eigenspaces of the β_I 's and the solutions V above, sending the eigenvalue of β_I to $\Delta_I(V)$.*
- (iii) If $t_1, \dots, t_{d(m-d)} \geq 0$, then the β_I 's are positive semidefinite.*

$$\beta_I := \sum_{\substack{X \subseteq \{1, \dots, n\}, \\ |X| = |\lambda(I)|}} \left(\prod_{i \notin X} t_i \right) \sum_{\pi \in \mathfrak{S}_X} \chi^{\lambda(I)}(\pi) \pi \in \mathbb{C}[\mathfrak{S}_n] \quad (n = d(m-d))$$

Example: $d = 2$, $m = 4$, and $t_3, t_4 \rightarrow \infty$

$$\beta_I := \sum_{\substack{X \subseteq \{1, \dots, n\}, \\ |X| = |\lambda(I)|}} \left(\prod_{i \notin X} t_i \right) \sum_{\pi \in \mathfrak{S}_X} \chi^{\lambda(I)}(\pi) \pi \in \mathbb{C}[\mathfrak{S}_n] \quad (n = 2)$$

- Write $\mathfrak{S}_2 = \{e, \sigma\}$, where e is the identity and $\sigma = (1 \ 2)$. We have

$$\beta_{1,2} \stackrel{\emptyset}{=} t_1 t_2 e, \quad \beta_{1,3} \stackrel{\square}{=} (t_1 + t_2) e, \quad \beta_{1,4} \stackrel{\square\square}{=} e + \sigma, \quad \beta_{2,3} \stackrel{\square}{=} e - \sigma,$$

and $\beta_I = 0$ if $|\lambda(I)| > 2$. The β_I 's commute and satisfy the Plücker relation

$$\beta_{1,3} \beta_{2,4} = \beta_{1,2} \beta_{3,4} + \beta_{1,4} \beta_{2,3} \rightsquigarrow 0 = 0 + (e + \sigma)(e - \sigma).$$

- On the eigenspace $\langle e - \sigma \rangle$, the eigenvalues are

$$\beta_{1,2} \rightsquigarrow t_1 t_2, \quad \beta_{1,3} \rightsquigarrow t_1 + t_2, \quad \beta_{1,4} \rightsquigarrow 0, \quad \beta_{2,3} \rightsquigarrow 2,$$

which are the Plücker coordinates of

$$V = \begin{bmatrix} \frac{t_1+t_2}{2} & 1 & 0 & 0 \\ -t_1 t_2 & 0 & 2 & 0 \end{bmatrix} \in \text{Gr}_{2,4}(\mathbb{C}).$$

Proof 1: KP hierarchy

- The key to the proof is showing that the β_I 's satisfy the Plücker relations.
- The *KP equation*

$$\frac{\partial}{\partial x} \left(-4 \frac{\partial u}{\partial t} + 6u \frac{\partial u}{\partial x} + \frac{\partial^3 u}{\partial x^3} \right) + 3 \frac{\partial^2 u}{\partial y^2} = 0$$

models shallow water waves. It is the first equation in the *KP hierarchy*.

- The solutions to the KP hierarchy are symmetric functions $\tau(\mathbf{x})$ in the variables $\mathbf{x} = (x_1, x_2, \dots)$ satisfying *Hirota's bilinear identity*:

$$[t^{-1}] \exp \left(\sum_{k \geq 1} \frac{t^k}{k} (p_k(\mathbf{x}) - p_k(\mathbf{y})) \right) \exp \left(\sum_{k \geq 1} -t^{-k} \left(\frac{\partial}{\partial p_k(\mathbf{x})} - \frac{\partial}{\partial p_k(\mathbf{y})} \right) \right) \tau(\mathbf{x}) \tau(\mathbf{y}) = 0.$$

- Sato (1981): $\tau(\mathbf{x})$ satisfies Hirota's bilinear identity if and only if its coefficients in the Schur basis $s_\lambda(\mathbf{x})$ satisfy the Plücker relations.
- Karp, Purbhoo (2023): $\sum_I \beta_I s_{\lambda(I)}(\mathbf{x})$ satisfies Hirota's bilinear identity.



M. Griffon, CC BY 3.0 Deed

Proof 2: higher Gaudin Hamiltonians

- The *higher Gaudin Hamiltonian* associated to the partition λ is

$$T_\lambda = (t_1 + d_1) \cdots (t_n + d_n) s_\lambda(h) \in \text{End}((\mathbb{C}^k)^{\otimes n}),$$

where:

- h is a $k \times k$ matrix;
- $s_\lambda(h)$ is the Schur polynomial evaluated at the eigenvalues of h ; and
- d_i is the derivative with respect to h^T acting in the i th tensor factor.

Theorem (Alexandrov, Leurent, Tsuboi, Zabrodin (2014))

The T_λ 's pairwise commute and satisfy the Plücker relations.

Theorem (Karp, Mukhin, Tarasov (2024))

(i) If $t_1, \dots, t_n \geq 0$ and h is positive semidefinite, then so is T_λ .

(ii) We have $\beta_I = T_{\lambda(I)}|_{h=0}$.

- Part (i) gives a positivity theorem for spaces $\langle e^{h_1 \times} f_1, \dots, e^{h_d \times} f_d \rangle$, where h_1, \dots, h_d are the eigenvalues of h and the f_i 's are polynomials.

Future directions

- Further explore the connection to the KP hierarchy.
- Find necessary and sufficient inequalities on the Plücker coordinates of V for all complex zeros of $\text{Wr}(V)$ to be nonpositive, generalizing the Aissen–Schoenberg–Whitney theorem in the case $\dim(V) = 1$. (The positivity conjecture implies that the inequalities $\Delta_I(V) \geq 0$ are necessary.)
- What happens to the higher Gaudin Hamiltonian T_λ if s_λ is replaced by a different symmetric function?
- Address generalizations and variations of the Shapiro–Shapiro conjecture: the discriminant conjecture, the general form of the secant conjecture, the monotone conjecture, the total reality conjecture for convex curves, ...

Thank you!