

Class09

Natalie Ogg A91030809

The PDB Database

```
stats <- read.csv("Data Export Summary.csv", row.names=1)
stats
```

	X.ray	EM	NMR	Multiple.methods	Neutron	Other
Protein (only)	158,844	11,759	12,296	197	73	32
Protein/Oligosaccharide	9,260	2,054	34	8	1	0
Protein/NA	8,307	3,667	284	7	0	0
Nucleic acid (only)	2,730	113	1,467	13	3	1
Other	164	9	32	0	0	0
Oligosaccharide (only)	11	0	6	1	0	4
Total						
Protein (only)	183,201					
Protein/Oligosaccharide	11,357					
Protein/NA	12,265					
Nucleic acid (only)	4,327					
Other	205					
Oligosaccharide (only)	22					

Q1: What percentage of structures in the PDB are solved by X-Ray and Electron Microscopy.

```
x <- stats$X.ray
x
```

```
[1] "158,844" "9,260" "8,307" "2,730" "164" "11"
```

```
as.numeric(gsub(",", "", x))
```

```
[1] 158844  9260  8307  2730  164  11
```

```
rm.comma <- function(x) {
  as.numeric(gsub(",", "", x))
}
```

```
rm.comma(stats$EM)
```

```
[1] 11759 2054 3667 113 9 0
```

```
pdb.data <- apply(stats, 2, rm.comma)
rownames(pdb.data) <- rownames(stats)
pdb.data
```

	X.ray	EM	NMR	Multiple.methods	Neutron	Other
Protein (only)	158844	11759	12296	197	73	32
Protein/Oligosaccharide	9260	2054	34	8	1	0
Protein/NA	8307	3667	284	7	0	0
Nucleic acid (only)	2730	113	1467	13	3	1
Other	164	9	32	0	0	0
Oligosaccharide (only)	11	0	6	1	0	4
Total						
Protein (only)	183201					
Protein/Oligosaccharide	11357					
Protein/NA	12265					
Nucleic acid (only)	4327					
Other	205					
Oligosaccharide (only)	22					

```
round(totals)
```

```
totals <- apply(pdb.data, 2, sum)
round(totals/totals["Total"]*100,2)
```

X.ray	EM	NMR	Multiple.methods
84.83	8.33	6.68	0.11
Neutron	Other	Total	
0.04	0.02	100.00	

Q2: What proportion of structures in the PDB are protein?

```
round(pdb.data[,"Total"] / sum(pdb.data[,"Total"])) *100, 2)
```

Protein (only)	Protein/Oligosaccharide	Protein/NA
86.67	5.37	5.80
Nucleic acid (only)	Other	Oligosaccharide (only)
2.05	0.10	0.01

Q3: Type HIV in the PDB website search box on the home page and determine how many HIV-1 protease structures are in the current PDB?

(skip)

Q4: Water molecules normally have 3 atoms. Why do we see just one atom per water molecule in this structure?


In MolStar, we can't see hydrogens because they are smaller than the resolution.

Q5: There is a critical "conserved" water molecule in the binding site. Can you identify this water molecule? What residue number does this water molecule have

Yes, it is water number 308.

Q6:

Here is a lovely figure of HIP-Pr with the catalytic ASP residues, the MK1 compound and the all-important water 308. The Asp residues are in ball and stick representation.



Reading PDB file data in R

We will use the Bio3D package for structural bioinformatics.

```
library(bio3d)

pdb <- read.pdb("1hsg")
```

Note: Accessing on-line PDB file

```
pdb
```

```
Call: read.pdb(file = "1hsg")
```

```
Total Models#: 1
```

```
Total Atoms#: 1686, XYZs#: 5058 Chains#: 2 (values: A B)
```

```
Protein Atoms#: 1514 (residues/Calpha atoms#: 198)
```

```
Nucleic acid Atoms#: 0 (residues/phosphate atoms#: 0)
```

```
Non-protein/nucleic Atoms#: 172 (residues: 128)
```

```
Non-protein/nucleic resid values: [ HOH (127), MK1 (1) ]
```

```
Protein sequence:
```

```
PQITLWQRPLVTIKIGGQLKEALLDTGADDTVLEEMSLPGRWKPKMIGGIGGFIKVRQYD
QILIEICGHKAIGTVLVGPTPVNIIGRNLLTQIGCTLNFPQITLWQRPLVTIKIGGQLKE
ALLDTGADDTVLEEMSLPGRWKPKMIGGIGGFIKVRQYDQILIEICGHKAIGTVLVGPTP
VNIIGRNLLTQIGCTLNF
```

```
+ attr: atom, xyz, seqres, helix, sheet,
      calpha, remark, call
```

```
head(pdb$atom)
```

	type	eleno	elety	alt	resid	chain	resno	insert	x	y	z	o	b
1	ATOM	1	N	<NA>	PRO	A	1	<NA>	29.361	39.686	5.862	1	38.10
2	ATOM	2	CA	<NA>	PRO	A	1	<NA>	30.307	38.663	5.319	1	40.62
3	ATOM	3	C	<NA>	PRO	A	1	<NA>	29.760	38.071	4.022	1	42.64
4	ATOM	4	O	<NA>	PRO	A	1	<NA>	28.600	38.302	3.676	1	43.40
5	ATOM	5	CB	<NA>	PRO	A	1	<NA>	30.508	37.541	6.342	1	37.87
6	ATOM	6	CG	<NA>	PRO	A	1	<NA>	29.296	37.591	7.162	1	38.40
	segid	elesy	charge										
1	<NA>	N	<NA>										
2	<NA>	C	<NA>										
3	<NA>	C	<NA>										
4	<NA>	O	<NA>										
5	<NA>	C	<NA>										
6	<NA>	C	<NA>										

Q7: How many amino acid residues are there in this pdb object?

198

Q8: Name one of the two non-protein residues?

MK1, the Merck drug.

Q9: How many protein chains are in this structure?

2: A and B.

We are finishing part 1 today with a calculation...

Predicting functional motions of a single structure

```
adk <- read.pdb("6s36")
```

Note: Accessing on-line PDB file


PDB has ALT records, taking A only, rm.alt=TRUE

```
m <- nma(adk)
```

Building Hessian... Done in 0.014 seconds.

Diagonalizing Hessian... Done in 0.283 seconds.

```
plot(m)
```



```
mktrj(m, file="adk_m7.pdb")
```