

Imperial College of Science, Technology and Medicine  
Department of Mathematics

# **Sparse Spectral Methods on Disk-Slices, Trapeziums and Spherical Caps**

Ben Snowball

Submitted in part fulfilment of the requirements for the degree of  
Doctor of Philosophy in Mathematics of Imperial College, May 2021



## Abstract

This thesis develops sparse spectral methods for solving partial differential equations (PDEs) on various multidimensional domains, with a specific focus on the disk-slice and trapezium in 2D, and the spherical cap as a surface in 3D. For the latter scenario, the PDEs are surface PDEs involving Laplace-Beltrami operators, spherical gradients and other spherical operators.

Throughout this thesis we wish to utilise our understanding of the orthogonal polynomials and sparse spectral methods to build up to new methods for subdomains of the sphere surface. To this end, we begin with an introduction to sparse spectral methods via the long established Spherical Harmonics on the whole sphere surface. We present a framework for how the spherical harmonics can be thought of as a set of multidimensional orthogonal polynomials in  $x$ ,  $y$  and  $z$ , and can be used to expand functions defined on the sphere. We explain how differential operators can be applied as banded-block-banded matrix operators to coefficient vectors for the function's expansion. This way of thinking about the spherical harmonics differs from the traditional sense (where they are considered as orthogonal functions in spherical coordinates) but we do so as to be able to generalise the techniques to spectral methods on other domains using multidimensional OPs. Further, we demonstrate how the Vector Spherical Harmonics can be used as an orthogonal basis for vector valued functions lying in the tangent space of the sphere, and thus how one can additionally derive gradient and divergence operators.

We move on working in 2D, where in recent years sparse spectral methods for solving PDEs have been derived using hierarchies of classical orthogonal polynomials on intervals, disks, and triangles. Presenting a new framework for choosing a suitable orthogonal polynomial basis for more general 2D domains defined via

an algebraic curve as a boundary, this work builds on the observation that sparsity is guaranteed due to this definition of the boundary, and that the entries of partial differential operators can be determined using formulae in terms of (non-classical) univariate orthogonal polynomials. Triangles and the full disk are then special cases of our framework, which we formalise for the disk-slice and trapezium cases.

Finally, we return to the surface of the sphere, namely the subdomain we call the spherical cap. Piecing together techniques and OPs from both the Spherical Harmonics and disk-slice work, we once again present a new orthogonal polynomial basis and sparse spectral method for the spherical cap, complete with the same observation about the guaranteed sparsity of differential and other operators. The motivation is for one to use spherical caps and spherical bands (which are a simple extension) as elements in a spectral element method for the sphere, with many applications in meteorology and astrophysics – in particular, as a potential replacement of the Spherical Harmonics approach that is currently in use at the European Centre for Medium-range Weather Forecasts (ECMWF) which is predicted in the future to be too costly due to a parallel scalability bottleneck arising from the global spectral transform.

## Foreword

Text of the foreword.



## Declaration

Declaration here.





## Acknowledgements

Firstly, I would like to express my thanks to ESPRC for their financial support, and to the Mathematics of Planet Earth Centre for Doctoral Training (MPE CDT) for providing me with the incredible opportunities that the course has brought.

I would like to thank the MPE CDT staff for any knowledge and wisdom they have bestowed, and in particular say a huge thank you to Colin Cotter for his guidance, understanding and general enthusiasm (it meant a lot). And to all my fellow 'Charlies' – I truly mean it when I say you are all such great, smart and kind people, and you all deserve the absolute best for the future.

I thank my family and friends for all their support and words of encouragement – it all helped a lot. An extra thanks to all my undergrad mates for many well needed fun times too.

A special thanks goes to my partner Jessica for putting up with me and helping me when things got tough during my 4 years at Imperial College London.

Finally, to Dr. Sheehan Olver – I cannot thank you enough for all your help, wisdom, advice, kindness, understanding and patience you have given me. I may not have expressed my gratitude enough over the past few years, but you have truly been a great supervisor to me.

“If we hit that bullseye, the rest of the dominoes will fall like a house of cards.  
Checkmate.”

*Captain Zapp Brannigan*

# Contents

<b>Abstract</b>	<b>i</b>
<b>Foreword</b>	<b>ii</b>
<b>Declaration</b>	<b>iii</b>
<b>Acknowledgements</b>	<b>vii</b>
<b>1 Introduction</b>	<b>1</b>
<b>2 Spherical harmonics as orthogonal polynomials</b>	<b>8</b>
2.1 Defining the spherical harmonics in three variables . . . . .	10
2.2 Jacobi matrices . . . . .	14
2.3 Three-term recurrence relation for $\mathbb{P}$ . . . . .	24
2.4 Computational aspects . . . . .	28
2.4.1 Obtaining coefficients . . . . .	28
2.4.2 Function evaluation . . . . .	29
2.4.3 Obtaining operator matrices for variable coefficients . . . .	29
2.5 Vector spherical harmonics as orthogonal vectors in three variables	30

2.5.1	Jacobi matrices . . . . .	32
2.5.2	Three-term recurrence relation for $\tilde{T}(x, y, z)$ . . . . .	55
2.5.3	Computational aspects . . . . .	58
2.5.4	Sparse partial differential operators . . . . .	61
2.6	Examples . . . . .	67
2.6.1	Heat equation . . . . .	67
2.6.2	Linearised shallow water equations . . . . .	69
2.7	Conclusion . . . . .	74
<b>3</b>	<b>Disk slices and trapeziums</b>	<b>76</b>
3.1	Orthogonal polynomials on the disk-slice and the trapezium . . .	80
3.1.1	Explicit construction . . . . .	80
3.1.2	Jacobi matrices . . . . .	84
3.1.3	Building the OPs . . . . .	91
3.2	Sparse partial differential operators . . . . .	93
3.3	Computational aspects . . . . .	108
3.3.1	Constructing $R_n^{(a,b,c)}(x)$ . . . . .	108
3.3.2	Quadrature rule on the disk-slice . . . . .	115
3.3.3	Obtaining the coefficients for expansion of a function on the disk-slice . . . . .	118
3.3.4	Calculating non-zero entries of the operator matrices . . .	119
3.4	Examples on the disk-slice with zero Dirichlet conditions . . . . .	120
3.4.1	Poisson . . . . .	121
3.4.2	Inhomogeneous variable-coefficient Helmholtz . . . . .	124

3.4.3	Biharmonic equation . . . . .	126
3.5	Other domains . . . . .	126
3.5.1	End-Disk-Slice . . . . .	126
3.5.2	Trapeziums . . . . .	128
3.6	P-finite element methods using sparse operators . . . . .	130
3.7	Conclusions . . . . .	135
<b>4</b>	<b>Spherical caps</b>	<b>137</b>
4.1	The circle arc . . . . .	142
4.2	Orthogonal polynomials on algebraic curves . . . . .	144
4.3	Orthogonal polynomials on spherical caps . . . . .	144
4.3.1	Explicit construction . . . . .	144
4.3.2	Jacobi matrices . . . . .	147
4.3.3	Building the OPs . . . . .	158
4.4	Sparse partial differential operators . . . . .	161
4.4.1	Further partial differential operators . . . . .	174
4.4.2	Stability of the Laplacian operator . . . . .	177
4.5	Computational aspects . . . . .	179
4.5.1	Constructing $R_n^{(a,b)}(x)$ . . . . .	179
4.5.2	Quadrature rule on the spherical cap . . . . .	179
4.5.3	Obtaining the coefficients for expansion of a function on the spherical cap . . . . .	183
4.5.4	Function evaluation . . . . .	184
4.5.5	Calculating non-zero entries of the operator matrices . . .	186

4.5.6	Obtaining operator matrices for variable coefficients . . . .	186
4.6	Examples on spherical caps with zero Dirichlet conditions . . . .	187
4.6.1	Poisson . . . . .	189
4.6.2	Inhomogeneous variable-coefficient Helmholtz . . . . .	190
4.6.3	Biharmonic equation . . . . .	195
4.6.4	Other boundary conditions . . . . .	195
4.7	Conclusions . . . . .	197
<b>5</b>	<b>Summary and future directions</b>	<b>199</b>
5.1	Summary . . . . .	199
5.2	Future directions . . . . .	200
	<b>Appendix</b>	<b>201</b>
.1	Vector-valued functions in the tangent space . . . . .	201
	<b>Bibliography</b>	<b>202</b>

# Chapter 1

## Introduction

Univariate orthogonal polynomials (hereon also referred to as OPs) have been extensively involved in the development of multiple fields of computational and applied mathematics. For example, univariate OPs have been used to derive spectral methods to numerically solve one-dimensional differential equations (see e.g. [63, 8, 37, 52, 44] ). While there are many famous examples of univariate OPs – such as the Jacobi polynomials, the Legendre polynomials and the Chebyshev polynomials but to name a few of the classical families [42, Section 18.3] – the area of multivariate orthogonal polynomials has a smaller array of research.

add more  
from the refer-  
ences in these  
papers

One could say this is surprising, given that there is a long history of around 150 years associated with multivariate OPs, beginning with Hermite first presenting the multivariate Hermite polynomials in 1865 [1, 28]. Zernike polynomials [70] were first introduced in 1934, as another example, that are a group of bivariate polynomials orthogonal on a unit circle. Koornwinder in 1975 described a method

for constructing two-variable OPs from univariate OPs [30]. However, few books have been published over the years on the topic of multivariate orthogonal polynomials [17].

This branch of mathematics has an encouraging future though, not least as a basis for sparse spectral methods for solving partial differential equations (PDEs) on multidimensional domains. Spectral methods have been established on the triangle [45] using OPs inspired by the Koornwinder approach for defining them, and on the disk [65] using Zernike polynomials. Applications include solving Volterra integral equations [22].

There are various interpretations of what a “spectral method” is. It could be described as one that achieves spectral convergence of its solutions, or one that uses eigenfunctions as basis functions, or one that uses OPs as basis polynomials

cite these as  
interpretations

[1]. It is the latter that we refer to for our purposes as a spectral method. By utilising OPs as basis functions, we can develop sparse spectral methods (SSMs), meaning that the naturally sparse relationships between the basis OPs lead to sparse operator matrices that represent the differential operations in the equation to solve. Sparse spectral methods for one dimensional problems have been shown to lead to “almost banded” matrices [44].

In this thesis we expand upon this knowledge by providing frameworks for similar sparse spectral methods for solving PDEs on other multidimensional domains (notably including the disk-slice and trapezium in 2D, and the spherical cap surface in 3D) that also yield matrices that are what is defined as “banded-block-banded”. We take inspiration from the work established for the triangle [45] and the unit



disk [65], which can be seen as special cases of the disk-slice and trapezium in the framework we present here. The spherical cap work serves to lay a foundation for using spherical caps and spherical bands as elements in a spectral element method for solving PDEs on the whole sphere, as an alternative to the Spherical Harmonic transform approach that the European Centre for Medium-range Weather Forecasts (ECMWF) use in their weather and climate model [13].

Spherical Harmonics are of course a long-established and famous group of functions defined on the surface of a sphere with certain useful properties (for example, they are orthogonal to each other on the unit sphere and form a basis for expanding functions defined on the sphere) and as a result are widely used in many fields including computer graphics (e.g. [39, 57]), astrophysics (e.g. [66]), quantum theory (e.g. [64]), geosciences (e.g. [25]) and meteorology (e.g. [18, 50]). Notably, they are also used as basis functions for the spectral transform method that makes up part of the model in the Integrated Forecasting System (IFS), which is used by ECMWF for their forecasts [67]. While the whole sphere spectral method using the Spherical Harmonics has been successful for numerous years [68], there is a drawback in the parallel scalability bottleneck that arises from the global spectral transform, which is expected to inhibit future performance of the IFS [3, 67, 15].

cite applications of SHs

Many implementations of an algorithm to compute the spectral transform (or spherical harmonic transform) exist (see e.g. [55, 60]). For the IFS, the spherical harmonic transform in fact uses two transforms – a Fourier transform (using the well-established Fast Fourier Transform (FFT) [14]) in the longitudinal direction

and a Legendre transform in the latitudinal direction – and it is the Legendre transform that has been identified as inhibiting future performance due to its computational cost. While a Fast Legendre Transform (FLT) [67] has been incorporated into the model, along with new grid types [36], to help to extend the lifespan of the spectral method for numerical weather prediction (NWP), it may not be sufficient for certain desired cases and resolutions [15].

The motivation for this project was to help address this problem while still utilising a spectral approach. More precisely, we aim to develop a sparse spectral method for solving PDEs on the spherical cap as a surface in 3D, with a simple extension to a spherical band. Together, these frameworks can be pieced together to create a spectral element method for the whole sphere, or further developed to investigate spectral methods on other spherical subdomains. This, however, is future work beyond the scope of this thesis. By spectral element method, we mean a finite element method (FEM) that uses high degree basis polynomials for its elements (this could also be referred to as a  $p$ -FEM with large  $p$ ). In other words, we can use our spectral methods developed for the spherical band and cap elements as part of a finite element framework. By using this approach, one would be able to avoid having to complete the global spectral transform (in particular, the global Legendre transform) and instead be able to simply apply the local element transforms in parallel. Moreover, by still using a spectral approach, one can maintain the high accuracy and excellent error properties that such methods bring.

There are also potential applications in physics too where solving equations on

the sphere and working in spherical geometries is also desirable (e.g. [6, 64, 56]), particularly in astrophysics [66, 49].

cite more, see  
p2 of Vasil  
and the refer-  
ences

Recently, a method for computing tensor fields in spherical coordinates using Jacobi polynomials has been proposed [66]. In this work the authors present the method for both the surface of the unit sphere and the three-dimensional generalisation of the unit ball. Their method involves using a spectral basis to represent functions too, choosing for the angular part of the basis to be the spin-weighted spherical harmonics (of which the spherical harmonics are a special case, with spin 0) in spherical coordinates. By using these as their basis functions, they are able to derive sparse relations for the Laplacian operator, as well as for multiplication of the basis by  $\cos \varphi$  and  $\sin \varphi$  (where  $\varphi$  is the polar angle from the  $z$  axis in cartesian coordinates) which lead to operators for operations with angular dependencies involving these.

On the other hand, we aim to propose a strictly orthogonal *polynomial* basis, and derive sparse operators for multiplication by the cartesian coordinate axes, which can in turn lead to operators for multiplication by trigonometric functions too. By utilising orthogonal polynomials, we can develop a sparse spectral method on subdomains of the sphere surface (e.g. the spherical cap) using our knowledge of other geometries and methods involving OPs developed for them.

The structure of this thesis is as follows.

Chapter 2 of this thesis provides an introduction to sparse spectral methods via the Spherical Harmonics on the whole sphere surface. Here, we think of the Spherical

Harmonics in a non-traditional way and write them as a group of multidimensional orthogonal polynomials in  $(x, y, z)$  as opposed to functions of spherical coordinates. We present an OP framework for how the Spherical Harmonics can be used to expand functions defined on the sphere as multidimensional polynomials in  $x, y, z$ , and how differential operators can be applied as banded-block-banded matrix operators to coefficient vectors for a function's expansion. Further, we demonstrate how the Vector Spherical Harmonics can be used as an orthogonal basis for vector valued functions lying in the tangent space of the sphere, and thus how one can additionally derive gradient and divergence operators.

In Chapter 3 we move on to working in 2D, where in recent years sparse spectral methods for solving PDEs have been derived using hierarchies of classical orthogonal polynomials on intervals, disks, and triangles. Presenting a new framework for choosing a suitable orthogonal polynomial basis for more general 2D domains defined via an algebraic curve as a boundary, this work builds on the observation that sparsity is guaranteed due to this definition of the boundary, and that the entries of partial differential operators can be determined using formulae in terms of (non-classical) univariate orthogonal polynomials, which we define. Triangles and the full disk are then special cases of our framework, which we formalise for the disk-slice and trapezium cases.

With a greater knowledge base in our quiver, we can adapt the techniques learnt from the founding of the disk-slice formulation to surfaces in 3D in Chapter 4. Using the same family of (non-classical) 1D OPs, we present a suitable orthogonal polynomial basis for the spherical cap, a subdomain of the surface of a unit

sphere, complete with sparse differential operators. A relatively simple adaption permits this framework to be extended to a spherical band. From here, a spectral element method could be devised for the whole sphere using the aforementioned as elements.

Finally, we summarise the work we have presented and detail avenues for future directions that one could take it in Chapter 5.

## Chapter 2

# Spherical harmonics as orthogonal polynomials

To introduce ourselves to the world of multidimensional orthogonal polynomials for solving PDEs on the sphere, we can naturally choose to look at the famous spherical harmonics. Our aim here is to express the Spherical Harmonics as polynomials in three variables  $x, y, z$  to evaluate functions and solve PDEs on the whole sphere. More precisely, we desire the solution to partial differential equations on the domain

$$\Omega := \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 = \rho(z)^2\}$$

where

$$\rho(z) := \sqrt{1 - z^2}.$$

While it may seem somewhat odd, it should hopefully become apparent why this is a useful way to define our domain.

In this chapter, we demonstrate how one can write the Spherical Harmonics as polynomials in  $(x, y, z)$ , and how the relationships between them can be used to obtain sparse “Jacobi operators” for multiplication by  $x, y, z$ . We demonstrate how these in turn lead to an algorithm for evaluating the Spherical Harmonics, an efficient algorithm for evaluating functions when expanded in the SH OP basis, and sparse “banded-block-banded” differential operator matrices. The techniques used here are done so with the aim of applying them to other OP families of other domains (and we do so in later chapters). We will finally demonstrate a proof-of-concept via the examples of the heat equation and the linearised shallow water equations on the unit sphere.

The code that allows one to produce the numerical examples in this chapter is publicly available as a Julia package<sup>1</sup> that partners the ApproxFun package [43] – however, this package is purely experimental at this stage.

---

<sup>1</sup><https://github.com/snowball13/SphericalHarmonics.jl>

## 2.1 Defining the spherical harmonics in three variables

Before we proceed further, let's build up to our definition of the spherical harmonics. We first need to introduce a few classical orthogonal polynomials. “Orthogonal polynomials” (OPs) are defined as such by the condition that each polynomial is orthogonal to each other, with respect to a given inner product. In particular, they are orthogonal with respect to all lower degree polynomials.

On the unit interval,  $[-1, 1]$ , we note that there is a hierarchy of OPs in the sense that [42, table 18.3.1, eqn 18.9.15]:

$$\begin{aligned} \frac{d}{dx} P_l^{(a,b)}(x) &= \frac{1}{2} (l + a + b + 1) P_{l-1}^{(a+1,b+1)}(x) \\ \implies \frac{d^m}{dx^m} P_l(x) &= \frac{(l+m)!}{2^m l!} P_{l-m}^{(m,m)}(x) \end{aligned}$$

where  $P_l^{(a,b)}(x)$  is the  $l$  degree *Jacobi polynomial*, and  $P_l(x) := P_l^{(0,0)}(x)$  is simply the *Legendre polynomial* of degree  $l$ . Jacobi polynomials are orthogonal with weight  $w(x) = (1-x)^a(1+x)^b$ ; that is for  $l, l' \in \mathbb{N}_0$ ,

$$\begin{aligned} \int_{-1}^1 P_l^{(a,b)}(x)^2 (1-x)^a(1+x)^b dx &=: \omega_{P,l}^{(a,b)} \\ \int_{-1}^1 P_l^{(a,b)}(x) P_{l'}^{(a,b)}(x) (1-x)^a(1+x)^b dx &= \omega_{P,l}^{(a,b)} \delta_{l,l'}. \end{aligned}$$

Legendre polynomials are special cases of *Legendre functions* when  $l$  is a non-negative integer. Legendre functions are a class of univariate functions that are



solutions to the differential equation

$$(1 - x^2) \frac{d^2}{dx^2} y - 2x \frac{d}{dx} y + l(l + 1)y = 0$$

which is helpfully known as the *Legendre equation* [42, 14.2.1]. Further, the *associated Legendre polynomials* are a set of polynomials orthogonal with respect to unit weight on the unit interval, and can be defined as the  $m^{\text{th}}$  derivative of a Legendre polynomial [4, p5]:

$$P_l^m(x) := (-1)^m (1 - x^2)^{\frac{m}{2}} \frac{d^m}{dx^m} P_l(x) = (-1)^m \frac{(l + m)!}{2^m l!} (1 - x^2)^{\frac{m}{2}} P_{l-m}^{(m,m)}(x),$$

for  $m = 0, 1, 2, \dots, l$ . It is also standard to include a further definition for the case of  $m$  negative – that is we also define:

$$P_l^m(x) := (-1)^m \frac{(l + m)!}{(l - m)!} P_l^{|m|}(x) = \frac{(l + m)!}{2^{|m|} l!} (1 - x^2)^{\frac{|m|}{2}} P_{l-|m|}^{(|m|,|m|)}(x)$$

for  $m = -l, \dots, -1$ .

Once again, associated Legendre polynomials are special cases of *associated Legendre functions* when  $l$  is a non-negative integer and  $m \in \{0, \dots, l\}$ , which are also a class of univariate functions that are solutions to a modified version of the Legendre equation

$$(1 - x^2) \frac{d^2}{dx^2} y - 2x \frac{d}{dx} y + \left[ l(l + 1) - \frac{m^2}{1 - x^2} \right] y = 0$$

which is also helpfully known as the *associated Legendre equation* ([4, p2], [42,

14.2.2]). We can see that Legendre functions (and polynomials) are just special cases of the associated versions when  $m = 0$ .

Although often referred to as the associated Legendre “polynomials” as we do here when  $l$  is a non-negative integer and  $m \in \{0, \dots, l\}$ ,  $P_l^m$  are not strictly polynomials when  $m$  is odd, and as such some authors refer to them as associated Legendre functions regardless instead. Moreover, it should be noted too that the inclusion of the  $(-1)^m$  factor in the definition for  $m > 0$ , known as the *Condon-Shortley phase* in physics, is sometimes omitted.

These relationships for the Jacobi polynomials and associated Legendre polynomials allow us to explicitly see where the normalising constants that we shall be using for our definition of the spherical harmonics come from.

On that note, we can now write down the spherical harmonics. Let  $(x, y, z) \in \Omega$ . It is useful to be able to transform between these cartesian coordinates and the spherical coordinates  $(\varphi, \theta)$ . Throughout this document we will use the convention that the spherical coordinate angles be defined by

$$x = \sin \varphi \cos \theta = \rho(z) \cos \theta$$

$$y = \sin \varphi \sin \theta = \rho(z) \sin \theta$$

$$z = \cos \varphi.$$

where  $\rho(z) := \sqrt{1 - z^2}$ . We will use the standard definition – that is, the spherical

harmonics, orthonormal on the unit sphere, are [42, 14.30.1]:

$$\begin{aligned}
Y_l^m(\varphi, \theta) &:= \left( \frac{(2l+1)(l-m)!}{4\pi(l+m)!} \right)^{\frac{1}{2}} e^{im\theta} P_l^m(\cos \varphi) \\
&= c_l^m (1 - (\cos \varphi)^2)^{\frac{|m|}{2}} e^{im\theta} P_{l-|m|}^{(|m|, |m|)}(\cos \varphi) \\
&= c_l^m P_{l-|m|}^{(|m|, |m|)}(z) \rho(z)^{|m|} e^{im\theta}
\end{aligned} \tag{2.1}$$

for  $0 \leq |m| \leq l$ ,  $l \in \mathbb{N}_0$ , where

$$c_l^m := \left( \frac{(2l+1)(l-m)!}{4\pi(l+m)!} \right)^{\frac{1}{2}} \frac{(l+m)!}{2^{|m|} l!} \begin{cases} (-1)^m & \text{if } m \geq 0 \\ 1 & \text{if } m < 0 \end{cases} \tag{2.2}$$

which can be simplified to

$$c_l^m = \left( \frac{(2l+1)(l+m)!(l-m)!}{\pi} \right)^{\frac{1}{2}} \frac{1}{2^{|m|+1} l!} \begin{cases} (-1)^m & \text{if } m \geq 0 \\ 1 & \text{if } m < 0 \end{cases}.$$

The spherical harmonics as defined are then orthonormal with respect to the inner product

$$\begin{aligned}
&\int_0^{2\pi} \int_0^\pi Y_l^m(\varphi, \theta) Y_{l'}^{m'}(\varphi, \theta)^* \sin \varphi \, d\varphi \, d\theta \\
&= 2\pi \delta_{m,m'} c_l^m c_{l'}^{m'} \int_{-1}^1 P_{l-|m|}^{(|m|, |m|)}(z) P_{l'-|m|}^{(|m|, |m|)}(z) \rho(z)^{2|m|} \, dz \\
&= \delta_{l,l'} \delta_{m,m'}
\end{aligned}$$

where  $\alpha^*$  denotes the complex conjugate of  $\alpha \in \mathbb{C}$ . Note how we can express the

spherical harmonics  $Y_l^m$  in terms of  $x, y, z$  instead of  $\varphi, \theta$  by noting that  $\rho(z)^{|m|}e^{im\theta}$  can be expressed in terms of  $x, y, z$  for any  $m \in \mathbb{Z}$ . Indeed, they are polynomials in  $x, y, z$  which we denote  $Y_l^m(x, y, z)$ . They span all polynomials modulo the vanishing ideal associated by the zero set of  $x^2 + y^2 + z^2 - 1$  in  $\mathbb{R}^3$ .

## 2.2 Jacobi matrices

Jacobi operators are matrices that correspond to multiplication of the orthogonal polynomial basis, in this case the spherical harmonics, by our cartesian coordinates  $x, y, z$ . We can obtain the entries to these operator matrices by determining the coefficients of  $x Y_l^m(x, y, z)$ ,  $y Y_l^m(x, y, z)$ , and  $z Y_l^m(x, y, z)$  in terms of  $Y_{l'}^{m'}(x, y, z)$  for any point  $(x, y, z)$  on the unit sphere. Relations concerning the products of Spherical Harmonics have already been established, and so the expressions we desire here could just be seen as special cases of these. More specifically, the coefficients of the expansion in the SH basis of the product of two SHs are composed of Clebsch-Gordan coefficients [51, p231]. Clebsch-Gordan coefficients are important in quantum mechanics and angular momentum, with tables and algorithms existing to calculate them [41]. By simply choosing appropriate combinations, one could gain the desired coefficients for the expansion of each of  $x Y_l^m(x, y, z)$ ,  $y Y_l^m(x, y, z)$ , and  $z Y_l^m(x, y, z)$ .

However we reiterate our goal here is to use techniques that can generalise to spectral methods on other domains using multidimensional OPs, and thus we proceed by exploiting the three-term recurrences and other relations of the univariate OPs

that make up the Spherical Harmonics as we have defined them. With that being said, we require a few results concerning the incrementing and decrementing of the parameters for the Jacobi polynomials, and the three-term recurrence for the Jacobi polynomials, that will also prove useful later on in this chapter.

**Lemma 1.** *Let  $\alpha, \beta \in \mathbb{R}$  be general parameters. Let  $z \in [-1, 1]$  and define  $P_{-2}^{(\alpha+1, \beta+1)}(z) \equiv P_{-1}^{(\alpha+1, \beta+1)}(z) \equiv 0$ . Then, for any  $n \in \mathbb{N}_0$  the following identities hold:*

$$\begin{aligned} 1) \quad P_n^{(\alpha, \beta)}(z) = & \frac{1}{2n + \alpha + \beta + 1} \left\{ -\frac{(n + \alpha)(n + \beta)}{2n + \alpha + \beta} P_{n-2}^{(\alpha+1, \beta+1)}(z) \right. \\ & + (n + \alpha + \beta + 1) \left[ \frac{n + \alpha}{2n + \alpha + \beta} - \frac{n + \beta + 1}{2n + \alpha + \beta + 2} \right] P_{n-1}^{(\alpha+1, \beta+1)}(z) \\ & \left. + \frac{(n + \alpha + \beta + 1)(n + \alpha + \beta + 2)}{2n + \alpha + \beta + 2} P_n^{(\alpha+1, \beta+1)}(z) \right\}, \end{aligned}$$

$$\begin{aligned} 2) \quad (1 - z^2) P_n^{(\alpha, \beta)}(z) = & \frac{4}{2n + \alpha + \beta + 1} \left\{ \frac{(n + \alpha)(n + \beta)}{2n + \alpha + \beta} P_n^{(\alpha-1, \beta-1)}(z) \right. \\ & + (n + 1) \left[ \frac{n + \alpha + 1}{2n + \alpha + \beta + 2} - \frac{n + \beta}{2n + \alpha + \beta} \right] P_{n+1}^{(\alpha-1, \beta-1)}(z) \\ & \left. - \frac{(n + 1)(n + 2)}{2n + \alpha + \beta + 2} P_{n+2}^{(\alpha-1, \beta-1)}(z) \right\}, \end{aligned}$$

$$\begin{aligned} 3) \quad z P_n^{(\alpha, \beta)}(z) = & \frac{2(n + \alpha)(n + \beta)}{(2n + \alpha + \beta)(2n + \alpha + \beta + 1)} P_{n-1}^{(\alpha, \beta)}(z) \\ & + \frac{(\beta^2 - \alpha^2)}{(2n + \alpha + \beta)(2n + \alpha + \beta + 2)} P_n^{(\alpha, \beta)}(z) \\ & + \frac{2(n + 1)(n + \alpha + \beta + 1)}{2(2n + \alpha + \beta + 1)(2n + \alpha + \beta + 2)} P_{n+1}^{(\alpha, \beta)}(z). \end{aligned}$$

*Proof.* The first two results are simply consequences of the relationships estab-

lished for incrementing and decrementing the parameters of Jacobi polynomials detailed in [42, 18.9.5, 18.9.6], along with the symmetry relation that  $P_n^{(\alpha,\beta)}(-z) \equiv (-1)^n P_n^{(\beta,\alpha)}(z)$ . The final one is simply a rearranging of the established three-term recurrence detailed in [42, 18.9.1, 18.9.2]  $\square$

We thus can present a useful corollary specific for our needs.

**Corollary 1.** *Let  $l, m \in \mathbb{N}_0$  s.t.  $0 \leq m \leq l$  and let  $z \in [-1, 1]$ . Then:*

$$\begin{aligned} (1 - z^2) P_{l-m}^{(m,m)}(z) &= \tilde{\alpha}_{l,m,1} P_{l-m}^{(m-1,m-1)}(z) + \tilde{\alpha}_{l,m,3} P_{l-m+2}^{(m-1,m-1)}(z) \\ P_{l-m}^{(m,m)}(z) &= \tilde{\alpha}_{l,m,2} P_{l-m-2}^{(m+1,m+1)}(z) + \tilde{\alpha}_{l,m,4} P_{l-m}^{(m+1,m+1)}(z), \\ z P_{l-m}^{(m,m)}(z) &= \tilde{\gamma}_{l,m,1} P_{l-m-1}^{(m,m)}(z) + \tilde{\gamma}_{l,m,2} P_{l-m+1}^{(m,m)}(z), \end{aligned}$$

where

$$\begin{aligned} \tilde{\alpha}_{l,m,1} &:= \frac{2l}{2l+1}, & \tilde{\alpha}_{l,m,2} &:= \begin{cases} -\frac{l}{2(2l+1)} & \text{if } l-m \geq 2 \\ 0 & \text{otherwise} \end{cases}, \\ \tilde{\alpha}_{l,m,3} &:= -\frac{2(l-m+2)(l-m+1)}{(2l+1)(l+1)}, & \tilde{\alpha}_{l,m,4} &:= \frac{(l+m+2)(l+m+1)}{2(2l+1)(l+1)}, \\ \tilde{\gamma}_{l,m,1} &:= \begin{cases} \frac{l}{2l+1} & \text{if } l-m \geq 1 \\ 0 & \text{otherwise} \end{cases}, & \tilde{\gamma}_{l,m,2} &:= \frac{(l-m+1)(l+m+1)}{(2l+1)(l+1)}. \end{aligned}$$

Corollary 1 allows us to easily determine the aforementioned coefficients, that we will formalise in the following Lemma.

**Lemma 2.** *For  $l \in \mathbb{N}_0$ ,  $m \in \mathbb{Z}$  s.t.  $0 \leq |m| \leq l$ , the spherical harmonics satisfy*

the relationships:

$$\begin{aligned} x Y_l^m(x, y, z) &= \alpha_{l,m,1} Y_{l-1}^{m-1}(x, y, z) + \alpha_{l,m,2} Y_{l-1}^{m+1}(x, y, z) \\ &\quad + \alpha_{l,m,3} Y_{l+1}^{m-1}(x, y, z) + \alpha_{l,m,4} Y_{l+1}^{m+1}(x, y, z), \end{aligned} \quad (2.3)$$

$$\begin{aligned} y Y_l^m(x, y, z) &= \beta_{l,m,1} Y_{l-1}^{m-1}(x, y, z) + \beta_{l,m,2} Y_{l-1}^{m+1}(x, y, z) \\ &\quad + \beta_{l,m,3} Y_{l+1}^{m-1}(x, y, z) + \beta_{l,m,4} Y_{l+1}^{m+1}(x, y, z), \end{aligned} \quad (2.4)$$

$$z Y_l^m(x, y, z) = \gamma_{l,m,1} Y_{l-1}^m(x, y, z) + \gamma_{l,m,2} Y_{l+1}^m(x, y, z), \quad (2.5)$$

where

$$\begin{aligned} \alpha_{l,m,1} &:= \frac{c_l^m}{2c_{l-1}^{m-1}} \begin{cases} \tilde{\alpha}_{l,m,1} & \text{if } m > 0 \\ \tilde{\alpha}_{l,|m|,2} & \text{if } m \leq 0 \\ 0 & \text{otherwise} \end{cases}, & \alpha_{l,m,2} &:= \frac{c_l^m}{2c_{l-1}^{m+1}} \begin{cases} \tilde{\alpha}_{l,m,2} & \text{if } m \geq 0 \\ \tilde{\alpha}_{l,|m|,1} & \text{if } m < 0 \\ 0 & \text{otherwise} \end{cases}, \\ \alpha_{l,m,3} &:= \frac{c_l^m}{2c_{l+1}^{m-1}} \begin{cases} \tilde{\alpha}_{l,m,3} & \text{if } m > 0 \\ \tilde{\alpha}_{l,|m|,4} & \text{if } m \leq 0 \end{cases}, & \alpha_{l,m,4} &:= \frac{c_l^m}{2c_{l+1}^{m+1}} \begin{cases} \tilde{\alpha}_{l,m,4} & \text{if } m \geq 0 \\ \tilde{\alpha}_{l,|m|,3} & \text{if } m < 0 \end{cases}, \\ \gamma_{l,m,1} &:= \frac{c_l^m}{c_{l-1}^m} \tilde{\gamma}_{l,m,1}, & \gamma_{l,m,2} &:= \frac{c_l^m}{c_{l+1}^m} \tilde{\gamma}_{l,m,2}, \\ \beta_{l,m,j} &:= (-1)^{j+1} i \alpha_{l,m,j}, \quad j = 1, 2, 3, 4. \end{aligned}$$

**Remark:** The results in Lemma 2 are fairly involved and can be derived via the relationships established for products of Spherical Harmonics and Clebsch-Gordan coefficients as mentioned earlier [51, p231] but we include this proof so as to generalise to other OPs later on in later chapters.

*Proof of Lemma 2.* The expression for multiplication of the spherical harmonic  $Y_l^m$  by  $z$  is simply a consequence of the third result in Corollary 1:

$$\begin{aligned}
z Y_l^m(x, y, z) &= c_l^m e^{im\theta} \rho(z)^{|m|} z P_{l-|m|}^{(|m|, |m|)}(z) \\
&= c_l^m e^{im\theta} \rho(z)^{|m|} [\tilde{\gamma}_{l,m,1} P_{l-|m|-1}^{(|m|, |m|)}(z) + \tilde{\gamma}_{l,m,2} P_{l-|m|+1}^{(|m|, |m|)}(z)] \\
&= \gamma_{l,m,1} Y_{l-1}^m(x, y, z) + \gamma_{l,m,2} Y_{l+1}^m(x, y, z).
\end{aligned}$$

For multiplication by  $x$  and  $y$ , note that the complex exponential satisfies

$$\begin{aligned}
\cos \theta e^{im\theta} &= \frac{1}{2}(e^{i\theta} + e^{-i\theta})e^{im\theta} = \frac{1}{2}(e^{i(m+1)\theta} + e^{i(m-1)\theta}) \\
\sin \theta e^{im\theta} &= \frac{1}{2i}(e^{i\theta} - e^{-i\theta})e^{im\theta} = \frac{-i}{2}(e^{i(m+1)\theta} - e^{i(m-1)\theta})
\end{aligned}$$

Combining this with the results in Corollary 1, the expressions for multiplication of the spherical harmonic  $Y_l^m$  by  $x$  and  $y$  are then:

$$\begin{aligned}
x Y_l^m(x, y, z) &= c_l^m \cos \theta e^{im\theta} \rho(z)^{|m|+1} P_{l-|m|}^{(|m|, |m|)}(z) \\
&= \frac{1}{2} c_l^m (e^{i(m+1)\theta} + e^{i(m-1)\theta}) \rho(z)^{|m|+1} P_{l-|m|}^{(|m|, |m|)}(z) \\
&= \frac{1}{2} c_l^m e^{i(m+1)\theta} \rho(z)^{|m|+1} [\tilde{\alpha}_{l,m,4} P_{l-|m|}^{(|m|+1, |m|+1)}(z) + \tilde{\alpha}_{l,m,2} P_{l-|m|-2}^{(|m|+1, |m|+1)}(z)] \\
&\quad + \frac{1}{2} c_l^m e^{i(m-1)\theta} \rho(z)^{|m|-1} [\tilde{\alpha}_{l,m,3} P_{l-|m|+2}^{(|m|-1, |m|-1)}(z) + \tilde{\alpha}_{l,m,1} P_{l-|m|}^{(|m|-1, |m|-1)}(z)] \\
&= \alpha_{l,m,1} Y_{l-1}^{m-1}(x, y, z) + \alpha_{l,m,2} Y_{l-1}^{m+1}(x, y, z) \\
&\quad + \alpha_{l,m,3} Y_{l+1}^{m-1}(x, y, z) + \alpha_{l,m,4} Y_{l+1}^{m+1}(x, y, z),
\end{aligned}$$



and

$$\begin{aligned}
y Y_l^m(x, y, z) &= c_l^m \sin \theta e^{im\theta} \rho(z)^{|m|+1} P_{l-|m|}^{(|m|, |m|)}(z) \\
&= -\frac{1}{2} i c_l^m (e^{i(m+1)\theta} - e^{i(m-1)\theta}) \rho(z)^{|m|+1} P_{l-|m|}^{(|m|, |m|)}(z) \\
&= i \left\{ \alpha_{l,m,1} Y_{l-1}^{m-1}(x, y, z) - \alpha_{l,m,2} Y_{l-1}^{m+1}(x, y, z) \right. \\
&\quad \left. + \alpha_{l,m,3} Y_{l+1}^{m-1}(x, y, z) - \alpha_{l,m,4} Y_{l+1}^{m+1}(x, y, z) \right\}.
\end{aligned}$$

□

These recurrences lead to Jacobi operators that correspond to multiplication by  $x, y, z$  of the OPs. Define for  $l \in \mathbb{N}_0$ :

$$\mathbb{P}_l := \begin{pmatrix} Y_l^{-l} \\ \vdots \\ Y_l^l \end{pmatrix} \in \mathbb{C}^{2l+1}, \quad \mathbb{P} := \begin{pmatrix} \mathbb{P}_0 \\ \mathbb{P}_1 \\ \mathbb{P}_2 \\ \vdots \end{pmatrix}.$$

Note that these OP vectors we have defined are grouped and ordered according to the degree  $l$ , and within each degree ordered by the Fourier mode  $m$ . Let the Jacobi matrices  $J_x, J_y, J_z$  hence be given by

$$J_x \mathbb{P}(x, y, z) = x \mathbb{P}(x, y, z), \quad J_y \mathbb{P}(x, y, z) = y \mathbb{P}(x, y, z), \quad J_z \mathbb{P}(x, y, z) = z \mathbb{P}(x, y, z) \quad (2.6)$$

for  $(x, y, z) \in \Omega$ . These Jacobi matrices can act on the coefficients vector of a

function's expansion in the spherical harmonic basis to implement the operation for multiplication by  $x$ ,  $y$  or  $z$ . For example, given sufficiently large  $n \in \mathbb{N}$  let the function  $f(x, y, z) : \Omega \rightarrow \mathbb{C}$  be given by its expansion

$$f(x, y, z) = \sum_{l=0}^N \sum_{m=-l}^l f_{l,m} Y_l^m(x, y, z) = \mathbb{P}(x, y, z)^\top \mathbf{f}$$

where


$$\mathbf{f}_l := \begin{pmatrix} f_{l,-l} \\ \vdots \\ f_{l,l} \end{pmatrix} \in \mathbb{C}^{2l+1}, \quad \mathbf{f} := \begin{pmatrix} \mathbf{f}_0 \\ \vdots \\ \mathbf{f}_N \end{pmatrix}.$$

We say that  $\mathbf{f}$  is the *coefficients vector* of the function  $f$ . Then  $xf(x, y, z)$  is approximated by  $\mathbb{P}(x, y, z)^\top J_x^\top \mathbf{f}$ . In other words,  $J_x^\top \mathbf{f}$  is the coefficients vector for the expansion of the function  $(x, y, z) \mapsto xf(x, y, z)$  in the spherical harmonics basis.

An important property of the Jacobi matrices is that they are sparse – specifically they are *banded-block-banded matrices*.

**Definition 1.** A block matrix  $A$  with blocks  $A_{i,j}$  has *block-bandwidths*  $(L, U)$  if  $A_{i,j} = 0$  for  $-L \leq j - i \leq U$ , and *subblock-bandwidths*  $(\lambda, \mu)$  if all blocks are banded with bandwidths  $(\lambda, \mu)$ . A matrix where the block-bandwidths and subblock-bandwidths are small compared to the dimensions is referred to as a *banded-block-banded matrix*.

Using equations (2.3–2.5), we have that the Jacobi matrices take the following



**Figure 2.1:** “Spy” plots of the Jacobi operator matrices, showing their sparsity and structure, up to order  $N = 10$ .

block-tridiagonal form (i.e. have block-bandwidths  $(1, 1)$ ):

$$J_x = \begin{pmatrix} B_{x,0} & A_{x,0} & & & \\ C_{x,1} & B_{x,1} & A_{x,1} & & \\ & C_{x,2} & B_{x,2} & A_{x,2} & \\ & & C_{x,3} & \ddots & \ddots \\ & & & \ddots & \ddots & \ddots \end{pmatrix},$$

$$J_y = \begin{pmatrix} B_{y,0} & A_{y,0} & & & \\ C_{y,1} & B_{y,1} & A_{y,1} & & \\ & C_{y,2} & B_{y,2} & A_{y,2} & \\ & & C_{y,3} & \ddots & \ddots \\ & & & \ddots & \ddots & \ddots \end{pmatrix},$$

$$J_z = \begin{pmatrix} B_{z,0} & A_{z,0} & & & \\ C_{z,1} & B_{z,1} & A_{z,1} & & \\ & C_{z,2} & B_{z,2} & A_{z,2} & \\ & & C_{z,3} & \ddots & \ddots \\ & & & \ddots & \ddots & \ddots \end{pmatrix}.$$

$J_x$  has subblock-bandwidths  $(2, 2)$ , where the blocks for  $l \in \mathbb{N}_0$  are given by:

$$A_{x,l} := \begin{pmatrix} \alpha_{l,-l,3} & 0 & \alpha_{l,-l,4} & & \\ & \ddots & \ddots & \ddots & \\ & & \alpha_{l,l,3} & 0 & \alpha_{l,l,4} \end{pmatrix} \in \mathbb{R}^{(2l+1) \times (2l+3)},$$

$$B_{x,l} := 0 \in \mathbb{R}^{(2l+1) \times (2l+1)}$$

$$C_{x,l} := \begin{pmatrix} \alpha_{l,-l,2} & & & & \\ 0 & \ddots & & & \\ \alpha_{l,-l+2,1} & \ddots & \alpha_{l,l-2,2} & & \\ & \ddots & 0 & & \\ & & & \alpha_{l,l,1} & \end{pmatrix} \in \mathbb{R}^{(2l+1) \times (2l-1)} \quad (l \neq 0).$$

$J_y$  also has subblock-bandwidths  $(2, 2)$ , where the blocks for  $l \in \mathbb{N}_0$  are given by:

$$A_{y,l} := \begin{pmatrix} \beta_{l,-l,3} & 0 & \beta_{l,-l,4} & & \\ & \ddots & \ddots & \ddots & \\ & & \beta_{l,l,3} & 0 & \beta_{l,l,4} \end{pmatrix} \in \mathbb{C}^{(2l+1) \times (2l+3)},$$

$$B_{y,l} := 0 \in \mathbb{R}^{(2l+1) \times (2l+1)}$$

$$C_{y,l} := \begin{pmatrix} \beta_{l,-l,2} & & & & \\ 0 & \ddots & & & \\ \beta_{l,-l+2,1} & \ddots & \beta_{l,l-2,2} & & \\ & \ddots & 0 & & \\ & & & \beta_{l,l,1} & \end{pmatrix} \in \mathbb{C}^{(2l+1) \times (2l-1)} \quad (l \neq 0).$$

Finally,  $J_z$  has subblock-bandwidths  $(1, 1)$ , where the blocks for  $l \in \mathbb{N}_0$  are given by:

$$A_{z,l} := \begin{pmatrix} 0 & \gamma_{l,-l,2} & 0 & & \\ & \ddots & \ddots & \ddots & \\ & & 0 & \gamma_{l,l,2} & 0 \end{pmatrix} \in \mathbb{R}^{(2l+1) \times (2l+3)},$$

$$B_{z,l} := 0 \in \mathbb{R}^{(2l+1) \times (2l+1)}$$

$$C_{z,l} := \begin{pmatrix} 0 & & & & \\ \gamma_{l,-l+1,1} & \ddots & & & \\ 0 & \ddots & 0 & & \\ & \ddots & \gamma_{l,l-1,1} & & \\ & & & 0 & \end{pmatrix} \in \mathbb{R}^{(2l+1) \times (2l-1)} \quad (l \neq 0).$$

The sparsity and structure of the Jacobi matrices as we have defined them can be seen visually in Figure 2.2.

### 2.3 Three-term recurrence relation for $\mathbb{P}$

Three-term recurrence relations for orthogonal polynomials are well established (see for example [42, Section 18.9]), including for multidimensional OPs [17, Section 3.2]. In a similar vein, we can obtain a recurrence relation for the Spherical Harmonics by combining each system in equation (2.6) in a specific way. It should be emphasised at this point that this is not necessarily the optimal way of computing the Spherical Harmonics; various efficient methods and algorithms have

been proposed (see e.g. [23, 57, 20]). We present this method in order to be able to generalise it to other multidimensional OP families later on in this thesis.

Rewriting equation (2.6), we have that:

$$(J_x - xI)\mathbb{P}(x, y, z) = (J_y - yI)\mathbb{P}(x, y, z) = (J_z - zI)\mathbb{P}(x, y, z) = \mathbf{0}.$$

Combining these relationships into a single system, complete with initial (degree 0) value, results in:

$$\begin{pmatrix} 1 & & & & \\ B_0 - G_0(x, y, z) & A_0 & & & \\ C_1 & B_1 - G_1(x, y, z) & A_1 & & \\ & C_2 & B_2 - G_2(x, y, z) & \ddots & \\ & & \ddots & \ddots & \end{pmatrix} \mathbb{P}(x, y, z) = \begin{pmatrix} Y_0 \\ 0 \\ 0 \\ 0 \\ \vdots \end{pmatrix},$$

where we note  $Y_0^0(x, y, z) \equiv Y_0 := c_0^0 P_0^{(0,0)} \equiv \frac{1}{2} \frac{1}{\sqrt{\pi}}$ , and for each  $l = 0, 1, 2, \dots$ ,

$$A_l := \begin{pmatrix} A_{x,l} \\ A_{y,l} \\ A_{z,l} \end{pmatrix} \in \mathbb{C}^{3(2l+1) \times (2l+3)}, \quad C_l := \begin{pmatrix} C_{x,l} \\ C_{y,l} \\ C_{z,l} \end{pmatrix} \in \mathbb{C}^{3(2l+1) \times (2l-1)} \quad (n \neq 0), \quad (2.7)$$

$$B_l := \begin{pmatrix} B_{x,l} \\ B_{y,l} \\ B_{z,l} \end{pmatrix} \in \mathbb{C}^{3(2l+1) \times (2l+1)}, \quad G_n(x, y) := \begin{pmatrix} xI_{2l+1} \\ yI_{2l+1} \\ zI_{2l+1} \end{pmatrix} \in \mathbb{C}^{3(2l+1) \times (2l+1)}. \quad (2.8)$$

For each  $l = 0, 1, 2, \dots$  let  $D_l^\top$  be any matrix that is a left inverse of  $A_l$ , i.e. such that  $D_l^\top A_l = I_{2l+3}$ . Multiplying our system by the preconditioner matrix that is given by the block diagonal matrix of the  $D_l^\top$ 's, we obtain a lower triangular system [17, Section 3.2], which can be expanded to obtain the three-term recurrence:

$$\begin{cases} \mathbb{P}_{-1}(x, y, z) := 0 \\ \mathbb{P}_0(x, y, z) := Y_0 \\ \mathbb{P}_{l+1}(x, y, z) = -D_l^\top (B_l - G_l(x, y, z)) \mathbb{P}_l(x, y, z) - D_l^\top C_l \mathbb{P}_{l-1}(x, y, z), \quad l = 0, 1, 2, \dots \end{cases}$$

Since the above holds for any  $D_l^\top$  that is a left inverse of  $A_l$ , we are free to choose the  $D_l^\top$  matrices in the following way. For  $l \in \mathbb{N}$ , we can set

$$D_l^\top = \begin{pmatrix} & & & \boldsymbol{\eta}_0^\top & & \\ 0 & & 0 & & \frac{1}{\gamma_{l,-l,2}} & \\ & \ddots & & \ddots & & \ddots \\ & & 0 & 0 & & \frac{1}{\gamma_{l,l,2}} \\ & & & \boldsymbol{\eta}_1^\top & & \end{pmatrix} \in \mathbb{R}^{(2n+3) \times 3(2n+1)}, \quad (2.9)$$



where  $\boldsymbol{\eta}_0, \boldsymbol{\eta}_1 \in \mathbb{R}^{3(2n+1)}$  are vectors with entries given by

$$(\boldsymbol{\eta}_0)_j = \begin{cases} \frac{1}{\alpha_{l,-l,3}} & j = 1 \\ \frac{-\alpha_{l,-l,4}}{\alpha_{l,-l,3} \gamma_{l,1-l,2}} & j = 2(2n+1) + 2, \\ 0 & o/w \end{cases}$$

$$(\boldsymbol{\eta}_1)_j = \begin{cases} \frac{1}{\alpha_{l,l,4}} & j = 2l + 1 \\ \frac{-\alpha_{l,l,3}}{\alpha_{l,l,4} \gamma_{l,l-1,2}} & j = 3(2n+1) - 1. \\ 0 & o/w \end{cases}$$

For  $l = 0$  we can simply set

$$D_0^T = \begin{pmatrix} \frac{\beta_{0,0,4}}{\alpha_{0,0,3}\beta_{0,0,4}-\alpha_{0,0,4}\beta_{0,0,3}} & \frac{-\alpha_{0,0,4}}{\alpha_{0,0,3}\beta_{0,0,4}-\alpha_{0,0,4}\beta_{0,0,3}} & 0 \\ 0 & 0 & \frac{1}{\gamma_{0,0,2}} \\ \frac{\beta_{0,0,3}}{\alpha_{0,0,4}\beta_{0,0,3}-\alpha_{0,0,3}\beta_{0,0,4}} & \frac{-\alpha_{0,0,3}}{\alpha_{0,0,4}\beta_{0,0,3}-\alpha_{0,0,3}\beta_{0,0,4}} & 0 \end{pmatrix}. \quad (2.10)$$

It will be useful for us to give a formal name for these coefficient matrices above for a family of multidimensional orthogonal polynomials.

**Definition 2.** The matrices  $-D_l^\top(B_l - G_l(x, y, z)), D_l^\top C_l$  for  $l \in \mathbb{N}_0$  defined via equations (2.7–2.10) are called the **recurrence coefficient matrices** for a given family of multidimensional orthogonal polynomials.

## 2.4 Computational aspects

Once again, let  $f(x, y, z)$  be a function on the unit sphere  $\Omega$  be approximated by its expansion

$$f(x, y, z) \approx \mathbb{P}(x, y, z)^\top \mathbf{f} = \sum_{l=0}^N \mathbb{P}_l(x, y, z)^\top \mathbf{f}_l = \sum_{l=0}^N \sum_{m=-l}^l f_{l,m} Y_l^m(x, y, z),$$

where  $\mathbb{P}_l(x, y, z), \mathbf{f}_l \in \mathbb{C}^{2l+1}$  for each  $l \in \{0, \dots, N\}$ , for some coefficients vector  $\mathbf{f} = (f_{l,m})$  up to degree order  $N \in \mathbb{N}$ .

### 2.4.1 Obtaining coefficients

In spectral space, we wish to work only with vectors of coefficients for the expansion of a function, to which we can apply operator matrices to that represent differential or other operations. Naturally, we of course need a way to obtain the coefficients  $f_{l,m}$ . The coefficients can be calculated via the integral

$$f_{l,m} = \int_{\Omega} f(\cos \theta \sin \varphi, \sin \theta \sin \varphi, \cos \varphi) Y_l^m(\varphi, \theta)^* \sin \varphi \, d\varphi \, d\theta$$

using the orthonormality of the spherical harmonics. As discussed in Section ??, methods to calculate these coefficients exist known as spectral transforms, and are well established for the spherical harmonics (see e.g. [38, 60]). Additionally, packages are available for public use that implement efficient spherical harmonics transforms, for example the FastTransforms package [55].

### 2.4.2 Function evaluation

We can use the Clenshaw algorithm to evaluate this function at a given point  $(x, y, z)$  on  $\Omega$  [48]. The *Clenshaw algorithm* is then as follows:

1) Set  $\boldsymbol{\xi}_{N+2} = \mathbf{0}$ ,  $\boldsymbol{\xi}_{N+1} = \mathbf{0}$ .

2) For  $n = N : -1 : 0$

$$\text{set } \boldsymbol{\xi}_n^T = \mathbf{f}_n^T - \boldsymbol{\xi}_{n+1}^T D_n^T (B_n - G_n(x, y, z)) - \boldsymbol{\xi}_{n+2}^T D_{n+1}^T C_{n+1}$$

3) Output:  $f(x, y, z) \approx \boldsymbol{\xi}_0^T \mathbb{P}_0(x, y, z) \equiv \xi_0 Y_0$ .

### 2.4.3 Obtaining operator matrices for variable coefficients

The Clenshaw algorithm presented in Section 2.4.2 can also be used with the Jacobi matrices replacing the point  $(x, y, z)$ , to yield an operator matrix.

Let's explain what we mean. Suppose  $v : \Omega \rightarrow \mathbb{C}$  is a function, and we encounter the problem of finding the coefficients of the expansion for  $v(x, y, z)f(x, y, z)$ . We wish to therefore find an operator  $V$  for  $v$  so that

$$v(x, y, z)f(x, y, z) = v(x, y, z)\mathbf{f}^\top \mathbb{P}(x, y, z) = (V\mathbf{f})^\top \mathbb{P}(x, y, z),$$

i.e.  $V\mathbf{f}$  is the coefficients vector for the expansion of the function  $(x, y, z) \mapsto v(x, y, z)f(x, y, z)$  in the spherical harmonic basis.

Let  $\mathbf{v}$  be the coefficients of the expansion of  $v$  up to order  $N$ . The operator  $V$

would then be the result of the following *operator Clenshaw algorithm*:

1) Set  $\boldsymbol{\xi}_{N+2} = \mathbf{0}$ ,  $\boldsymbol{\xi}_{N+1} = \mathbf{0}$ .

2) For  $n = N : -1 : 0$

$$\text{set } \boldsymbol{\xi}_n^T = \mathbf{v}_n^T - \boldsymbol{\xi}_{n+1}^T D_n^T (B_n - G_n(J_x, J_y, J_z)) - \boldsymbol{\xi}_{n+2}^T D_{n+1}^T C_{n+1}$$

3) Output:  $f(x, y, z) \approx \boldsymbol{\xi}_0^T \mathbb{P}_0(x, y, z) \equiv \xi_0 Y_0$

where at each iteration,  $\boldsymbol{\xi}_n$  is a vector of matrices (note that we are abusing notation here a bit, however it is the simplest way to present the algorithm without introducing yet more matrices!).

## 2.5 Vector spherical harmonics as orthogonal vectors in three variables

We have established that the Spherical Harmonics can be viewed as a basis of orthonormal polynomials on the sphere. Specifically, given a function  $f : \Omega \rightarrow \mathbb{C}$ , we can write  $f = \sum_{l=0}^N \sum_{m=-l}^l f_{l,m} Y_l^m$  where  $\{f_{l,m}\}$  are coefficients and  $N \in \mathbb{N}$  is sufficiently large. But what about vector-valued functions?

**Definition 3.** Let  $(x, y, z) \in \Omega$ . We refer to the **tangent space** of the point  $(x, y, z)$  as the space of all vectors that are orthogonal to the vector  $\hat{\mathbf{r}} := [x, y, z]^\top$ . We further define the **tangent bundle** of  $\Omega$  as the set of all such spaces for every point in  $\Omega$ .

For the purposes of this discussion, we will only consider the tangent bundle, and thus wish to write down a basis for vector-valued functions with values in the tangent bundle. Fortunately, there is a simple solution known as the *vector spherical harmonics* (VSHs) [2]:

$$\begin{aligned}\Psi_l^m &:= \nabla Y_l^m = \frac{\partial}{\partial \phi} Y_l^m \hat{\phi} + \frac{1}{\sin \phi} \frac{\partial}{\partial \theta} Y_l^m \hat{\theta}, \\ \Phi_l^m &:= \nabla^\perp Y_l^m \equiv \hat{\mathbf{r}} \times \nabla Y_l^m = \frac{\partial}{\partial \phi} Y_l^m \hat{\theta} - \frac{1}{\sin \phi} \frac{\partial}{\partial \theta} Y_l^m \hat{\phi}.\end{aligned}\quad (2.11)$$

Here,  $\hat{\mathbf{r}}$  is simply the outward unit normal vector to the surface of the sphere at the point  $(x, y, z)$ , while  $\hat{\phi}, \hat{\theta}$  are the standard spherical coordinate unit vectors, i.e.  $\hat{\phi} := [\cos \theta \cos \phi, \sin \theta \cos \phi, -\sin \phi]^\top$ ,  $\hat{\theta} := [-\sin \theta, \cos \theta, 0]^\top$ . They are orthogonal with respect to the inner product  $\langle \mathbf{A}, \mathbf{B} \rangle = \iint_\Omega \mathbf{A} \cdot \mathbf{B}^* \sin \phi \, d\phi \, d\theta$  – in particular, for any  $l, l' \in \mathbb{N}$ ,  $m, m' \in \mathbb{Z}$  s.t.  $-l \leq m \leq l$ ,  $-l' \leq m' \leq l'$ ,

$$\begin{aligned}\langle \Psi_l^m, \Psi_{l'}^{m'} \rangle &= \iint_\Omega \Psi_l^m \cdot \Psi_{l'}^{m'*} \sin \phi \, d\phi \, d\theta = \delta_{l,l'} \delta_{m,m'} l(l+1), \\ \langle \Phi_l^m, \Phi_{l'}^{m'} \rangle &= \iint_\Omega \Phi_l^m \cdot \Phi_{l'}^{m'*} \sin \phi \, d\phi \, d\theta = \delta_{l,l'} \delta_{m,m'} l(l+1), \\ \langle \Psi_l^m, \Phi_{l'}^{m'} \rangle &= \iint_\Omega \Psi_l^m \cdot \Phi_{l'}^{m'*} \sin \phi \, d\phi \, d\theta = 0.\end{aligned}$$

Let  $\mathbf{u}$  be a vector-valued function on  $\Omega$  whose values are of the form

$$\mathbf{u}(x, y, z) = u_\phi(x, y, z) \hat{\phi} + u_\theta(x, y, z) \hat{\theta}, \quad (x, y, z) \in \Omega$$

for some scalar functions  $u^\phi, u^\theta$ . Such a function  $\mathbf{u}$  we refer to as a *vector-valued function in the tangent bundle* of the sphere. Importantly, the set  $\{\Psi_l^m, \Phi_l^m\}$  form

a complete and orthogonal basis for such functions [2]. That is, given  $\mathbf{u}$ , there exist coefficients  $\{u_{l,m}^{\Psi}, u_{l,m}^{\Phi}\}$  such that for large enough  $N$ :

$$\mathbf{u} = \sum_{l=0}^N \sum_{m=-l}^l u_{l,m}^{\Psi} \Psi_l^m + u_{l,m}^{\Phi} \Phi_l^m.$$

Vector spherical harmonics have been widely used in electrostatics (e.g. [2]), electrodynamics (e.g. [11]), and of course fluid dynamics including weather and climate modelling (e.g. [40, 19, 62]). Other definitions for VSHs are used (e.g. [24]), however it is convenient for deriving explicit sparse relations and operators to use the ones described here. In particular, we can note that computing the gradient or divergence of a function is merely a matter of conversion between the SH and the VSH bases.

A simple calculation shows that such orthogonal vectors must still have block-tridiagonal Jacobi operators, as multiplication by  $x, y$ , or  $z$  remains inside the ideal. Thus, we can use the same techniques that we used for the scalar SHs to derive Jacobi matrices, function evaluation and transforms.

### 2.5.1 Jacobi matrices

We start as before for the scalar case by finding  $x \Psi_l^m(x, y, z)$ ,  $y \Psi_l^m(x, y, z)$ ,  $z \Psi_l^m(x, y, z)$ , and  $x \Phi_l^m(x, y, z)$ ,  $y \Phi_l^m(x, y, z)$ ,  $z \Phi_l^m(x, y, z)$  in terms of  $\Psi_{l'}^{m'}(x, y, z)$ ,  $\Phi_{l'}^{m'}(x, y, z)$ . A lovely consequence of our definitions of the scalar and vector spherical harmonics is that we can once again find explicit expressions for the coefficients in the aforementioned relations. To this end, we recall and derive some important

relations of the complex exponential and Jacobi polynomials that are a part of the spherical harmonics definitions. In particular we first note that:

$$\int_0^{2\pi} e^{im\theta} e^{-im'\theta} \cos(\theta) d\theta = \pi(\delta_{m',m-1} + \delta_{m',m+1}) \quad (2.12)$$

$$\int_0^{2\pi} e^{im\theta} e^{-im'\theta} \sin(\theta) d\theta = i \pi(\delta_{m',m-1} - \delta_{m',m+1}) \quad (2.13)$$

and, for  $l \in \mathbb{N}_0, m \in \mathbb{Z}$  s.t.  $-l \leq m \leq l$ ,

$$\int_{-1}^1 P_{l-|m|}^{(|m|,|m|)}(z) P_{l'-|m|}^{(|m|,|m|)}(z) \rho(z)^{2|m|} dz = \delta_{l',l} \frac{1}{2\pi(c_l^m)^2} \quad (2.14)$$

$$P_{l-|m|}^{(|m|,|m|)'}(z) = d_{l,m} P_{l-(|m|+1)}^{(|m|+1,|m|+1)} \quad \text{where } d_{l,m} := \begin{cases} \frac{1}{2}(l + |m| + 1) & l > |m| \\ 0 & l = |m| \end{cases}. \quad (2.15)$$

Equation (2.14) and equation (2.15) are simply consequences of the definition of the Jacobi polynomials – see [42, Table 18.3.1] and [42, 18.9.15].

The expressions for the VSHs as given in equation (2.11) together with the above allow us to be able to find the coefficients for our desired expressions for  $x\Psi_l^m$ ,  $y\Psi_l^m$ ,  $z\Psi_l^m$ ,  $x\Phi_l^m$ ,  $y\Phi_l^m$  and  $z\Phi_l^m$ .

**Lemma 3.** *For  $l \in \mathbb{N}$  and  $m \in \mathbb{Z}$  s.t.  $-l \leq m \leq l$ , the vector spherical harmonics*

satisfy the relationships:

$$\begin{aligned}
x\Psi_l^m &= A_{l,m,1}\Psi_{l-1}^{m-1} + A_{l,m,2}\Psi_{l-1}^{m+1} + A_{l,m,3}\Psi_{l+1}^{m-1} + A_{l,m,4}\Psi_{l+1}^{m+1} \\
&\quad + A_{l,m,5}\Phi_l^{m-1} + A_{l,m,6}\Phi_l^{m+1}, \\
x\Phi_l^m &= A_{l,m,1}\Phi_{l-1}^{m-1} + A_{l,m,2}\Phi_{l-1}^{m+1} + A_{l,m,3}\Phi_{l+1}^{m-1} + A_{l,m,4}\Phi_{l+1}^{m+1} \\
&\quad - A_{l,m,5}\Psi_l^{m-1} - A_{l,m,6}\Psi_l^{m+1}, \\
y\Psi_l^m &= B_{l,m,1}\Psi_{l-1}^{m-1} + B_{l,m,2}\Psi_{l-1}^{m+1} + B_{l,m,3}\Psi_{l+1}^{m-1} + B_{l,m,4}\Psi_{l+1}^{m+1} \\
&\quad + B_{l,m,5}\Phi_l^{m-1} + B_{l,m,6}\Phi_l^{m+1}, \\
y\Phi_l^m &= B_{l,m,1}\Phi_{l-1}^{m-1} + B_{l,m,2}\Phi_{l-1}^{m+1} + B_{l,m,3}\Phi_{l+1}^{m-1} + B_{l,m,4}\Phi_{l+1}^{m+1} \\
&\quad - B_{l,m,5}\Psi_l^{m-1} - B_{l,m,6}\Psi_l^{m+1}, \\
z\Psi_l^m &= \Gamma_{l,m,1}\Psi_{l-1}^m + \Gamma_{l,m,2}\Psi_{l+1}^m + \Gamma_{l,m,3}\Phi_l^m, \\
z\Phi_l^m &= \Gamma_{l,m,1}\Phi_{l-1}^m + \Gamma_{l,m,2}\Phi_{l+1}^m - \Gamma_{l,m,3}\Psi_l^m
\end{aligned}$$



where,

$$\begin{aligned}
A_{l,m,1} &:= \frac{c_l^m c_{l-1}^{m-1}}{2l(l-1)} \begin{cases} \tilde{A}_{l,m,1} & \text{if } m > 0 \\ \tilde{A}_{l,|m|,2} & \text{if } m \leq 0 \end{cases} \\
A_{l,m,2} &:= \frac{c_l^m c_{l-1}^{m+1}}{2l(l-1)} \begin{cases} \tilde{A}_{l,m,2} & \text{if } m \geq 0 \\ \tilde{A}_{l,|m|,1} & \text{if } m < 0 \end{cases} \\
A_{l,m,3} &:= \frac{c_l^m c_{l+1}^{m-1}}{2(l+1)(l+2)} \begin{cases} \tilde{A}_{l,m,3} & \text{if } m > 0 \\ \tilde{A}_{l,|m|,4} & \text{if } m \leq 0 \end{cases} \\
A_{l,m,4} &:= \frac{c_l^m c_{l+1}^{m+1}}{2(l+1)(l+2)} \begin{cases} \tilde{A}_{l,m,4} & \text{if } m \geq 0 \\ \tilde{A}_{l,|m|,3} & \text{if } m < 0 \end{cases} \\
A_{l,m,5} &:= \frac{i}{2l(l+1)} \begin{cases} -i d_{l,m-1} \frac{c_l^{m-1}}{2c_l^m} & \text{if } m > 0 \\ i d_{l,|m|} \frac{c_l^m}{2c_l^{m-1}} & \text{if } m \leq 0 \end{cases} \\
A_{l,m,6} &:= \frac{i}{2l(l+1)} \begin{cases} -d_{l,m} \frac{c_l^m}{c_l^{m+1}} & \text{if } m \geq 0 \\ d_{l,|m|-1} \frac{c_l^{m+1}}{c_l^m} & \text{if } m < 0 \end{cases} \\
B_{l,m,j} &:= i(-1)^{j+1} A_{l,m,j} \quad \text{for } j = 1, \dots, 6 \\
\Gamma_{l,m,1} &:= \frac{c_l^m c_{l-1}^m}{l(l-1)} \left[ |m| (|m| + 2) \frac{\tilde{\gamma}_{l,|m|,1}}{(c_{l-1}^{|m|})^2} + d_{l,|m|} d_{l-1,|m|} \frac{\tilde{\gamma}_{l,|m|+1,1}}{(c_{l-1}^{|m|+1})^2} \right] \\
\Gamma_{l,m,2} &:= \frac{c_l^m c_{l+1}^m}{(l+1)(l+2)} \left[ |m| (|m| + 2) \frac{\tilde{\gamma}_{l,|m|,2}}{(c_{l+1}^{|m|})^2} + d_{l,|m|} d_{l+1,|m|} \frac{\tilde{\gamma}_{l,|m|+1,2}}{(c_{l+1}^{|m|+1})^2} \right] \\
\Gamma_{l,m,3} &:= \frac{i m}{l(l+1)}
\end{aligned}$$

where the  $c_l^{m'}$ ,  $d_{l,m'}$ ,  $\tilde{\gamma}_{l,m',j}$  are defined in equation (2.2), equation (2.15) and

Corollary 1 respectively, and

$$\begin{aligned}\tilde{A}_{l,m,1} &:= \left( (m^2 - 1)\tilde{\alpha}_{l-1,m-1,4} - d_{l-1,m-1}\tilde{\gamma}_{l-1,m,2} \right) \frac{1}{(c_l^m)^2} + d_{l,m}d_{l-1,m-1} \frac{\tilde{\alpha}_{l,m+1,1}}{(c_{l-1}^m)^2} \\ \tilde{A}_{l,m,2} &:= \left( m(m+2)\tilde{\alpha}_{l,m,2} - d_{l,m}\tilde{\gamma}_{l,m+1,1} \right) \frac{1}{(c_{l-1}^{m+1})^2} + d_{l,m}d_{l-1,m+1} \frac{\tilde{\alpha}_{l-1,m+2,3}}{(c_l^{m+1})^2} \\ \tilde{A}_{l,m,3} &:= \left( (m^2 - 1)\tilde{\alpha}_{l+1,m-1,2} - d_{l+1,m-1}\tilde{\gamma}_{l+1,m,1} \right) \frac{1}{(c_l^m)^2} + d_{l,m}d_{l+1,m-1} \frac{\tilde{\alpha}_{l,m+1,3}}{(c_{l+1}^m)^2} \\ \tilde{A}_{l,m,4} &:= \left( m(m+2)\tilde{\alpha}_{l,m,4} - d_{l,m}\tilde{\gamma}_{l,m+1,2} \right) \frac{1}{(c_{l+1}^{m+1})^2} + d_{l,m}d_{l+1,m+1} \frac{\tilde{\alpha}_{l+1,m+2,1}}{(c_l^{m+1})^2}.\end{aligned}$$

where again the  $\tilde{\alpha}_{l',m',j}$  are defined in Corollary 1 too.

**Remark:** We emphasise again that it is due to the unique relationships that the Jacobi polynomials possess that we are able to explicitly write down these coefficients.

*Proof of Lemma 3.* Fix  $l \in \mathbb{N}$  and  $m \in \mathbb{Z}$  s.t.  $-l \leq m \leq l$ . The proof we will use is to directly calculate the nonzero coefficients in each expansion using inner products. To this end, note that

$$x\Psi_l^m = \sum_{l'=1}^{l+1} \sum_{m'=-l'}^{l'} \mathcal{A}_{l',m'} \Psi_{l'}^{m'} + \mathcal{A}_{l',m'}^\perp \Phi_{l'}^{m'}$$

where the coefficients are given by

$$\begin{aligned}\mathcal{A}_{l',m'} &= \frac{\langle x\Psi_l^m, \Psi_{l'}^{m'} \rangle}{\|\Psi_{l'}^{m'}\|^2} = \frac{1}{l'(l'+1)} \langle x\Psi_l^m, \Psi_{l'}^{m'} \rangle \\ \mathcal{A}_{l',m'}^\perp &= \frac{\langle x\Psi_l^m, \Phi_{l'}^{m'} \rangle}{\|\Phi_{l'}^{m'}\|^2} = \frac{1}{l'(l'+1)} \langle x\Psi_l^m, \Phi_{l'}^{m'} \rangle.\end{aligned}$$

The aim will be to show that the only nonzero coefficients  $\mathcal{A}_{l',m'}, \mathcal{A}_{l',m'}^\perp$  match the  $A_{l,m,j}$  for  $j = 1, \dots, 6$  in the Lemma. Now, using a change of variable, we have that

$$\begin{aligned}
 & \langle x \Psi_l^m, \Psi_{l'}^{m'} \rangle \\
 &= \iint_{\Omega} \cos \theta \sin \varphi \Psi_l^m \cdot \Psi_{l'}^{m'*} \sin \varphi \, d\phi \, d\theta \\
 &= c_l^m c_{l'}^{m'} \left( \int_0^{2\pi} e^{im\theta} e^{im'\theta} \cos \theta \, d\theta \right) \\
 & \quad \cdot \left( \int_{-1}^1 \left\{ [|m| z P_{l-|m|}^{(|m|,|m|)}(z) - \rho(z)^2 P_{l-|m|}'^{(|m|,|m|)}(z)] [|m'| z P_{l'-|m'|}^{(|m'|,|m'|)}(z) - \rho(z)^2 P_{l'-|m'|}'^{(|m'|,|m'|)}(z)] \right. \right. \\
 & \quad \left. \left. + mm' P_{l-|m|}^{(|m|,|m|)}(z) P_{l'-|m'|}^{(|m'|,|m'|)}(z) \right\} \rho(z)^{|m|+|m'|-1} \, dz \right) \\
 &= c_l^m c_{l'}^{m'} \pi (\delta_{m',m-1} + \delta_{m',m+1}) \\
 & \quad \cdot \left( \int_{-1}^1 \left\{ P_{l-|m|}^{(|m|,|m|)} P_{l'-|m'|}^{(|m'|,|m'|)} \rho^{|m|+|m'|-1} (mm' + |m| |m'| z^2) \right. \right. \\
 & \quad - |m| z P_{l-|m|}^{(|m|,|m|)} P_{l'-|m'|}'^{(|m'|,|m'|)} \rho^{|m|+|m'|+1} \\
 & \quad - |m'| z P_{l-|m|}'^{(|m|,|m|)} P_{l'-|m'|}^{(|m'|,|m'|)} \rho^{|m|+|m'|+1} \\
 & \quad \left. \left. + P_{l-|m|}'^{(|m|,|m|)} P_{l'-|m'|}'^{(|m'|,|m'|)} \rho^{|m|+|m'|+3} \right\} \, dz \right). \tag{2.16}
 \end{aligned}$$

We can conclude that the only possible nonzero coefficients are when  $m' = m \pm 1$ , and so it is left to evaluate the integral that is left at equation (2.16) when  $m'$  takes either value. Since we need to take account of the fact that  $m$  can be negative, to simplify our argument we make use of the fact that  $m' = m - 1$  for  $m < 0$  is analogous to the case  $m' = m + 1$  for  $m > 0$ , and vice versa.

On that note, we will first consider the case that  $m > 0$  and  $m' = m - 1$ , and

calculate the integral in equation (2.16):

$$\begin{aligned}
& \int_{-1}^1 \left\{ m(m-1) P_{l-m}^{(m,m)} P_{l'-(m-1)}^{(m-1,m-1)} \rho^{2m-2} (1+z^2) - m z P_{l-m}^{(m,m)} P_{l'-(m-1)}^{(m-1,m-1)'} \rho^{2m} \right. \\
& \quad \left. - (m-1) z P_{l-m}^{(m,m)'} P_{l'-(m-1)}^{(m-1,m-1)} \rho^{2m} + P_{l-m}^{(m,m)'} P_{l'-(m-1)}^{(m-1,m-1)'} \rho^{2m+2} \right\} dz \\
&= \int_{-1}^1 \left\{ m(m-1) P_{l-m}^{(m,m)} P_{l'-(m-1)}^{(m-1,m-1)} \rho^{2m-2} (1+z^2) - m z P_{l-m}^{(m,m)} P_{l'-(m-1)}^{(m-1,m-1)'} \rho^{2m} \right. \\
& \quad + (m-1) P_{l-m}^{(m,m)} \rho^{2m-2} \left( -2m z^2 P_{l'-(m-1)}^{(m-1,m-1)} + P_{l'-(m-1)}^{(m-1,m-1)} \rho^2 - z P_{l'-(m-1)}^{(m-1,m-1)'} \rho^2 \right) \\
& \quad \left. + P_{l-m}^{(m,m)'} P_{l'-(m-1)}^{(m-1,m-1)'} \rho^{2m+2} \right\} dz \\
&= \int_{-1}^1 \left\{ (m-1)(m+1) P_{l-m}^{(m,m)} P_{l'-(m-1)}^{(m-1,m-1)} \rho^{2m} - z P_{l-m}^{(m,m)} P_{l'-(m-1)}^{(m-1,m-1)'} \rho^{2m} \right. \\
& \quad \left. + P_{l-m}^{(m,m)'} P_{l'-(m-1)}^{(m-1,m-1)'} \rho^{2m+2} \right\} dz \\
&= \int_{-1}^1 \left\{ (m-1)(m+1) P_{l-m}^{(m,m)} P_{l'-(m-1)}^{(m-1,m-1)} \rho^{2m} - d_{l',m-1} z P_{l-m}^{(m,m)} P_{l'-m}^{(m,m)} \rho^{2m} \right. \\
& \quad \left. + d_{l,m} d_{l',m-1} P_{l-(m+1)}^{(m+1,m+1)} P_{l'-m}^{(m,m)} \rho^{2m+2} \right\} dz \\
&= \int_{-1}^1 \left\{ (m^2-1) P_{l-m}^{(m,m)} (\tilde{\alpha}_{l',m-1,2} P_{l'-1-m}^{(m,m)} + \tilde{\alpha}_{l',m-1,4} P_{l'+1-m}^{(m,m)}) \rho^{2m} \right. \\
& \quad - d_{l',m-1} P_{l-m}^{(m,m)} (\tilde{\gamma}_{l',m,1} P_{l'-1-m}^{(m,m)} + \tilde{\gamma}_{l',m,2} P_{l'+1-m}^{(m,m)}) \rho^{2m} \\
& \quad \left. + d_{l,m} d_{l',m-1} P_{l'-m}^{(m,m)} (\tilde{\alpha}_{l,m+1,1} P_{l-1-m}^{(m,m)} + \tilde{\alpha}_{l,m+1,3} P_{l+1-m}^{(m,m)}) \rho^{2m} \right\} dz \\
&= \delta_{l',l-1} \left( [(m^2-1) \tilde{\alpha}_{l-1,m-1,4} - d_{l-1,m-1} \tilde{\gamma}_{l-1,m,2}] \frac{1}{2\pi(c_l^m)^2} + d_{l,m} d_{l-1,m-1} \frac{\tilde{\alpha}_{l-1,m-1,1}}{2\pi(c_{l-1}^m)^2} \right) \\
& \quad + \delta_{l',l+1} \left( [(m^2-1) \tilde{\alpha}_{l+1,m-1,2} - d_{l+1,m-1} \tilde{\gamma}_{l+1,m,1}] \frac{1}{2\pi(c_l^m)^2} + d_{l,m} d_{l+1,m-1} \frac{\tilde{\alpha}_{l-1,m-1,3}}{2\pi(c_{l+1}^m)^2} \right).
\end{aligned} \tag{2.17}$$

As we can see, the coefficients are then also only possibly nonzero for  $l' = l \pm 1$ .

Combining the above with the equation (2.16), we have that our two coefficients

for  $\mathcal{A}_{l-1,m-1}, \mathcal{A}_{l+1,m-1}$  are those stated in the Lemma as  $A_{l,m,1}$  and  $A_{l,m,3}$ , for  $m > 0$ .

Next, assume  $m \geq 0$  and consider the case  $m' = m + 1$ . The integral in equation (2.16) is then:

$$\begin{aligned}
 & \int_{-1}^1 \left\{ m(m+1) P_{l-m}^{(m,m)} P_{l'-(m+1)}^{(m+1,m+1)} \rho^{2m} (1+z^2) - m z P_{l-m}^{(m,m)} P_{l'-(m+1)}^{(m+1,m+1)'} \rho^{2m+2} \right. \\
 & \quad \left. - (m+1) z P_{l-m}^{(m,m)'} P_{l'-(m+1)}^{(m+1,m+1)} \rho^{2m+2} + P_{l-m}^{(m,m)'} P_{l'-(m+1)}^{(m+1,m+1)'} \rho^{2m+4} \right\} dz \\
 &= \int_{-1}^1 \left\{ m(m+1) P_{l-m}^{(m,m)} P_{l'-(m+1)}^{(m+1,m+1)} \rho^{2m} (1+z^2) - (m+1) z P_{l-m}^{(m,m)'} P_{l'-(m+1)}^{(m+1,m+1)} \rho^{2m+2} \right. \\
 & \quad \left. + m P_{l'-(m+1)}^{(m+1,m+1)} \rho^{2m} (-2(m+1) z^2 P_{l-m}^{(m,m)} + P_{l-m}^{(m,m)} \rho^2 - z P_{l-m}^{(m,m)'} \rho^2) \right. \\
 & \quad \left. + P_{l-m}^{(m,m)'} P_{l'-(m+1)}^{(m+1,m+1)'} \rho^{2m+2} \right\} dz \\
 &= \int_{-1}^1 \left\{ m(m+2) P_{l-m}^{(m,m)} P_{l'-(m+1)}^{(m+1,m+1)} \rho^{2m+2} - z P_{l-m}^{(m,m)'} P_{l'-(m+1)}^{(m+1,m+1)} \rho^{2m+2} \right. \\
 & \quad \left. + P_{l-m}^{(m,m)'} P_{l'-(m+1)}^{(m+1,m+1)'} \rho^{2m+4} \right\} dz \\
 &= \int_{-1}^1 \left\{ m(m+2) P_{l-m}^{(m,m)} P_{l'-(m+1)}^{(m+1,m+1)} \rho^{2m+2} - d_{l,m} z P_{l-(m+1)}^{(m+1,m+1)} P_{l'-(m+1)}^{(m+1,m+1)} \rho^{2m+2} \right. \\
 & \quad \left. + d_{l,m} d_{l',m+1} P_{l-(m+1)}^{(m+1,m+1)} P_{l'-(m+2)}^{(m+2,m+2)} \rho^{2m+4} \right\} dz \\
 &= \int_{-1}^1 \left\{ m(m+2) P_{l'-(m+1)}^{(m+1,m+1)} (\tilde{\alpha}_{l,m,2} P_{l-1-(m+1)}^{(m+1,m+1)} + \tilde{\alpha}_{l,m,4} P_{l+1-(m+1)}^{(m+1,m+1)}) \rho^{2m+2} \right. \\
 & \quad \left. - d_{l,m} P_{l'-(m+1)}^{(m+1,m+1)} (\tilde{\gamma}_{l,m+1,1} P_{l-1-(m+1)}^{(m+1,m+1)} + \tilde{\gamma}_{l,m+1,2} P_{l+1-(m+1)}^{(m+1,m+1)}) \rho^{2m+2} \right. \\
 & \quad \left. + d_{l,m} d_{l',m+1} P_{l-(m+1)}^{(m+1,m+1)} (\tilde{\alpha}_{l',m+2,1} P_{l'-1-(m+1)}^{(m+1,m+1)} + \tilde{\alpha}_{l',m+2,3} P_{l'+1-(m+1)}^{(m+1,m+1)}) \rho^{2m+2} \right\} dz \\
 &= \delta_{l',l-1} \left( [m(m+2) \tilde{\alpha}_{l,m,2} - d_{l,m} \tilde{\gamma}_{l,m+1,1}] \frac{1}{2\pi (c_{l-1}^{m+1})^2} + d_{l,m} d_{l-1,m+1} \frac{\tilde{\alpha}_{l-1,m+2,3}}{2\pi (c_l^{m+1})^2} \right) \\
 & \quad + \delta_{l',l+1} \left( [m(m+2) \tilde{\alpha}_{l,m,4} - d_{l,m} \tilde{\gamma}_{l,m+1,2}] \frac{1}{2\pi (c_{l-1}^{m+1})^2} + d_{l,m} d_{l+1,m+1} \frac{\tilde{\alpha}_{l+1,m+2,1}}{2\pi (c_l^{m+1})^2} \right).
 \end{aligned} \tag{2.18}$$

Again, we can see that the coefficients are then also only possibly nonzero for  $l' = l \pm 1$ . Combining the above with the equation (2.16), we have that our two coefficients for  $\mathcal{A}_{l-1,m-1}, \mathcal{A}_{l+1,m-1}$  are those stated in the Lemma as  $A_{l,m,2}$  and  $A_{l,m,4}$ , for  $m \geq 0$ .

We still have to account for the remaining  $m = 0$  cases and  $m$  negative cases. Thankfully, as noted earlier, the case of  $m \leq 0, m' = m - 1$  is the same as replacing  $m$  with  $|m|$  in equation (2.18). Thus, our two coefficients for  $\mathcal{A}_{l-1,m-1}, \mathcal{A}_{l+1,m-1}$  are those stated in the Lemma as  $A_{l,m,1}$  and  $A_{l,m,3}$ , for  $m \leq 0$ . Similarly the case of  $m < 0, m' = m + 1$  is the same as replacing  $m$  with  $|m|$  in equation (2.17), and thus our two coefficients for  $\mathcal{A}_{l-1,m+1}, \mathcal{A}_{l+1,m+1}$  are those stated in the Lemma as  $A_{l,m,2}$  and  $A_{l,m,4}$  for  $m < 0$ .

The remaining coefficients we need to determine are  $\mathcal{A}_{l',m'}^\perp$ . To this end, we have

that

$$\begin{aligned}
& \langle x \Psi_l^m, \Phi_{l'}^{m'} \rangle \\
&= \iint_{\Omega} \cos \theta \sin \varphi \Psi_l^m \cdot \Phi_{l'}^{m'*} \sin \varphi \, d\phi \, d\theta \\
&= c_l^m c_{l'}^{m'} \left( \int_0^{2\pi} e^{im\theta} e^{im'\theta} \cos \theta \, d\theta \right) \\
&\quad \cdot \left( \int_{-1}^1 \left\{ im' P_{l'-|m'|}^{(|m'|, |m'|)}(z) \rho(z)^{|m|+|m'|-1} [ |m| z P_{l-|m|}^{(|m|, |m|)}(z) - \rho(z)^2 P_{l-|m|}^{(|m|, |m|)'}(z) \right. \right. \\
&\quad \left. \left. + im P_{l-|m|}^{(|m|, |m|)}(z) \rho(z)^{|m|+|m'|-1} [ |m'| z P_{l'-|m'|}^{(|m'|, |m'|)}(z) - \rho(z)^2 P_{l'-|m'|}^{(|m'|, |m'|)'}(z) ] \right\} dz \right) \\
&= c_l^m c_{l'}^{m'} \pi (\delta_{m', m-1} + \delta_{m', m+1}) \\
&\quad \cdot \left( \int_{-1}^1 \left\{ \operatorname{sgn} m 2i m m' z P_{l-|m|}^{(|m|, |m|)} P_{l'-|m'|}^{(|m'|, |m'|)} \rho^{|m|+|m'|-1} \right. \right. \\
&\quad \left. \left. - im P_{l-|m|}^{(|m|, |m|)'} P_{l'-|m'|}^{(|m'|, |m'|)} \rho^{|m|+|m'|+1} \right. \right. \\
&\quad \left. \left. - im' P_{l-|m|}^{(|m|, |m|)} P_{l'-|m'|}^{(|m'|, |m'|)'} \rho^{|m|+|m'|+1} \right\} dz \right), \tag{2.19}
\end{aligned}$$

since when the above is nonzero,  $m, m'$  will have the same sign (or one is zero) and so  $|m| m' + |m'| m = \operatorname{sgn} m m m'$ . Once again, we can conclude that  $\mathcal{A}_{l', m'}^\perp$  are zero for  $m' \notin \{m-1, m+1\}$ . It is left to evaluate the integral that is left at equation (2.19) when  $m'$  takes either  $m \pm 1$  value. First, consider the case that

$m > 0$  and  $m' = m - 1$ . The integral is then, using integration by parts:

$$\begin{aligned}
&= \int_{-1}^1 \left\{ 2i m(m-1) z P_{l-m}^{(m,m)} P_{l'-(m-1)}^{(m-1,m-1)} \rho^{2m-2} - im P_{l-m}^{(m,m)} P_{l'-(m-1)}^{(m-1,m-1)'} \rho^{2m} \right. \\
&\quad \left. - i(m-1) P_{l-m}^{(m,m)'} P_{l'-(m-1)}^{(m-1,m-1)} \rho^{2m} \right\} dz \\
&= \int_{-1}^1 \left\{ 2i m(m-1) z P_{l-m}^{(m,m)} P_{l'-(m-1)}^{(m-1,m-1)} \rho^{2m-2} - im P_{l-m}^{(m,m)} P_{l'-(m-1)}^{(m-1,m-1)'} \rho^{2m} \right. \\
&\quad \left. + i(m-1) P_{l-m}^{(m,m)} \rho^{2m-2} [P_{l'-(m-1)}^{(m-1,m-1)'} \rho^2 - 2mz P_{l'-(m-1)}^{(m-1,m-1)}] \right\} dz \\
&= - \int_{-1}^1 i P_{l-m}^{(m,m)} P_{l'-(m-1)}^{(m-1,m-1)'} \rho^{2m} dz \\
&= - \int_{-1}^1 i d_{l',m-1} P_{l-m}^{(m,m)} P_{l'-m}^{(m,m)} \rho^{2m} dz \\
&= -i \frac{d_{l,m-1}}{2\pi(c_l^m)^2} \delta_{l',l}. \tag{2.20}
\end{aligned}$$

Combining equation (2.20) with equation (2.19) means we retrieve the correct values for our coefficients  $\mathcal{A}_{l,m-1}^\perp$  when  $m > 0$ , i.e. they are  $A_{l,m,5}$  as stated in the Lemma for  $m > 0$ . The case of  $m < 0$  and  $m' = m + 1$  is very similar with the



integral becoming:

$$\begin{aligned}
&= \int_{-1}^1 \left\{ -2i m(m+1) z P_{l-|m|}^{(|m|,|m|)} P_{l'-(|m|-1)}^{(|m|-1,|m|-1)} \rho^{2|m|-2} - im P_{l-|m|}^{(|m|,|m|)} P_{l'-(|m|-1)}^{(|m|-1,|m|-1)} \rho^{2|m|} \right. \\
&\quad \left. - i(m+1) P_{l-|m|}^{(|m|,|m|)} P_{l'-(|m|-1)}^{(|m|-1,|m|-1)} \rho^{2|m|} \right\} dz \\
&= \int_{-1}^1 \left\{ -2i m(m+1) z P_{l-|m|}^{(|m|,|m|)} P_{l'-(|m|-1)}^{(|m|-1,|m|-1)} \rho^{2|m|-2} - im P_{l-|m|}^{(|m|,|m|)} P_{l'-(|m|-1)}^{(|m|-1,|m|-1)} \rho^{2|m|} \right. \\
&\quad \left. + i(m+1) P_{l-|m|}^{(|m|,|m|)} \rho^{2|m|-2} [P_{l'-(|m|-1)}^{(|m|-1,|m|-1)} \rho^2 - 2|m| z P_{l'-(|m|-1)}^{(|m|-1,|m|-1)}] \right\} dz \\
&= \int_{-1}^1 i P_{l-|m|}^{(|m|,|m|)} P_{l'-(|m|-1)}^{(|m|-1,|m|-1)} \rho^{2|m|} dz \\
&= \int_{-1}^1 i d_{l',|m|-1} P_{l-|m|}^{(|m|,|m|)} P_{l'-(|m|-1)}^{(|m|,|m|)} \rho^{2|m|} dz \\
&= i \frac{d_{l,|m|-1}}{2\pi(c_l^{|m|})^2} \delta_{l',l}. \tag{2.21}
\end{aligned}$$

Combining equation (2.21) with equation (2.19) means we retrieve the correct values for our coefficients  $\mathcal{A}_{l,m+1}^\perp$  when  $m < 0$ , i.e. they are  $A_{l,m,6}$  as stated in the Lemma for  $m < 0$ .

Next, assume  $m \geq 0$  and  $m' = m + 1$ . Then the integral in equation (2.19)

becomes:

$$\begin{aligned}
&= \int_{-1}^1 \left\{ 2i m(m+1) z P_{l-m}^{(m,m)} P_{l'-(m+1)}^{(m+1,m+1)} \rho^{2m} - im P_{l-m}^{(m,m)} P_{l'-(m+1)}^{(m+1,m+1)'} \rho^{2m+2} \right. \\
&\quad \left. - i(m+1) P_{l-m}^{(m,m)'} P_{l'-(m+1)}^{(m+1,m+1)} \rho^{2m+2} \right\} dz \\
&= \int_{-1}^1 \left\{ 2i m(m+1) z P_{l-m}^{(m,m)} P_{l'-(m+1)}^{(m+1,m+1)} \rho^{2m} - i(m+1) P_{l-m}^{(m,m)'} P_{l'-(m+1)}^{(m+1,m+1)} \rho^{2m+2} \right. \\
&\quad \left. + im P_{l'-(m+1)}^{(m+1,m+1)} \rho^{2m} [P_{l-m}^{(m,m)'} \rho^2 - 2(m+1) z P_{l-m}^{(m,m)}] \right\} dz \\
&= - \int_{-1}^1 i P_{l-m}^{(m,m)'} P_{l'-(m+1)}^{(m+1,m+1)} \rho^{2m+2} dz \\
&= - \int_{-1}^1 i d_{l,m} P_{l-(m+1)}^{(m+1,m+1)} P_{l'-(m+1)}^{(m+1,m+1)} \rho^{2m+2} dz \\
&= -i \frac{d_{l,m}}{2\pi(c_l^{m+1})^2} \delta_{l',l}. \tag{2.22}
\end{aligned}$$

Again, combining equation (2.22) with equation (2.19) means we retrieve the correct values for our coefficients  $\mathcal{A}_{l,m+1}^\perp$  when  $m \geq 0$ , i.e. they are  $A_{l,m,6}$  as stated in the Lemma for  $m \geq 0$ . In a similar vein, the integral for the case  $m \leq 0$

and  $m' = m - 1$  is:

$$\begin{aligned}
&= \int_{-1}^1 \left\{ -2i m(m-1) z P_{l-|m|}^{(|m|,|m|)} P_{l'-(|m|+1)}^{(|m|+1,|m|+1)} \rho^{2|m|} - im P_{l-|m|}^{(|m|,|m|)} P_{l'-(|m|+1)}^{(|m|+1,|m|+1)} \rho^{2|m|+2} \right. \\
&\quad \left. - i(m-1) P_{l-|m|}^{(|m|,|m|)} P_{l'-(|m|+1)}^{(|m|+1,|m|+1)} \rho^{2|m|+2} \right\} dz \\
&= \int_{-1}^1 \left\{ -2i m(m-1) z P_{l-|m|}^{(|m|,|m|)} P_{l'-(|m|+1)}^{(|m|+1,|m|+1)} \rho^{2|m|} - i(m-1) P_{l-|m|}^{(|m|,|m|)} P_{l'-(|m|+1)}^{(|m|+1,|m|+1)} \rho^{2|m|+2} \right. \\
&\quad \left. + im P_{l'-(|m|+1)}^{(|m|+1,|m|+1)} \rho^{2|m|} [P_{l-|m|}^{(|m|,|m|)} \rho^2 - 2(|m|+1)z P_{l-|m|}^{(|m|,|m|)}] \right\} dz \\
&= \int_{-1}^1 i P_{l-|m|}^{(|m|,|m|)} P_{l'-(|m|+1)}^{(|m|+1,|m|+1)} \rho^{2|m|+2} dz \\
&= \int_{-1}^1 i d_{l,|m|} P_{l-|m|}^{(|m|+1,|m|+1)} P_{l'-(|m|+1)}^{(|m|+1,|m|+1)} \rho^{2|m|+2} dz \\
&= i \frac{d_{l,|m|}}{2\pi(c_l^{|m|+1})^2} \delta_{l',l}, \tag{2.23}
\end{aligned}$$

and so combining equation (2.23) with equation (2.19), we have that our coefficients for  $\mathcal{A}_{l,m-1}^\perp$  are those stated in the Lemma as  $A_{l,m,5}$  for  $m \leq 0$ , concluding the argument for the expansion of  $x\Psi_l^m$ .

An almost identical argument holds for multiplication by  $y$ , since

$$\int_0^{2\pi} e^{im\theta} e^{im'\theta} \sin \theta d\theta = i\pi(\delta_{m',m-1} - \delta_{m',m+1})$$

replaces the integrals over  $\theta$  in equation (2.16) and equation (2.19).

For multiplication by  $z$ , we will use the same method to directly calculate the nonzero coefficients in each expansion using inner products. To this end, note

that

$$z\Psi_l^m = \sum_{l'=1}^{l+1} \sum_{m'=-l'}^{l'} \mathcal{C}_{l',m'} \Psi_{l'}^{m'} + \mathcal{C}_{l',m'}^\perp \Phi_{l'}^{m'}$$

where the coefficients are given by

$$\begin{aligned} \mathcal{C}_{l',m'} &= \frac{\langle z\Psi_l^m, \Psi_{l'}^{m'} \rangle}{\|\Psi_{l'}^{m'}\|^2} = \frac{1}{l'(l'+1)} \langle z\Psi_l^m, \Psi_{l'}^{m'} \rangle \\ \mathcal{C}_{l',m'}^\perp &= \frac{\langle z\Psi_l^m, \Phi_{l'}^{m'} \rangle}{\|\Phi_{l'}^{m'}\|^2} = \frac{1}{l'(l'+1)} \langle z\Psi_l^m, \Phi_{l'}^{m'} \rangle. \end{aligned}$$

The aim once again will be to show that the only nonzero coefficients  $\mathcal{C}_{l',m'}$ ,  $\mathcal{C}_{l',m'}^\perp$  match the  $\Gamma_{l,m,j}$  for  $j = 1, 2, 3$  in the Lemma. Now, using a change of variable and integration by parts, we have that

$$\begin{aligned} &\langle z\Psi_l^m, \Psi_{l'}^{m'} \rangle \\ &= \iint_{\Omega} \cos \varphi \Psi_l^m \cdot \Psi_{l'}^{m'*} \sin \varphi \, d\varphi \, d\theta \\ &= c_l^m c_{l'}^{m'} \left( \int_0^{2\pi} e^{im\theta} e^{im'\theta} \, d\theta \right) \\ &\quad \cdot \left( \int_{-1}^1 \left\{ [|m| z P_{l-|m|}^{(|m|,|m|)}(z) - \rho(z)^2 P_{l-|m|}^{(|m|,|m|)'}(z)] [|m'| z P_{l'-|m'|}^{(|m'|,|m'|)}(z) - \rho(z)^2 P_{l'-|m'|}^{(|m'|,|m'|)'}(z)] \right. \right. \\ &\quad \left. \left. + mm' P_{l-|m|}^{(|m|,|m|)}(z) P_{l'-|m'|}^{(|m'|,|m'|)}(z) \right\} z \rho(z)^{|m|+|m'|-2} \, dz \right) \end{aligned} \quad (2.24)$$

The integral over  $\theta$  in equation (2.24) is simply  $2\pi\delta_{m',m}$  showing the above is zero when  $m' \neq m$ , and thus so will the coefficients  $\mathcal{C}_{l',m'}$  be when this is the case.

Focusing on the second integral (over  $z$ ) when  $m' = m$ , this becomes:

$$\begin{aligned}
& \int_{-1}^1 \left\{ m^2 z P_{l-|m|}^{(|m|,|m|)} P_{l'-|m|}^{(|m|,|m|)} \rho^{2|m|-2} (1+z^2) \right. \\
& \quad - |m| z^2 P_{l-|m|}^{(|m|,|m|)} P_{l'-|m|}^{(|m|,|m|)} \rho^{2|m|} \\
& \quad - |m| z^2 P_{l-|m|}^{(|m|,|m|)} P_{l'-|m|}^{(|m|,|m|)} \rho^{2|m|} \\
& \quad \left. + z P_{l-|m|}^{(|m|,|m|)} P_{l'-|m|}^{(|m|,|m|)} \rho^{2|m|+2} \right\} dz \\
&= \int_{-1}^1 \left\{ m^2 z P_{l-|m|}^{(|m|,|m|)} P_{l'-|m|}^{(|m|,|m|)} \rho^{2|m|-2} (1+z^2) \right. \\
& \quad + |m| P_{l'-|m|}^{(|m|,|m|)} \rho^{2|m|-2} [z^2 P_{l-|m|}^{(|m|,|m|)} \rho^2 + 2z P_{l'-|m|}^{(|m|,|m|)} - 2|m| z^3 P_{l'-|m|}^{(|m|,|m|)} \rho^2] \\
& \quad - |m| z^2 P_{l-|m|}^{(|m|,|m|)} P_{l'-|m|}^{(|m|,|m|)} \rho^{2|m|} \\
& \quad \left. + z P_{l-|m|}^{(|m|,|m|)} P_{l'-|m|}^{(|m|,|m|)} \rho^{2|m|+2} \right\} dz \\
&= \int_{-1}^1 \left\{ |m| (|m| + 2) z P_{l-|m|}^{(|m|,|m|)} P_{l'-|m|}^{(|m|,|m|)} \rho^{2|m|} \right. \\
& \quad \left. + z d_{l,|m|} d_{l',|m|} P_{l-(|m|+1)}^{(|m|+1,|m|+1)} P_{l'-(|m|+1)}^{(|m|+1,|m|+1)} \rho^{2|m|+2} \right\} dz \\
&= \int_{-1}^1 \left\{ |m| (|m| + 2) P_{l'-|m|}^{(|m|,|m|)} \rho^{2|m|} [\tilde{\gamma}_{l,|m|,1} P_{l-1-|m|}^{(|m|,|m|)} + \tilde{\gamma}_{l,|m|,2} P_{l+1-|m|}^{(|m|,|m|)}] \right. \\
& \quad \left. + d_{l,|m|} d_{l',|m|} P_{l'-(|m|+1)}^{(|m|+1,|m|+1)} \rho^{2|m|+2} [\tilde{\gamma}_{l,|m|+1,1} P_{l-1-(|m|+1)}^{(|m|+1,|m|+1)} + \tilde{\gamma}_{l,|m|+1,2} P_{l+1-(|m|+1)}^{(|m|+1,|m|+1)}] \right\} dz \\
&= c_{l-1}^m \delta_{l',l-1} \left[ |m| (|m| + 2) \frac{\tilde{\gamma}_{l,|m|,1}}{2\pi (c_{l-1}^{|m|})^2} + d_{l,|m|} d_{l-1,|m|} \frac{\tilde{\gamma}_{l,|m|+1,1}}{2\pi (c_{l-1}^{|m|+1})^2} \right] \\
& \quad + c_{l+1}^m \delta_{l',l+1} \left[ |m| (|m| + 2) \frac{\tilde{\gamma}_{l,|m|,2}}{2\pi (c_{l+1}^{|m|})^2} + d_{l,|m|} d_{l+1,|m|} \frac{\tilde{\gamma}_{l,|m|+1,2}}{2\pi (c_{l+1}^{|m|+1})^2} \right] \\
&= c_{l-1}^m \delta_{l',l-1} \left[ |m| (|m| + 2) \frac{\tilde{\gamma}_{l,|m|,1}}{(c_{l-1}^{|m|})^2} + d_{l,|m|} d_{l-1,|m|} \frac{\tilde{\gamma}_{l,|m|+1,1}}{(c_{l-1}^{|m|+1})^2} \right] \\
& \quad + c_{l+1}^m \delta_{l',l+1} \left[ |m| (|m| + 2) \frac{\tilde{\gamma}_{l,|m|,2}}{(c_{l+1}^{|m|})^2} + d_{l,|m|} d_{l+1,|m|} \frac{\tilde{\gamma}_{l,|m|+1,2}}{(c_{l+1}^{|m|+1})^2} \right]. \tag{2.25}
\end{aligned}$$

Combining equation (2.25) with equation (2.24), we obtain that

$$\begin{aligned} \langle z \Psi_l^m, \Psi_{l'}^{m'} \rangle &= c_l^m \delta_{m',m} \left\{ c_{l-1}^m \delta_{l',l-1} \left[ |m| (|m| + 2) \frac{\tilde{\gamma}_{l,|m|,1}}{(c_{l-1}^{|m|})^2} + d_{l,|m|} d_{l-1,|m|} \frac{\tilde{\gamma}_{l,|m|+1,1}}{(c_{l-1}^{|m|+1})^2} \right] \right. \\ &\quad \left. + c_{l+1}^m \delta_{l',l+1} \left[ |m| (|m| + 2) \frac{\tilde{\gamma}_{l,|m|,2}}{(c_{l+1}^{|m|})^2} + d_{l,|m|} d_{l+1,|m|} \frac{\tilde{\gamma}_{l,|m|+1,2}}{(c_{l+1}^{|m|+1})^2} \right] \right\}. \end{aligned} \quad (2.26)$$

We can see here in equation (2.26) that the nonzero coefficients  $\mathcal{C}_{l',m'}$  are those described in the Lemma as  $\Gamma_{l,m,1}$  and  $\Gamma_{l,m,2}$ .

Next, we also have that

$$\begin{aligned} &\langle z \Psi_l^m, \Phi_{l'}^{m'} \rangle \\ &= \iint_{\Omega} \cos \varphi \Psi_l^m \cdot \Phi_{l'}^{m'*} \sin \varphi \, d\varphi \, d\theta \\ &= c_l^m c_{l'}^{m'} \left( \int_0^{2\pi} e^{im\theta} e^{im'\theta} \, d\theta \right) \\ &\quad \cdot \left( \int_{-1}^1 \left\{ im' z P_{l'-|m'|}^{(|m'|,|m'|)}(z) \rho(z)^{|m|+|m'|-2} \left[ |m| z P_{l-|m|}^{(|m|,|m|)}(z) - \rho(z)^2 P_{l-|m|}^{(|m|,|m|)'}(z) \right] \right. \right. \\ &\quad \left. \left. + im z P_{l-|m|}^{(|m|,|m|)}(z) \rho(z)^{|m|+|m'|-2} \left[ |m'| z P_{l'-|m'|}^{(|m'|,|m'|)}(z) - \rho(z)^2 P_{l'-|m'|}^{(|m'|,|m'|)'}(z) \right] \right\} \, dz \right) \\ &= c_l^m c_{l'}^{m'} 2\pi \delta_{m',m} \int_{-1}^1 \left\{ 2im |m| z^2 P_{l-|m|}^{(|m|,|m|)}(z) P_{l'-|m|}^{(|m|,|m|)}(z) \rho(z)^{2|m|-2} \right. \\ &\quad - im z P_{l-|m|}^{(|m|,|m|)}(z) P_{l'-|m|}^{(|m|,|m|)'}(z) \rho(z)^{2|m|} \\ &\quad \left. - im z P_{l-|m|}^{(|m|,|m|)'}(z) P_{l'-|m|}^{(|m|,|m|)}(z) \rho(z)^{2|m|} \right\} \, dz \\ &= c_l^m c_{l'}^{m'} 2\pi \delta_{m',m} \int_{-1}^1 im P_{l-|m|}^{(|m|,|m|)}(z) P_{l'-|m|}^{(|m|,|m|)}(z) \rho(z)^{2|m|} \, dz \\ &= \delta_{l',l} \delta_{m',m} im. \end{aligned} \quad (2.27)$$

Again, we can see here in equation (2.27) that the coefficients  $\mathcal{C}_{l',m'}^\perp$  are zero when  $l' \neq l$  or  $m' \neq m$ , and that the nonzero coefficients are those described in the Lemma as  $\Gamma_{l,m,3}$ .

Finally, to complete our proof, for multiplication by  $x, y, z$  of the other VSHs  $\Phi_l^m$ , simply take the cross product by  $\hat{\mathbf{r}}$  of  $x\Psi_l^m$ , etc. noting that  $\hat{\mathbf{r}} \times \Psi_l^m = \Phi_l^m$  and  $\hat{\mathbf{r}} \times \Phi_l^m = -\|\hat{\mathbf{r}}\| \Psi_l^m = -\Psi_l^m$  by the triple vector product and orthogonality.  $\square$


As for the scalar case, these coefficients detailed above are then the nonzero entries of the Jacobi matrices for the VSHs. Define  $\tilde{\mathbb{T}}$  by

$$\tilde{\mathbb{T}} = \begin{pmatrix} \tilde{\mathbb{T}}_0 \\ \tilde{\mathbb{T}}_1 \\ \vdots \end{pmatrix}, \quad \text{where} \quad \tilde{\mathbb{T}}_l = \begin{pmatrix} (\Psi_l^{-l})^\top \\ (\Phi_l^{-l})^\top \\ \vdots \\ (\Psi_l^l)^\top \\ (\Phi_l^l)^\top \end{pmatrix} \in \mathbb{C}^{2(2l+1) \times 3} \quad \forall l \in \mathbb{N}_0. \quad (2.28)$$

The Jacobi matrices are then defined according to the matrices  ${}^TJ_x, {}^TJ_y, {}^TJ_z$  satisfying

$$\begin{aligned} {}^TJ_x \tilde{\mathbb{T}}(x, y, z) &= x \tilde{\mathbb{T}}(x, y, z) \\ {}^TJ_y \tilde{\mathbb{T}}(x, y, z) &= y \tilde{\mathbb{T}}(x, y, z) \\ {}^TJ_z \tilde{\mathbb{T}}(x, y, z) &= z \tilde{\mathbb{T}}(x, y, z) \end{aligned} \quad (2.29)$$

for each  $(x, y, z) \in \Omega$ . The VSH Jacobi matrices are banded-block-banded due



**Figure 2.2:** “Spy” plots of the Jacobi operator matrices for the Vector Spherical Harmonic OP basis, showing their sparsity and structure, up to order  $N = 10$ .



to the sparse relationships we obtain from Lemma 3. Specifically, they each have block-bandwidths  $(1, 1)$ :

$${}^T J_{x/y/z} = \begin{pmatrix} B_{x/y/z,0} & A_{x/y/z,0} & & & \\ C_{x/y/z,1} & B_{x/y/z,1} & A_{x/y/z,1} & & \\ & C_{x/y/z,2} & B_{x/y/z,2} & A_{x/y/z,2} & \\ & & C_{x/y/z,3} & \ddots & \ddots \\ & & & \ddots & \ddots & \ddots \end{pmatrix}.$$

Let  $0_2 \in \mathbb{R}^{2 \times 2}$  be the 2x2 zero matrix.  ${}^T J_x$  has subblock-bandwidths  $(4, 4)$  since

for  $l \in \mathbb{N}_0$ :

$$\begin{aligned}
 A_{x,l} &:= \begin{pmatrix} \mathcal{A}_{l,-l,3} & 0_2 & \mathcal{A}_{l,-l,4} & & \\ & \ddots & \ddots & \ddots & \\ & & \mathcal{A}_{l,l,3} & 0_2 & \mathcal{A}_{l,l,4} \end{pmatrix} \in \mathbb{R}^{2(2l+1) \times 2(2l+3)}, \\
 B_{x,l} &:= \begin{pmatrix} & 0_2 & & \mathcal{A}_{l,-l,6} & & \\ \mathcal{A}_{l,-l+1,5} & & \ddots & \ddots & & \\ & & \ddots & \ddots & \mathcal{A}_{l,l-1,6} & \\ & & & \mathcal{A}_{l,l,5} & 0_2 & \end{pmatrix} \in \mathbb{R}^{2(2l+1) \times 2(2l+1)}, \\
 C_{x,l} &:= \begin{pmatrix} \mathcal{A}_{l,-l,2} & & & & & \\ & 0_2 & & \ddots & & \\ \mathcal{A}_{l,-l+2,1} & & \ddots & \ddots & & \\ & & \ddots & \ddots & \mathcal{A}_{l,l-2,2} & \\ & & & \ddots & 0_2 & \\ & & & & \mathcal{A}_{l,l,1} & \end{pmatrix} \in \mathbb{R}^{2(2l+1) \times 2(2l-1)} \quad (l \neq 0),
 \end{aligned}$$

where

$$\mathcal{A}_{l,m,j} := \begin{pmatrix} A_{l,m,j} & 0 \\ 0 & A_{l,m,j} \end{pmatrix} \text{ for } j = 1, 2, 3, 4, \quad \mathcal{A}_{l,m,j} := \begin{pmatrix} 0 & A_{l,m,j} \\ -A_{l,m,j} & 0 \end{pmatrix} \text{ for } j = 5, 6.$$

Similarly,  ${}^T J_y$  also has subblock-bandwidths  $(4, 4)$  since for  $l \in \mathbb{N}_0$ :

$$\begin{aligned}
 A_{y,l} &:= \begin{pmatrix} \mathcal{B}_{l,-l,3} & 0_2 & \mathcal{B}_{l,-l,4} & & \\ & \ddots & \ddots & \ddots & \\ & & \mathcal{B}_{l,l,3} & 0_2 & \mathcal{B}_{l,l,4} \end{pmatrix} \in \mathbb{R}^{2(2l+1) \times 2(2l+3)}, \\
 B_{y,l} &:= \begin{pmatrix} & 0_2 & \mathcal{B}_{l,-l,6} & & \\ \mathcal{B}_{l,-l+1,5} & \ddots & \ddots & & \\ & \ddots & \ddots & \mathcal{B}_{l,l-1,6} & \\ & & \mathcal{B}_{l,l,5} & 0_2 & \end{pmatrix} \in \mathbb{R}^{2(2l+1) \times 2(2l+1)}, \\
 C_{y,l} &:= \begin{pmatrix} \mathcal{B}_{l,-l,2} & & & & \\ & 0_2 & \ddots & & \\ \mathcal{B}_{l,-l+2,1} & \ddots & \ddots & & \\ & \ddots & \ddots & \mathcal{B}_{l,l-2,2} & \\ & & \ddots & 0_2 & \\ & & & \mathcal{B}_{l,l,1} & \end{pmatrix} \in \mathbb{R}^{2(2l+1) \times 2(2l-1)} \quad (l \neq 0),
 \end{aligned}$$

where

$$\mathcal{B}_{l,m,j} := \begin{pmatrix} B_{l,m,j} & 0 \\ 0 & B_{l,m,j} \end{pmatrix} \text{ for } j = 1, 2, 3, 4, \quad \mathcal{B}_{l,m,j} := \begin{pmatrix} 0 & B_{l,m,j} \\ -B_{l,m,j} & 0 \end{pmatrix} \text{ for } j = 5, 6.$$

Finally,  ${}^tJ_z$  has subblock-bandwidths  $(2, 2)$ , as for  $l \in \mathbb{N}_0$ :

$$\begin{aligned}
 A_l^z &:= \begin{pmatrix} 0 & 0 & \Gamma_{l,-l,2} & & & \\ & & \Gamma_{l,-l,2} & & & \\ & & & \ddots & & \\ & & & & \Gamma_{l,l,2} & \\ & & & & & \Gamma_{l,l,2} & 0 & 0 \end{pmatrix} \in \mathbb{R}^{2(2l+1) \times 2(2l+3)}, \\
 B_l^z &:= \begin{pmatrix} 0 & \Gamma_{l,-l,3} & & & \\ -\Gamma_{l,-l,3} & 0 & & & \\ & & \ddots & & \\ & & & 0 & \Gamma_{l,l,3} \\ & & & -\Gamma_{l,l,3} & 0 \end{pmatrix} \in \mathbb{R}^{2(2l+1) \times 2(2l+1)}, \\
 C_l^z &:= \begin{pmatrix} 0 & & & & & \\ 0 & & & & & \\ \Gamma_{l,-l+1,1} & & & & & \\ & \Gamma_{l,-l+1,1} & & & & \\ & & \ddots & & & \\ & & & \Gamma_{l,l-1,1} & & \\ & & & & \Gamma_{l,l-1,1} & \\ & & & & & 0 \\ & & & & & & 0 \end{pmatrix} \in \mathbb{R}^{2(2l+1) \times 2(2l-1)} \quad (l \neq 0).
 \end{aligned}$$

The sparsity and structure of the Jacobi matrices for the VSHs as we have defined them can be seen visually in Figure ??.

### 2.5.2 Three-term recurrence relation for $\tilde{\mathbb{T}}(x, y, z)$

We can combine each system in equation (2.29) into a block-tridiagonal system for any  $(x, y, z) \in \Omega$ :

$$\begin{pmatrix} I_6 & & & & & \\ B_1 - G_1(x, y, z) & A_1 & & & & \\ & C_2 & B_2 - G_2(x, y, z) & A_3 & & \\ & & C_3 & B_3 - G_3(x, y, z) & \ddots & \\ & & & \ddots & \ddots & \\ & & & & \ddots & \ddots \end{pmatrix} \tilde{\mathbb{T}}(x, y, z) = \begin{pmatrix} \tilde{\mathbb{T}}_1(x, y, z) \\ \mathbf{0} \\ \mathbf{0} \\ \vdots \end{pmatrix},$$

where we note that

$$\tilde{\mathbb{T}}_1(x, y, z) := \begin{pmatrix} \Psi_1^{-1}(x, y, z)^\top \\ \Phi_1^{-1}(x, y, z)^\top \\ \Psi_1^0(x, y, z)^\top \\ \Phi_1^0(x, y, z)^\top \\ \Psi_1^1(x, y, z)^\top \\ \Phi_1^1(x, y, z)^\top \end{pmatrix},$$

can be calculated explicitly, and for each  $l \in \mathbb{N}$ ,

$$A_l := \begin{pmatrix} A_{x,l} \\ A_{y,l} \\ A_{z,l} \end{pmatrix} \in \mathbb{C}^{6(2l+1) \times 2(2l+3)}, \quad C_l := \begin{pmatrix} C_{x,l} \\ C_{y,l} \\ C_{z,l} \end{pmatrix} \in \mathbb{C}^{6(2l+1) \times 2(2l-1)} \quad (n \neq 0), \quad (2.30)$$

$$B_l := \begin{pmatrix} B_{x,l} \\ B_{y,l} \\ B_{z,l} \end{pmatrix} \in \mathbb{C}^{6(2l+1) \times 2(2l+1)}, \quad G_n(x, y) := \begin{pmatrix} xI_{2l+1} \\ yI_{2l+1} \\ zI_{2l+1} \end{pmatrix} \in \mathbb{C}^{6(2l+1) \times 2(2l+1)}. \quad (2.31)$$

Just like for the scalar case, for each  $l = 0, 1, 2 \dots$  let  $D_l^\top$  be any matrix that is a left inverse of  $A_l$ , i.e. such that  $D_l^\top A_l = I_{2(2l+3)}$ . Multiplying our system by the preconditioner matrix that is given by the block diagonal matrix of the  $D_l^\top$ 's, we obtain a lower triangular system [17, p78], which can be expanded to obtain the recurrence:

$$\begin{cases} \tilde{T}_0(x, y, z) := 0_{2,3} \\ \tilde{T}_{l+1}(x, y, z) = -D_l^\top (B_l - G_l(x, y, z)) \tilde{T}_l(x, y, z) - D_l^\top C_l \tilde{T}_{l-1}(x, y, z), \quad l \in \mathbb{N}. \end{cases}$$

For  $l \in \mathbb{N}$ , we can set

$$D_l^\top = \begin{pmatrix} \boldsymbol{\eta}_0^\top \\ \boldsymbol{\eta}_1^\top \\ 0_{2(2l+1)} & 0_{2(2l+1)} & \mathcal{D}_l \\ \boldsymbol{\eta}_2^\top \\ \boldsymbol{\eta}_3^\top \end{pmatrix} \in \mathbb{R}^{2(2l+3) \times 6(2l+1)}, \quad (2.32)$$

where  $\mathcal{D}_l \in \mathbb{R}^{2(2l+1) \times 2(2l+1)}$  is the matrix

$$\mathcal{D}_l := \begin{pmatrix} \frac{1}{\Gamma_{l,-l,2}} & & & & \\ & \frac{1}{\Gamma_{l,-l,2}} & & & \\ & & \ddots & & \\ & & & \frac{1}{\Gamma_{l,l,2}} & \\ & & & & \frac{1}{\Gamma_{l,l,2}} \end{pmatrix}$$

and where  $\boldsymbol{\eta}_0, \boldsymbol{\eta}_1, \boldsymbol{\eta}_2, \boldsymbol{\eta}_3 \in \mathbb{R}^{6(2l+1)}$  are vectors with entries given by

$$\begin{aligned}
 (\boldsymbol{\eta}_0)_j &= \begin{cases} \frac{1}{A_{l,-l,3}} & j = 1 \\ \frac{-A_{l,-l,4}}{A_{l,-l,3} \Gamma_{l,1-l,2}} & j = 4(2l+1) + 3, \\ 0 & o/w \end{cases} \\
 (\boldsymbol{\eta}_1)_j &= \begin{cases} (\boldsymbol{\eta}_0)_{j-1} & j = 2, 4(2l+1) + 4 \\ 0 & o/w \end{cases}, \\
 (\boldsymbol{\eta}_2)_j &= \begin{cases} \frac{1}{A_{l,l,4}} & j = 2(2l+1) - 1 \\ \frac{-A_{l,l,3}}{A_{l,l,4} \Gamma_{l,l-1,2}} & j = 6(2l+1) - 3, \\ 0 & o/w \end{cases} \\
 (\boldsymbol{\eta}_3)_j &= \begin{cases} (\boldsymbol{\eta}_2)_{j-1} & j = 2(2l+1), 6(2l+1) - 2 \\ 0 & o/w \end{cases}.
 \end{aligned}$$

### 2.5.3 Computational aspects

It will be useful to be able to describe the coefficients of a vector-valued function in the tangent bundle of  $\Omega$  in multiple ways, and as such we will introduce some more notation for the “splitting” of the basis vectors into the two groups stemming from the gradient and perpendicular gradient directions. Let us explain what we mean by this.



Define  $\mathbb{T}^\Psi, \mathbb{T}^\Phi$  by

$$\mathbb{T}^\Psi := \begin{pmatrix} \mathbb{T}_0^\Psi \\ \mathbb{T}_1^\Psi \\ \vdots \end{pmatrix}, \quad \text{where} \quad \mathbb{T}_l^\Psi := \begin{pmatrix} (\Psi_l^{-l})^\top \\ \vdots \\ (\Psi_l^l)^\top \end{pmatrix} \in \mathbb{C}^{(2l+1) \times 3} \quad \forall l \in \mathbb{N}_0, \quad (2.33)$$

$$\mathbb{T}^\Phi := \begin{pmatrix} \mathbb{T}_0^\Phi \\ \mathbb{T}_1^\Phi \\ \vdots \end{pmatrix}, \quad \text{where} \quad \mathbb{T}_l^\Phi := \begin{pmatrix} (\Phi_l^{-l})^\top \\ \vdots \\ (\Phi_l^l)^\top \end{pmatrix} \in \mathbb{C}^{(2l+1) \times 3} \quad \forall l \in \mathbb{N}_0. \quad (2.34)$$

If  $\mathbf{u}$  is a vector-valued function on the tangent bundle of  $\Omega$ , it will have the coefficients vectors  $\mathbf{u}^c, \mathbf{u}^{\Psi^c}, \mathbf{u}^{\Phi^c}$  for its expansion in the VSH basis up to order  $N \in \mathbb{N}$ , defined such that

$$\begin{aligned} \mathbf{u}(x, y, z) &\approx \tilde{\mathbb{T}}(x, y, z)^\top \mathbf{u}^c \\ &\equiv \mathbb{T}^\Psi(x, y, z)^\top \mathbf{u}^{\Psi^c} + \mathbb{T}^\Phi(x, y, z)^\top \mathbf{u}^{\Phi^c} \\ &:= \sum_{l=0}^N \sum_{m=-l}^l [u_{l,m}^\Psi \Psi_l^m(x, y, z) + u_{l,m}^\Phi \Phi_l^m(x, y, z)]. \end{aligned}$$

Of course, just like for the scalar case, we can use transforms to compute the coefficients of such a known function, and further evaluate, and apply differential operators to, a function knowing just its coefficients.

## Obtaining coefficients

As eluded to, we would like to be able to obtain the coefficients for the expansion of a vector valued function in the VSH basis up to a given degree, similar to how we approach the scalar case. To do this, we can use a vector spherical harmonic transform. The development of vector spherical harmonic transforms is well established [61], and continues today (see [21]).

## Function evaluation

The Clenshaw algorithm presented in Section 2.4.2 can also be used to evaluate a vector-valued function in the tangent bundle of  $\Omega$ , where the matrices  $C_n, B_n, D_n^\top, G_n(x, y, z)$  are all defined in Section 2.5.2:

1) Set  $\boldsymbol{\xi}_{N+2} = \mathbf{0}, \boldsymbol{\xi}_{N+1} = \mathbf{0}$ .

2) For  $n = N : -1 : 1$

$$\text{set } \boldsymbol{\xi}_n^\top = (\mathbf{u}_n^c)^\top - \boldsymbol{\xi}_{n+1}^\top D_n^\top (B_n - G_n(x, y, z)) - \boldsymbol{\xi}_{n+2}^\top D_{n+1}^\top C_{n+1}$$

3) Output:  $\mathbf{u}(x, y, z) \approx \tilde{\mathbb{T}}_1(x, y, z)^\top \boldsymbol{\xi}_1$

$$\begin{aligned} &= (\boldsymbol{\xi}_1)_1 \boldsymbol{\Psi}_1^{-1} + (\boldsymbol{\xi}_1)_2 \boldsymbol{\Phi}_1^{-1} + (\boldsymbol{\xi}_1)_3 \boldsymbol{\Psi}_1^0 \\ &\quad + (\boldsymbol{\xi}_1)_4 \boldsymbol{\Phi}_1^0 + (\boldsymbol{\xi}_1)_5 \boldsymbol{\Psi}_1^1 + (\boldsymbol{\xi}_1)_6 \boldsymbol{\Phi}_1^1. \end{aligned}$$

Here, at each iteration, the vector  $\mathbf{u}_n^c$  is given by the merging of the two order  $n$  subvector of  $\mathbf{u}^c$ :

$$\mathbf{u}_n^c := \begin{pmatrix} u_{n,-n}^\Psi \\ u_{n,-n}^\Phi \\ \vdots \\ u_{n,n}^\Psi \\ u_{n,n}^\Phi \end{pmatrix}.$$

Notice how the iteration also ends at  $n = 1$  (not  $n = 0$ ) here, since the VSHs (and hence the coefficients) for order  $n = 0$  are zero.

#### 2.5.4 Sparse partial differential operators

The framework we have outlined for the spherical harmonics and the vector spherical harmonics allows us to easily and explicitly derive sparse operator matrices for partial differential operators such as the spherical gradient, divergence and Laplacian, as well as other key operators useful in examples such as the linearised shallow water equations. These operators will take coefficients of a function's expansion in either the SSH or VSH basis, to coefficients in either basis.

Let  $\mathbf{u}$  be a vector-valued function on the tangent bundle of  $\Omega$  with coefficients vectors  $\mathbf{u}^{\Psi^c}, \mathbf{u}^{\Phi^c}$  for its expansion in the VSH basis up to order  $N \in \mathbb{N}$ , and let  $f$  be a scalar function on  $\Omega$  with coefficients vector  $\mathbf{f}^c$  for its expansion in the SSH

basis also up to order  $N$ , i.e.

$$\begin{aligned}
\mathbf{u}(x, y, z) &\approx \tilde{\mathbb{T}}(x, y, z)^\top \mathbf{u}^c \\
&\equiv \mathbb{T}^\Psi(x, y, z)^\top \mathbf{u}^{\Psi^c} + \mathbb{T}^\Phi(x, y, z)^\top \mathbf{u}^{\Phi^c} \\
&:= \sum_{l=0}^N \sum_{m=-l}^l \left[ u_{l,m}^\Psi \Psi_l^m(x, y, z) + u_{l,m}^\Phi \Phi_l^m(x, y, z) \right], \\
f(x, y, z) &\approx \mathbb{P}(x, y, z)^\top \mathbf{f}^c := \sum_{l=0}^N \sum_{m=-l}^l f_{l,m} Y_l^m(x, y, z).
\end{aligned}$$

In order to fully exploit the sparsity of the relationships and ensure we are using banded-block-banded operators, in problems where we have an interdependency of scalar and vector-valued functions it can be useful to “expand” the scalar function’s coefficients vector  $\mathbf{f}^c$  to be the length of the vector-valued one  $\mathbf{u}^c$ , with zeros being inserted after each entry, i.e. define the *expanded coefficients for a scalar function on  $\Omega$  in the SH basis* by

$$\tilde{\mathbf{f}}^c := \begin{pmatrix} \tilde{\mathbf{f}}^c_0 \\ \vdots \\ \tilde{\mathbf{f}}^c_N \end{pmatrix} \in \mathbb{R}^{2(N+1)^2}, \quad \tilde{\mathbf{f}}^c_l := \begin{pmatrix} f_{l,-l} \\ 0 \\ f_{l,-l+1} \\ 0 \\ \vdots \\ f_{l,l} \\ 0 \end{pmatrix} \in \mathbb{R}^{4(l+1)}.$$

To help with the definitions of the operators we will discuss, let  $\mathcal{E}_s, \mathcal{E}_v$  be (severely)

rectangular matrices defined such that  $\mathcal{E}_s \tilde{\mathbf{f}}^c = \mathbf{f}^c$  and  $\mathcal{E}_v \mathbf{f}^c = \tilde{\mathbf{f}}^c$ . We are now in a position to derive the sparse differential operators that can be applied to the functions  $\mathbf{u}, f$ .

**Definition 4.** We define the operator matrices  $\mathcal{D}, \tilde{\mathcal{D}}, \mathcal{G}, \tilde{\mathcal{G}}, \mathcal{L}$  according to:

$$\begin{aligned}\nabla \cdot \mathbf{u}(x, y, z) &= \mathbb{P}(x, y, z)^\top \mathcal{D} \mathbf{u}^{\Psi^c}, \\ \nabla \cdot \mathbf{u}(x, y, z) &= \mathbb{P}(x, y, z)^\top \mathcal{E}_s (\tilde{\mathcal{D}} \mathbf{u}^c), \\ \nabla_S f(x, y, z) &= \mathbb{T}^\Psi(x, y, z)^\top \mathcal{G} \mathbf{f}^c, \\ \nabla_S f(x, y, z) &= \tilde{\mathbb{T}}(x, y, z)^\top \tilde{\mathcal{G}} \tilde{\mathbf{f}}^c, \\ \Delta_S f(x, y, z) &= \mathbb{P}(x, y, z)^\top \mathcal{L} \mathbf{f}^c.\end{aligned}$$

**Remark:** The operators  $\tilde{\mathcal{D}}, \tilde{\mathcal{G}}$  would be used when vectors of the resulting length are required for the problem you are wishing to solve.

**Theorem 1.** The operator matrices defined in Definition 4 are diagonal, and are

given by:

$$\mathcal{D} \equiv \mathcal{L} := \begin{pmatrix} \mathcal{L}_0 & & \\ & \ddots & \\ & & \mathcal{L}_N \end{pmatrix} \in \mathbb{R}^{(N+1)^2 \times (N+1)^2}, \quad \tilde{\mathcal{D}} = \begin{pmatrix} \tilde{\mathcal{D}}_0 & & \\ & \ddots & \\ & & \tilde{\mathcal{D}}_N \end{pmatrix} \in \mathbb{R}^{2(N+1)^2 \times 2(N+1)^2},$$

$$\mathcal{G} = I_{(N+1)^2}, \quad \tilde{\mathcal{G}} = \begin{pmatrix} 1 & & & \\ & 0 & & \\ & & \ddots & \\ & & & 1 \\ & & & & 0 \end{pmatrix} \in \mathbb{R}^{2(N+1)^2 \times 2(N+1)^2},$$

where

$$\mathcal{L}_l := -l(l+1) I_{2l+1}, \quad \tilde{\mathcal{D}}_l := \begin{pmatrix} -l(l+1) & & & \\ & 0 & & \\ & & \ddots & \\ & & & -l(l+1) \\ & & & & 0 \end{pmatrix} \in \mathbb{R}^{2(2l+1) \times 2(2l+1)}.$$

*Proof.* The arguments for  $\mathcal{G}$  and  $\tilde{\mathcal{G}}$  are trivial by definition. The spherical harmonics, as the name eludes to, satisfy a harmonic relationship  $\Delta_{\mathbb{S}} Y_l^m = l(l+1) Y_l^m$

(see e.g. [26]). Thus, for the proof for  $\mathcal{D}$  and  $\tilde{\mathcal{D}}$ , consider


$$\begin{aligned}\nabla \cdot \mathbf{u} &= \nabla \cdot \left( \sum_{l=0}^N \sum_{m=-l}^l [u_{l,m}^{\Psi} \Psi_l^m + u_{l,m}^{\Phi} \Phi_l^m] \right) \\ &= \sum_{l=0}^N \sum_{m=-l}^l u_{l,m}^{\Psi} \Delta_S Y_l^m \\ &= \sum_{l=0}^N \sum_{m=-l}^l -u_{l,m}^{\Psi} l(l+1) Y_l^m,\end{aligned}$$

using the fact that  $\nabla \cdot (\nabla^\perp f) \equiv 0$  for any function  $f$ . We similarly have for  $\mathcal{L}$ :

$$\begin{aligned}\Delta_S f &= \Delta_S \left( \sum_{l=0}^N \sum_{m=-l}^l f_{l,m} Y_l^m \right) \\ &= \sum_{l=0}^N \sum_{m=-l}^l f_{l,m} \Delta_S Y_l^m \\ &= \sum_{l=0}^N \sum_{m=-l}^l -f_{l,m} l(l+1) Y_l^m.\end{aligned}$$

□

Now that we have our differential operators, let's apply them in some examples.



**Figure 2.3:** Four snapshots of the solution  $u$  of the heat equation with  $\alpha = 1/42$  and a step size of  $\Delta t = 0.01$ , solved using the BDF2 timestepping method. Top Left: Initial conditions (0 timesteps). Top Right: After 25 timesteps. Bottom Left: After 50 timesteps. Bottom Right: After 75 timesteps.



## 2.6 Examples

### 2.6.1 Heat equation

To demonstrate proof of concept, we can apply the framework we have proposed to modelling the heat equation on the unit sphere. In doing so, we can demonstrate how our method allows us to expand and evaluate functions, and turn the differential equation into a matrix-vector problem at each timestep.

Let  $u : \Omega \rightarrow \mathbb{C}$  represent the temperature at a point  $(x, y, z)$  on the sphere and suppose that it is modelled by the equation

$$\frac{\partial}{\partial t} u = \alpha \Delta_S u \quad (2.35)$$

for some  $\alpha \in \mathbb{R}$ . A simple timestepping method we can use to iterate with is the BDF2 method [27, 2.3]. Let  $u_n$  for  $n = 0, 1, 2, \dots$  be the solution at time  $n\Delta t$  where  $\Delta t$  is the timestep. The are timestepping method is

$$\begin{aligned} \frac{u_1 - u_0}{\Delta t} &= \alpha \Delta_S u_1 \\ \frac{3u_{n+1} - 4u_n + u_{n-1}}{2\Delta t} &= \alpha \Delta_S u_{n+1}, \quad n = 1, 2, \dots \end{aligned}$$

$\Longleftrightarrow$

$$\begin{aligned} [1 - \Delta t \alpha \Delta_S] u_1 &= u_0 \\ [3 - 2\Delta t \alpha \Delta_S] u_{n+1} &= 4u_n - u_{n-1}, \quad n = 1, 2, \dots \end{aligned}$$

Let us write this in coefficient space. Firstly, our coefficients vectors for the function  $u$  we will denote  $\mathbf{u}^c$  – that is:

$$u(x, y, z) \approx \mathbb{P}(x, y, z)^\top \mathbf{u}^c.$$

Our system then becomes

$$\begin{aligned} [I - \Delta t \alpha \mathcal{L}] \mathbf{u}_1^c &= u_0 \\ [3I - 2\Delta t \alpha \mathcal{L}] \mathbf{u}_{n+1}^c &= 4\mathbf{u}_n^c - \mathbf{u}_{n-1}^c, \quad n = 1, 2, \dots \end{aligned}$$

From here, it is clear that at each timestep we simply have a fairly trivial linear system to solve, with the matrices  $I - \Delta t \alpha \mathcal{L}$ ,  $3I - 2\Delta t \alpha \mathcal{L}$  being banded-block-banded (in fact, simply diagonal here). We demonstrate this theory in Figure 2.3, where we see four snapshots of the solution to the heat equation in equation (2.35) on the unit sphere with our order  $N$  set large enough (here,  $N = 20$ ). We set  $\alpha = \frac{1}{42}$  and a step size of  $\Delta t = 0.01$ , and take the initial condition for the heat  $u$  to be a “football function”<sup>2</sup> – that is we take

$$u_0 := Y_6^0 + \sqrt{\frac{14}{11}} Y_6^6.$$

The snapshots of the solution  $u$  in `bsreffigfig:heateqn` are taken of the initial conditions, plus after 25, 50 and 75 timesteps respectively.

---

<sup>2</sup>These initial conditions are taken from an example found at <https://www.chebfun.org/examples/sphere/SphereHeatConduction.html>

### 2.6.2 Linearised shallow water equations

The linearised shallow water equations allow us to showcase more of the differential operators that we defined in Definition 4 (namely the divergence and gradient operators) and demonstrate how the natural sparsity that our framework brings to the problem leads to a simple sparse linear system to solve.

Let  $\mathbf{u}(x, y, z)$  be the tangential velocity of a flow and  $h(x, y, z)$  be the height deviation of the flow from some constant reference height  $\mathcal{H}$ . Define  $\hat{\mathbf{r}}$  as the unit outward normal vector at the point on the sphere  $(x, y, z)$ , so that

$$\hat{\mathbf{r}} = \begin{pmatrix} x \\ y \\ z \end{pmatrix}.$$

The linear SWEs are


$$\begin{cases} \frac{\partial \mathbf{u}}{\partial t} + f \hat{\mathbf{r}} \times \mathbf{u} - \nabla_S h = \mathbf{0} \\ \frac{\partial h}{\partial t} + \mathcal{H} \nabla_S \cdot \mathbf{u} = 0 \end{cases} \quad (2.36)$$

where  $f = 2\tilde{\Omega} \cos(\varphi) = 2\tilde{\Omega}z$  is the Coriolis parameter<sup>3</sup>, and  $\tilde{\Omega}$  is the rotation rate of the sphere surface (that is, the amount the body rotates per unit time)<sup>4</sup>.

A keen eye may notice that – while we have operators for the multiplication by  $z$ , the spherical gradient and divergence – we require one more operator not yet

<sup>3</sup>Normally, the Coriolis parameter is expressed in terms of the latitude: latitude =  $\frac{\pi}{2} - \varphi$

<sup>4</sup>On a full scale Earth, the rotation rate is  $\tilde{\Omega} = 7.292110^{-5}$  rad s<sup>-1</sup>.



**Figure 2.4:** Four snapshots of the solution for the height  $h$  of the linearised shallow water equations with  $\tilde{\Omega} = \mathcal{H} = 1$  and a step size of  $\Delta t = 0.01$ , solved using the backwards Euler timestepping method. Top Left: Initial conditions (0 timesteps). Top Right: After 50 timesteps. Bottom Left: After 75 timesteps. Bottom Right: After 100 timesteps.

defined for the unit vector cross product  $\hat{\mathbf{r}} \times \mathbf{u}$ .

**Definition 5.** Define the operator matrix  $\mathcal{R}$  according to:

$$\hat{\mathbf{r}}(x, y, z) \times \mathbf{u}(x, y, z) = \tilde{\mathbb{T}}(x, y, z)^\top \mathcal{R} \mathbf{u}^c.$$

**Lemma 4.** The operator  $\mathcal{R}$  defined in Definition 5 is sparse with banded-block-banded structure. More specifically,  $\mathcal{R}$  has block-bandwidths  $(0, 0)$ , and sub-block-bandwidths  $(1, 1)$  (i.e. is tridiagonal).

*Proof.* Using the relations that  $\Psi_l^m = \nabla_S Y_l^m$  and  $\Phi_l^m = \hat{\mathbf{r}} \times \nabla_S Y_l^m$  for each  $l \in \mathbb{N}$ ,  $m \in \mathbb{Z}$  s.t.  $-l \leq m \leq l$ , we have that:

$$\begin{aligned} \hat{\mathbf{r}} \times \mathbf{u} &= \hat{\mathbf{r}} \times \sum_{l=0}^N \sum_{m=-l}^l [u_{l,m}^\Psi \Psi_l^m + u_{l,m}^\Phi \Phi_l^m] \\ &= \sum_{l=0}^N \sum_{m=-l}^l [u_{l,m}^\Psi \hat{\mathbf{r}} \times \Psi_l^m + u_{l,m}^\Phi \hat{\mathbf{r}} \times (\hat{\mathbf{r}} \times \Psi_l^m)] \\ &= \sum_{l=0}^N \sum_{m=-l}^l [u_{l,m}^\Psi \Phi_l^m - u_{l,m}^\Phi \Psi_l^m]. \end{aligned}$$

We can then write down the operator  $\mathcal{R}$  as follows:

$$\mathcal{R} = \begin{pmatrix} R & & \\ & R & \\ & & \ddots \end{pmatrix} \in \mathbb{R}^{2(N+1)^2 \times 2(N+1)^2},$$

where the blocks  $R$  are simply given by

$$R := \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \in \mathbb{R}^{2 \times 2}.$$

□

For simplicity, we will implement a backward Euler timestepping method to solve the linear SWEs with timestep  $\Delta t$ . Of course, other timestepping methods are equally implementable (for example, the method used in the IFS model that ECMWF uses is a semi-implicit, semi-Lagrangian (SISL) timestepping method [15]), but we simplify things here to show proof of concept. Using  $\mathbf{u}_n, h_n$  to represent the solution at time  $n\Delta t$ :

$$\mathbf{u}_{n+1} = \mathbf{u}_n + \Delta t (\nabla_S h_{n+1} - f \hat{\mathbf{r}} \times \mathbf{u}_{n+1})$$

$$h_{n+1} = h_n - \Delta t \mathcal{H} \nabla_S \cdot \mathbf{u}_{n+1}$$

Let us write this in coefficient space. Firstly, our coefficients vectors for the functions  $\mathbf{u}, h$  we will denote  $\mathbf{u}^c, \mathbf{h}^c$  – that is:

$$\mathbf{u}(x, y, z) \approx \tilde{\mathbb{T}}(x, y, z)^\top \mathbf{u}^c$$

$$h(x, y, z) \approx \mathbb{P}(x, y, z)^\top \mathbf{h}^c.$$

As eluded to earlier, we will “extend” the scalar function’s coefficients vector, writing  $\tilde{\mathbf{h}}^c = E_v \mathbf{h}^c$ , as described in Section 2.5.4. Thus, our system can be written

as a matrix-vector system:

$$\begin{aligned}\mathbf{u}_{n+1}^c &= \mathbf{u}_n^c + \Delta t (\tilde{\mathcal{G}}\tilde{\mathbf{h}}_{n+1}^c - 2\tilde{\Omega}^T J_z \mathcal{R} \mathbf{u}_{n+1}^c) \\ \tilde{\mathbf{h}}_{n+1}^c &= \tilde{\mathbf{h}}_n^c - \Delta t \mathcal{H} \tilde{\mathcal{D}} \mathbf{u}_{n+1}^c\end{aligned}$$

$\Longleftrightarrow$

$$\begin{aligned}\mathbf{u}_{n+1}^c &= \mathbf{u}_n^c + \Delta t (\tilde{\mathcal{G}}\tilde{\mathbf{h}}_n^c - \Delta t \mathcal{H} \tilde{\mathcal{G}} \tilde{\mathcal{D}} \mathbf{u}_{n+1}^c - 2\tilde{\Omega}^T J_z \mathcal{R} \mathbf{u}_{n+1}^c) \\ \tilde{\mathbf{h}}_{n+1}^c &= \tilde{\mathbf{h}}_n^c - \Delta t \mathcal{H} \tilde{\mathcal{D}} \mathbf{u}_{n+1}^c\end{aligned}$$

$\Longleftrightarrow$

$$\begin{aligned}\left[ I + \Delta t^2 \mathcal{H} \tilde{\mathcal{G}} \tilde{\mathcal{D}} + 2\tilde{\Omega} \Delta t^T J_z \mathcal{R} \right] \mathbf{u}_{n+1}^c &= \mathbf{u}_n^c + \Delta t \tilde{\mathcal{G}} \tilde{\mathbf{h}}_n^c \\ \tilde{\mathbf{h}}_{n+1}^c &= \tilde{\mathbf{h}}_n^c - \Delta t \mathcal{H} \tilde{\mathcal{D}} \mathbf{u}_{n+1}^c.\end{aligned}$$

Here, it is clear that the matrix  $I + \Delta t^2 \mathcal{H} \tilde{\mathcal{G}} \tilde{\mathcal{D}} + 2\tilde{\Omega} \Delta t^T J_z \mathcal{R}$  is sparse with banded-block-banded structure, and hence this system will be efficient to solve. Doing so at each iteration provides us with the coefficients vectors  $\mathbf{u}_{n+1}^c, \tilde{\mathbf{h}}_{n+1}^c$  if we know  $\mathbf{u}_n^c, \tilde{\mathbf{h}}_n^c$ . Of course, when evaluating the function  $h$ , we can simply gather back the coefficients vector  $\mathbf{h}^c$  by  $\mathbf{h}^c = E_s \tilde{\mathbf{h}}^c$ .

As a demonstration, in Figure 2.4 we see four snapshots of the solution to the system in equation (2.36) on the unit sphere, with  $\tilde{\Omega} = \mathcal{H} = 1$  and initial conditions

given by

$$\mathbf{u}_0 := \hat{\phi} \mu \sin \varphi, \quad h_0 := \mu \sin^2 \varphi,$$

where  $\mu = \frac{\pi}{6}$ , with the solution order  $N$  set large enough (here,  $N = 20$ ). This example is very similar to “test case 2” in the well-known standard test set for shallow water equations in spherical geometries [69]. The snapshots of the wave height  $h$  are taken of the initial conditions, plus after 50, 75 and 100 timesteps respectively, using a step size of  $\Delta t = 0.01$ .

## 2.7 Conclusion

Spherical Harmonics have been widely used to solve PDEs, in particular a Spherical Harmonics approach is used as part of the Integrated Forecasting System (IFS) that ECMWF uses for their forecasts. In this chapter, we have chosen to view the Spherical Harmonics as orthogonal (in fact orthonormal) polynomials in  $x$ ,  $y$  and  $z$  in order to illustrate and understand more about how the framework presented here for using multivariate OPs in a spectral method translate to the sphere. We have derived the entries of *Jacobi matrices* that represent multiplication by the coordinates, showing how they are *banded-block-banded* in structure. We have set out how one can find coefficients for the expansion of a function in the Spherical Harmonics basis using established transforms, and how one can evaluate a function given its coefficients vector using a multidimensional version of the Clenshaw algorithm. We extended our framework to include the *tangent*



*bundle* of the sphere – that is, the space of vectors orthogonal to the unit outward normal for each point on the sphere – where we can use the *Vector Spherical Harmonics* as vector-valued orthogonal polynomials to be a basis, demonstrating how similar such Jacobi matrices, function expansion and evaluation can be derived. Further, by exploiting the natural sparsity of the relationships between the OPs in the bases, we have shown how matrices for differential and other operators will be sparse and banded-block-banded in structure too. In fact, for the Spherical Harmonics, these are simply diagonal. Finally, we presented a couple of simple examples of how one can put this all into practice, namely solving the heat equation and the linearised shallow water equations on the unit sphere.

# Chapter 3

## Disk slices and trapeziums

The contents of the previous chapter on spherical harmonics can allow us to use similar techniques for developing sparse spectral methods for solving linear partial differential equations on other domains. Before we can return to the 3D surface realm, let us first achieve a foundation in two-dimensions by looking at a special class of geometries that includes disk slices and trapeziums.

In recent years, there has been some interesting work in the area of multivariate orthogonal polynomials in two-dimensional domains (see [17]), such as for a Chebyshev-based spectral method on the rectangle [29] . In particular, spectral methods have been established on the disk [10, 65] and the triangle [45]. Jacobi polynomials are extremely useful for solving a number of problems in not just 1D but 2D also, since one can utilise the sparsity of the relations that they hold to yield operator matrices that are efficient to solve [16]. In a similar vein to the ultraspherical method for the unit interval [44? ], for the triangle a choice of OPs

list some domains and cite references for them

are the Jacobi polynomials on the triangle, which are a three-parameter family of polynomials in  $(x, y)$  with differential operators acting in such a way so as to increment or decrement the parameter values [17, 30, 45]:

$$P_{n,k}^{(a,b,c)}(x, y) := \tilde{P}_{n-k}^{(2k+b+c+1,a)}(x) (1-x)^k \tilde{P}_k^{(c,b)}\left(\frac{y}{1-x}\right) \quad (3.1)$$

for  $(x, y) \in \{(x, y) \in \mathbb{R}^2 \mid 0 < x < 1, 0 < y < 1-x\}$ , where  $\tilde{P}_k^{(a,b)}$  is the degree  $k$  *shifted Jacobi polynomial*<sup>1</sup> with  $a, b, c$  being parameter values.

For the disk, a review of spectral methods on the disk for solving the Poisson equation has been completed [10], that looks at methods involving various bases for expanding functions in the disk including Zernike polynomials and Chebyshev polynomials. It is concluded that using Chebyshev polynomials yields mixed results, but that there is also no single advised basis to use. While Zernike polynomials yield accurate results, there is a lack of radial fast transform, which is a disadvantage. However, ideas for how to overcome this have progressed [54? ?]. One solution proposed is a radial basis defined in terms of Jacobi polynomials [65]:

$$Q_n^{k,m}(r) := C_n^{k,m} r^m P_n^{(k,m)}(2r^2 - 1), \quad (3.2)$$

where  $C_n^{k,m}$  is some normalising constant so that they are orthonormal with respect to  $\int_0^1 Q_n^{k,m}(r) Q_{n'}^{k,m}(r) (1-r^2)^k r \, dr = \delta_{n,n'}$ .  $\{Q_n^{k,m}\}$  are therefore a two-

---

<sup>1</sup>The shifted Jacobi polynomials are orthogonal on the interval  $[0, 1]$  and given by  $\tilde{P}_k^{(a,b)}(x) = P_k^{(a,b)}(2x-1)$  where  $P_k^{(a,b)}$  is the standard degree  $k$  Jacobi polynomial.

parameter family of orthogonal polynomials. Again, the idea here is that differential and other operators, as well as basis (parameter) conversion operators, can be represented as banded matrices.

In this chapter, we aim to present a basis and develop a sparse spectral method on generalisations of these domains, namely the disk-slice and trapezium. The full unit disk and triangle cases are special cases of the spectral methods we investigate here too. We will show that the operators for multiplication by  $x$  and  $y$ , as well as differential and parameter conversion, will be banded-block-banded.

More precisely, we will consider the solution of partial differential equations on the domain

$$\Omega := \{(x, y) \in \mathbb{R}^2 \mid \alpha \leq x \leq \beta, \gamma\rho(x) \leq y \leq \delta\rho(x)\}$$

where either of the following conditions hold:

**Condition 1.**  $\rho$  is a degree 1 polynomial.

**Condition 2.**  $\rho$  is the square root of a non-negative degree  $\leq 2$  polynomial,  $-\gamma = \delta > 0$ .

For simplicity of presentation we focus on the disk-slice in this chapter, where  $\rho(x) = \sqrt{1 - x^2}$ ,  $(\alpha, \beta) \subset (0, 1)$ , and  $(\gamma, \delta) = (-1, 1)$ . However, we will discuss an extension to other geometries, including the half-disk and trapeziums, by choosing the function  $\rho$  and constants  $\alpha, \beta, \gamma, \delta$  appropriately (for example, choosing  $\rho(x) = 1 - \xi x$  and  $(\alpha, \beta) = (\gamma, \delta) = (0, 1)$  for some constant  $\xi > 0$  returns the trapezium case). Moreover, one should be able to see that the full disk and triangle can also

be obtained in such a way.

We will show that partial differential equations become sparse linear systems when viewed as acting on expansions involving a family of orthogonal polynomials (OPs) that generalise Jacobi polynomials, mirroring the ultraspherical spectral method for ordinary differential equations on the interval [44], and its analogues on the disk [65] and triangle [46, 45]. On the disk-slice the family of weights we consider are of the form

$$W^{(a,b,c)}(x, y) = (\beta - x)^a (x - \alpha)^b (1 - x^2 - y^2)^c$$

for  $\alpha \leq x \leq \beta$ ,  $-\rho(x) \leq y \leq \rho(x)$ . The corresponding OPs denoted  $H_{n,k}^{(a,b,c)}(x, y)$ , where  $n$  denotes the polynomial degree, and  $0 \leq k \leq n$ . We define these to be orthogonalised lexicographically, that is,

$$H_{n,k}^{(a,b,c)}(x, y) = C_{n,k} x^{n-k} y^k + (\text{lower order terms})$$

where  $C_{n,k} \neq 0$  and “lower order terms” includes degree  $n$  polynomials of the form  $x^{n-j} y^j$  where  $j < k$ . The precise normalization arises from their definition in terms of one-dimensional OPs we will see in Definition 6.

Sparsity comes from expanding the domain and range of an operator using different choices of the parameters  $a$ ,  $b$  and  $c$ . Whereas the sparsity pattern and entries derived for equations on the triangle [46, 45] and for equations on the disk [65] with the frameworks discussed earlier both result from manipulations of Jacobi polynomials, in the present work we use a more general integration-by-parts

argument to deduce the sparsity structure, alongside careful use of the Christoffel-Darboux formula [42, 18.2.2] and quadrature rules to determine the entries. In particular, by exploiting the connection with one-dimensional orthogonal polynomials we can construct discretizations of general partial differential operators of size  $p(p-1)/2 \times p(p-1)/2$  in  $O(p^3)$  operations, where  $p$  is the total polynomial degree. This compares favourably to  $O(p^6)$  operations if one proceeds naïvely. Furthermore, we use this framework to derive sparse  $p$ -finite element methods that are analogous to those of Beuchler and Schöberl on tetrahedra [5], see also work by Li and Shen [32].

### 3.1 Orthogonal polynomials on the disk-slice and the trapezium

We can mirror the approach for the spherical harmonics, in that we will outline the construction and some basic properties of our 2D OPs on  $\Omega$  that we denote by  $H_{n,k}^{(a,b,c,d)}(x,y)$ . The symmetry in the weight allows us to express the polynomials in terms of 1D OPs, and deduce certain properties such as recurrence relationships.

#### 3.1.1 Explicit construction

We can construct 2D orthogonal polynomials on  $\Omega$  from 1D orthogonal polynomials on the intervals  $[\alpha, \beta]$  and  $[\gamma, \delta]$ .

**Proposition 1** ([17, p55–56]). *Let  $w_1 : (\alpha, \beta) \rightarrow \mathbb{R}$ ,  $w_2 : (\gamma, \delta) \rightarrow \mathbb{R}$  be weight functions with  $\alpha, \beta, \gamma, \delta \in \mathbb{R}$ , and let  $\rho : (\alpha, \beta) \rightarrow (0, \infty)$  be such that either Condition 1 or Condition 2 with  $w_2$  being an even function hold.  $\forall, n = 0, 1, 2, \dots$ , let  $\{p_{n,k}\}$  be polynomials orthogonal with respect to the weight  $\rho(x)^{2k+1}w_1(x)$  where  $0 \leq k \leq n$ , and  $\{q_n\}$  be polynomials orthogonal with respect to the weight  $w_2(x)$ . Then, for  $n = 0, 1, 2, \dots$ ,  $k = 0, \dots, n$ , the 2D polynomials defined on  $\Omega$  given by*

$$H_{n,k}(x, y) := p_{n-k,k}(x) \rho(x)^k q_k\left(\frac{y}{\rho(x)}\right)$$

*are orthogonal polynomials with respect to the weight  $W(x, y) := w_1(x) w_2\left(\frac{y}{\rho(x)}\right)$  on  $\Omega$ .*

For disk slices and trapeziums, we specialise Proposition 1 in the following definitions. First we introduce notation for two families of univariate OPs.

**Definition 6.** *Let  $w_R^{(a,b,c)}(x)$  and  $w_P^{(a,b)}(x)$  be two weight functions on the intervals  $(\alpha, \beta)$  and  $(\gamma, \delta)$  respectively, given by:*

$$\begin{cases} w_R^{(a,b,c)}(x) &:= (\beta - x)^a (x - \alpha)^b \rho(x)^c \\ w_P^{(a,b)}(x) &:= (\delta - x)^a (x - \gamma)^b \end{cases}$$

*and define the associated inner products by:*

$$\langle p, q \rangle_{w_R^{(a,b,c)}} := \frac{1}{\omega_R^{(a,b,c)}} \int_{\alpha}^{\beta} p(x) q(x) w_R^{(a,b,c)}(x) dx \quad (3.3)$$

$$\langle p, q \rangle_{w_P^{(a,b)}} := \frac{1}{\omega_P^{(a,b)}} \int_{\gamma}^{\delta} p(y) q(y) w_P^{(a,b)}(y) dy \quad (3.4)$$

where

$$\omega_R^{(a,b,c)} := \int_{\alpha}^{\beta} w_R^{(a,b,c)}(x) dx, \quad \omega_P^{(a,b)} := \int_{\gamma}^{\delta} w_P^{(a,b)}(y) dy. \quad (3.5)$$

Denote the three-parameter family of orthonormal polynomials on  $[\alpha, \beta]$  by  $\{R_n^{(a,b,c)}\}$ , orthonormal with respect to the inner product defined in (3.3), and the two-parameter family of orthonormal polynomials on  $[\gamma, \delta]$  by  $\{P_n^{(a,b)}\}$ , orthonormal with respect to the inner product defined in (3.4).

We can now write down our 2D OP family.

**Definition 7.** For parameters  $a, b, c, d \in \mathbb{R}$ , define the four-parameter 2D orthogonal polynomials as the set  $\{H_{n,k}^{(a,b,c,d)}\}$  where:

$$H_{n,k}^{(a,b,c,d)}(x, y) := R_{n-k}^{(a,b,c+d+2k+1)}(x) \rho(x)^k P_k^{(d,c)}\left(\frac{y}{\rho(x)}\right), \quad (x, y) \in \Omega,$$

The comparison with the construction of the 3D spherical harmonics is evident here (the  $R$  OPs are in place of the Jacobi polynomials, and the  $P$  OPs are in place of the complex exponentials, which are just Chebyshev polynomials). By defining our OPs in this way, we will be able to yield sparse and banded relations for the necessary operators involved in PDEs, just as we did before.

$\{H_{n,k}^{(a,b,c,d)}\}$  are orthogonal with respect to the weight

$$W^{(a,b,c,d)}(x, y) := w_R^{(a,b,c+d)}(x) w_P^{(d,c)}\left(\frac{y}{\rho(x)}\right), \quad (x, y) \in \Omega,$$

assuming that either Condition 1 holds, or Condition 2 holds with  $w_P^{(a,b)}$  being



an even function (i.e.  $a = b$ , and we can hence denote the weight as  $w_P^{(a)}(x) = w_P^{(a,a)}(x) = (\delta - x^2)^a$ ). That is,

$$\left\langle H_{n,k}^{(a,b,c,d)}, H_{m,j}^{(a,b,c,d)} \right\rangle_{W^{(a,b,c,d)}} = \omega_R^{(a,b,c,d+2k+1)} \omega_P^{(d,c)} \delta_{n,m} \delta_{k,j}$$

where for  $f, g : \Omega \rightarrow \mathbb{R}$  the inner product is defined as

$$\langle f, g \rangle_{W^{(a,b,c,d)}} := \iint_{\Omega} f(x, y) g(x, y) W^{(a,b,c,d)}(x, y) dy dx.$$

We can see that they are indeed orthogonal using the change of variable  $t = \frac{y}{\rho(x)}$ , for the following normalisation:

$$\left\langle H_{n,k}^{(a,b,c,d)}, H_{m,j}^{(a,b,c,d)} \right\rangle_{W^{(a,b,c,d)}} \quad (3.6)$$

$$\begin{aligned} &= \iint_{\Omega} \left[ R_{n-k}^{(a,b,c,d+2k+1)}(x) R_{m-j}^{(a,b,c,d+2j+1)}(x) \rho(x)^{k+j} \right. \\ &\quad \cdot P_k^{(d,c)}\left(\frac{y}{\rho(x)}\right) P_j^{(d,c)}\left(\frac{y}{\rho(x)}\right) W^{(a,b,c,d)}(x, y) \Big] dy dx \\ &= \left( \int_{\alpha}^{\beta} R_{n-k}^{(a,b,c,d+2k+1)}(x) R_{m-j}^{(a,b,c,d+2j+1)}(x) w_R^{(a,b,c,d+k+j+1)}(x) dx \right) \\ &\quad \cdot \left( \int_{\gamma}^{\delta} P_k^{(d,c)}(t) P_j^{(d,c)}(t) w_P^{(d,c)}(t) dt \right) \\ &= \omega_P^{(d,c)} \delta_{k,j} \int_{\alpha}^{\beta} R_{n-k}^{(a,b,c,d+2k+1)}(x) R_{m-k}^{(a,b,c,d+2k+1)}(x) w_R^{(a,b,c,d+2k+1)}(x) dx \\ &= \omega_R^{(a,b,c,d+2k+1)} \omega_P^{(d,c)} \delta_{n,m} \delta_{k,j}. \end{aligned} \quad (3.7)$$

For the disk-slice, the weight  $W^{(a,b,c)}(x, y) = (\beta - x)^a (x - \alpha)^b (1 - x^2 - y^2)^c$  results

from setting:

$$\begin{cases} (\alpha, \beta) & \subset (0, 1) \\ (\gamma, \delta) & := (-1, 1) \\ \rho(x) & := (1 - x^2)^{\frac{1}{2}} \end{cases}$$

so that

$$\begin{cases} w_R^{(a,b,c)}(x) := (\beta - x)^a (x - \alpha)^b \rho(x)^c \\ w_P^{(c)}(x) := (1 - x)^c (1 + x)^c = (1 - x^2)^c. \end{cases}$$

Note here we can simply remove the need for including a fourth parameter  $d$ . The 2D OPs orthogonal with respect to the weight above on the disk-slice  $\Omega$  are then given by:

$$H_{n,k}^{(a,b,c)}(x, y) := R_{n-k}^{(a,b,2c+2k+1)}(x) \rho(x)^k P_k^{(c,c)}\left(\frac{y}{\rho(x)}\right), \quad (x, y) \in \Omega \quad (3.8)$$

In this case the weight  $w_P(x)$  is an ultraspherical weight, and the corresponding OPs are the normalized Jacobi polynomials  $\{P_n^{(b,b)}\}$ , while the weight  $w_R(x)$  is non-classical (it is in fact semi-classical, and is equivalent to a generalized Jacobi weight [35, §5]).

### 3.1.2 Jacobi matrices

Recall that Jacobi matrices are an important piece of the puzzle for using these orthogonal polynomials to practically solve PDEs in a sparse spectral method. To obtain the entries for them, we first need to establish the three-term recurrences associated with  $R_n^{(a,b,c)}$  and  $P_n^{(d,c)}$ . These can be expressed as:

$$xR_n^{(a,b,c)}(x) = \beta_n^{(a,b,c)}R_{n+1}^{(a,b,c)}(x) + \alpha_n^{(a,b,c)}R_n^{(a,b,c)}(x) + \beta_{n-1}^{(a,b,c)}R_{n-1}^{(a,b,c)}(x) \quad (3.9)$$

$$yP_n^{(d,c)}(y) = \delta_n^{(d,c)}P_{n+1}^{(d,c)}(y) + \gamma_n^{(d,c)}P_n^{(d,c)}(y) + \delta_{n-1}^{(d,c)}P_{n-1}^{(d,c)}(y). \quad (3.10)$$

Of course, for the disk-slice case, we have that  $c = d$  and  $\gamma_n^{(c,c)} = 0 \forall n = 0, 1, 2, \dots$

We can use (3.9) and (3.10) to determine the 2D recurrences for  $H_{n,k}^{(a,b,c,d)}(x, y)$ .

Importantly, we can deduce sparsity in the recurrence relationships.

**Lemma 5.**  $H_{n,k}^{(a,b,c,d)}(x, y)$  satisfy the following 3-term recurrences:

$$\begin{aligned} xH_{n,k}^{(a,b,c,d)}(x, y) &= \alpha_{n,k,1}^{(a,b,c,d)}H_{n-1,k}^{(a,b,c,d)}(x, y) + \alpha_{n,k,2}^{(a,b,c,d)}H_{n,k}^{(a,b,c,d)}(x, y) + \alpha_{n+1,k,1}^{(a,b,c,d)}H_{n+1,k}^{(a,b,c,d)}(x, y), \\ yH_{n,k}^{(a,b,c,d)}(x, y) &= \beta_{n,k,1}^{(a,b,c,d)}H_{n-1,k-1}^{(a,b,c,d)}(x, y) + \beta_{n,k,2}^{(a,b,c,d)}H_{n-1,k}^{(a,b,c,d)}(x, y) + \beta_{n,k,3}^{(a,b,c,d)}H_{n-1,k+1}^{(a,b,c,d)}(x, y) \\ &\quad + \beta_{n,k,4}^{(a,b,c,d)}H_{n,k-1}^{(a,b,c,d)}(x, y) + \beta_{n,k,5}^{(a,b,c,d)}H_{n,k}^{(a,b,c,d)}(x, y) + \beta_{n,k,6}^{(a,b,c,d)}H_{n,k+1}^{(a,b,c,d)}(x, y) \\ &\quad + \beta_{n,k,7}^{(a,b,c,d)}H_{n+1,k-1}^{(a,b,c,d)}(x, y) + \beta_{n,k,8}^{(a,b,c,d)}H_{n+1,k}^{(a,b,c,d)}(x, y) + \beta_{n,k,9}^{(a,b,c,d)}H_{n+1,k+1}^{(a,b,c,d)}(x, y), \end{aligned}$$

for  $(x, y) \in \Omega$ , where

$$\begin{aligned}
\alpha_{n,k,1}^{(a,b,c,d)} &:= \beta_{n-k-1}^{(a,b,c,d+2k+1)}, & \alpha_{n,k,2}^{(a,b,c,d)} &:= \alpha_{n-k}^{(a,b,c,d+2k+1)} \\
\beta_{n,k,1}^{(a,b,c,d)} &:= \delta_{k-1}^{(d,c)} \left\langle R_{n-k}^{(a,b,c,d+2k+1)}, R_{n-k}^{(a,b,c,d+2k-1)} \right\rangle_{w_R^{(a,b,c,d+2k+1)}} \\
\beta_{n,k,2}^{(a,b,c,d)} &:= \gamma_k^{(d,c)} \left\langle R_{n-k}^{(a,b,c,d+2k+1)}, \rho(x) R_{n-k-1}^{(a,b,c,d+2k+1)} \right\rangle_{w_R^{(a,b,c,d+2k+1)}} \\
\beta_{n,k,3}^{(a,b,c,d)} &:= \delta_k^{(d,c)} \left\langle R_{n-k}^{(a,b,c,d+2k+1)}, R_{n-k-2}^{(a,b,c,d+2k+3)} \right\rangle_{w_R^{(a,b,c,d+2k+3)}} \\
\beta_{n,k,4}^{(a,b,c,d)} &:= \delta_{k-1}^{(d,c)} \left\langle R_{n-k}^{(a,b,c,d+2k+1)}, R_{n-k+1}^{(a,b,c,d+2k-1)} \right\rangle_{w_R^{(a,b,c,d+2k+1)}} \\
\beta_{n,k,5}^{(a,b,c,d)} &:= \gamma_k^{(d,c)} \left\langle R_{n-k}^{(a,b,c,d+2k+1)}, \rho(x) R_{n-k}^{(a,b,c,d+2k+1)} \right\rangle_{w_R^{(a,b,c,d+2k+1)}} \\
\beta_{n,k,6}^{(a,b,c,d)} &:= \delta_k^{(d,c)} \left\langle R_{n-k}^{(a,b,c,d+2k+1)}, R_{n-k-1}^{(a,b,c,d+2k+3)} \right\rangle_{w_R^{(a,b,c,d+2k+3)}} \\
\beta_{n,k,7}^{(a,b,c,d)} &:= \delta_{k-1}^{(d,c)} \left\langle R_{n-k}^{(a,b,c,d+2k+1)}, R_{n-k+2}^{(a,b,c,d+2k-1)} \right\rangle_{w_R^{(a,b,c,d+2k+1)}} \\
\beta_{n,k,8}^{(a,b,c,d)} &:= \gamma_k^{(d,c)} \left\langle R_{n-k}^{(a,b,c,d+2k+1)}, \rho(x) R_{n-k+1}^{(a,b,c,d+2k+1)} \right\rangle_{w_R^{(a,b,c,d+2k+1)}} \\
\beta_{n,k,9}^{(a,b,c,d)} &:= \delta_k^{(d,c)} \left\langle R_{n-k}^{(a,b,c,d+2k+1)}, R_{n-k}^{(a,b,c,d+2k+3)} \right\rangle_{w_R^{(a,b,c,d+2k+3)}}.
\end{aligned}$$

*Proof.* The 3-term recurrence for multiplication by  $x$  follows from equation (3.9).

For the recurrence for multiplication by  $y$ , since  $\{H_{m,j}^{(a,b,c,d)}\}$  for  $m = 0, \dots, n+1$ ,  $j = 0, \dots, m$  is an orthogonal basis for any degree  $n+1$  polynomial, we can expand  $y H_{n,k}^{(a,b,c,d)}(x, y) = \sum_{m=0}^{n+1} \sum_{j=0}^m c_{m,j} H_{m,j}^{(a,b,c,d)}(x, y)$ . These coefficients are given by

$$c_{m,j} = \left\langle y H_{n,k}^{(a,b,c,d)}, H_{m,j}^{(a,b,c,d)} \right\rangle_{W^{(a,b,c,d)}} \left\| H_{m,j}^{(a,b,c,d)} \right\|_{W^{(a,b,c,d)}}^{-2}.$$

Recall from equation (3.7) that  $\left\| H_{m,j}^{(a,b,c,d)} \right\|_{W^{(a,b,c,d)}}^2 = \omega_R^{(a,b,c,d+2j+1)} \omega_P^{(d,c)}$ . Then

for  $m = 0, \dots, n+1$ ,  $j = 0, \dots, m$ , using the change of variable  $t = \frac{y}{\rho(x)}$ :

$$\begin{aligned}
& \left\langle y H_{n,k}^{(a,b,c,d)}, H_{m,j}^{(a,b,c,d)} \right\rangle_{W^{(a,b,c,d)}} \\
&= \iint_{\Omega} H_{n,k}^{(a,b,c,d)}(x, y) H_{m,j}^{(a,b,c,d)}(x, y) y W^{(a,b,c,d)}(x, y) dy dx \\
&= \left( \int_{\alpha}^{\beta} R_{n-k}^{(a,b,c+d+2k+1)}(x) R_{m-j}^{(a,b,c+d+2j+1)}(x) \rho(x)^{k+j+2} w_R^{(a,b,c+d)}(x) dx \right) \\
&\quad \cdot \left( \int_{\gamma}^{\delta} P_k^{(d,c)}(t) P_j^{(d,c)}(t) t w_P^{(d,c)}(t) dt \right) \\
&= \left( \int_{\alpha}^{\beta} R_{n-k}^{(a,b,c+d+2k+1)}(x) R_{m-j}^{(a,b,c+d+2j+1)}(x) w_R^{(a,b,c+d+k+j+2)}(x) dx \right) \\
&\quad \cdot \left( \int_{\gamma}^{\delta} P_k^{(d,c)}(t) P_j^{(d,c)}(t) t w_P^{(d,c)}(t) dt \right) \\
&= \begin{cases} \delta_k^{(d,c)} \omega_P^{(d,c)} \omega_R^{(a,b,c+d+2k+3)} \left\langle R_{n-k}^{(a,b,c+d+2k+1)}, R_{m-k-1}^{(a,b,c+d+2k+3)} \right\rangle_{w_R^{(a,b,c+d+2k+3)}} & \text{if } j = k+1 \\ \gamma_k^{(d,c)} \omega_P^{(d,c)} \omega_R^{(a,b,c+d+2k+1)} \left\langle R_{n-k}^{(a,b,c+d+2k+1)}, \rho(x) R_{m-k}^{(a,b,c+d+2k+1)} \right\rangle_{w_R^{(a,b,c+d+2k+1)}} & \text{if } j = k \\ \delta_{k-1}^{(d,c)} \omega_P^{(d,c)} \omega_R^{(a,b,c+d+2k-1)} \left\langle R_{n-k}^{(a,b,c+d+2k-1)}, \rho(x)^2 R_{m-k+1}^{(a,b,c+d+2k-1)} \right\rangle_{w_R^{(a,b,c+d+2k-1)}} & \text{if } j = k-1 \\ 0 & \text{otherwise} \end{cases}
\end{aligned}$$

where, by orthogonality,

$$\begin{aligned}
& \left\langle R_{n-k}^{(a,b,c+d+2k+1)}, R_{m-k-1}^{(a,b,c+d+2k+3)} \right\rangle_{w_R^{(a,b,c+d+2k+3)}} = 0 \quad \text{for } m < n-1, \\
& \left\langle R_{n-k}^{(a,b,c+d+2k+1)}, \rho(x)^2 R_{m-k+1}^{(a,b,c+d+2k-1)} \right\rangle_{w_R^{(a,b,c+d+2k-1)}} = 0 \quad \text{for } m < n-1.
\end{aligned}$$

Finally, if Condition 1 holds we have that

$$\left\langle R_{n-k}^{(a,b,c+d+2k+1)}, \rho(x) R_{m-k}^{(a,b,c+d+2k+1)} \right\rangle_{w_R^{(a,b,c+d+2k+1)}} = 0 \quad \text{for } m < n - 1.$$

If Condition 2 holds we have that  $\gamma_k^{(d,c)} = \gamma_k^{(c,c)} \equiv 0$  for any  $k$ .

□

Three-term recurrences lead to Jacobi operators that correspond to multiplication by  $x$  and  $y$ . Define, for  $n = 0, 1, 2, \dots$ :

$$\mathbb{H}_n^{(a,b,c,d)} := \begin{pmatrix} H_{n,0}^{(a,b,c,d)}(x, y) \\ \vdots \\ H_{n,n}^{(a,b,c,d)}(x, y) \end{pmatrix} \in \mathbb{R}^{n+1}, \quad \mathbb{H}^{(a,b,c,d)} := \begin{pmatrix} \mathbb{H}_0^{(a,b,c,d)} \\ \mathbb{H}_1^{(a,b,c,d)} \\ \mathbb{H}_2^{(a,b,c,d)} \\ \vdots \end{pmatrix}$$

and set  $J_x^{(a,b,c,d)}, J_y^{(a,b,c,d)}$  as the Jacobi matrices corresponding to

$$J_x^{(a,b,c,d)} \mathbb{H}^{(a,b,c,d)}(x, y) = x \mathbb{H}^{(a,b,c,d)}(x, y), \quad J_y^{(a,b,c,d)} \mathbb{H}^{(a,b,c,d)}(x, y) = y \mathbb{H}^{(a,b,c,d)}(x, y). \quad (3.11)$$

The matrices  $J_x^{(a,b,c,d)}, J_y^{(a,b,c,d)}$  act on the coefficients vector of a function's expansion in the  $\{H_{n,k}^{(a,b,c,d)}\}$  basis. For example, let  $a, b$  be general parameters and a function  $f(x, y)$  defined on  $\Omega$  be approximated by its expansion  $f(x, y) = \mathbb{H}^{(a,b,c,d)}(x, y)^\top \mathbf{f}$ . Then  $x f(x, y)$  is approximated by  $\mathbb{H}^{(a,b,c,d)}(x, y)^\top J_x^{(a,b,c,d)} \mathbf{f}$ . In other words,  $J_x^{(a,b,c,d)} \mathbf{f}$  is the coefficients vector for the expansion of the function

$(x, y) \mapsto x f(x, y)$  in the  $\{H_{n,k}^{(a,b,c,d)}\}$  basis.

Further, note that  $J_x^{(a,b,c,d)}, J_y^{(a,b,c,d)}$  are banded-block-banded matrices (see Definition 1). They are block-tridiagonal (block-bandwidths  $(1, 1)$ ):

$$J_{x/y}^{(a,b,c,d)} = \begin{pmatrix} B_0^{x/y} & A_0^{x/y} & & & \\ C_1^{x/y} & B_1^{x/y} & A_1^{x/y} & & \\ & C_2^{x/y} & B_2^{x/y} & A_2^{x/y} & \\ & & C_3^{x/y} & \ddots & \ddots \\ & & & \ddots & \ddots & \ddots \end{pmatrix}$$

where the blocks themselves are diagonal for  $J_x^{(a,b,c,d)}$  (sub-block-bandwidths  $(0, 0)$ ),

$$A_n^x := \begin{pmatrix} \alpha_{n+1,0,1}^{(a,b,c,d)} & 0 & \dots & 0 \\ & \ddots & & \vdots \\ & & \alpha_{n+1,n,1}^{(a,b,c,d)} & 0 \end{pmatrix} \in \mathbb{R}^{(n+1) \times (n+2)}, \quad n = 0, 1, 2, \dots$$

$$B_n^x := \begin{pmatrix} \alpha_{n,0,2}^{(a,b,c,d)} & & \\ & \ddots & \\ & & \alpha_{n,n,2}^{(a,b,c,d)} \end{pmatrix} \in \mathbb{R}^{(n+1) \times (n+1)} \quad n = 0, 1, 2, \dots$$

$$C_n^x := (A_n^x)^\top \in \mathbb{R}^{(n+1) \times n}, \quad n = 1, 2, \dots$$

and tridiagonal for  $J_y^{(a,b,c,d)}$  (sub-block-bandwidths  $(1, 1)$ ),

$$\begin{aligned}
 A_n^y &:= \begin{pmatrix} \beta_{n,0,8}^{(a,b,c,d)} & \beta_{n,0,9}^{(a,b,c,d)} & & & \\ \beta_{n,1,7}^{(a,b,c,d)} & \ddots & \ddots & & \\ & \ddots & \ddots & \ddots & \\ & & \beta_{n,n,7}^{(a,b,c,d)} & \beta_{n,n,8}^{(a,b,c,d)} & \beta_{n,n,9}^{(a,b,c,d)} \end{pmatrix} \in \mathbb{R}^{(n+1) \times (n+2)}, \quad n = 0, 1, 2, \dots \\
 B_n^y &:= \begin{pmatrix} \beta_{n,0,5}^{(a,b,c,d)} & \beta_{n,0,6}^{(a,b,c,d)} & & & \\ \beta_{n,1,4}^{(a,b,c,d)} & \ddots & \ddots & & \\ & \ddots & \ddots & \beta_{n,n-1,6}^{(a,b,c,d)} & \\ & & \beta_{n,n,4}^{(a,b,c,d)} & \beta_{n,n,5}^{(a,b,c,d)} & \end{pmatrix} \in \mathbb{R}^{(n+1) \times (n+1)} \quad n = 0, 1, 2, \dots \\
 C_n^y &:= \begin{pmatrix} \beta_{n,0,2}^{(a,b,c,d)} & \beta_{n,0,3}^{(a,b,c,d)} & & & \\ \beta_{n,1,1}^{(a,b,c,d)} & \ddots & \ddots & & \\ & \ddots & \ddots & \beta_{n,n-2,3}^{(a,b,c,d)} & \\ & & \ddots & \beta_{n,n-1,2}^{(a,b,c,d)} & \\ & & & \beta_{n,n,1}^{(a,b,c,d)} & \end{pmatrix} \in \mathbb{R}^{(n+1) \times n}, \quad n = 1, 2, \dots
 \end{aligned}$$

Note that the sparsity of the Jacobi matrices (in particular the sparsity of the sub-blocks) comes from the natural sparsity of the three-term recurrences of the 1D OPs, meaning that the sparsity is not limited to the specific disk-slice case.



### 3.1.3 Building the OPs

We can combine each system in (3.11) into a block-tridiagonal system:

$$\begin{pmatrix} 1 & & & & \\ B_0 - G_0(x, y) & A_0 & & & \\ C_1 & B_1 - G_1(x, y) & A_1 & & \\ & C_2 & B_2 - G_2(x, y) & \ddots & \\ & & \ddots & \ddots & \ddots \end{pmatrix} \mathbb{H}^{(a,b,c,d)}(x, y) = \begin{pmatrix} H_{0,0}^{(a,b,c,d)} \\ 0 \\ 0 \\ 0 \\ \vdots \end{pmatrix},$$

where we note  $H_{0,0}^{(a,b,c,d)}(x, y) \equiv R_0^{(a,b,c+d+1)} P_0^{(d,c)}$ , and for each  $n = 0, 1, 2, \dots$ ,

$$\begin{aligned} A_n &:= \begin{pmatrix} A_n^x \\ A_n^y \end{pmatrix} \in \mathbb{R}^{2(n+1) \times (n+2)}, & C_n &:= \begin{pmatrix} C_n^x \\ C_n^y \end{pmatrix} \in \mathbb{R}^{2(n+1) \times n} \quad (n \neq 0), \\ B_n &:= \begin{pmatrix} B_n^x \\ B_n^y \end{pmatrix} \in \mathbb{R}^{2(n+1) \times (n+1)}, & G_n(x, y) &:= \begin{pmatrix} xI_{n+1} \\ yI_{n+1} \end{pmatrix} \in \mathbb{R}^{2(n+1) \times (n+1)}. \end{aligned}$$

For each  $n = 0, 1, 2, \dots$  let  $D_n^\top$  be any matrix that is a left inverse of  $A_n$ , i.e. such that  $D_n^\top A_n = I_{n+2}$ . Multiplying our system by the preconditioner matrix that is given by the block diagonal matrix of the  $D_n^\top$ 's, we obtain a lower triangular

system [17, p78], which can be expanded to obtain the recurrence:

$$\begin{cases} \mathbb{H}_{-1}^{(a,b,c,d)}(x, y) := 0 \\ \mathbb{H}_0^{(a,b,c,d)}(x, y) := H_{0,0}^{(a,b,c,d)} \\ \mathbb{H}_{n+1}^{(a,b,c,d)}(x, y) = -D_n^\top (B_n - G_n(x, y)) \mathbb{H}_n^{(a,b,c,d)}(x, y) - D_n^\top C_n \mathbb{H}_{n-1}^{(a,b,c,d)}(x, y), \quad n = 0, 1, 2, \dots \end{cases}$$


Note that we can define an explicit  $D_n^\top$  as follows:

$$D_n^\top := \begin{pmatrix} \frac{1}{\alpha_{n+1,0,1}^{(a,b,c,d)}} & & & & & & & \\ & \ddots & & & & & & \\ & & \ddots & & & & & \\ & & & \ddots & & & & \\ & & & & \frac{1}{\alpha_{n+1,n,1}^{(a,b,c,d)}} & & & \\ \eta_0 & 0 & \dots & 0 & \eta_1 & \dots & \dots & \eta_{n+1} \end{pmatrix},$$

where

$$\begin{aligned} \eta_{n+1} &= \frac{1}{\beta_{n,n,9}^{(a,b,c,d)}}, \\ \eta_n &= -\frac{1}{\beta_{n,n-1,9}^{(a,b,c,d)}} (\beta_{n,n,8}^{(a,b,c,d)} \eta_{n+1}), \\ \eta_j &= -\frac{1}{\beta_{n,j-1,9}^{(a,b,c,d)}} (\beta_{n,n+j+1,7}^{(a,b,c,d)} \eta_{j+2} + \beta_{n,n+j,8}^{(a,b,c,d)} \eta_{j+1}) \quad \text{for } j = n-1, n-2, \dots, 1, \\ \eta_0 &= -\frac{1}{\alpha_{n+1,0,1}^{(a,b,c,d)}} (\beta_{n,1,7}^{(a,b,c,d)} \eta_2 + \beta_{n,0,8}^{(a,b,c,d)} \eta_1). \end{aligned}$$

It follows that we can apply  $D_n^\top$  in  $O(n)$  complexity, and thereby calculate  $\mathbb{H}_0^{(a,b,c,d)}(x, y)$  through  $\mathbb{H}_n^{(a,b,c,d)}(x, y)$  in optimal  $O(n^2)$  complexity.



**Figure 3.1:** The Laplace operator acting on vectors of  $H_{n,k}^{(0,0,0)}$  coefficients has a sparse matrix representation if the range is represented as vectors of  $H_{n,k}^{(2,2,2)}$  coefficients. Here, the arrows indicate that the corresponding operation has a sparse matrix representation when the domain is  $H_{n,k}^{(a,b,c)}$  coefficients, where  $(a,b,c)$  is at the tail of the arrow, and the range is  $H_{n,k}^{(\tilde{a},\tilde{b},\tilde{c})}$  coefficients, where  $(\tilde{a},\tilde{b},\tilde{c})$  is at the head of the arrow.

For the disk-slice,  $\beta_{n,k,2}^{(a,b,c)} = \beta_{n,k,5}^{(a,b,c)} = \beta_{n,k,8}^{(a,b,c)} \equiv 0$  for any  $n, k$ .

## 3.2 Sparse partial differential operators

In this section, we concentrate on the disk-slice case, and simply note that similar arguments apply for the trapezium case. Recall that, for the disk-slice,

$$\Omega := \{(x, y) \in \mathbb{R}^2 \mid \alpha < x < \beta, \gamma\rho(x) < y < \delta\rho(x)\}$$

where

$$\begin{cases} (\alpha, \beta) & \subset (0, 1) \\ (\gamma, \delta) & := (-1, 1) \\ \rho(x) & := (1 - x^2)^{\frac{1}{2}} \end{cases}.$$

The 2D OPs on the disk-slice  $\Omega$ , orthogonal with respect to the weight

$$\begin{aligned} W^{(a,b,c)}(x, y) &:= w_R^{(a,b,2c)}(x) w_P^{(c)}\left(\frac{y}{\rho(x)}\right) \\ &= (\beta - x)^a (x - \alpha)^b (1 - x^2 - y^2)^c, \quad (x, y) \in \Omega, \end{aligned}$$

are then given by:

$$H_{n,k}^{(a,b,c)}(x, y) := R_{n-k}^{(a,b,2c+2k+1)}(x) \rho(x)^k P_k^{(c,c)}\left(\frac{y}{\rho(x)}\right), \quad (x, y) \in \Omega$$

where the 1D OPs  $\{R_n^{(a,b,c)}\}$  are orthogonal on the interval  $(\alpha, \beta)$  with respect to the weight

$$w_R^{(a,b,c)}(x) := (\beta - x)^a (x - \alpha)^b \rho(x)^c$$

and the 1D OPs  $\{P_n^{(c,c)}\}$  are orthogonal on the interval  $(\gamma, \delta) = (-1, 1)$  with respect to the weight

$$w_P^{(c)}(x) := (1 - x)^c (1 + x)^c = (1 - x^2)^c.$$

Denote the weighted OPs by

$$\mathbb{W}^{(a,b,c)}(x, y) := W^{(a,b,c)}(x, y) \mathbb{H}^{(a,b,c)}(x, y),$$

and recall that a function  $f(x, y)$  defined on  $\Omega$  is approximated by its expansion  $f(x, y) = \mathbb{H}^{(a,b,c)}(x, y)^\top \mathbf{f}$ .

**Definition 8.** Define the operator matrices  $D_x^{(a,b,c)}$ ,  $D_y^{(a,b,c)}$ ,  $W_x^{(a,b,c)}$ ,  $W_y^{(a,b,c)}$  according to:

$$\begin{aligned} \frac{\partial f}{\partial x} &= \mathbb{H}^{(a+1,b+1,c+1)}(x, y)^\top D_x^{(a,b,c)} \mathbf{f}, \\ \frac{\partial f}{\partial y} &= \mathbb{H}^{(a,b,c+1)}(x, y)^\top D_y^{(a,b,c)} \mathbf{f}, \\ \frac{\partial}{\partial x} [W^{(a,b,c)}(x, y) f(x, y)] &= \mathbb{W}^{(a-1,b-1,c-1)}(x, y)^\top W_x^{(a,b,c)} \mathbf{f}, \\ \frac{\partial}{\partial y} [W^{(a,b,c)}(x, y) f(x, y)] &= \mathbb{W}^{(a,b,c-1)}(x, y)^\top W_y^{(a,b,c)} \mathbf{f}. \end{aligned}$$

The incrementing and decrementing of parameters as seen here is analogous to other well known orthogonal polynomial families' derivatives, for example the Jacobi polynomials on the interval, as seen in the DLMF [42, (18.9.3)], and on the triangle [46]. An illustration of how the non-weighted differential operators increment the parameters (a,b,c) is seen in Figure 3.1.

**Theorem 2.** The operator matrices  $D_x^{(a,b,c)}$ ,  $D_y^{(a,b,c)}$ ,  $W_x^{(a,b,c)}$ ,  $W_y^{(a,b,c)}$  from Definition 8 are sparse, with banded-block-banded structure. More specifically:

- $D_x^{(a,b,c)}$  has block-bandwidths  $(-1, 3)$ , and sub-block-bandwidths  $(0, 2)$ .
- $D_y^{(a,b,c)}$  has block-bandwidths  $(-1, 1)$ , and sub-block-bandwidths  $(-1, 1)$ .

- $W_x^{(a,b,c)}$  has block-bandwidths  $(3, -1)$ , and sub-block-bandwidths  $(2, 0)$ .
- $W_y^{(a,b,c)}$  has block-bandwidths  $(1, -1)$ , and sub-block-bandwidths  $(1, -1)$ .

*Proof.* First, note that:

$$w_R^{(a,b,c)'}(x) = -a w_R^{(a-1,b,c)}(x) + b w_R^{(a,b-1,c)}(x) + c \rho(x) \rho'(x) w_R^{(a,b,c-2)}(x), \quad (3.12)$$

$$w_P^{(c)'}(y) = -2c y w_P^{(c-1)}(y), \quad (3.13)$$

$$\rho(x) \rho'(x) = -x. \quad (3.14)$$

We proceed with the case for the operator  $D_y^{(a,b,c)}$  for partial differentiation by  $y$ . Since  $\{H_{m,j}^{(a,b,c+1)}\}$  for  $m = 0, \dots, n-1$ ,  $j = 0, \dots, m$  is an orthogonal basis for any degree  $n-1$  polynomial, we can expand  $\frac{\partial}{\partial y} H_{n,k}^{(a,b,c)} = \sum_{m=0}^{n-1} \sum_{j=0}^m c_{m,j}^y H_{m,j}^{(a,b,c+1)}$ . The coefficients of the expansion are then the entries of the relevant operator matrix. We can use an integration-by-parts argument to show that the only non-zero coefficient of this expansion is when  $m = n-1$ ,  $j = k-1$ . First, note that

$$c_{m,j}^y = \left\langle \frac{\partial}{\partial y} H_{n,k}^{(a,b,c)}, H_{m,j}^{(a,b,c+1)} \right\rangle_{W^{(a,b,c+1)}} \left\| H_{m,j}^{(a,b,c+1)} \right\|_{W^{(a,b,c+1)}}^{-2}.$$

Then, using the change of variable  $t = \frac{y}{\rho(x)}$ , we have that

$$\begin{aligned}
& \left\langle \frac{\partial}{\partial y} H_{n,k}^{(a,b,c)}, H_{m,j}^{(a,b,c+1)} \right\rangle_{W^{(a,b,c+1)}} \\
&= \iint_{\Omega} \left[ R_{n-k}^{(a,b,2c+2k+1)}(x) \rho(x)^{k-1} P_k^{(c,c)'} \left( \frac{y}{\rho(x)} \right) \right. \\
&\quad \cdot R_{m-j}^{(a,b,2c+2j+3)}(x) \rho(x)^j P_j^{(c+1,c+1)} \left( \frac{y}{\rho(x)} \right) \left. \right] dy dx \\
&= \omega_R^{(a,b,2c+2k+1)} \left\langle R_{n-k}^{(a,b,2c+2k+1)}, \rho(x)^{j-k+1} R_{m-j}^{(a,b,2c+2j+3)} \right\rangle_{w_R^{(a,b,2c+2k+1)}} \\
&\quad \cdot \omega_P^{(c+1)} \left\langle P_k^{(c,c)'}, P_j^{(c+1,c+1)} \right\rangle_{w_P^{(c+1)}}
\end{aligned}$$

Now, using (3.13), integration-by-parts, and noting that the weight  $w_P^{(c)}$  is a polynomial of degree  $2c$  and vanishes at the limits of the integral for positive parameter  $c$ , we have that

$$\begin{aligned}
\omega_P^{(c+1)} \left\langle P_k^{(c,c)'}, P_j^{(c+1,c+1)} \right\rangle_{w_P^{(c+1)}} &= \int_{\gamma}^{\delta} P_k^{(c,c)'}(y) P_j^{(c+1,c+1)}(y) w_P^{(c+1)}(y) dy \\
&= - \int_{-1}^1 P_k^{(c,c)}(y) \frac{d}{dy} [w_P^{(c+1)}(y) P_j^{(c+1,c+1)}(y)] dy \\
&= - \int_{-1}^1 P_k^{(c,c)} [P_j^{(c+1,c+1)'} w_P^{(c+1)} - 2c y P_j^{(c+1,c+1)} w_P^{(c)}] dy \\
&= - \omega_P^{(c)} \left\langle P_k^{(c,c)}, w_P^{(1)} P_j^{(c+1,c+1)'} - 2c y P_j^{(c+1,c+1)} \right\rangle_{w_P^{(c)}}
\end{aligned}$$

which is zero for  $j < k - 1$  by orthogonality. Further, when  $j = k - 1$ , we have

that

$$\begin{aligned}
& \omega_R^{(a,b,2c+2k+1)} \left\langle R_{n-k}^{(a,b,2c+2k+1)}, \rho(x)^{j-k+1} R_{m-j}^{(a,b,2c+2j+3)} \right\rangle_{w_R^{(a,b,2c+2k+1)}} \\
&= \omega_R^{(a,b,2c+2k+1)} \left\langle R_{n-k}^{(a,b,2c+2k+1)}, R_{m-j}^{(a,b,2c+2k+1)} \right\rangle_{w_R^{(a,b,2c+2k+1)}} \\
&= \omega_R^{(a,b,2c+2k+1)} \delta_{n,m+1},
\end{aligned}$$

showing that the only possible non-zero coefficient is when  $m = n - 1, j = k - 1$ .

Finally,

$$c_{n-1,k-1}^y = \left\langle P_k^{(c,c)'}, P_{k-1}^{(c+1,c+1)} \right\rangle_{w_P^{(c+1)}}.$$

We next consider the case for the operator  $D_x^{(a,b,c)}$  for partial differentiation by  $x$ .

Since  $\{H_{m,j}^{(a+1,b+1,c+1)}\}$  for  $m = 0, \dots, n-1, j = 0, \dots, m$  is an orthogonal basis for any degree  $n-1$  polynomial, we can expand  $\frac{\partial}{\partial x} H_{n,k}^{(a,b,c)} = \sum_{m=0}^{n-1} \sum_{j=0}^m c_{m,j}^x H_{m,j}^{(a+1,b+1,c+1)}$ .

The coefficients of the expansion are then the entries of the relevant operator matrix. As before, we can use an integration-by-parts argument to show that the only non-zero coefficients of this expansion are when  $m = n - 1, n - 2, n - 3, j = k, k - 1, k - 2$  and  $0 \leq j \leq m$ . First, note that

$$c_{m,j}^x = \left\langle \frac{\partial}{\partial x} H_{n,k}^{(a,b,c)}, H_{m,j}^{(a+1,b+1,c+1)} \right\rangle_{W^{(a+1,b+1,c+1)}} \left\| H_{m,j}^{(a+1,b+1,c+1)} \right\|_{W^{(a+1,b+1,c+1)}}^{-2}.$$



Now, again using the change of variable  $t = \frac{y}{\rho(x)}$ , we have that

$$\begin{aligned}
& \left\langle \frac{\partial}{\partial x} H_{n,k}^{(a,b,c)}, H_{m,j}^{(a+1,b+1,c+1)} \right\rangle_{W^{(a+1,b+1,c+1)}} \\
&= \left( \int_{\alpha}^{\beta} R_{n-k}^{(a,b,2c+2k+1)'} R_{m-j}^{(a+1,b+1,2c+2j+3)} \rho^{k+j+1} w_R^{(a+1,b+1,2c+2)} dx \right) \\
&\quad \cdot \left( \int_{\gamma}^{\delta} P_k^{(c,c)} P_j^{(c+1,c+1)} w_P^{(c+1)} dt \right) \\
&\quad + k \left( \int_{\alpha}^{\beta} R_{n-k}^{(a,b,2c+2k+1)} R_{m-j}^{(a+1,b+1,2c+2j+3)} \rho^{k+j} \rho' w_R^{(a+1,b+1,2c+2)} dx \right) \\
&\quad \cdot \left( \int_{\gamma}^{\delta} P_k^{(c,c)} P_j^{(c+1,c+1)} w_P^{(c+1)} dt \right) \\
&\quad - \left( \int_{\alpha}^{\beta} R_{n-k}^{(a,b,2c+2k+1)} R_{m-j}^{(a+1,b+1,2c+2j+3)} \rho^{k+j} \rho' w_R^{(a+1,b+1,2c+2)} dx \right) \\
&\quad \cdot \left( \int_{\gamma}^{\delta} t P_k^{(c,c)'} P_j^{(c+1,c+1)} w_P^{(c+1)} dt \right). \tag{3.15}
\end{aligned}$$

We will first show that the second factor of each term in (3.15) are zero for  $j < k - 2$  and also for  $j = k - 1$ . To this end, observe that, for any integer  $c$ ,  $P^{(c,c)}(-t) = (-1)^k P^{(c,c)}(t)$  and so  $P_k^{(c,c)}$  is an even polynomial for even  $k$ , and an odd polynomial for odd  $k$ . Thus,  $P_k^{(c,c)} P_{k-1}^{(c+1,c+1)}$  is an odd polynomial for any  $k$ . Hence

$$\int_{\gamma}^{\delta} P_k^{(c,c)} P_j^{(c+1,c+1)} w_P^{(c+1)} dt = \int_{-\delta}^{\delta} P_k^{(c,c)} P_j^{(c+1,c+1)} w_P^{(1)} w_P^{(c)} dt$$

is zero for  $j < k - 2$  by orthogonality, and is zero for  $j = k - 1$  due to symmetry over the domain. Moreover,  $t P_k^{(c,c)'}(t) P_j^{(c+1,c+1)}(t)$  is also an odd polynomial for

any  $k$  and so

$$\int_{\gamma}^{\delta} t P_k^{(c,c)'}(t) P_j^{(c+1,c+1)}(t) w_P^{(c+1)}(t) dt$$

is zero for  $j = k - 1$  due to symmetry over the domain, and

$$\begin{aligned} & \int_{\gamma}^{\delta} t P_k^{(c,c)'} P_j^{(c+1,c+1)} w_P^{(c+1)} dt \\ &= - \int_{-\delta}^{\delta} P_k^{(c,c)} \frac{d}{dt} [t P_j^{(c+1,c+1)} w_P^{(c+1)}] dt \\ &= - \int_{-\delta}^{\delta} P_k^{(c,c)} [P_j^{(c+1,c+1)} w_P^{(c+1)} + t P_j^{(c+1,c+1)'} w_P^{(c+1)} - 2c t^2 P_j^{(c+1,c+1)} w_P^{(c)}] dt \\ &= - \omega_P^{(c)} \left\langle P_k^{(c,c)}, P_j^{(c+1,c+1)} w_P^{(1)} + t P_j^{(c+1,c+1)'} w_P^{(1)} - 2c t^2 P_j^{(c+1,c+1)} \right\rangle_{w_P^{(c)}} \end{aligned}$$

which is zero for  $j < k - 2$  by orthogonality. Thus, (3.15) is zero for  $j \notin \{k - 2, k\}$ .

Now, using (3.12), integration-by-parts, and noting that the weight  $w_R^{(a,b,2c)}$  is a polynomial degree  $a + b + 2c$  and vanishes at the limits of the integral for positive

parameters  $a, b, c$ , we have that

$$\begin{aligned}
& \int_{\alpha}^{\beta} R_{n-k}^{(a,b,2c+2k+1)'} R_{m-j}^{(a+1,b+1,2c+2j+3)} \rho^{k+j+1} w_R^{(a+1,b+1,2c+2)} dx \\
&= \int_{\alpha}^{\beta} R_{n-k}^{(a,b,2c+2k+1)'} R_{m-j}^{(a+1,b+1,2c+2j+3)} w_R^{(a+1,b+1,2c+k+j+3)} dx \\
&= - \int_{\alpha}^{\beta} R_{n-k}^{(a,b,2c+2k+1)} \frac{d}{dx} \left[ R_{m-j}^{(a+1,b+1,2c+2j+3)} w_R^{(a+1,b+1,2c+k+j+3)} \right] dx \\
&= - \int_{\alpha}^{\beta} R_{n-k}^{(a,b,2c+2k+1)} \left\{ R_{m-j}^{(a+1,b+1,2c+2j+3)'} w_R^{(a+1,b+1,2c+k+j+3)} \right. \\
&\quad + (2c+k+j+3) \rho \rho' w_R^{(a+1,b+1,2c+k+j+1)} R_{m-j}^{(a+1,b+1,2c+2j+3)} \\
&\quad + (b+1) w_R^{(a+1,b,2c+k+j+3)} R_{m-j}^{(a+1,b+1,2c+2j+3)} \\
&\quad \left. - (a+1) w_R^{(a,b+1,2c+k+j+3)} R_{m-j}^{(a+1,b+1,2c+2j+3)} \right\} dx \\
&= - \omega_R^{(a,b,2c+2k+1)} \left\{ \left\langle R_{n-k}^{(a,b,2c+2k+1)}, w_R^{(1,1,j-k+2)} R_{m-j}^{(a+1,b+1,2c+2j+3)'} \right\rangle_{w_R^{(a,b,2c+2k+1)}} \right. \\
&\quad + (2c+k+j+3) \left\langle R_{n-k}^{(a,b,2c+2k+1)}, \rho \rho' w_R^{(1,1,j-k)} R_{m-j}^{(a+1,b+1,2c+2j+3)} \right\rangle_{w_R^{(a,b,2c+2k+1)}} \\
&\quad + (b+1) \left\langle R_{n-k}^{(a,b,2c+2k+1)}, w_R^{(1,0,j-k+2)} R_{m-j}^{(a+1,b+1,2c+2j+3)} \right\rangle_{w_R^{(a,b,2c+2k+1)}} \\
&\quad \left. - (a+1) \left\langle R_{n-k}^{(a,b,2c+2k+1)}, w_R^{(0,1,j-k+2)} R_{m-j}^{(a+1,b+1,2c+2j+3)} \right\rangle_{w_R^{(a,b,2c+2k+1)}} \right\}.
\end{aligned} \tag{3.16}$$

By recalling (3.14) and noting that  $j-k$  is even by the earlier argument, we can see  $\rho \rho' w_R^{(1,1,j-k)}$ ,  $w_R^{(1,0,j-k+2)}$  and  $w_R^{(0,1,j-k+2)}$  are all polynomials, and further that

$$\deg(\rho \rho' w_R^{(1,1,j-k)}) = \deg(w_R^{(1,0,j-k+2)}) = \deg(w_R^{(0,1,j-k+2)}) = 3 + j - k.$$

Hence, by orthogonality, each term in (3.16) is zero for  $m-j+3+j-k <$

$$n - k \iff m < n - 3.$$

Finally,

$$\begin{aligned} & \int_{\alpha}^{\beta} R_{n-k}^{(a,b,2c+2k+1)} R_{m-j}^{(a+1,b+1,2c+2j+3)} \rho^{k+j} \rho' w_R^{(a+1,b+1,2c+2)} dx \\ &= \omega_R^{(a,b,2c+2k+1)} \left\langle R_{n-k}^{(a,b,2c+2k+1)}, \rho \rho' w_R^{(1,1,j-k)} R_{m-j}^{(a+1,b+1,2c+2j+3)} \right\rangle_{w_R^{(a,b,2c+2k+1)}} \end{aligned}$$

which is also zero for  $m < n - 3$ . Thus

$$\left\langle \frac{\partial}{\partial x} H_{n,k}^{(a,b,c)}, H_{m,j}^{(a+1,b+1,c+1)} \right\rangle_{W^{(a+1,b+1,c+1)}} = 0 \quad \text{for } m < n - 3, j \notin \{k - 2, k\},$$

showing that the only possible non-zero coefficients  $c_{m,j}^x$  are when  $m = n - 3, n - 2, n - 1$  and  $j = k - 2, k$  ( $j \leq m$ ).

We can gain the non-zero entries of the weighted differential operators similarly, by noting that for the disk-slice

$$\frac{\partial}{\partial x} W^{(a,b,c)}(x, y) = -a W^{(a-1,b,c)}(x, y) + b W^{(a,b,c)}(x, y) + 2c \rho(x) \rho'(x) W^{(a,b,c-1)} \quad (3.17)$$

$$\frac{\partial}{\partial y} W^{(a,b,c)}(x, y) = -2c y W^{(a,b,c-1)}(x, y) \quad (3.18)$$

and also that

$$\left\langle W^{(a,b,c)} H_{n,k}^{(a,b,c)}, W^{(\tilde{a},\tilde{b},\tilde{c})} H_{m,j}^{(\tilde{a},\tilde{b},\tilde{c})} \right\rangle_{W^{(-\tilde{a},-\tilde{b},-\tilde{c})}} = \left\langle H_{n,k}^{(a,b)}, H_{m,j}^{(\tilde{a},\tilde{b})} \right\rangle_{W^{(a,b,c)}}.$$

□

There exist conversion matrix operators that increment/decrement the parameters, transforming the OPs from one (weighted or non-weighted) parameter space to another.

**Definition 9.** *Define the operator matrices*

$$T^{(a,b,c) \rightarrow (a+1,b+1,c)}, \quad T^{(a,b,c) \rightarrow (a,b,c+1)} \quad \text{and} \quad T^{(a,b,c) \rightarrow (a+1,b+1,c+1)}$$

*for conversion between non-weighted spaces, and*

$$T_W^{(a,b,c) \rightarrow (a-1,b-1,c)}, \quad T_W^{(a,b,c) \rightarrow (a,b,c-1)} \quad \text{and} \quad T_W^{(a,b,c) \rightarrow (a-1,b-1,c-1)}$$

*for conversion between weighted spaces, according to:*

$$\begin{aligned} \mathbb{H}^{(a,b,c)}(x, y) &= \left( T^{(a,b,c) \rightarrow (a+1,b+1,c)} \right)^\top \mathbb{H}^{(a+1,b+1,c)}(x, y) \\ \mathbb{H}^{(a,b,c)}(x, y) &= \left( T^{(a,b,c) \rightarrow (a,b,c+1)} \right)^\top \mathbb{H}^{(a,b,c+1)}(x, y) \\ \mathbb{H}^{(a,b,c)}(x, y) &= \left( T^{(a,b,c) \rightarrow (a+1,b+1,c+1)} \right)^\top \mathbb{H}^{(a+1,b+1,c+1)}(x, y) \\ \mathbb{W}^{(a,b,c)}(x, y) &= \left( T_W^{(a,b,c) \rightarrow (a-1,b-1,c)} \right)^\top \mathbb{W}^{(a-1,b-1,c)}(x, y) \\ \mathbb{W}^{(a,b,c)}(x, y) &= \left( T_W^{(a,b,c) \rightarrow (a,b,c-1)} \right)^\top \mathbb{W}^{(a,b,c-1)}(x, y) \\ \mathbb{W}^{(a,b,c)}(x, y) &= \left( T_W^{(a,b,c) \rightarrow (a-1,b-1,c-1)} \right)^\top \mathbb{W}^{(a-1,b-1,c-1)}(x, y). \end{aligned}$$

**Lemma 6.** *The operator matrices in Definition 9 are sparse, with banded-block-banded structure. More specifically:*

- $T^{(a,b,c) \rightarrow (a+1,b+1,c)}$  has block-bandwidth  $(0, 2)$ , with diagonal blocks.
- $T^{(a,b,c) \rightarrow (a,b,c+1)}$  has block-bandwidth  $(0, 2)$  and sub-block-bandwidth  $(0, 2)$ .
- $T^{(a,b,c) \rightarrow (a+1,b+1,c+1)}$  has block-bandwidth  $(0, 4)$  and sub-block-bandwidth  $(0, 2)$ .
- $T_W^{(a,b,c) \rightarrow (a-1,b-1,c)}$  has block-bandwidth  $(2, 0)$  with diagonal blocks.
- $T_W^{(a,b,c) \rightarrow (a,b,c-1)}$  has block-bandwidth  $(2, 0)$  and sub-block-bandwidth  $(2, 0)$ .
- $T_W^{(a,b,c) \rightarrow (a-1,b-1,c-1)}$  has block-bandwidth  $(4, 0)$  and sub-block-bandwidth  $(2, 0)$ .

*Proof.* We proceed with the case for the non-weighted operators  $T^{(a,b) \rightarrow (a+\tilde{a},b+\tilde{b},c+\tilde{c})}$ , where  $\tilde{a}, \tilde{b}, \tilde{c} \in \{0, 1\}$ . Since  $\{H_{m,j}^{(a+\tilde{a},b+\tilde{b},c+\tilde{c})}\}$  for  $m = 0, \dots, n$ ,  $j = 0, \dots, m$  is an orthogonal basis for any degree  $n$  polynomial, we can expand  $H_{n,k}^{(a,b,c)} = \sum_{m=0}^n \sum_{j=0}^m c_{m,j} H_{m,j}^{(a+\tilde{a},b+\tilde{b},c+\tilde{c})}$ . The coefficients of the expansion are then the entries of the relevant operator matrix. We will show that the only non-zero coefficients are for  $m \geq n - \tilde{a} - \tilde{b} - 2\tilde{c}$ ,  $j \geq k - 2\tilde{c}$  and  $0 \leq j \leq m$ . First, note that

$$c_{m,j} = \left\langle H_{n,k}^{(a,b,c)}, H_{m,j}^{(a+\tilde{a},b+\tilde{b},c+\tilde{c})} \right\rangle_{W^{(a+\tilde{a},b+\tilde{b},c+\tilde{c})}} \left\| H_{m,j}^{(a+\tilde{a},b+\tilde{b},c+\tilde{c})} \right\|_{W^{(a+\tilde{a},b+\tilde{b},c+\tilde{c})}}^{-2}.$$

Then, using the change of variable  $t = \frac{y}{\rho(x)}$ , we have that


$$\begin{aligned}
& \left\langle H_{n,k}^{(a,b,c)}, H_{m,j}^{(a+\tilde{a},b+\tilde{b},c+\tilde{c})} \right\rangle_{W^{(a+\tilde{a},b+\tilde{b},c+\tilde{c})}} \\
&= \omega_R^{(a+\tilde{a},b+\tilde{b},2c+2\tilde{c})} \left\langle R_{n-k}^{(a,b,2c+2k+1)}, \rho(x)^{k+j+1} R_{m-j}^{(a+\tilde{a},b+\tilde{b},2c+2\tilde{c}+2j+1)} \right\rangle_{w_R^{(a+\tilde{a},b+\tilde{b},2c+2\tilde{c})}} \\
&\quad \cdot \omega_P^{(c,\tilde{c})} \left\langle P_k^{(c,c)}, P_j^{(c+\tilde{c},c+\tilde{c})} \right\rangle_{w_P^{(c,\tilde{c})}} \\
&= \omega_R^{(a,b,2c+2k+1)} \left\langle R_{n-k}^{(a,b,2c+2k+1)}, w_R^{(\tilde{a},\tilde{b},2\tilde{c}+j-k)} R_{m-j}^{(a+\tilde{a},b+\tilde{b},2c+2\tilde{c}+2j+1)} \right\rangle_{w_R^{(a,b,2c+2k+1)}} \\
&\quad \cdot \omega_P^{(c)} \left\langle P_k^{(c,c)}, w_P^{(\tilde{c})} P_j^{(c+\tilde{c},c+\tilde{c})} \right\rangle_{w_P^{(c)}}.
\end{aligned}$$

Since  $w_P^{(\tilde{c})}$  is a polynomial degree  $2\tilde{c}$ , we have that the above is then zero for  $j < k - 2\tilde{c}$ . Further, since  $w_R^{(\tilde{a},\tilde{b},2\tilde{c}+j-k)}$  is a polynomial of degree  $\tilde{a} + \tilde{b} + 2\tilde{c} + j - k$ , we have that the above is zero for  $m - j + \tilde{a} + \tilde{b} + 2\tilde{c} + j - k < n - k \iff m < n - \tilde{a} - \tilde{b} - 2\tilde{c}$ .


The sparsity argument for the weighted parameter transformation operators follows similarly.  $\square$

General linear partial differential operators with polynomial variable coefficients can be constructed by composing the sparse representations for partial derivatives, conversion between bases, and Jacobi operators. As a canonical example, we can obtain the matrix operator for the Laplacian  $\Delta$ , that will take us from coefficients for expansion in the weighted space

$$\mathbb{W}^{(1,1,1)}(x, y) = W^{(1,1,1)}(x, y) \mathbb{H}^{(1,1,1)}(x, y)$$



(a) The Laplacian operator  $\Delta_W^{(1,1,1) \rightarrow (1,1,1)}$  (b) A variable coefficient Helmholtz operator



(c) The biharmonic operator  ${}_2\Delta_W^{(2,2,2) \rightarrow (2,2,2)}$

**Figure 3.2:** "Spy" plots of (differential) operator matrices, showing their sparsity. For (1) is the variable coefficient Helmholtz operator  $\Delta_W^{(1,1,1) \rightarrow (1,1,1)}$ .



to coefficients in the non-weighted space  $\mathbb{H}^{(1,1,1)}(x, y)$ . Note that this construction will ensure the imposition of the Dirichlet zero boundary conditions on  $\Omega$ . The matrix operator for the Laplacian we denote  $\Delta_W^{(1,1,1) \rightarrow (1,1,1)}$  acting on the coefficients vector is then given by

$$\Delta_W^{(1,1,1) \rightarrow (1,1,1)} := D_x^{(0,0,0)} W_x^{(1,1,1)} + T^{(0,0,1) \rightarrow (1,1)} D_y^{(0,0,0)} T_W^{(1,1,0) \rightarrow (0,0,0)} W_y^{(1,1,1)}.$$

Importantly, this operator will have banded-block-banded structure, and hence will be sparse, as seen in Figure 3.2.

Another important example is the Biharmonic operator  $\Delta^2$ , where we assume zero Dirichlet and Neumann conditions. To construct this operator, we first note that we can obtain the matrix operator for the Laplacian  $\Delta$  that will take us from coefficients for expansion in the space  $\mathbb{H}^{(0,0,0)}(x, y)$  to coefficients in the space  $\mathbb{H}^{(2,2,2)}(x, y)$ . We denote this matrix operator that acts on the coefficients vector as  $\Delta^{(0,0,0) \rightarrow (2,2,2)}$ , and is given by

$$\Delta^{(0,0,0) \rightarrow (2,2,2)} := D_x^{(1,1,1)} D_x^{(0,0,0)} + T^{(1,1,2) \rightarrow (2,2,2)} D_y^{(1,1,1)} T^{(0,0,1) \rightarrow (1,1,1)} D_y^{(0,0,0)}.$$

Further, we can represent the Laplacian as a map from coefficients in the space  $\mathbb{W}^{(2,2)}$  to coefficients in the space  $\mathbb{H}^{(0,0,0)}$ . Note that a function expanded in the  $\mathbb{W}^{(2,2)}$  basis will satisfy both zero Dirichlet and Neumann boundary conditions on  $\Omega$ . We denote this matrix operator as  $\Delta_W^{(2,2,2) \rightarrow (0,0,0)}$ , and is given by

$$\Delta_W^{(2,2,2) \rightarrow (0,0,0)} := W_x^{(1,1,1)} W_x^{(2,2,2)} + T_W^{(1,1,0) \rightarrow (0,0,0)} W_y^{(1,1,1)} T_W^{(2,2,1) \rightarrow (1,1,1)} W_y^{(2,2,2)}.$$

We can then construct a matrix operator for  $\Delta^2$  that will take coefficients in the space  $\mathbb{W}^{(2,2,2)}$  to coefficients in the space  $\mathbb{H}^{(2,2,2)}$ . Note that any function expanded in the  $\mathbb{W}^{(2,2,2)}$  basis will satisfy both zero Dirichlet and zero Neumann boundary conditions on  $\Omega$ . The matrix operator for the Biharmonic operator is then given by

$${}_2\Delta_W^{(2,2,2)\rightarrow(2,2,2)} = \Delta^{(0,0,0)\rightarrow(2,2,2)} \Delta_W^{(2,2,2)\rightarrow(0,0,0)}.$$

The sparsity and structure of this biharmonic operator are seen in Figure 3.2.

### 3.3 Computational aspects

In this section we discuss how to take advantage of the proposed basis and sparsity structure in partial differential operators in practical computational applications.

#### 3.3.1 Constructing $R_n^{(a,b,c)}(x)$

It is possible to obtain the recurrence coefficients for the  $\{R_n^{(a,b,c)}\}$  OPs in (3.9), by careful application of the Christoffel–Darboux formula [42, 18.2.12]. We explain the process here for the disk-slice case, however we note that a similar but simpler argument holds for the trapezium case. We thus first need to define a new set of ‘interim’ 1D OPs.

**Definition 10.** Let  $w_{\tilde{R}}^{(a,b,c,d)}(x) := (\beta - x)^a (x - \alpha)^b (1 - x)^c (1 + x)^d$  be a weight

function on the interval  $(\alpha, \beta)$ , and define the associated inner product by:

$$\langle p, q \rangle_{w_{\tilde{R}}^{(a,b,c,d)}} := \frac{1}{\omega_{\tilde{R}}^{(a,b,c,d)}} \int_{\alpha}^{\beta} p(x) q(x) w_{\tilde{R}}^{(a,b,c,d)}(x) dx \quad (3.19)$$

where

$$\omega_{\tilde{R}}^{(a,b,c,d)} := \int_{\alpha}^{\beta} w_{\tilde{R}}^{(a,b,c,d)}(x) dx \quad (3.20)$$

Denote the four-parameter family of orthonormal polynomials on  $[\alpha, \beta]$  by  $\{\tilde{R}_n^{(a,b,c,d)}\}$ , orthonormal with respect to the inner product defined in (3.19).

Note that the OPs  $\{R_n^{(a,b,2c)}\}$  are then equivalent to the OPs  $\{\tilde{R}_n^{(a,b,c,c)}\}$ . Let the recurrence coefficients for the OPs  $\{\tilde{R}_n^{(a,b,c,d)}\}$  be given by:

$$x \tilde{R}_n^{(a,b,c,d)}(x) = \tilde{\beta}_n^{(a,b,c,d)} \tilde{R}_{n+1}^{(a,b,c,d)}(x) + \tilde{\alpha}_n^{(a,b,c,d)} \tilde{R}_n^{(a,b,c,d)}(x) + \tilde{\beta}_{n-1}^{(a,b,c,d)} \tilde{R}_{n-1}^{(a,b,c,d)}(x) \quad (3.21)$$

**Proposition 2.** *There exist constants  $\mathcal{C}_n^{(a,b,c,d)}$ ,  $\mathcal{D}_n^{(a,b,c,d)}$  such that*

$$\tilde{R}_n^{(a,b,c+1,d)}(x) = \mathcal{C}_n^{(a,b,c,d)} \sum_{k=0}^n \tilde{R}_k^{(a,b,c,d)}(1) \tilde{R}_k^{(a,b,c,d)}(x) \quad (3.22)$$

$$\tilde{R}_n^{(a,b,c,d+1)}(x) = \mathcal{D}_n^{(a,b,c,d)} \sum_{k=0}^n \tilde{R}_k^{(a,b,c,d)}(-1) \tilde{R}_k^{(a,b,c,d)}(x) \quad (3.23)$$

*Proof.* Fix  $n, m \in \{0, 1, \dots\}$  and without loss of generality, assume  $m \leq n$ . First

recall that

$$\int_{\alpha}^{\beta} \tilde{R}_n^{(a,b,c+1,d)}(x) \tilde{R}_m^{(a,b,c+1,d)}(x) w_{\tilde{R}}^{(a,b,c+1,d)}(x) dx = \delta_{n,m} \omega_{\tilde{R}}^{(a,b,c+1,d)}$$

and define

$$\mathcal{C}_n^{(a,b,c,d)} = \left( \frac{\omega_{\tilde{R}}^{(a,b,c+1,d)}}{\omega_{\tilde{R}}^{(a,b,c,d)} \tilde{R}_n^{(a,b,c,d)}(1) \tilde{R}_{n+1}^{(a,b,c,d)}(1) \tilde{\beta}_n^{(a,b,c,d)}} \right)^{\frac{1}{2}}, \quad (3.24)$$

$$\mathcal{D}_n^{(a,b,c,d)} = (-1)^n \left( \frac{-\omega_{\tilde{R}}^{(a,b,c,d+1)}}{\omega_{\tilde{R}}^{(a,b,c,d)} \tilde{R}_n^{(a,b,c,d)}(-1) \tilde{R}_{n+1}^{(a,b,c,d)}(-1) \tilde{\beta}_n^{(a,b,c,d)}} \right)^{\frac{1}{2}}. \quad (3.25)$$

Now, by the Christoffel–Darboux formula [42, 18.2.12], we have that for any

$x, y \in \mathbb{R}$ ,

$$\sum_{k=0}^n \tilde{R}_k^{(a,b,c,d)}(y) \tilde{R}_k^{(a,b,c,d)}(x) = \tilde{\beta}_n^{(a,b,c,d)} \frac{\tilde{R}_n^{(a,b,c,d)}(x) \tilde{R}_{n+1}^{(a,b,c,d)}(y) - \tilde{R}_{n+1}^{(a,b,c,d)}(x) \tilde{R}_n^{(a,b,c,d)}(y)}{y - x}. \quad (3.26)$$

Then,

$$\begin{aligned}
& \int_{\alpha}^{\beta} \left( [\mathcal{C}_n^{(a,b,c,d)} \sum_{k=0}^n \tilde{R}_k^{(a,b,c,d)}(1) \tilde{R}_k^{(a,b,c,d)}(x)] \right. \\
& \quad \cdot [\mathcal{C}_m^{(a,b,c,d)} \sum_{k=0}^m \tilde{R}_k^{(a,b,c,d)}(1) \tilde{R}_k^{(a,b,c,d)}(x)] w_{\tilde{R}}^{(a,b,c+1,d)}(x) \Big) dx \\
&= \mathcal{C}_n^{(a,b,c,d)} \mathcal{C}_m^{(a,b,c,d)} \tilde{\beta}_n^{(a,b,c,d)} \\
& \quad \cdot \sum_{k=0}^m \int_{\alpha}^{\beta} \left( \tilde{R}_k^{(a,b,c,d)}(1) \tilde{R}_k^{(a,b,c,d)}(x) \right. \\
& \quad \cdot [\tilde{R}_n^{(a,b,c,d)}(x) \tilde{R}_{n+1}^{(a,b,c,d)}(1) - \tilde{R}_{n+1}^{(a,b,c,d)}(x) \tilde{R}_n^{(a,b,c,d)}(1)] w_{\tilde{R}}^{(a,b,c,d)}(x) \Big) dx \\
&= \delta_{m,n} \mathcal{C}_n^{(a,b,c,d)^2} \tilde{\beta}_n^{(a,b,c,d)} \omega_{\tilde{R}}^{(a,b,c,d)} \tilde{R}_n^{(a,b,c,d)}(1) \tilde{R}_{n+1}^{(a,b,c,d)}(1) \\
&= \delta_{m,n} \omega_{\tilde{R}}^{(a,b,c+1,d)}
\end{aligned}$$

using (3.24) and (3.26), showing that the RHS and LHS of (3.22) are equivalent.

Further,

$$\begin{aligned}
& \int_{\alpha}^{\beta} \left( [\mathcal{D}_n^{(a,b,c,d)} \sum_{k=0}^n \tilde{R}_k^{(a,b,c,d)}(-1) \tilde{R}_k^{(a,b,c,d)}(x)] \right. \\
& \quad \cdot [\mathcal{D}_m^{(a,b,c,d)} \sum_{k=0}^m \tilde{R}_k^{(a,b,c,d)}(-1) \tilde{R}_k^{(a,b,c,d)}(x)] w_{\tilde{R}}^{(a,b,c,d+1)}(x) \Big) dx \\
&= -\mathcal{D}_n^{(a,b,c,d)} \mathcal{D}_m^{(a,b,c,d)} \tilde{\beta}_n^{(a,b,c,d)} \\
& \quad \cdot \sum_{k=0}^m \int_{\alpha}^{\beta} \left( \tilde{R}_k^{(a,b,c,d)}(-1) \tilde{R}_k^{(a,b,c,d)}(x) \right. \\
& \quad \cdot [\tilde{R}_n^{(a,b,c,d)}(x) \tilde{R}_{n+1}^{(a,b,c,d)}(-1) - \tilde{R}_{n+1}^{(a,b,c,d)}(x) \tilde{R}_n^{(a,b,c,d)}(-1)] w_{\tilde{R}}^{(a,b,c,d)}(x) \Big) dx \\
&= -\delta_{m,n} \mathcal{D}_n^{(a,b,c,d)^2} \tilde{\beta}_n^{(a,b,c,d)} \omega_{\tilde{R}}^{(a,b,c,d)} \tilde{R}_n^{(a,b,c,d)}(-1) \tilde{R}_{n+1}^{(a,b,c,d)}(-1) \\
&= \delta_{m,n} \omega_{\tilde{R}}^{(a,b,c,d+1)}
\end{aligned}$$

using (3.25) and (3.26), showing that the RHS and LHS of (3.23) are also equivalent.  $\square$

**Proposition 3.** *The recurrence coefficients for the OPs  $\{\tilde{R}_n^{(a,b,c+1,d)}\}$  are given by:*

$$\tilde{\alpha}_n^{(a,b,c+1,d)} = \frac{\tilde{R}_{n+2}^{(a,b,c,d)}(1)}{\tilde{R}_{n+1}^{(a,b,c,d)}(1)} \tilde{\beta}_{n+1}^{(a,b,c,d)} - \frac{\tilde{R}_{n+1}^{(a,b,c,d)}(1)}{\tilde{R}_n^{(a,b,c,d)}(1)} \tilde{\beta}_n^{(a,b,c,d)} + \tilde{\alpha}_{n+1}^{(a,b,c,d)}, \quad (3.27)$$

$$\tilde{\beta}_n^{(a,b,c+1,d)} = \frac{\mathcal{C}_n^{(a,b,c,d)}}{\mathcal{C}_{n+1}^{(a,b,c,d)}} \frac{\tilde{R}_n^{(a,b,c,d)}(1)}{\tilde{R}_{n+1}^{(a,b,c,d)}(1)} \tilde{\beta}_n^{(a,b,c,d)}. \quad (3.28)$$

*The recurrence coefficients for the OPs  $\{\tilde{R}_n^{(a,b,c,d+1)}\}$  are given by:*

$$\tilde{\alpha}_n^{(a,b,c,d+1)} = \frac{\tilde{R}_{n+2}^{(a,b,c,d)}(-1)}{\tilde{R}_{n+1}^{(a,b,c,d)}(-1)} \tilde{\beta}_{n+1}^{(a,b,c,d)} - \frac{\tilde{R}_{n+1}^{(a,b,c,d)}(-1)}{\tilde{R}_n^{(a,b,c,d)}(-1)} \tilde{\beta}_n^{(a,b,c,d)} + \tilde{\alpha}_{n+1}^{(a,b,c,d)}, \quad (3.29)$$

$$\tilde{\beta}_n^{(a,b,c,d+1)} = \frac{\mathcal{D}_n^{(a,b,c,d)}}{\mathcal{D}_{n+1}^{(a,b,c,d)}} \frac{\tilde{R}_n^{(a,b,c,d)}(-1)}{\tilde{R}_{n+1}^{(a,b,c,d)}(-1)} \tilde{\beta}_n^{(a,b,c,d)}. \quad (3.30)$$

*Proof.* First, using (3.22) and (3.26) we have that

$$\begin{aligned} & (1-x) x \tilde{R}_n^{(a,b,c+1,d)}(x) \\ &= \mathcal{C}_n^{(a,b,c,d)} \tilde{\beta}_n^{(a,b,c,d)} x \left[ \tilde{R}_n^{(a,b,c,d)}(x) \tilde{R}_{n+1}^{(a,b,c,d)}(1) - \tilde{R}_{n+1}^{(a,b,c,d)}(x) \tilde{R}_n^{(a,b,c,d)}(1) \right] \\ &= \mathcal{C}_n^{(a,b,c,d)} \tilde{\beta}_n^{(a,b,c,d)} \\ & \quad \cdot \left[ \left( \tilde{\beta}_n^{(a,b,c,d)} \tilde{R}_{n+1}^{(a,b,c,d)}(x) + \tilde{\alpha}_n^{(a,b,c,d)} \tilde{R}_n^{(a,b,c,d)}(x) + \tilde{\beta}_{n-1}^{(a,b,c,d)} \tilde{R}_{n-1}^{(a,b,c,d)}(x) \right) \tilde{R}_{n+1}^{(a,b,c,d)}(1) \right. \\ & \quad \left. - \left( \tilde{\beta}_{n+1}^{(a,b,c,d)} \tilde{R}_{n+2}^{(a,b,c,d)}(x) + \tilde{\alpha}_{n+1}^{(a,b,c,d)} \tilde{R}_{n+1}^{(a,b,c,d)}(x) + \tilde{\beta}_n^{(a,b,c,d)} \tilde{R}_n^{(a,b,c,d)}(x) \right) \tilde{R}_n^{(a,b,c,d)}(1) \right] \end{aligned} \quad (3.31)$$

Next, note that the recurrence coefficients for  $\tilde{R}_n^{(a,b,c+1,d)}(x)$  satisfy

$$\begin{aligned}
& (1-x) x \tilde{R}_n^{(a,b,c+1,d)}(x) \\
&= (1-x) \left[ \tilde{\beta}_n^{(a,b,c+1,d)} \tilde{R}_{n+1}^{(a,b,c+1,d)}(x) + \tilde{\alpha}_n^{(a,b,c+1,d)} \tilde{R}_n^{(a,b,c+1,d)}(x) + \tilde{\beta}_{n-1}^{(a,b,c+1,d)} \tilde{R}_{n-1}^{(a,b,c+1,d)}(x) \right] \\
&= \mathcal{C}_{n+1}^{(a,b,c,d)} \tilde{\beta}_n^{(a,b,c+1,d)} \tilde{\beta}_{n+1}^{(a,b,c,d)} \left( \tilde{R}_{n+1}^{(a,b,c,d)}(x) \tilde{R}_{n+2}^{(a,b,c,d)}(1) - \tilde{R}_{n+2}^{(a,b,c,d)}(x) \tilde{R}_{n+1}^{(a,b,c,d)}(1) \right) \\
&\quad + \mathcal{C}_n^{(a,b,c,d)} \tilde{\alpha}_n^{(a,b,c+1,d)} \tilde{\beta}_n^{(a,b,c,d)} \left( \tilde{R}_n^{(a,b,c,d)}(x) \tilde{R}_{n+1}^{(a,b,c,d)}(1) - \tilde{R}_{n+1}^{(a,b,c,d)}(x) \tilde{R}_n^{(a,b,c,d)}(1) \right) \\
&\quad + \mathcal{C}_{n-1}^{(a,b,c,d)} \tilde{\beta}_{n-1}^{(a,b,c+1,d)} \tilde{\beta}_{n-1}^{(a,b,c,d)} \left( \tilde{R}_{n-1}^{(a,b,c,d)}(x) \tilde{R}_n^{(a,b,c,d)}(1) - \tilde{R}_n^{(a,b,c,d)}(x) \tilde{R}_{n-1}^{(a,b,c,d)}(1) \right)
\end{aligned} \tag{3.32}$$

We can set  $\tilde{\beta}_{-1}^{(a,b,c+1,d)} = 0$ . By comparing coefficients of  $\tilde{R}_{n+2}^{(a,b,c,d)}(x)$  and  $\tilde{R}_{n+1}^{(a,b,c,d)}(x)$  in both (3.31) and (3.32) we obtain the desired recurrence coefficients for the OP  $\tilde{R}_n^{(a,b,c+1,d)}(x)$ . The recurrence coefficients for the OPs  $\tilde{R}_n^{(a,b,c,d+1)}(x)$  are found similarly.  $\square$

**Corollary 2.** *The recurrence coefficients for the OPs  $\{\tilde{R}_n^{(a,b,c+1,d)}\}$  can be written as:*

$$\tilde{\alpha}_n^{(a,b,c+1,d)} = \frac{\tilde{\beta}_{n-1}^{(a,b,c,d)}}{\chi_{n-1}^{(a,b,c,d)}(1)} - \frac{\tilde{\beta}_n^{(a,b,c,d)}}{\chi_n^{(a,b,c,d)}(1)} + \tilde{\alpha}_n^{(a,b,c,d)}, \tag{3.33}$$

$$\tilde{\beta}_n^{(a,b,c+1,d)} = \left( \frac{1 - \tilde{\alpha}_{n+1}^{(a,b,c,d)} - \frac{\tilde{\beta}_n^{(a,b,c,d)}}{\chi_n^{(a,b,c,d)}(1)}}{1 - \tilde{\alpha}_n^{(a,b,c,d)} - \frac{\tilde{\beta}_{n-1}^{(a,b,c,d)}}{\chi_{n-1}^{(a,b,c,d)}(1)}} \right)^{\frac{1}{2}} \tilde{\beta}_n^{(a,b,c,d)}. \tag{3.34}$$

The recurrence coefficients for the OPs  $\{\tilde{R}_n^{(a,b,c,d+1)}\}$  can be written as:

$$\tilde{\alpha}_n^{(a,b,c,d+1)} = \frac{\tilde{\beta}_{n-1}^{(a,b,c,d)}}{\chi_{n-1}^{(a,b,c,d)}(-1)} - \frac{\tilde{\beta}_n^{(a,b,c,d)}}{\chi_n^{(a,b,c,d)}(-1)} + \tilde{\alpha}_n^{(a,b,c,d)}, \quad (3.35)$$

$$\tilde{\beta}_n^{(a,b,c,d+1)} = \left( \frac{-1 + \tilde{\alpha}_{n+1}^{(a,b,c,d)} + \frac{\tilde{\beta}_n^{(a,b,c,d)}}{\chi_n^{(a,b,c,d)}(-1)}}{-1 + \tilde{\alpha}_n^{(a,b,c,d)} + \frac{\tilde{\beta}_{n-1}^{(a,b,c,d)}}{\chi_{n-1}^{(a,b,c,d)}(-1)}} \right)^{\frac{1}{2}} \tilde{\beta}_n^{(a,b,c,d)}. \quad (3.36)$$

where

$$\chi_n^{(a,b,c,d)}(y) := \frac{\tilde{R}_{n+1}^{(a,b,c,d)}(y)}{\tilde{R}_n^{(a,b,c,d)}(y)} \quad (3.37)$$

$$= \frac{1}{\tilde{\beta}_n^{(a,b,c,d)}} \left( y - \tilde{\alpha}_n^{(a,b,c,d)} - \frac{\tilde{\beta}_{n-1}^{(a,b,c,d)}}{\chi_{n-1}^{(a,b,c,d)}(y)} \right), \quad y \in \{-1, 1\}. \quad (3.38)$$

These two propositions allow us to recursively obtain the recurrence coefficients for the OPs  $\{R_{n-k}^{(a,b,2c+2k+1)}\}$  as  $k$  increases to be large.

**Remark:** The Corollary demonstrates that in order to obtain the recurrence coefficients  $\{\alpha_m^{(a,b,2c+2k+1)}\}$ ,  $\{\beta_m^{(a,b,2c+2k+1)}\}$  for some  $m$  and  $k$ , we require that we obtain the recurrence coefficients  $\{\alpha_{m+2}^{(a,b,2c+2(k-1)+1)}\}$ ,  $\{\beta_{m+2}^{(a,b,2c+2(k-1)+1)}\}$ . Thus, for large  $N$ , this recursive method of obtaining the recurrence coefficients requires a large initialisation (i.e. using the Lanczos algorithm to compute the recurrence coefficients  $\{\alpha_n^{(a,b,2c+1)}\}$ ,  $\{\beta_n^{(a,b,2c+1)}\}$  – however, we only need to compute these once, and can store and save this initialisation to disk once computed, for the given values of  $a, b, c$ ).



### 3.3.2 Quadrature rule on the disk-slice

In this section we construct a quadrature rule exact for polynomials in the disk-slice  $\Omega$  that can be used to expand functions in  $H_{n,k}^{(a,b,c)}(x,y)$  when  $\Omega$  is a disk-slice.

**Theorem 3.** *Denote the Gauss quadrature nodes and weight on  $[\alpha, \beta]$  with weight  $(\beta - s)^a (s - \alpha)^b \rho(s)^{2c+1}$  as  $(s_k, w_k^{(s)})$ , and on  $[-1, 1]$  with weight  $(1 - t^2)^c$  as  $(t_k, w_k^{(t)})$ . Define*

$$\begin{aligned} x_{i+(j-1)N} &:= s_j, \quad i, j = 1, \dots, \left\lceil \frac{N+1}{2} \right\rceil, \\ y_{i+(j-1)N} &:= \rho(s_j) t_i, \quad i, j = 1, \dots, \left\lceil \frac{N+1}{2} \right\rceil, \\ w_{i+(j-1)N} &:= w_j^{(s)} w_i^{(t)}, \quad i, j = 1, \dots, \left\lceil \frac{N+1}{2} \right\rceil. \end{aligned}$$

Let  $f(x, y)$  be a polynomial on  $\Omega$ . The quadrature rule is then

$$\iint_{\Omega} f(x, y) W^{(a,b)}(x, y) \, dA \approx \frac{1}{2} \sum_{j=1}^M w_j [f(x_j, y_j) + f(x_j, -y_j)],$$

where  $M = \left\lceil \frac{1}{2}(N+1) \right\rceil^2$ , and the quadrature rule is exact if  $f(x, y)$  is a polynomial of degree  $\leq N$ .

*Proof.* We will use the substitution that

$$x = s, \quad y = \rho(s) t.$$

First, note that, for  $(x, y) \in \Omega$ ,

$$\begin{aligned}
 W^{(a,b,c)}(x, y) &= w_R^{(a,b,2c)}(x) w_P^{(c)}\left(\frac{y}{\rho(x)}\right) \\
 &= w_R^{(a,b,c2)}(s) w_P^{(c)}(t) \\
 &=: V^{(a,b,c)}(s, t), \quad \text{for } (s, t) \in [\alpha, \beta] \times [-1, 1].
 \end{aligned}$$

Let  $f : \Omega \rightarrow \mathbb{R}$ . Define the functions  $f_e, f_o : \Omega \rightarrow \mathbb{R}$  by

$$\begin{aligned}
 f_e(x, y) &:= \frac{1}{2} \left( f(x, y) + f(x, -y) \right), \quad \forall (x, y) \in \Omega \\
 f_o(x, y) &:= \frac{1}{2} \left( f(x, y) - f(x, -y) \right), \quad \forall (x, y) \in \Omega
 \end{aligned}$$

so that  $y \mapsto f_e(x, y)$  for fixed  $x$  is an even function, and  $y \mapsto f_o(x, y)$  for fixed  $x$  is an odd function. Note that if  $f$  is a polynomial, then  $f_e(s, \rho(s)t)$  is a polynomial in  $s \in [\alpha, \beta]$  for fixed  $t$ .

Now, we have that

$$\begin{aligned}
\iint_{\Omega} f_e(x, y) W^{(a,b,c)}(x, y) dy dx &= \int_{\alpha}^{\beta} \int_{-1}^1 f_e(s, \rho(s)t) V^{(a,b,c)}(s, t) \rho(s) dt ds \\
&= \int_{\alpha}^{\beta} w_R^{(a,b,2c+1)}(s) \left( \int_{-1}^1 f_e(s, \rho(s)t) w_P^{(c)}(t) dt \right) ds \\
&\approx \int_{\alpha}^{\beta} w_R^{(a,b,2c+1)}(s) \sum_{i=1}^{M_2} \left( w_i^{(t)} f_e(s, \rho(s)t_i) \right) ds \quad (\star) \\
&\approx \sum_{j=1}^{M_1} \left( w_j^{(s)} \sum_{i=1}^{M_2} \left( w_i^{(t)} f_e(s_j, \rho(s_j)t_i) \right) \right) \quad (\star\star) \\
&= \sum_{k=1}^{M_1 M_2} w_k f_e(x_k, y_k).
\end{aligned}$$

Suppose  $f$  is a polynomial in  $x$  and  $y$  of degree  $N$ , and hence that  $f_e$  is a degree  $\leq N$  polynomial. First, note that the degree of the polynomial given by  $x \mapsto f_e(x, y)$  for fixed  $y$  is  $\leq N$  and the degree of the polynomial given by  $y \mapsto f_e(x, y)$  for fixed  $x$  is  $\leq N$ . Also note that  $s \mapsto f_e(s, \rho(s)t)$  for fixed  $t$  is then a degree  $N$  polynomial (since  $\rho$  is a degree 1 polynomial). Hence, we achieve equality at  $(\star)$  if  $2M_2 - 1 \geq N$  and we achieve equality at  $(\star\star)$  if also  $2M_1 - 1 \geq N$ .

Next, note that

$$\begin{aligned}
\iint_{\Omega} f_o(x, y) W^{(a,b,c)}(x, y) dy dx &= \int_{\alpha}^{\beta} \int_{-1}^1 f_o(s, \rho(s)t) V^{(a,b,c)}(s, t) \rho(s) dt ds \\
&= \int_{\alpha}^{\beta} w_R^{(a,b,2c+1)}(s) \left( \int_{-1}^1 f_o(s, \rho(s)t) w_P^{(c)}(t) dt \right) ds \quad (\dagger) \\
&= 0
\end{aligned}$$

since the inner integral at  $(\dagger)$  over  $t$  is zero, due to the symmetry over the domain.

Hence, for a polynomial  $f$  in  $x$  and  $y$  of degree  $N$ ,

$$\begin{aligned} \iint_{\Omega} f(x, y) W^{(a,b,c)}(x, y) \, dy \, dx &= \iint_{\Omega} \left( f_e(x, y) + f_o(x, y) \right) W^{(a,b,c)}(x, y) \, dy \, dx \\ &= \iint_{\Omega} f_e(x, y) W^{(a,b,c)}(x, y) \, dy \, dx \\ &= \sum_{j=1}^M w_j f_e(x_j, y_j), \end{aligned}$$

where  $M = \left\lceil \frac{1}{2}(N+1) \right\rceil^2$ . □

### 3.3.3 Obtaining the coefficients for expansion of a function on the disk-slice

Fix  $a, b, c \in \mathbb{R}$ . Then for any function  $f : \Omega \rightarrow \mathbb{R}$  we can express  $f$  by

$$f(x, y) \approx \sum_{n=0}^N \mathbb{H}_n^{(a,b,c)}(x, y)^\top \mathbf{f}_n$$

for  $N$  sufficiently large, where

$$\mathbb{H}_n^{(a,b,c)}(x, y) := \begin{pmatrix} H_{n,0}^{(a,b,c)}(x, y) \\ \vdots \\ H_{n,n}^{(a,b,c)}(x, y) \end{pmatrix} \in \mathbb{R}^{n+1} \quad \forall n = 0, 1, 2, \dots, N,$$

and where

$$\mathbf{f}_n := \begin{pmatrix} f_{n,0} \\ \vdots \\ f_{n,n} \end{pmatrix} \in \mathbb{R}^{n+1} \quad \forall n = 0, 1, 2, \dots, N, \quad f_{n,k} := \frac{\langle f, H_{n,k}^{(a,b,c)} \rangle_{W^{(a,b,c)}}}{\|H_{n,k}^{(a,b,c)}\|_{W^{(a,b,c)}}}$$


Recall from (3.7) that  $\|H_{n,k}^{(a,b,c)}\|_{W^{(a,b,c)}}^2 = \omega_R^{(a,b,2c+2k+1)} \omega_P^{(c)}$ . Using the quadrature rule detailed in Section 4.2 for the inner product, we can calculate the coefficients  $f_{n,k}$  for each  $n = 0, \dots, N$ ,  $k = 0, \dots, n$ :

$$f_{n,k} = \frac{1}{2 \omega_R^{(a,b,2c+2k+1)} \omega_P^{(c)}} \sum_{j=1}^M w_j [f(x_j, y_j) H_{n,k}^{(a,b,c)}(x_j, y_j) + f(x_j, -y_j) H_{n,k}^{(a,b,c)}(x_j, -y_j)]$$

where  $M = \lceil \frac{1}{2}(N+1) \rceil^2$ .

### 3.3.4 Calculating non-zero entries of the operator matrices


The proofs of Theorem 2 and Lemma 6 provide a way to calculate the non-zero entries of the operator matrices given in Definition 8 and Definition 9. We can simply use quadrature to calculate the 1D inner products, which has a complexity of  $\mathcal{O}(N^3)$ . This proves much cheaper computationally than using the 2D quadrature rule to calculate the 2D inner products, which has a complexity of  $\mathcal{O}(N^4)$ .



**Figure 3.3:** Left: The computed solution to  $\Delta u = f$  with zero boundary conditions with  $f(x, y) = 1 + \text{erf}(5(1 - 10((x - 0.5)^2 + y^2)))$ . Right: The norms of each block of the computed solution of the Poisson equation with the given right hand side functions. This demonstrates algebraic convergence with the rate dictated by the decay at the corners, with spectral convergence observed when the right-hand side vanishes to all orders.

### 3.4 Examples on the disk-slice with zero Dirichlet conditions

We now demonstrate how the sparse linear systems constructed as above can be used to efficiently solve PDEs with zero Dirichlet conditions. We consider Poisson, inhomogeneous variable coefficient Helmholtz equation and the Biharmonic equation, demonstrating the versatility of the approach.



**Figure 3.4:** The computed solution to  $\Delta u = f$  with zero boundary conditions compared with the exact solution  $u(x, y) = W^{(1,1,1)}(x, y)y^3 \exp(x)$ . Left: Computed. Centre: Exact. Right: Plot of the error (colourbar is shown to demonstrate magnitude of the error is of the order  $10^{-17}$ )


### 3.4.1 Poisson

The Poisson equation is the classic problem of finding  $u(x, y)$  given a function  $f(x, y)$  such that:

$$\begin{cases} \Delta u(x, y) = f(x, y) & \text{in } \Omega \\ u(x, y) = 0 & \text{on } \partial\Omega \end{cases}. \quad (3.39)$$

noting the imposition of zero Dirichlet boundary conditions on  $u$ .

We can tackle the problem as follows. Denote the coefficient vector for expansion of  $u$  in the  $\mathbb{W}^{(1,1,1)}$  OP basis up to degree  $N$  by  $\mathbf{u}$ , and the coefficient vector for




**Figure 3.5:** Left: The computed solution to  $\Delta u + k^2 v u = f$  with zero boundary conditions with  $f(x, y) = x(1 - x^2 - y^2)e^x$ ,  $v(x, y) = 1 - (3(x - 1)^2 + 5y^2)$  and  $k = 100$ . Right: The norms of each block of the computed solution of the Helmholtz equation with the given right hand side functions, with  $k = 20$  and  $v(x, y) = 1 - (3(x - 1)^2 + 5y^2)$ .

expansion of  $f$  in the  $\mathbb{H}^{(1,1,1)}$  OP basis up to degree  $N$  by  $\mathbf{f}$ . Since  $f$  is known, we can obtain  $\mathbf{f}$  using the quadrature rule above. In matrix-vector notation, our system hence becomes:

$$\Delta_W^{(1,1,1) \rightarrow (1,1,1)} \mathbf{u} = \mathbf{f}$$

which can be solved to find  $\mathbf{u}$ . In Figure 3.3 we see the solution to the Poisson equation with zero boundary conditions given in (3.39) in the disk-slice  $\Omega$ . In Figure 3.3 we also show the norms of each block of calculated coefficients of the approximation for four right-hand sides of the Poisson equation with  $N = 990$ , that is, 491,536 unknowns. The rate of decay in the coefficients is a proxy for





**Figure 3.6:** Left: The computed solution to  $\Delta^2 u = f$  with zero Dirichlet and Neumann boundary conditions with  $f(x, y) = 1 + \text{erf}(5(1 - 10((x - 0.5)^2 + y^2)))$ . Right: The norms of each block of the computed solution of the biharmonic equation with the given right hand side functions.

the rate of convergence of the computed solution: as typical of spectral methods, we expect the numerical scheme to converge at the same rate as the coefficients decay. We see that we achieve algebraic convergence for the first three examples, noting that for right hand-sides that vanish at the corners of our disk-slice ( $x \in \{\alpha, \beta\}$ ,  $y = \pm \rho(x)$ ) we observe faster convergence.

In Figure 3.4 we see an example where the solution calculated to the Poisson equation is shown together with a plot of the exact solution and the error. The example was chosen so that the exact solution was  $u(x, y) = W^{(1,1,1)}(x, y)y^3 \exp(x)$ , and thus the RHS function  $f$  would be  $f(x, y) = \Delta[W^{(1,1,1)}(x, y)y^3 \exp(x)]$ . We see that the computed solution is almost exact.

### 3.4.2 Inhomogeneous variable-coefficient Helmholtz

Find  $u(x, y)$  given functions  $v, f : \Omega \rightarrow \mathbb{R}$  such that:

$$\begin{cases} \Delta u(x, y) + k^2 v(x, y) u(x, y) = f(x, y) & \text{in } \Omega \\ u(x, y) = 0 & \text{on } \partial\Omega \end{cases} \quad (3.40)$$

where  $k \in \mathbb{R}$ , noting the imposition of zero Dirichlet boundary conditions on  $u$ .

We can tackle the problem as follows. Denote the coefficient vector for expansion of  $u$  in the  $\mathbb{W}^{(1,1,1)}$  OP basis up to degree  $N$  by  $\mathbf{u}$ , and the coefficient vector for expansion of  $f$  in the  $\mathbb{H}^{(1,1,1)}$  OP basis up to degree  $N$  by  $\mathbf{f}$ . Since  $f$  is known, we can obtain the coefficients  $\mathbf{f}$  using the quadrature rule above. We can obtain the matrix operator for the variable-coefficient function  $v(x, y)$  by using the Clenshaw algorithm with matrix inputs as the Jacobi matrices  $J_x^{(0,0,0)\top}, J_y^{(0,0,0)\top}$ , yielding an operator matrix of the same dimension as the input Jacobi matrices a la the procedure introduced in [45]. We can denote the resulting operator acting on coefficients in the  $\mathbb{H}^{(0,0,0)}$  space by  $V(J_x^{(0,0,0)\top}, J_y^{(0,0,0)\top})$ . In matrix-vector notation, our system hence becomes:

$$(\Delta_W^{(1,1,1) \rightarrow (1,1,1)} + k^2 T^{(0,0,0) \rightarrow (1,1,1)} V(J_x^{(0,0,0)\top}, J_y^{(0,0,0)\top}) T_W^{(1,1,1) \rightarrow (0,0,0)}) \mathbf{u} = \mathbf{f}$$

which can be solved to find  $\mathbf{u}$ . We can see the sparsity and structure of this matrix system in Figure 3.2 with  $v(x, y) = xy^2$  as an example. In Figure 3.5 we see the

solution to the inhomogeneous variable-coefficient Helmholtz equation with zero boundary conditions given in (3.40) in the half-disk  $\Omega$ , with  $k = 100$ ,  $v(x, y) = 1 - (3(x-1)^2 + 5y^2)$  and  $f(x, y) = x(1 - x^2 - y^2)e^x$ . In Figure 3.5 we also show the norms of each block of calculated coefficients of the approximation for four right-hand sides of the inhomogeneous variable-coefficient Helmholtz equation with  $k = 20$  and  $v(x, y) = 1 - (3(x-1)^2 + 5y^2)$  using  $N = 500$ , that is, 125,751 unknowns. The rate of decay in the coefficients is a proxy for the rate of convergence of the computed solution. We see that we achieve algebraic convergence for the first three examples, noting that for right hand sides that vanish at the corners of our disk-slice ( $x \in \{\alpha, \beta\}$ ,  $y = \pm\rho(x)$ ) we see faster convergence.

We can extend this to constant non-zero boundary conditions by simply noting that the problem

$$\begin{cases} \Delta u(x, y) + k^2 v(x, y) u(x, y) = f(x, y) & \text{in } \Omega \\ u(x, y) = c \in \mathbb{R} & \text{on } \partial\Omega \end{cases}$$

is equivalent to letting  $u = \tilde{u} + c$  and solving

$$\begin{cases} \Delta \tilde{u}(x, y) + k^2 v(x, y) \tilde{u}(x, y) = f(x, y) - c k^2 v(x, y) =: g(x, y) & \text{in } \Omega \\ \tilde{u}(x, y) = 0 & \text{on } \partial\Omega. \end{cases}.$$

### 3.4.3 Biharmonic equation

Find  $u(x, y)$  given a function  $f(x, y)$  such that:

$$\begin{cases} \Delta^2 u(x, y) = f(x, y) & \text{in } \Omega \\ u(x, y) = 0, \quad \frac{\partial u}{\partial n}(x, y) = 0 & \text{on } \partial\Omega \end{cases}. \quad (3.41)$$

where  $\Delta^2$  is the Biharmonic operator, noting the imposition of zero Dirichlet and Neumann boundary conditions on  $u$ . In Figure 3.6 we see the solution to the Biharmonic equation (3.41) in the disk-slice  $\Omega$ . In Figure 3.6 we also show the norms of each block of calculated coefficients of the approximation for four right-hand sides of the biharmonic equation with  $N = 500$ , that is, 125,751 unknowns. We see that we achieve algebraic convergence for the first three examples, noting that for right hand sides that vanish at the corners of our disk-slice ( $x \in \{\alpha, \beta\}$ ,  $y = \pm\rho(x)$ ) we see faster convergence.

## 3.5 Other domains

### 3.5.1 End-Disk-Slice

The work in this paper on the disk-slice can be easily transferred to the special-case domain of the end-disk-slice, such as half disks, by which we mean

$$\Omega := \{(x, y) \in \mathbb{R}^2 \mid \alpha < x < \beta, \gamma\rho(x) < y < \delta\rho(x)\}$$

with

$$\left\{ \begin{array}{ll} \alpha & \in (0, 1) \\ \beta & := 1 \\ (\gamma, \delta) & := (-1, 1) \\ \rho(x) & := (1 - x^2)^{\frac{1}{2}}. \end{array} \right.$$

Our 1D weight functions on the intervals  $(\alpha, \beta)$  and  $(\gamma, \delta)$  respectively are then given by:

$$\left\{ \begin{array}{ll} w_R^{(a,b)}(x) & := (x - \alpha)^a \rho(x)^b \\ w_P^{(a)}(x) & := (1 - x^2)^b. \end{array} \right.$$

Note here how we can remove the need for third parameter, which is why we consider this a special case. This will make some calculations easier, and the operator matrices more sparse. The weight  $w_P^{(b)}(x)$  is a still the same ultraspherical weight (and the corresponding OPs are the Jacobi polynomials  $\{P_n^{(b,b)}\}$ ).  $w_R^{(a,b)}(x)$  is the (non-classical) weight for the OPs denoted  $\{R_n^{(a,b)}\}$ . Thus we arrive at the two-parameter family of 2D orthogonal polynomials  $\{H_{n,k}^{(a,b)}\}$  on  $\Omega$  given by, for  $0 \leq k \leq n$ ,  $n = 0, 1, 2, \dots$ ,

$$H_{n,k}^{(a,b)}(x, y) := R_{n-k}^{(a, 2b+2k+1)}(x) \rho(x)^k P_k^{(b,b)}\left(\frac{y}{\rho(x)}\right), \quad (x, y) \in \Omega,$$

orthogonal with respect to the weight

$$\begin{aligned}
 W^{(a,b)}(x, y) &:= w_R^{(a,2b)}(x)w_P^{(b)}\left(\frac{y}{\rho(x)}\right) \\
 &= (x - \alpha)^a (\rho(x)^2 - y^2)^b \\
 &= (x - \alpha)^a (1 - x^2 - y^2)^b, \quad (x, y) \in \Omega.
 \end{aligned}$$

The sparsity of operator matrices for partial differentiation by  $x, y$  as well as for parameter transformations generalise to such end-disk-slice domains. For instance, if we inspect the proof of Lemma 2, we see that it can easily generalise to the weights and domain  $\Omega$  for an end-disk-slice.

In Figure 3.7 we see the solution to the Poisson equation with zero boundary conditions in the half-disk  $\Omega$  with  $(\alpha, \beta) := (0, 1)$ .

### 3.5.2 Trapeziums

We can further extend this work to trapezium shaped domains. Note that for any trapezium there exists an affine map to the canonical trapezium domain that we consider here, given by

$$\Omega := \{(x, y) \in \mathbb{R}^2 \mid \alpha < x < \beta, \gamma\rho(x) < y < \delta\rho(x)\}$$

with


$$\left\{ \begin{array}{ll} (\alpha, \beta) & := (0, 1) \\ (\gamma, \delta) & := (0, 1) \\ \rho(x) & := 1 - \xi x, \quad \xi \in (0, 1) \\ w_R^{(a,b,c)}(x) & := (\beta - x)^a (x - \alpha)^b \rho(x)^c = (1 - x)^a x^b (1 - \xi x)^c \\ w_P^{(a,b)}(x) & := (\delta - x)^a (x - \gamma)^b = (1 - x)^a x^b. \end{array} \right.$$

The weight  $w_P^{(a,b)}(x)$  is the weight for the shifted Jacobi polynomials on the interval  $[0, 1]$ , and hence the corresponding OPs are the shifted Jacobi polynomials  $\{\tilde{P}_n^{(a,b)}\}$ . We note that the shifted Jacobi polynomials relate to the normal Jacobi polynomials by the relationship  $\tilde{P}_n^{(a,b)}(x) = P_n^{(a,b)}(2x - 1)$  for any degree  $n = 0, 1, 2, \dots$  and  $x \in [0, 1]$ .  $w_R^{(a,b,c)}(x)$  is the (non-classical) weight for the OPs we denote  $\{R_n^{(a,b,c)}\}$ . Thus we arrive at the four-parameter family of 2D orthogonal polynomials  $\{H_{n,k}^{(a,b,c,d)}\}$  on  $\Omega$  given by, for  $0 \leq k \leq n$ ,  $n = 0, 1, 2, \dots$ ,

$$H_{n,k}^{(a,b,c,d)}(x, y) := R_{n-k}^{(a,b,c+d+2k+1)}(x) \rho(x)^k \tilde{P}_k^{(d,c)}\left(\frac{y}{\rho(x)}\right), \quad (x, y) \in \Omega,$$

orthogonal with respect to the weight

$$\begin{aligned} W^{(a,b,c,d)}(x, y) &:= w_R^{(a,b,c+d)}(x) w_P^{(d,c)}\left(\frac{y}{\rho(x)}\right) \\ &= (1 - x)^a x^b y^c (1 - \xi x - y)^d, \quad (x, y) \in \Omega. \end{aligned}$$



**Figure 3.7:** Left: The computed solution to  $\Delta u = f$  with zero boundary conditions with  $f(x, y) = W^{(1,1,1)}(x, y)y \cos(x)$  in the disk-slice using the  $p$ -FEM approach with a single element. Centre: The computed solution to  $\Delta u = f$  with zero boundary conditions with  $f(x, y) = 1 + \text{erf}(5(1 - 10((x - 0.5)^2 + y^2)))$  in the half-disk. Right: The computed solution to  $\Delta u + k^2 v u = f$  with zero boundary conditions with  $f(x, y) = (1 - x) x y (1 - \frac{1}{2}x - y) e^x$ ,  $v(x, y) = 1 - (3(x - 1)^2 + 5y^2)$  and  $k = 100$ . in the trapezium.

In Figure 3.7 we see the solution to the Helmholtz equation with zero boundary conditions in the trapezium  $\Omega$  with  $\xi := \frac{1}{2}$ .

### 3.6 P-finite element methods using sparse operators

It is possible for our framework to be applied to a  $p$ -finite element method – that is, one where we can vary the polynomial degree  $p$  of the basis functions used in each element (compare this to a normal  $h$ -FEM, where we can tune the element



size  $h$ ). For example, one could discretise the disk into disk-slice elements, and apply a  $p$ -finite element method to solve PDEs on the disk. As a precursor to this, in this section we limit our discretisation to a single element. Specifically, we follow the method of [5] to construct a sparse  $p$ -finite element method in terms of the operators constructed above, with the benefit of ensuring that the resulting discretisation is symmetric. Consider the 2D Dirichlet problem on a domain  $\Omega$ :

$$\begin{cases} -\Delta u(x, y) = f(x, y) & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega \end{cases}$$

This has the weak formulation for any test function  $v \in V := H_0^1(\Omega) = \{v \in H^1(\Omega) \mid v|_{\partial\Omega} = 0\}$ ,

$$L(v) := \int_{\Omega} f v \, d\mathbf{x} = \int_{\Omega} \nabla u \cdot \nabla v \, d\mathbf{x} =: a(u, v).$$

As eluded to, in general we would let  $\mathcal{T}$  be the set of elements  $\tau$  that make up our finite element discretisation of the domain, where each  $\tau$  is a trapezium or disk slice for example. However, here we simply consider our domain to be a disk-slice and our discretisation to be a single element – that is we let  $\tau = \Omega$  for a disk-slice domain. We can choose our finite dimensional space  $V_p = \{v_p \in V \mid \deg(v_p|_{\tau}) \leq p\}$  for some  $p \in \mathbb{N}$ .

We seek  $u_p \in V_p$  s.t.

$$L(v_p) = a(u_p, v_p) \quad \forall v_p \in V_p. \quad (3.42)$$

Recall that the OPs  $\mathbb{H}^{(a,b,c)}$  are orthogonal with respect to the weight  $W^{(a,b,c)}$  on  $\Omega$ , and define the matrix  $\Lambda^{(a,b,c)} := \left\langle \mathbb{H}^{(a,b,c)}, \mathbb{H}^{(a,b,c)\top} \right\rangle_{W^{(a,b,c)}}$ . Note that due to orthogonality this is a diagonal matrix. We can choose a basis for  $V_p$  by using the weighted orthogonal polynomials on  $\tau$  with parameters  $a = b = 1$ :

$$\mathbb{W}^{(1,1,1)}(x, y) := \begin{pmatrix} \mathbb{W}^{(1,1,1)}_0(x, y) \\ \mathbb{W}^{(1,1,1)}_1(x, y) \\ \mathbb{W}^{(1,1,1)}_2(x, y) \\ \vdots \\ \mathbb{W}^{(1,1,1)}_p(x, y) \end{pmatrix},$$

$$\mathbb{W}^{(1,1,1)}_n(x, y) := W^{(1,1,1)}(x, y) \begin{pmatrix} H_{n,0}^{(1,1,1)}(x, y) \\ \vdots \\ H_{n,n}^{(1,1,1)}(x, y) \end{pmatrix} \in \mathbb{R}^{n+1} \quad \forall n = 0, 1, 2, \dots, p,$$

and rewrite (3.42) in matrix form:

$$\begin{aligned}
a(u_p, v_p) &= \int_{\tau} \nabla u_p \cdot \nabla v_p \, d\mathbf{x} \\
&= \int_{\tau} \begin{pmatrix} \partial_x v_p \\ \partial_y v_p \end{pmatrix}^{\top} \begin{pmatrix} \partial_x u_p \\ \partial_y u_p \end{pmatrix} \, d\mathbf{x} \\
&= \int_{\tau} \begin{pmatrix} \mathbb{H}^{(0,0,0)\top} W_x^{(1,1,1)} \mathbf{v} \\ \mathbb{H}^{(0,0,0)\top} T_W^{(1,1,0) \rightarrow (0,0,0)} W_y^{(1,1,1)} \mathbf{v} \end{pmatrix}^{\top} \begin{pmatrix} \mathbb{H}^{(0,0,0)\top} W_x^{(1,1,1)} \mathbf{u} \\ \mathbb{H}^{(0,0,0)\top} T_W^{(1,1,0) \rightarrow (0,0,0)} W_y^{(1,1,1)} \mathbf{u} \end{pmatrix} \, d\mathbf{x} \\
&= \int_{\tau} \left( \mathbf{v}^{\top} W_x^{(1,1,1)\top} \mathbb{H}^{(0,0,0)} \mathbb{H}^{(0,0,0)\top} W_x^{(1,1,1)} \mathbf{u} \right. \\
&\quad \left. + \mathbf{v}^{\top} (T_W^{(1,1,0) \rightarrow (0,0,0)} W_y^{(1,1,1)})^{\top} \mathbb{H}^{(0,0,0)} \mathbb{H}^{(0,0,0)\top} T_W^{(1,1,0) \rightarrow (0,0,0)} W_y^{(1,1,1)} \mathbf{u} \right) \, d\mathbf{x} \\
&= \mathbf{v}^{\top} \left( W_x^{(1,1,1)\top} \Lambda^{(0,0,0)} W_x^{(1,1,1)} \right. \\
&\quad \left. + (T_W^{(1,1,0) \rightarrow (0,0,0)} W_y^{(1,1,1)})^{\top} \Lambda^{(0,0,0)} T_W^{(1,1,0) \rightarrow (0,0,0)} W_y^{(1,1,1)} \right) \mathbf{u}
\end{aligned}$$

where  $\mathbf{u}, \mathbf{v}$  are the coefficient vectors of the expansions of  $u_p, v_p \in V_p$  respectively in the  $V_p$  basis ( $\mathbb{W}^{(1,1,1)}$  OPs), and

$$\begin{aligned}
L(v_p) &= \int_{\tau} v_p f \, d\mathbf{x} \\
&= \int_{\tau} \mathbf{v}^{\top} \mathbb{W}^{(1,1,1)} \mathbb{H}^{(1,1,1)\top} \mathbf{f} \, d\mathbf{x} \\
&= \mathbf{v}^{\top} \left\langle \mathbb{H}^{(1,1,1)}, \mathbb{H}^{(1,1,1)\top} \right\rangle_{W^{(1,1,1)}} \, d\mathbf{x} \\
&= \mathbf{v}^{\top} \Lambda^{(1,1,1)} \mathbf{f},
\end{aligned}$$

where  $\mathbf{f}$  is the coefficient vector for the expansion of the function  $f(x, y)$  in the  $\mathbb{H}^{(1,1,1)}$  OP basis.

Since (3.42) is equivalent to stating that

$$L(W^{(1,1,1)} H_{n,k}^{(1,1,1)}) = a(u_p, W^{(1,1,1)} H_{n,k}^{(1,1,1)}) \quad \forall n = 0, \dots, p, k = 0, \dots, n,$$

(i.e. holds for all basis functions of  $V_p$ ) by choosing  $v_p$  as each basis function, we can equivalently write the linear system for our finite element problem as:

$$A\mathbf{u} = \tilde{\mathbf{f}}.$$

where the (element) stiffness matrix  $A$  is defined by

$$A = W_x^{(1,1,1)\top} \Lambda^{(0,0,0)} W_x^{(1,1,1)} + (T_W^{(1,1,0) \rightarrow (0,0,0)} W_y^{(1,1,1)})^\top \Lambda^{(0,0,0)} T_W^{(1,1,0) \rightarrow (0,0,0)} W_y^{(1,1,1)}$$

and the load vector  $\tilde{\mathbf{f}}$  is given by

$$\tilde{\mathbf{f}} = \Lambda^{(1,1,1)} \mathbf{f}.$$

Note that since we have sparse operator matrices for partial derivatives and basis-transform, we obtain a symmetric sparse (element) stiffness matrix, as well as a sparse operator matrix for calculating the load vector (rhs).

## 3.7 Conclusions

In this chapter we have shown that bivariate orthogonal polynomials can lead to sparse discretizations of general linear PDEs on specific domains whose boundary is specified by an algebraic curve – notably here the disk-slice – with Dirichlet boundary conditions. This work extends the triangle case [5, 32, 45] and whole disk case [ ] to non-classical geometries, and forms a building block in developing an *hp*-finite element method to solve PDEs on other polygonal domains by using suitably shaped elements (for example, by dividing the disk or a section of the disk into disk-slice elements, which has applications in turbulent pipe flow [ ]).

cite

cite

We have demonstrated how one can construct the three-or-four-parameter OP families (depending on the domain) that form the basis for our sparse spectral methods, and presented a procedure, utilising the Christoffel-Darboux formula [42, 18.2.2], of explicitly calculating the recurrence coefficients for the univariate OPs that are part of the construction. These coefficients contribute to calculations of the entries in the Jacobi matrices and other differential operators. Moreover, we have defined a quadrature rule that can be used for expanding functions in our OP basis and for calculating said entries of important operators. We have looked at a few mathematical examples including the Poisson, variable coefficient Helmholtz, and biharmonic equations and shown that our method performs well. Importantly, all operator matrices used are shown to be sparse, and in fact banded-block-banded in structure.

Looking ahead, this work serves as a stepping stone to constructing similar meth-

ods to solve partial differential equations on sub-domains of the 2-sphere surface, such as spherical caps that we will discuss in the next chapter.

# Chapter 4

## Spherical caps

While the work in the previous chapter looked at developing a sparse spectral method inside a two dimensional domain, we move on to investigating the realm of a surface in three dimensional space. Specifically, we look to extend the methodology to a hierarchy of non-classical multivariate orthogonal polynomials on spherical caps. The entries of discretisations of partial differential operators can be effectively computed using formulae in terms of (non-classical) univariate orthogonal polynomials. We demonstrate the results on partial differential equations involving the spherical Laplacian and biharmonic operators, showing spectral convergence with discretisations that can be made well-conditioned using a simple preconditioner.

Our aim in this chapter is to develop a sparse spectral method for solving linear partial differential equations on certain subsets of the sphere – specifically spherical caps. More precisely, we consider the solution of partial differential equations

on the *spherical cap*  $\Omega$  defined by

$$\Omega := \{(x, y, z) \in \mathbb{R}^3 \mid \alpha < z < \beta, x^2 + y^2 + z^2 = 1\}$$

where  $\alpha \in (-1, 1)$  and  $\beta := 1$ . Simply put, the region of the surface of the 2-sphere where the  $z$ -coordinate range is limited to a sub-interval of  $[-1, 1]$  is what we refer to as the spherical cap.

**Remark:** For simplicity we focus on the case of a spherical cap, though there is an extension to a spherical band by taking  $\beta \in (\alpha, 1)$ . The methods presented here translate to the spherical band case by including the necessary adjustments to the weights and recurrence relations we present in this paper. These adjustments make the construction more involved, which is why they are omitted here, but the approach is essentially the same.

For the spherical cap, we advocate using a basis that is polynomial in cartesian coordinates, that is, polynomial in  $x$ ,  $y$ , and  $z$ , and orthogonal with respect to a prescribed weight: that is, multivariate orthogonal polynomials, whose construction was considered in [47]. Equivalently, we can think of these as polynomials modulo the vanishing ideal  $\{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 + z^2 = 1\}$ , or simply as a linear recombination of spherical harmonics that are orthogonalised on a subset of the sphere. This is in contrast to more standard approaches based on mapping the geometry to a simpler one (e.g., a rectangle or disk) and using orthogonal polynomials in the mapped coordinates (e.g., a basis that is polynomial in the spherical coordinates  $\varphi$  and  $\theta$ ). The benefit of the new approach is that we do not



need to resolve Jacobians, and thereby we can achieve sparse discretisations for partial differential operators, including those with polynomial variable coefficients. Further, we avoid the singular nature at the poles or as  $\alpha$  approaches 0 that such a projection may give, since our new approach yields a smooth polynomial basis for all  $\alpha \in [-1, 1)$ .

There are of course other approaches for solving PDEs on surfaces, such as the closest point method [34, 33]. This involves recasting the PDE as one involving a “closest point” operator in a 3D volume that acts as a “shell” for the surface. While such an approach is useful for other geometries, it is of low order and does not achieve spectral convergence with sparse discretisations. Further, for our domain of interest, the closest point method does not take advantage of any rotational symmetry, and so is unable to achieve optimal complexity with a direct solver.

On the spherical cap, the family of weights we will consider are of the form

$$W^{(a)}(x, y, z) := (z - \alpha)^a, \quad \text{for } (x, y, z) \in \Omega,$$

noting that  $W^{(a)}(x, y, z) = 0$  for  $(x, y, z) \in \partial\Omega$  when  $a > 0$ . The corresponding OPs denoted  $Q_{n,k,i}^{(a)}(x, y, z)$ , where  $n$  denotes the polynomial degree,  $0 \leq k \leq n$  and  $i \in \{0, \min(1, k)\}$ . We define these to be orthogonalised lexicographically, that is,

$$Q_{n,k,i}^{(a)}(x, y, z) = C_{n,k,i} x^{k-i} y^i z^{n-k} + (\text{lower order terms})$$

where  $C_{n,k,i} \neq 0$  and “lower order terms” includes degree  $n$  polynomials of the form  $x^{j-i} y^i z^{n-j}$  where  $j < k$ . The precise normalization arises from their definition in terms of one-dimensional OPs that we will see in Definition ??.

We consider partial differential operators involving the spherical Laplacian (the Laplace–Beltrami operator): in spherical coordinates

$$z = \cos \varphi,$$

$$x = \sin \varphi \cos \theta = \rho(z) \cos \theta,$$

$$y = \sin \varphi \sin \theta = \rho(z) \sin \theta.$$

where  $\rho(z) := \sqrt{1 - z^2}$ , we have

$$\Delta_S = \frac{1}{\sin \varphi} \frac{\partial}{\partial \varphi} \left( \sin \varphi \frac{\partial}{\partial \varphi} \right) + \frac{1}{\sin^2 \varphi} \frac{\partial^2}{\partial \theta^2} = \frac{1}{\rho} \frac{\partial}{\partial \varphi} \left( \rho \frac{\partial}{\partial \varphi} \right) + \frac{1}{\rho^2} \frac{\partial^2}{\partial \theta^2}$$

i.e.  $\Delta_S f(\mathbf{x}) = \Delta f(\frac{\mathbf{x}}{\|\mathbf{x}\|})$  for  $\mathbf{x} := (x, y, z) \in \mathbb{R}^3$ . We do so by considering the component operators  $\rho \frac{\partial}{\partial \varphi}$  and  $\frac{\partial}{\partial \theta}$  applied to OPs with a specific choices of weight so that their discretisation is sparse, see Theorem 4. Sparsity comes from expanding the domain and range of an operator using different choices of the parameter  $a$ , a la the ultraspherical spectral method for intervals [44], triangles [45], disk-slices and trapeziums [58], and the related work on sparse discretisations involving Jacobi polynomials on disks [65] and spheres [66, 31]. Just as we proceeded in Chapter ?? for the disk-slice case in 2D (see also [58]), we will use an integration-by-parts argument to deduce the sparsity structure.

The three-dimensional orthogonal polynomials defined here involve the same non-classical (in fact, semi-classical [35, §5]) 1D OPs as those outlined for the disk-slice, and so methods for calculating these 1D OP recurrence coefficients and integrals have already been outlined in Chapter ?? (see also [58]). In particular, by exploiting the connection with these 1D OPs we can construct discretizations of general partial differential operators of size  $(p+1)^2 \times (p+1)^2$  in  $O(p^3)$  operations, where  $p$  is the total polynomial degree. This clearly compares favourably to proceeding in a naïve approach where one would require  $O(p^6)$  operations.

Note that we consider partial differential operators that are not necessarily rotational invariant: for example, one can use these techniques for Schrödinger operators  $\Delta_S + v(x, y, z)$  where  $v$  is first approximated by a polynomial. A nice feature is that if the partial differential operator is invariant with respect to rotation around the  $z$  axis (e.g., a Schrödinger operator with potential  $v(z)$ ) the discretisation decouples, and can be reordered as a block-diagonal matrix. This improves the complexity further to an optimal  $O(p^2)$ , which is demonstrated in Figure 4.5 with  $v(x, y, z) = \cos z$ .

The code that allows one to produce the numerical examples in this chapter is publicly available as a Julia package<sup>1</sup> to partner the ApproxFun package [43] – however, this package is purely experimental at this stage.

---

<sup>1</sup><https://github.com/snowball13/OrthogonalPolynomialFamilies.jl>

## 4.1 The circle arc

The spherical cap can be thought of as higher dimensional version of the circle arc (a one-dimensional “surface” in two-dimensional space). One can define OPs on the circle in terms of Fourier series, which we write here as orthogonal polynomials in  $x$  and  $y$ :

**Definition 11.** Define the unit circle  $\omega := \{\mathbf{x} = (x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 1\}$ , and define the parameter  $\theta$  for each  $(x, y) \in \omega$  by  $x = \cos \theta$ ,  $y = \sin \theta$ . Define the polynomials  $\{Y_{k,i}\}$  for  $k = 0, 1, \dots$ ,  $i = 0, 1$  on  $(x, y) \in \omega$  by

$$Y_{0,0}(\mathbf{x}) \equiv Y_{0,0}(x, y) := Y_0 =: Y_{0,0}(\theta)$$

$$Y_{k,0}(\mathbf{x}) \equiv Y_{k,0}(x, y) := T_k(x) = \cos k\theta =: Y_{k,0}(\theta)$$

$$Y_{k,1}(\mathbf{x}) \equiv Y_{k,1}(x, y) := y U_{k-1}(x) = \sin k\theta =: Y_{k,1}(\theta)$$

for  $k = 1, 2, 3, \dots$  where  $Y_0 := \frac{\sqrt{2}}{2}$  and  $T_k$ ,  $U_{k-1}$  are the standard Chebyshev polynomials on the interval  $[-1, 1]$ . The  $\{Y_{k,i}\}$  are orthonormal with respect to the inner product

$$\langle p, q \rangle_Y := \frac{1}{\pi} \int_0^{2\pi} p(\mathbf{x}(\theta)) q(\mathbf{x}(\theta)) d\theta.$$

Note that  $Y_0$  is defined so as to ensure orthonormality.

However, for an arc of the unit circle  $\{\mathbf{x} = (x, y) \in \mathbb{R}^2 \mid x \geq h, x^2 + y^2 = 1\}$  for some  $h \in (-1, 1)$  (note we can orientate the arc as we please without loss of generality), we would need to make a modification so that the polynomials are

orthonormal with respect to a different inner product accounting for the truncated domain of the arc:

$$\begin{aligned}\langle p, q \rangle_{Y^h} &:= c^h \int_{-\arccos h}^{\arccos h} p(\mathbf{x}(\theta)) q(\mathbf{x}(\theta)) d\theta \\ &= c^h \int_h^1 \left\{ p(x, \sqrt{1-x^2}) q(x, \sqrt{1-x^2}) \right. \\ &\quad \left. + p(x, -\sqrt{1-x^2}) q(x, -\sqrt{1-x^2}) \right\} \frac{dx}{\sqrt{1-x^2}}\end{aligned}$$

for some normalising constant  $c^h$ . In the same vein, we can choose to construct the arc OPs in the same way as the whole circle OPs – that is, define  $\{T_k^h\}$ ,  $\{U_k^h\}$  as two sets of univariate orthogonal polynomials on the interval  $[h, 1]$ , orthogonal with respect to the weight functions  $(1-x^2)^{-\frac{1}{2}}$  and  $(1-x^2)^{\frac{1}{2}}$  respectively, and then define

$$\begin{aligned}Y_{0,0}^h(\mathbf{x}) &\equiv Y_{0,0}^h(x, y) := Y_0^h =: Y_{0,0}^h(\theta) \\ Y_{k,0}^h(\mathbf{x}) &\equiv Y_{k,0}^h(x, y) := T_k^h(x) =: Y_{k,0}^h(\theta) \\ Y_{k,1}^h(\mathbf{x}) &\equiv Y_{k,1}^h(x, y) := y U_{k-1}^h(x) =: Y_{k,1}^h(\theta)\end{aligned}$$

where  $Y_0^h$  is chosen so that  $\langle Y_0^h, Y_0^h \rangle_{Y^h} = 1$ .

Expand on this. Read the Olver Xu "OPs on quadratic curves paper." where the arc might apply??? Either way, cite this paper in intro (CH1).

## 4.2 Orthogonal polynomials on algebraic curves

This idea of using 1D OPs on a shortened interval to create the modified OPs for a circle arc has been generalised to higher dimensional domains, where a way of constructing orthogonal polynomials on algebraic curves has been presented [47].

Expand on this. Summarise the "on" surfaces approach. Use references in the paper.

## 4.3 Orthogonal polynomials on spherical caps

### 4.3.1 Explicit construction

Using the approach presented for constructing OPs on algebraic curves, we can construct the 3D orthogonal polynomials on the spherical cap or band  $\Omega$  from 1D orthogonal polynomials on the interval  $[\alpha, \beta]$ , and from Fourier series written as orthogonal polynomials in  $x$  and  $y$ . Recall that the spherical cap is defined by

$$\Omega := \{(x, y, z) \in \mathbb{R}^3 \mid \alpha < z < \beta, x^2 + y^2 + z^2 = 1\}.$$

**Proposition 4** ([47]). *Let  $w : (\alpha, \beta) \rightarrow \mathbb{R}$  be a weight function. For  $n = 0, 1, 2, \dots$ , let  $\{r_{n,k}\}$  be polynomials orthogonal with respect to the weight  $\rho(x)^{2k}w(x)$*

where  $0 \leq k \leq n$ . Then the 3D polynomials defined on  $\Omega$  given by

$$Q_{n,k,i}(x, y, z) := r_{n-k,k}(z) \rho(z)^k Y_{k,i}\left(\frac{x}{\rho(z)}, \frac{y}{\rho(z)}\right)$$

for  $i \in 0, 1$ ,  $0 \leq k \leq n$ ,  $n = 0, 1, 2, \dots$  are orthogonal polynomials with respect to the inner product

$$\begin{aligned} \langle p, q \rangle &:= \int_{\Omega} p(x, y, z) q(x, y, z) w(z) dA \\ &= \int_0^{\cos^{-1}(\alpha)} \int_0^{2\pi} \left\{ p(\sin \varphi \cos \theta, \sin \varphi \sin \theta, \cos \varphi) q(\sin \varphi \cos \theta, \sin \varphi \sin \theta, \cos \varphi) \right. \\ &\quad \left. \cdot w(\cos \varphi) \sin \varphi \right\} d\theta d\varphi \\ &= \int_{\alpha}^1 \int_0^{2\pi} p(\rho(z) \cos \theta, \rho(z) \sin \theta, z) q(\rho(z) \cos \theta, \rho(z) \sin \theta, z) w(z) d\theta dz \end{aligned}$$

on  $\Omega$ , where  $dA = \sin \varphi d\theta d\varphi$  is the uniform spherical measure on  $\Omega$ .

For the spherical cap, we can use Proposition 4 to create our one-parameter family of OPs. The univariate OPs that we will choose for the  $r_{n,k}$  polynomials above will be the non-classical  $R^{(a,b)}$  OPs that we defined in Definition ???. Since there is only one boundary for the spherical cap, we will only need to use a two-parameter version<sup>2</sup>). For reference, the family of orthonormal polynomials on  $[\alpha, \beta]$  denoted  $\{R_n^{(a,b)}\}$  are defined such that they are orthonormal with respect to the inner product

$$\langle p, q \rangle_{w_R^{(a,b)}} := \frac{1}{\omega_R^{(a,b)}} \int_{\alpha}^1 p(x) q(x) w_R^{(a,b)}(x) dx \quad (4.1)$$

---

<sup>2</sup>For a spherical band with two boundaries, we would need the three-parameter version

where

$$w_R^{(a,b)}(x) := (x - \alpha)^a \rho(x)^b$$

is the weight function and

$$\omega_R^{(a,b)} := \int_{\alpha}^1 w_R^{(a,b)}(x) \, dx \quad (4.2)$$

is a normalising constant.

We can now define the 3D OPs for the spherical cap.

**Definition 12.** *Define the one-parameter 3D orthogonal polynomials via:*

$$Q_{n,k,i}^{(a)}(x, y, z) := R_{n-k}^{(a,2k)}(z) \rho(z)^k Y_{k,i}\left(\frac{x}{\rho(z)}, \frac{y}{\rho(z)}\right), \quad (x, y, z) \in \Omega. \quad (4.3)$$

By construction,  $\{Q_{n,k,i}^{(a)}\}$  are orthogonal with respect to the inner product

$$\begin{aligned} \langle p, q \rangle_{Q^{(a)}} &:= \int_{\Omega} p(\mathbf{x}, z) q(\mathbf{x}, z) w_R^{(a,0)}(z) \, dA \\ &= \int_{\alpha}^1 \int_0^{2\pi} p(\rho(z) \cos \theta, \rho(z) \sin \theta, z) q(\rho(z) \cos \theta, \rho(z) \sin \theta, z) \, d\theta w_R^{(a,0)}(z) \, dz, \end{aligned}$$

with

$$\left\| Q_{n,k,i}^{(a)} \right\|_{Q^{(a)}}^2 := \left\langle Q_{n,k,i}^{(a)}, Q_{n,k,i}^{(a)} \right\rangle_{Q^{(a)}} = \pi \omega_R^{(a,2k)}. \quad (4.4)$$

A method for obtaining explicit recurrence coefficients and evaluating integrals for



the weight function  $w_R^{(a,b)}(x)$  was established in Chapter ?? (see also [58]). The weight is in fact semi-classical, and is equivalent to a generalized Jacobi weight [35, §5].

### 4.3.2 Jacobi matrices

Recall that we can express the three-term recurrences associated with  $R_n^{(a,b)}$  as

$$xR_n^{(a,b)}(x) = \beta_n^{(a,b)}R_{n+1}^{(a,b)}(x) + \alpha_n^{(a,b)}R_n^{(a,b)}(x) + \beta_{n-1}^{(a,b)}R_{n-1}^{(a,b)}(x) \quad (4.5)$$

where the coefficients are calculable (see Chapter ??). We can use equation (4.5) to determine the 3D recurrences for  $Q_{n,k,i}^{(a)}(x, y, z)$ . Importantly, we can deduce sparsity in the recurrence relationships. We first require the following lemma.

**Lemma 7.** *The following identities hold for  $k = 2, 3, \dots$ ,  $j = 0, 1, \dots$  and  $i, h \in \{0, 1\}$ :*

- 1)  $\int_0^{2\pi} Y_0 Y_{j,h}(\theta) \cos \theta \, d\theta = Y_0 \pi \delta_{0,h} \delta_{1,j}$
- 2)  $\int_0^{2\pi} Y_0 Y_{j,h}(\theta) \sin \theta \, d\theta = Y_0 \pi \delta_{1,h} \delta_{1,j}$
- 3)  $\int_0^{2\pi} Y_{1,i}(\theta) Y_{j,h}(\theta) \cos \theta \, d\theta = \pi \delta_{i,h} (Y_0 \delta_{0,j} + \frac{1}{2} \delta_{2,j})$
- 4)  $\int_0^{2\pi} Y_{1,i}(\theta) Y_{j,h}(\theta) \sin \theta \, d\theta = \pi \delta_{|i-1|,h} ((-1)^{i+1} Y_0 \delta_{0,j} + (-1)^i \frac{1}{2} \delta_{2,j})$
- 5)  $\int_0^{2\pi} Y_{k,i}(\theta) Y_{j,h}(\theta) \cos \theta \, d\theta = \frac{1}{2} \pi \delta_{i,h} (\delta_{k-1,j} + \delta_{k+1,j})$
- 6)  $\int_0^{2\pi} Y_{k,i}(\theta) Y_{j,h}(\theta) \sin \theta \, d\theta = \frac{1}{2} \pi \delta_{|i-1|,h} ((-1)^{i+1} \delta_{k-1,j} + (-1)^i \delta_{k+1,j}).$

*Proof.* Each follows from the definitions of  $Y_{k,i}$  and  $Y_0$ , as well as the trigonometric relationships:

$$2 \cos k\theta \cos \theta = \cos(k-1)\theta + \cos(k+1)\theta$$

$$2 \sin k\theta \cos \theta = \sin(k-1)\theta + \sin(k+1)\theta$$

$$2 \cos k\theta \sin \theta = -\sin(k-1)\theta + \sin(k+1)\theta$$

$$2 \sin k\theta \sin \theta = \cos(k-1)\theta - \cos(k+1)\theta.$$

□

**Lemma 8.** *Define*

$$\eta_k := \begin{cases} 0 & \text{if } k < 0 \\ Y_0 & \text{if } k = 0 \\ \frac{1}{2} & \text{otherwise} \end{cases} \quad (4.6)$$

$Q_{n,k,i}^{(a)}(x, y, z)$  satisfy the following recurrences:

$$\begin{aligned} x Q_{n,k,i}^{(a)}(x, y, z) &= \alpha_{n,k,1}^{(a)} Q_{n-1,k-1,i}^{(a)}(x, y, z) + \alpha_{n,k,2}^{(a)} Q_{n-1,k+1,i}^{(a)}(x, y, z) \\ &\quad + \alpha_{n,k,3}^{(a)} Q_{n,k-1,i}^{(a)}(x, y, z) + \alpha_{n,k,4}^{(a)} Q_{n,k+1,i}^{(a)}(x, y, z) \\ &\quad + \alpha_{n,k,5}^{(a)} Q_{n+1,k-1,i}^{(a)}(x, y, z) + \alpha_{n,k,6}^{(a)} Q_{n+1,k+1,i}^{(a)}(x, y, z), \end{aligned}$$

$$\begin{aligned} y Q_{n,k,i}^{(a)}(x, y, z) &= \beta_{n,k,i,1}^{(a)} Q_{n-1,k-1,|i-1|}^{(a)}(x, y, z) + \beta_{n,k,i,2}^{(a)} Q_{n-1,k+1,|i-1|}^{(a)}(x, y, z) \\ &\quad + \beta_{n,k,i,3}^{(a)} Q_{n,k-1,|i-1|}^{(a)}(x, y, z) + \beta_{n,k,i,4}^{(a)} Q_{n,k+1,|i-1|}^{(a)}(x, y, z) \\ &\quad + \beta_{n,k,i,5}^{(a)} Q_{n+1,k-1,|i-1|}^{(a)}(x, y, z) + \beta_{n,k,i,6}^{(a)} Q_{n+1,k+1,|i-1|}^{(a)}(x, y, z), \end{aligned}$$

$$z Q_{n,k,i}^{(a)}(x, y, z) = \gamma_{n,k,1}^{(a)} Q_{n-1,k,i}^{(a)}(x, y, z) + \gamma_{n,k,2}^{(a)} Q_{n,k,i}^{(a)}(x, y, z) + \gamma_{n,k,3}^{(a)} Q_{n+1,k,i}^{(a)}(x, y, z),$$

for  $(x, y, z) \in \Omega$ , where

$$\begin{aligned}
\alpha_{n,k,1}^{(a)} &:= \eta_{k-1} \left\langle R_{n-k}^{(a,2k)}, R_{n-k}^{(a,2(k-1))} \right\rangle_{w_R^{(a,2k)}}, \\
\alpha_{n,k,2}^{(a)} &:= \eta_k \left\langle R_{n-k}^{(a,2k)}, R_{n-k-2}^{(a,2(k+1))} \right\rangle_{w_R^{(a,2(k+1))}}, \\
\alpha_{n,k,3}^{(a)} &:= \eta_{k-1} \left\langle R_{n-k}^{(a,2k)}, R_{n-k+1}^{(a,2(k-1))} \right\rangle_{w_R^{(a,2k)}}, \\
\alpha_{n,k,4}^{(a)} &:= \eta_k \left\langle R_{n-k}^{(a,2k)}, R_{n-k-1}^{(a,2(k+1))} \right\rangle_{w_R^{(a,2(k+1))}}, \\
\alpha_{n,k,5}^{(a)} &:= \eta_{k-1} \left\langle R_{n-k}^{(a,2k)}, R_{n-k+2}^{(a,2(k-1))} \right\rangle_{w_R^{(a,2k)}}, \\
\alpha_{n,k,6}^{(a)} &:= \eta_k \left\langle R_{n-k}^{(a,2k)}, R_{n-k}^{(a,2(k+1))} \right\rangle_{w_R^{(a,2(k+1))}}, \\
\beta_{n,k,i,j}^{(a)} &:= \begin{cases} -\alpha_{n,k,j}^{(a)} & \text{if } (i=0 \text{ and } j \text{ is odd}) \text{ or } (i=1 \text{ and } j \text{ is even}) \\ \alpha_{n,k,j}^{(a)} & \text{otherwise} \end{cases}, \\
\gamma_{n,k,1}^{(a)} &:= \beta_{n-k-1}^{(a,2k)}, \quad \gamma_{n,k,2}^{(a)} := \alpha_{n-k}^{(a,2k)}, \quad \gamma_{n,k,3}^{(a)} := \beta_{n-k}^{(a,2k)}.
\end{aligned}$$

**Remark:** For  $z$  multiplication, note that different Fourier modes do not interact. This is because multiplication by  $z$  is invariant with respect to rotation around the  $z$ -axis.

*Proof.* The 3-term recurrence for multiplication by  $z$  follows from equation (4.5). For the recurrence for multiplication by  $x$ , since  $\{Q_{m,j,h}^{(a)}\}$  for  $m = 0, \dots, n+1$ ,  $j = 0, \dots, m$ ,  $h = 0, 1$  is an orthogonal basis for any degree  $n+1$  polynomial on

$\Omega$ , we can expand

$$x Q_{n,k,i}^{(a)}(x, y, z) = \sum_{m=0}^{n+1} \sum_{j=0}^m \sum_{h=0}^1 c_{m,j} Q_{m,j,h}^{(a)}(x, y, z).$$

These coefficients are given by

$$c_{m,j} = \left\langle x Q_{n,k,i}^{(a)}, Q_{m,j,h}^{(a)} \right\rangle_{Q^{(a)}} \left\| Q_{m,j,h}^{(a)} \right\|_{Q^{(a)}}^{-2}$$

where we show the non-zero coefficients that result are the  $\alpha_{n,k,1}^{(a)}, \dots, \alpha_{n,k,6}^{(a)}$  in the lemma. Recall from equation (4.4) that  $\left\| Q_{m,j,h}^{(a)} \right\|_{Q^{(a)}}^2 = \pi \omega_R^{(a,2j)}$ . Then for  $m = 0, \dots, n+1, j = 0, \dots, m$ , using a change of variables  $(\cos \theta \sin \varphi, \sin \theta \sin \varphi, \cos \varphi) = (x, y, z)$ :

$$\begin{aligned} & \left\langle x Q_{n,k,i}^{(a)}, Q_{m,j,h}^{(a)} \right\rangle_{Q^{(a)}} \\ &= \int_{\Omega} Q_{n,k,i}^{(a)}(\mathbf{x}, z) Q_{m,j,h}^{(a)}(\mathbf{x}, z) x w_R^{(a,0)}(z) dA \\ &= \left( \int_{\alpha}^1 R_{n-k}^{(a,2k)}(z) R_{m-j}^{(a,2j)}(z) \rho(z)^{k+j+1} w_R^{(a,0)}(z) dz \right) \cdot \left( \int_0^{2\pi} Y_{k,i}(\theta) Y_{j,h}(\theta) \cos \theta d\theta \right) \\ &= \left( \int_{\alpha}^1 R_{n-k}^{(a,2k)}(z) R_{m-j}^{(a,2j)}(z) w_R^{(a,k+j+1)}(z) dz \right) \cdot \left( \int_0^{2\pi} Y_{k,i}(\theta) Y_{j,h}(\theta) \cos \theta d\theta \right) \\ &= \frac{1}{2} \pi \delta_{i,h} (\eta_{k-1} \delta_{k-1,j} + \eta_k \delta_{k+1,j}) \int_{\alpha}^1 R_{n-k}^{(a,2k)}(z) R_{m-j}^{(a,2j)}(z) w_R^{(a,k+j+1)}(z) dz. \end{aligned}$$

where  $\delta_{k,j}$  is the standard Kronecker delta function, using Lemma 7. Similarly, for the recurrence for multiplication by  $y$ , we can expand

$$y Q_{n,k,i}^{(a)}(x, y, z) = \sum_{m=0}^{n+1} \sum_{j=0}^m \sum_{h=0}^1 d_{m,j} Q_{m,j,h}^{(a)}(x, y, z).$$

These coefficients are given by

$$d_{m,j} = \left\langle y Q_{n,k,i}^{(a)}, Q_{m,j,h}^{(a)} \right\rangle_{Q^{(a)}} \left\| Q_{m,j,h}^{(a)} \right\|_{Q^{(a)}}^{-2}$$

where we show the non-zero coefficients that result are the  $\beta_{n,k,1}^{(a)}, \dots, \beta_{n,k,6}^{(a)}$  in the lemma:sc:

$$\begin{aligned} & \left\langle y Q_{n,k,i}^{(a)}, Q_{m,j,h}^{(a)} \right\rangle_{Q^{(a)}} \\ &= \int_{\Omega} Q_{n,k,i}^{(a)}(\mathbf{x}, z) Q_{m,j,h}^{(a)}(\mathbf{x}, z) y w_R^{(a,0)}(z) dA \\ &= \left( \int_{\alpha}^1 R_{n-k}^{(a,2k)}(z) R_{m-j}^{(a,2j)}(z) \rho(z)^{k+j+1} w_R^{(a,0)}(z) dz \right) \cdot \left( \int_0^{2\pi} Y_{k,i}(\theta) Y_{j,h}(\theta) \sin \theta d\theta \right) \\ &= \left( \int_{\alpha}^1 R_{n-k}^{(a,2k)}(z) R_{m-j}^{(a,2j)}(z) w_R^{(a,k+j+1)}(z) dz \right) \cdot \left( \int_0^{2\pi} Y_{k,i}(\theta) Y_{j,h}(\theta) \sin \theta d\theta \right) \\ &= \frac{1}{2} \pi \delta_{|i-1|,h} \left[ (-1)^{i+1} \eta_{k-1} \delta_{k-1,j} + (-1)^i \eta_k \delta_{k+1,j} \right] \int_{\alpha}^1 R_{n-k}^{(a,2k)}(z) R_{m-j}^{(a,2j)}(z) w_R^{(a,k+j+1)}(z) dz. \end{aligned}$$

where again  $\delta_{k,j}$  is the standard Kronecker delta function, and we have used Lemma 7.

□

The recurrences in Lemma 8 lead to (block) Jacobi matrices that correspond to multiplication by  $x$ ,  $y$  and  $z$ . In later sections we will use an ordering of the OPs so that they are grouped by Fourier mode  $k$ , which is convenient for the application of differential and other operators to the vector of coefficients of a given function's expansion (some operators will exploit this ordering for operators where Fourier modes do not interact, and thus will be block-diagonal). Before that, the

ordering we will use in the remainder of this section is convenient for establishing Jacobi operators for multiplication by  $x$ ,  $y$  and  $z$ , and hence building the OPs and importantly obtaining the associated *recurrence coefficient matrices* necessary for efficient function evaluation using the Clenshaw algorithm. In practice, it is simply a matter of converting coefficients between the two orderings. To this end, we define our OP-building ordering as follows. For  $n = 0, 1, 2, \dots$ :

$$\tilde{\mathbb{Q}}_n^{(a)} := \begin{pmatrix} Q_{n,0,0}^{(a)}(x, y, z) \\ Q_{n,1,0}^{(a)}(x, y, z) \\ Q_{n,1,1}^{(a)}(x, y, z) \\ \vdots \\ Q_{n,n,0}^{(a)}(x, y, z) \\ Q_{n,n,1}^{(a)}(x, y, z) \end{pmatrix} \in \mathbb{R}^{2n+1}, \quad \tilde{\mathbb{Q}}^{(a)} := \begin{pmatrix} \tilde{\mathbb{Q}}_0^{(a)} \\ \tilde{\mathbb{Q}}_1^{(a)} \\ \tilde{\mathbb{Q}}_2^{(a)} \\ \vdots \end{pmatrix}$$

and set  $J_x^{(a)}$ ,  $J_y^{(a)}$ ,  $J_z^{(a)}$  as the Jacobi matrices corresponding to

$$\begin{aligned} J_x^{(a)} \tilde{\mathbb{Q}}^{(a)}(x, y, z) &= x \tilde{\mathbb{Q}}^{(a)}(x, y, z), \\ J_y^{(a)} \tilde{\mathbb{Q}}^{(a)}(x, y, z) &= y \tilde{\mathbb{Q}}^{(a)}(x, y, z), \\ J_z^{(a)} \tilde{\mathbb{Q}}^{(a)}(x, y, z) &= z \tilde{\mathbb{Q}}^{(a)}(x, y, z). \end{aligned} \tag{4.7}$$

where

$$J_{x/y/z}^{(a)} = \begin{pmatrix} B_{x/y/z,0}^{(a)} & A_{x/y/z,0}^{(a)} & & & \\ C_{x/y/z,1}^{(a)} & B_{x/y/z,1}^{(a)} & A_{x/y/z,1}^{(a)} & & \\ & C_{x/y/z,2}^{(a)} & B_{x/y/z,2}^{(a)} & A_{x/y/z,2}^{(a)} & \\ & & C_{x/y/z,3}^{(a)} & \ddots & \ddots \\ & & & \ddots & \ddots & \ddots \end{pmatrix}.$$

While  $J_x^{(a)}, J_y^{(a)}, J_z^{(a)}$  are not Jacobi matrices in the classical sense, they are block Jacobi matrices. However, it is useful to simply label them as such. In fact, note that they are in fact *banded-block-banded matrices*:

**Definition 13.** A block matrix  $A$  with blocks  $A_{i,j}$  has block-bandwidths  $(L, U)$  if  $A_{i,j} = 0$  for  $-L \leq j - i \leq U$ , and sub-block-bandwidths  $(\lambda, \mu)$  if all blocks  $A_{i,j}$  are banded with bandwidths  $(\lambda, \mu)$ . A matrix where the block-bandwidths and sub-block-bandwidths are small compared to the dimensions is referred to as a *banded-block-banded matrix*.

Each of these Jacobi matrices are then block-tridiagonal (block-bandwidths  $(1, 1)$ ).



For  $J_x^{(a)}$ , the sub-blocks have sub-block-bandwidths  $(2, 2)$ :

$$\begin{aligned}
 A_{x,n}^{(a)} &:= \begin{pmatrix} 0 & A_{n,0,6}^{(a)} & 0 \\ A_{n,1,5}^{(a)} & \ddots & \ddots \\ & \ddots & \ddots & \ddots \\ & & A_{n,n,5}^{(a)} & 0 & A_{n,n,6}^{(a)} \end{pmatrix} \in \mathbb{R}^{(2n+1) \times (2n+3)}, \quad n = 0, 1, 2, \dots \\
 B_{x,n}^{(a)} &:= \begin{pmatrix} 0 & A_{n,0,4}^{(a)} \\ A_{n,1,3}^{(a)} & \ddots & \ddots \\ & \ddots & \ddots & A_{n,n-1,4}^{(a)} \\ & & A_{n,n,3}^{(a)} & 0 \end{pmatrix} \in \mathbb{R}^{(2n+1) \times (2n+1)} \quad n = 0, 1, 2, \dots \\
 C_{x,n}^{(a)} &:= \begin{pmatrix} 0 & A_{n,0,2}^{(a)} \\ A_{n,1,1}^{(a)} & \ddots & \ddots \\ & \ddots & \ddots & A_{n,n-2,2}^{(a)} \\ & & \ddots & 0 \\ & & & A_{n,n,1}^{(a)} \end{pmatrix} \in \mathbb{R}^{(2n+1) \times (2n-1)}, \quad n = 1, 2, \dots
 \end{aligned}$$

where for  $k = 1, \dots, N$ ,  $n = k, \dots, N$

$$\begin{aligned}
 A_{n,k,j}^{(a)} &:= \begin{pmatrix} \alpha_{n,k,j}^{(a)} & 0 \\ 0 & \alpha_{n,k,j}^{(a)} \end{pmatrix} \in \mathbb{R}^{2 \times 2}, (k \neq 1 \text{ for } j \text{ odd}) \\
 A_{n,0,j}^{(a)} &:= \begin{pmatrix} \alpha_{n,0,j}^{(a)} & 0 \end{pmatrix} \in \mathbb{R}^{1 \times 2}, j \text{ even} \\
 A_{n,1,j}^{(a)} &:= \begin{pmatrix} \alpha_{n,1,j}^{(a)} \\ 0 \end{pmatrix} \in \mathbb{R}^{2 \times 1}, j \text{ odd.}
 \end{aligned}$$

For  $J_y^{(a)}$ , the sub-blocks have sub-block-bandwidths  $(3, 3)$ :

$$\begin{aligned}
 A_{y,n}^{(a)} &:= \begin{pmatrix} 0 & B_{n,0,6}^{(a)} & 0 \\ B_{n,1,5}^{(a)} & \ddots & \ddots \\ & \ddots & \ddots & \ddots \\ & & B_{n,n,5}^{(a)} & 0 & B_{n,n,6}^{(a)} \end{pmatrix} \in \mathbb{R}^{(2n+1) \times (2n+3)}, \quad n = 0, 1, 2, \dots \\
 B_{y,n}^{(a)} &:= \begin{pmatrix} 0 & B_{n,0,4}^{(a)} \\ B_{n,1,3}^{(a)} & \ddots & \ddots \\ & \ddots & \ddots & B_{n,n-1,4}^{(a)} \\ & & B_{n,n,3}^{(a)} & 0 \end{pmatrix} \in \mathbb{R}^{(2n+1) \times (2n+1)} \quad n = 0, 1, 2, \dots \\
 C_{y,n}^{(a)} &:= \begin{pmatrix} 0 & B_{n,0,2}^{(a)} \\ B_{n,1,1}^{(a)} & \ddots & \ddots \\ & \ddots & \ddots & B_{n,n-2,2}^{(a)} \\ & & \ddots & 0 \\ & & & B_{n,n,1}^{(a)} \end{pmatrix} \in \mathbb{R}^{(2n+1) \times (2n-1)}, \quad n = 1, 2, \dots
 \end{aligned}$$

where for  $k = 1, \dots, N$ ,  $n = k, \dots, N$

$$\begin{aligned}
 B_{n,k,j}^{(a)} &:= \begin{pmatrix} 0 & \beta_{n,k,0,j}^{(a)} \\ \beta_{n,k,1,j}^{(a)} & 0 \end{pmatrix} \in \mathbb{R}^{2 \times 2}, (k \neq 1 \text{ for } j \text{ odd}) \\
 B_{n,0,j}^{(a)} &:= \begin{pmatrix} 0 & \beta_{n,0,0,j}^{(a)} \end{pmatrix} \in \mathbb{R}^{1 \times 2}, j \text{ even} \\
 B_{n,1,j}^{(a)} &:= \begin{pmatrix} 0 \\ \beta_{n,1,1,j}^{(a)} \end{pmatrix} \in \mathbb{R}^{2 \times 1}, j \text{ odd}.
 \end{aligned}$$

For  $J_z^{(a)}$ , the sub-blocks are diagonal, i.e. have sub-block-bandwidths  $(0, 0)$ :

$$A_{z,n}^{(a)} := \begin{pmatrix} \Gamma_{n,0,3}^{(a)} & 0 & & & \\ & 0 & \ddots & \ddots & \\ & & \ddots & \ddots & \ddots \\ & & & 0 & \Gamma_{n,n,3}^{(a)} & 0 \end{pmatrix} \in \mathbb{R}^{(2n+1) \times (2n+3)}, \quad n = 0, 1, 2, \dots$$

$$B_{z,n}^{(a)} := \begin{pmatrix} \Gamma_{n,0,2}^{(a)} & & & & \\ & \ddots & & & \\ & & \ddots & & \\ & & & \ddots & \\ & & & & \Gamma_{n,n,2}^{(a)} \end{pmatrix} \in \mathbb{R}^{(2n+1) \times (2n+1)} \quad n = 0, 1, 2, \dots$$

$$C_{z,n}^{(a)} := \begin{pmatrix} \Gamma_{n,0,1}^{(a)} & 0 & & & \\ & 0 & \ddots & \ddots & \\ & & \ddots & \ddots & 0 \\ & & & \ddots & \Gamma_{n,n-1,1}^{(a)} \\ & & & & 0 \end{pmatrix} \in \mathbb{R}^{(2n+1) \times (2n-1)}, \quad n = 1, 2, \dots$$

where for  $k = 1, \dots, N$ ,  $n = k, \dots, N$

$$\Gamma_{n,k,j}^{(a)} := \begin{pmatrix} \gamma_{n,k,j} & 0 \\ 0 & \gamma_{n,k,j} \end{pmatrix} \in \mathbb{R}^{2 \times 2}, \quad (4.8)$$

$$\Gamma_{n,0,j}^{(a)} := \gamma_{n,0,j}^{(a)}. \quad (4.9)$$

Note that the sparsity of the Jacobi matrices (in particular the sparsity of the sub-blocks) comes from the natural sparsity of the three-term recurrences of the

1D OPs and the circular harmonics, meaning that the sparsity is not limited to the specific spherical cap, and would extend to the spherical band.

### 4.3.3 Building the OPs

A multivariate analogue of Clenshaw's algorithm for recursive evaluation of the OPs has been established for methods on other domains in 2D such as the triangle [45] and the disk-slice [58], and we proceed similarly here. Combining each system in equation (4.7) into a block-tridiagonal system, for any  $(x, y, z) \in \Omega$ , yields:

$$\begin{pmatrix} 1 & & & & \\ B_0 - G_0(x, y, z) & A_0 & & & \\ C_1 & B_1 - G_1(x, y, z) & A_1 & & \\ & C_2 & B_2 - G_2(x, y, z) & \ddots & \\ & & \ddots & \ddots & \ddots \end{pmatrix} \tilde{\mathbb{Q}}^{(a)}(x, y, z) = \begin{pmatrix} Q_0^{(a)} \\ 0 \\ 0 \\ 0 \\ \vdots \end{pmatrix},$$

where we note  $Q_0^{(a)} := Q_{0,0,0}^{(a)}(x, y, z) \equiv R_0^{(a,0)} Y_0$ , and for each  $n = 0, 1, 2, \dots$ ,

$$A_n := \begin{pmatrix} A_{x,n}^{(a)} \\ A_{y,n}^{(a)} \\ A_{z,n}^{(a)} \end{pmatrix} \in \mathbb{R}^{3(2n+1) \times (2n+3)}, \quad C_n := \begin{pmatrix} C_{x,n}^{(a)} \\ C_{y,n}^{(a)} \\ C_{z,n}^{(a)} \end{pmatrix} \in \mathbb{R}^{3(2n+1) \times (2n-1)} \quad (n \neq 0),$$

$$B_n := \begin{pmatrix} B_{x,n}^{(a)} \\ B_{y,n}^{(a)} \\ B_{z,n}^{(a)} \end{pmatrix} \in \mathbb{R}^{3(2n+1) \times (2n+1)}, \quad G_n(x, y, z) := \begin{pmatrix} xI_{2n+1} \\ yI_{2n+1} \\ zI_{2n+1} \end{pmatrix} \in \mathbb{R}^{3(2n+1) \times (n+1)}.$$

For each  $n = 0, 1, 2, \dots$  let  $D_n^\top$  be any matrix that is a left inverse of  $A_n$ , i.e. such that  $D_n^\top A_n = I_{2n+3}$ . Multiplying our system by the preconditioner matrix that is given by the block diagonal matrix of the  $D_n^\top$ 's, we obtain a lower triangular system [17, p78], which can be expanded to obtain the recurrence:

$$\begin{cases} \tilde{Q}_{-1}^{(a)}(x, y, z) := 0 \\ \tilde{Q}_0^{(a)}(x, y, z) := Q_0^{(a)} \\ \tilde{Q}_{n+1}^{(a)}(x, y, z) = -D_n^\top (B_n - G_n(x, y, z)) \tilde{Q}_n^{(a)}(x, y, z) - D_n^\top C_n \tilde{Q}_{n-1}^{(a)}(x, y, z), \quad n = 0, 1, 2, \dots \end{cases}$$

Note that we can define an explicit  $D_n^\top$  as follows:

$$D_n^\top := \begin{pmatrix} 0 & 0 & (\Gamma_{n,0,3}^{(a)})^{-1} & & \\ & \ddots & \ddots & & \ddots \\ & & 0 & 0 & (\Gamma_{n,n,3}^{(a)})^{-1} \\ & & & \boldsymbol{\eta}_0^\top & \\ & & & \boldsymbol{\eta}_1^\top & \end{pmatrix} \in \mathbb{R}^{(2n+3) \times 3(2n+1)},$$

for  $n = 1, 2, \dots$  where again  $\Gamma_{n,k,3}^{(a)}$  are defined in equation (4.8) and equation (4.9) for  $k = 0, \dots, n$ , and where  $\boldsymbol{\eta}_0, \boldsymbol{\eta}_1 \in \mathbb{R}^{3(2n+1)}$  with entries given by

$$(\boldsymbol{\eta}_0)_j = \begin{cases} \frac{1}{\beta_{n,n,1,6}^{(a)}} & j = 2(2n+1) \\ \frac{-\beta_{n,n,1,5}^{(a)}}{\beta_{n,n,1,6}^{(a)} \gamma_{n,n-1,3}^{(a)}} & j = 3(2n+1) - 3 \\ 0 & o/w \end{cases}$$

$$(\boldsymbol{\eta}_1)_j = \begin{cases} \frac{1}{\alpha_{n,n,6}^{(a)}} & j = 2n+1 \\ \frac{-\alpha_{n,n,5}^{(a)}}{\alpha_{n,n,6}^{(a)} \gamma_{n,n-1,3}^{(a)}} & j = 3(2n+1) - 2 \text{ and } n > 1 \\ 0 & o/w \end{cases}$$

For  $n = 0$ , we can simply take

$$D_0^\top := \begin{pmatrix} 0 & 0 & \frac{1}{\gamma_{0,0,3}^{(a)}} \\ \frac{1}{\alpha_{0,0,6}^{(a)}} & 0 & 0 \\ 0 & \frac{1}{\beta_{0,0,6}^{(a)}} & 0 \end{pmatrix} \in \mathbb{R}^{3 \times 3}.$$

It follows that we can apply  $D_n^\top$  in  $O(n)$  complexity, and thereby calculate  $\tilde{\mathbb{Q}}_0^{(a)}(x, y, z)$  through  $\tilde{\mathbb{Q}}_n^{(a)}(x, y, z)$  in optimal  $O(n^2)$  complexity.

**Definition 14.** *The recurrence coefficient matrices associated with the OPs  $\{Q_{n,k,i}^{(a)}\}$  are given by the matrices  $A_n, B_n, C_n, D_n^\top$  for  $n = 0, 1, 2, \dots$  defined above.*

## 4.4 Sparse partial differential operators

In this section we will derive the entries of spherical partial differential operators applied to our basis, demonstrating their sparsity in the process. To this end, as alluded to in Section 4.3.2, we introduce new notation for a different ordering of the OP vector, in order to exploit the orthogonality the polynomials  $Y_{k,i}$  will bring

and thus ensure the operators will be block-diagonal. Let  $N \in \mathbb{N}$  and define:

$$\mathbb{Q}_{N,k}^{(a)} := \begin{pmatrix} Q_{k,k,0}^{(a)}(x, y, z) \\ Q_{k,k,1}^{(a)}(x, y, z) \\ \vdots \\ Q_{N,k,0}^{(a)}(x, y, z) \\ Q_{N,k,1}^{(a)}(x, y, z) \end{pmatrix} \in \mathbb{R}^{2(N-k+1)}, \quad k = 1, \dots, N, \quad (4.10)$$

$$\mathbb{Q}_{N,0}^{(a)} := \begin{pmatrix} Q_{0,0,0}^{(a)}(x, y, z) \\ \vdots \\ Q_{N,0,0}^{(a)}(x, y, z) \end{pmatrix} \in \mathbb{R}^{N+1}, \quad (4.11)$$

$$\mathbb{Q}_N^{(a)} := \begin{pmatrix} \mathbb{Q}_{N,0}^{(a)} \\ \vdots \\ \mathbb{Q}_{N,N}^{(a)} \end{pmatrix} \in \mathbb{R}^{(N+1)^2} \quad (4.12)$$

We further denote the weighted set of OPs on  $\Omega$  by

$$\mathbb{W}_N^{(a)}(x, y, z) := w_R^{(a,0)}(z) \mathbb{Q}_N^{(a)}(x, y, z),$$

The operator matrices we derive here act on coefficient vectors, that represent a function  $f(x, y, z)$  defined on  $\Omega$  in spectral space – such a function is approximated by its expansion up to degree  $N$ :

$$f(x, y, z) = \mathbb{Q}_N^{(a)}(x, y, z)^\top \mathbf{f} = \sum_{n=0}^N \sum_{k=0}^n \sum_{i=0}^1 f_{n,k,i} Q_{n,k,i}^{(a)}(x, y, z),$$



where  $\mathbf{f} = (f_{n,k,i})$  is the coefficients vector for the function  $f$ .

**Definition 15.** Let  $a$  be a nonnegative parameter, and  $\tilde{a} \geq 2$  be an integer. Define the operator matrices  $D_\varphi^{(a)}$ ,  $W_\varphi^{(a)}$ ,  $D_\theta$ ,  $\mathcal{L}^{(a) \rightarrow (a+\tilde{a})}$ ,  $\mathcal{L}_W^{(a) \rightarrow (a-\tilde{a})}$ ,  $\Delta_W^{(1)}$  according to:

$$\begin{aligned} \rho \frac{\partial f}{\partial \varphi}(x, y, z) &= \mathbb{Q}_N^{(a+1)}(x, y, z)^\top D_\varphi^{(a)} \mathbf{f}, \\ \rho \frac{\partial}{\partial \varphi}[w_R^{(a,0)}(z) f(x, y, z)] &= \mathbb{W}_N^{(a-1)}(x, y)^\top W_\varphi^{(a)} \mathbf{f}, \\ \frac{\partial f}{\partial \theta}(x, y, z) &= \mathbb{Q}_N^{(a)}(x, y, z)^\top D_\theta \mathbf{f}, \\ \Delta_S f(x, y, z) &= \mathbb{Q}_N^{(a+\tilde{a})}(x, y, z)^\top \mathcal{L}^{(a) \rightarrow (a+\tilde{a})} \mathbf{f}, \\ \Delta_S(w_R^{(a,0)}(z) f(x, y, z)) &= \mathbb{W}_N^{(a-\tilde{a})}(x, y, z)^\top \mathcal{L}_W^{(a) \rightarrow (a-\tilde{a})} \mathbf{f}, \quad (\text{for } a \geq 2 \text{ only}) \\ \Delta_S(w_R^{(1,0)}(z) f(x, y, z)) &= \mathbb{Q}_N^{(1)}(x, y, z)^\top \Delta_W^{(1)} \mathbf{f}, \quad (\text{for } a = 1 \text{ only}) \end{aligned}$$

The incrementing and decrementing of parameters as seen here is analogous to other well known orthogonal polynomial families' derivatives, for example the Jacobi polynomials on the interval, as seen in the DLMF [42, (18.9.3)], on the triangle [46], and on the disk-slice [58]. The operators we define here are for partial derivatives with respect to the spherical coordinates  $(\varphi, \theta)$ , so that we can more easily apply the operators to PDEs on the surface of a sphere (for example, surface Laplacian operator in the Poisson equation). With the OP ordering by Fourier mode  $k$  defined in equations (4.10–4.12) these rotationally invariant operators are block-diagonal, meaning simple and parallelisable practical application.

**Theorem 4.** The operator matrices  $D_\varphi^{(a)}$ ,  $W_\varphi^{(a)}$ ,  $D_\theta$ ,  $\mathcal{L}^{(a) \rightarrow (a+\tilde{a})}$ ,  $\mathcal{L}_W^{(a) \rightarrow (a-\tilde{a})}$ ,  $\Delta_W^{(1)}$  from Definition 15 are sparse, with banded-block-banded structure. More specifically:

- $D_\varphi^{(a)}$  is block-diagonal with sub-block-bandwidths  $(2, 4)$
- $W_\varphi^{(a)}$  is block-diagonal with sub-block-bandwidths  $(4, 2)$
- $D_\theta$  is block-diagonal with sub-block-bandwidths  $(1, 1)$
- $\mathcal{L}^{(a) \rightarrow (a+\tilde{a})}$  is block-diagonal with sub-block-bandwidths  $(0, 4)$
- $\mathcal{L}_W^{(a) \rightarrow (a-\tilde{a})}$  is block-diagonal with sub-block-bandwidths  $(4, 0)$
- $\Delta_W^{(1)}$  is block-diagonal with sub-block-bandwidths  $(2, 2)$

In order to show the last part of Theorem 4, we require the following short lemma.

**Lemma 9.** *For any general parameter  $a$  and any  $n = 0, 1, \dots$ ,  $k = 0, \dots, n$  we have that*

$$\begin{aligned}
 & \frac{d}{dz} [w_R^{(a+1, 2(k+1))} R_{n-k}^{(a, 2k)'}] \\
 &= w_R^{(a+1, 2(k+1))} R_{n-k}^{(a, 2k)''} - 2(k+1)z w_R^{(a+1, 2k)} R_{n-k}^{(a, 2k)'} + (a+1)w_R^{(a, 2(k+1))} R_{n-k}^{(a, 2k)'} \\
 &= \sum_{m=n-1}^{n+1} c_{m,k} w_R^{(a, 2k)} R_{m-k}^{(a, 2k)}
 \end{aligned}$$

where

$$c_{m,k} = -\frac{1}{\omega_R^{(a, 2k)}} \int_\alpha^1 R_{n-k}^{(a, 2k)'} R_{m-k}^{(a, 2k)'} w_R^{(a+1, 2(k+1))} dz$$

*Proof of Lemma 9.* Since  $\frac{d}{dz} [w_R^{(a+1, 2(k+1))} R_{n-k}^{(a, 2k)'}] = w_R^{(a, 2k)} r_{n-k+1}$  where  $r_{n-k+1}$  is

a degree  $n - k + 1$  polynomial, we have that

$$\frac{d}{dz} [w_R^{(a+1,2(k+1))} R_{n-k}^{(a,2k)'}] = \sum_{m=0}^{n-k+1} \tilde{c}_{\{n,k\},m} w_R^{(a,2k)} R_m^{(a,2k)}$$

for some coefficients  $\tilde{c}_{\{n,k\},m}$ . These coefficients are given by

$$\begin{aligned} \tilde{c}_{\{n,k\},m} &= \frac{1}{\omega_R^{(a,2k)}} \left\langle \frac{d}{dz} [w_R^{(a+1,2(k+1))} R_{n-k}^{(a,2k)'}], R_m^{(a,2k)} \right\rangle_{w_R^{(0,0)}} \\ &= -\frac{1}{\omega_R^{(a,2k)}} \int_{\alpha}^1 R_{n-k}^{(a,2k)'} R_m^{(a,2k)'} w_R^{(a+1,2(k+1))} dz \end{aligned}$$

We show that these are zero for  $m < n - k - 1$  by integrating twice by parts:

$$\begin{aligned} &\left\langle \frac{d}{dz} [w_R^{(a+1,2(k+1))} R_{n-k}^{(a,2k)'}], R_m^{(a,2k)} \right\rangle_{w_R^{(0,0)}} \\ &= -\int_{\alpha}^1 R_{n-k}^{(a,2k)'} R_{m-k}^{(a,2k)'} w_R^{(a+1,2(k+1))} dz \\ &= \int_{\alpha}^1 R_{n-k}^{(a,2k)'} [(a+1)R_m^{(a,2k)'} w_R^{(0,2)} \\ &\quad - 2(k+1)z R_m^{(a,2k)'} w_R^{(1,0)} + R_m^{(a,2k)''} w_R^{(1,2)}] w_R^{(a,2k)} dz \end{aligned}$$

which is indeed zero for  $m < n - k - 1$  by orthogonality.  $\square$

*Proof of Theorem 4.* For the operator  $D_{\theta}$  for partial differentiation by  $\theta$ , we sim-

ply have that

$$\begin{aligned} \frac{\partial}{\partial \theta} Q_{n,k,i}^{(a)}(x, y, z) &= R_{n-k}^{(a,2k)}(z) \rho(z)^k \frac{d}{d\theta} Y_{k,i}(\theta) \\ &= \begin{cases} (-1)^{i+1} k Q_{n,k,|i-1|}^{(a)}(x, y, z) & k > 0 \\ 0 & k = 0 \end{cases}. \end{aligned}$$

We now proceed with the case for the operator  $D_\varphi^{(a)}$  for partial differentiation by  $\varphi$ . The entries of the operator are given by the coefficients in the expansion

$$\rho \frac{\partial}{\partial \varphi} Q_{n,k,i}^{(a)} = \sum_{m=0}^{n+1} \sum_{j=0}^m \sum_{h=0}^1 c_{m,j,h} Q_{m,j,h}^{(a+1)},$$

where the coefficients are

$$c_{m,j,h} = \left\| Q_{m,j,h}^{(a+1)} \right\|_{Q^{(a+1)}}^{-2} \left\langle \rho \frac{\partial}{\partial \varphi} Q_{n,k,i}^{(a)}, Q_{m,j,h}^{(a+1)} \right\rangle_{Q^{(a+1)}}.$$

Now, note that:

$$\begin{aligned} w_R^{(a,b)'}(z) &= a w_R^{(a-1,b)}(z) + c \rho(z) \rho'(z) w_R^{(a,b-2)}(z), \\ \rho(z) \rho'(z) &= -z, \\ \frac{\partial}{\partial \varphi} Q_{n,k,i}^{(a)}(x, y, z) &= -\rho(z) \frac{d}{dz} \left[ \rho(z)^k R_{n-k}^{(a,2k)}(z) \right] Y_{k,i}(\theta), \\ \frac{\partial}{\partial \varphi} \left[ w_R^{(a,0)}(z) Q_{n,k,i}^{(a)}(x, y, z) \right] &= -\rho(z) \frac{d}{dz} \left[ w_R^{(a,k)}(z) R_{n-k}^{(a,2k)}(z) \right] Y_{k,i}(\theta). \end{aligned}$$

Then,

$$\begin{aligned}
& \left\langle \rho \frac{\partial}{\partial \varphi} Q_{n,k,i}^{(a)}, Q_{m,j,h}^{(a+1)} \right\rangle_{Q^{(a+1)}} \\
&= - \int_{\alpha}^1 \left( \int_0^{2\pi} \rho(z)^2 \frac{d}{dz} [R_{n-k}^{(a,2k)}(z) \rho(z)^k] R_{m-j}^{(a+1,2j)}(z) \rho(z)^j Y_{k,i}(\theta) Y_{j,h}(\theta) d\theta \right) w_R^{(a+1,0)} dz \\
&= \left( \int_0^{2\pi} Y_{k,i}(\theta) Y_{j,h}(\theta) d\theta \right) \left( \int_{\alpha}^1 R_{m-j}^{(a+1,2j)} [kz R_{n-k}^{(a,2k)} - \rho^2 R_{n-k}^{(a,2k)'}] w_R^{(a+1,k+j)} dz \right) \\
&= \pi \delta_{k,j} \delta_{i,h} \int_{\alpha}^1 R_{m-k}^{(a+1,2k)} [kz R_{n-k}^{(a,2k)} - \rho^2 R_{n-k}^{(a,2k)'}] w_R^{(a+1,2k)} dz \\
&= \pi \delta_{k,j} \delta_{i,h} \int_{\alpha}^1 R_{n-k}^{(a,2k)} \left\{ kz R_{m-k}^{(a+1,2k)} w_R^{(1,0)} + R_{m-k}^{(a+1,2k)'} w_R^{(1,2)} \right. \\
&\quad \left. + a \rho^2 R_{m-k}^{(a+1,2k)} - (2k+2)z R_{m-k}^{(a+1,2k)} w_R^{(1,0)} \right\} w_R^{(a,2k)} dz
\end{aligned}$$

which is zero for  $j \neq k$ ,  $h \neq i$ , and  $m < n - 2$  by orthogonality.

Similarly for the operator  $W_{\varphi}^{(a)}$  for partial differentiation by  $\varphi$  on the weighted space, the entries of the operator are given by the coefficients in the expansion  $\rho \frac{\partial}{\partial \varphi} (w_R^{(a,0)} Q_{n,k,i}^{(a)}) = \sum_{m=0}^{n+2} \sum_{j=0}^m \sum_{h=0}^1 c_{m,j,h} w_R^{(a-1,0)} Q_{m,j,h}^{(a-1)}$ , where the coefficients are

$$c_{m,j,h} = \left\| Q_{m,j,h}^{(a-1)} \right\|_{Q^{(a-1)}}^{-2} \left\langle \rho \frac{\partial}{\partial \varphi} (w_R^{(a,0)} Q_{n,k,i}^{(a)}), Q_{m,j,h}^{(a-1)} \right\rangle_{Q^{(0)}}.$$

Now,

$$\begin{aligned}
& \left\langle \rho \frac{\partial}{\partial \varphi} (w_R^{(a,0)} Q_{n,k,i}^{(a)}, Q_{m,j,h}^{(a-1)}) \right\rangle_{Q^{(0)}} \\
&= - \int_{\alpha}^1 \left( \int_0^{2\pi} \rho(z)^2 \frac{d}{dz} [R_{n-k}^{(a,2k)}(z) w_R^{(a,k)}(z)] R_{m-j}^{(a-1,2j)}(z) \rho(z)^j Y_{k,i}(\theta) Y_{j,h}(\theta) d\theta \right) dz \\
&= \left( \int_0^{2\pi} Y_{k,i}(\theta) Y_{j,h}(\theta) d\theta \right) \\
&\quad \cdot \left( \int_{\alpha}^1 R_{m-j}^{(a-1,2j)} [kz R_{n-k}^{(a,2k)} w_R^{(1,0)} - R_{n-k}^{(a,2k)'} w_R^{(1,2)} - a R_{n-k}^{(a,2k)} \rho^2] w_R^{(a-1,k+j)} dz \right) \\
&= \pi \delta_{k,j} \delta_{i,h} \int_{\alpha}^1 R_{m-k}^{(a-1,2k)} [kz R_{n-k}^{(a,2k)} w_R^{(1,0)} - R_{n-k}^{(a,2k)'} w_R^{(1,2)} - a R_{n-k}^{(a,2k)} \rho^2] w_R^{(a-1,2k)} dz \\
&= \pi \delta_{k,j} \delta_{i,h} \int_{\alpha}^1 R_{n-k}^{(a,2k)} \left\{ kz R_{m-k}^{(a-1,2k)} w_R^{(1,0)} - a \rho^2 R_{m-k}^{(a-1,2k)} + R_{m-k}^{(a-1,2k)'} w_R^{(1,2)} \right. \\
&\quad \left. + a \rho^2 R_{m-k}^{(a-1,2k)} - (2k+2)z R_{m-k}^{(a-1,2k)} w_R^{(1,0)} \right\} w_R^{(a-1,2k)} dz \\
&= \pi \delta_{k,j} \delta_{i,h} \int_{\alpha}^1 R_{n-k}^{(a,2k)} \left\{ kz R_{m-k}^{(a-1,2k)} + R_{m-k}^{(a-1,2k)'} \rho^2 - (2k+2)z R_{m-k}^{(a-1,2k)} \right\} w_R^{(a,2k)} dz
\end{aligned}$$

which is zero for  $j \neq k$ ,  $h \neq i$ , and  $m < n - 1$  by orthogonality.

We move on to the spherical Laplacian operators. Note that the Laplacian acting

on the weighted and non-weighted spherical cap OP  $Q_{n,k,i}^{(a)}$  yield

$$\begin{aligned}\Delta_S Q_{n,k,i}^{(a)} &= \frac{1}{\rho} \frac{\partial}{\partial \varphi} \left( \rho \frac{\partial}{\partial \varphi} [R_{n-k}^{(a,2k)}(\cos \varphi) \sin^k \varphi] \right) Y_{k,i}(\theta) \\ &\quad + R_{n-k}^{(a,2k)}(\cos \varphi) \sin^{k-2} \varphi \frac{\partial^2}{\partial \theta^2} Y_{k,i}(\theta) \\ &= Y_{k,i}(\theta) \rho(z)^k \left\{ -k(k+1) R_{n-k}^{(a,2k)}(z) - 2(k+1)z R_{n-k}^{(a,2k)'}(z) \right. \\ &\quad \left. + \rho(z)^2 R_{n-k}^{(a,2k)''}(z) \right\},\end{aligned}\tag{4.13}$$

$$\begin{aligned}\Delta_S (w_R^{(a,0)} Q_{n,k,i}^{(a)}) &= \frac{1}{\rho} \frac{\partial}{\partial \varphi} \left( \rho \frac{\partial}{\partial \varphi} [w_R^{(a,0)}(\cos \varphi) R_{n-k}^{(a,2k)}(\cos \varphi) \sin^k \varphi] \right) Y_{k,i}(\theta) \\ &\quad + w_R^{(a,0)}(\cos \varphi) R_{n-k}^{(a,2k)}(\cos \varphi) \sin^{k-2} \varphi \frac{\partial^2}{\partial \theta^2} Y_{k,i}(\theta) \\ &= Y_{k,i}(\theta) \left\{ R_{n-k}^{(a,2k)}(z) [-k(k+1)w_R^{(a,k)}(z) - 2a(k+1)z w_R^{(a-1,k)}(z)] \right. \\ &\quad + a(a-1)R_{n-k}^{(a,2k)}(z) w_R^{(a-2,k+1)}(z) \\ &\quad + R_{n-k}^{(a,2k)'}(z) [-2(k+1)z w_R^{(a,k)}(z) + 2a w_R^{(a-1,k+2)}(z)] \\ &\quad \left. + R_{n-k}^{(a,2k)''}(z) w_R^{(a,k+2)}(z) \right\}.\end{aligned}\tag{4.14}$$

For the operator  $\mathcal{L}^{(a) \rightarrow (a+\tilde{a})}$  for the surface Laplacian on a non-weighted space, the entries of the operator are given by the coefficients in the expansion

$$\Delta_S Q_{n,k,i}^{(a)} = \sum_{m=0}^n \sum_{j=0}^m \sum_{h=0}^1 c_{m,j,h} Q_{m,j,h}^{(a+\tilde{a})},$$

where the coefficients are

$$c_{m,j,h} = \left\| Q_{m,j,h}^{(a+\tilde{a})} \right\|_{Q^{(a+\tilde{a})}}^{-2} \left\langle \Delta_S Q_{n,k,i}^{(a)}, Q_{m,j,h}^{(a+\tilde{a})} \right\rangle_{Q^{(a+\tilde{a})}}.$$

Using equation (4.13), and integrating by parts twice, we then have that

$$\begin{aligned}
& \left\langle \Delta_S Q_{n,k,i}^{(a)}, Q_{m,j,h}^{(a+\tilde{a})} \right\rangle_{Q^{(a+\tilde{a})}} \\
&= \left( \int_0^{2\pi} Y_{k,i}(\theta) Y_{j,h}(\theta) d\theta \right) \\
&\quad \cdot \left( \int_\alpha^1 R_{m-j}^{(a+\tilde{a},2j)} w_R^{(a+\tilde{a}),k+j} \left\{ -k(k+1)R_{n-k}^{(a,2k)} - 2(k+1)z R_{n-k}^{(a,2k)'} \right. \right. \\
&\quad \quad \quad \left. \left. + \rho(z)^2 R_{n-k}^{(a,2k)''} \right\} dz \right) \\
&= \pi \delta_{k,j} \delta_{i,h} \int_\alpha^1 R_{m-k}^{(a+\tilde{a},2k)} w_R^{(a+\tilde{a},0)} \left( -k(k+1)R_{n-k}^{(a,2k)} \rho^{2k} + \frac{d}{dz} [R_{n-k}^{(a,2k)'} \rho^{2(k+1)}] \right) dz \\
&= \pi \delta_{k,j} \delta_{i,h} \int_\alpha^1 \left\{ -k(k+1)R_{m-k}^{(a+\tilde{a},2k)} R_{n-k}^{(a,2k)} w_R^{(a+\tilde{a},2k)} \right. \\
&\quad \quad \quad \left. - R_{n-k}^{(a,2k)'} w_R^{(a+\tilde{a}-1,2k)} [R_{m-k}^{(a+\tilde{a},2k)'} w_R^{(1,0)} + (a+\tilde{a})R_{m-k}^{(a+\tilde{a},2k)}] \right\} dz \\
&= \pi \delta_{k,j} \delta_{i,h} \int_\alpha^1 R_{n-k}^{(a,2k)} w_R^{(a,2k)} r_{m-k+\tilde{a}} dz
\end{aligned}$$

where  $r_{m-k+\tilde{a}}$  is a degree  $m-k+\tilde{a}$  polynomial in  $z$ , and so the above is zero for  $n-k > m-k+\tilde{a} \iff m < n-\tilde{a}$ .

For the operator  $\mathcal{L}_W^{(a) \rightarrow (a-\tilde{a})}$  for the surface Laplacian on a weighted space, the entries of the operator are given by the coefficients in the expansion

$$\Delta_S(w_R^{(a,0)} Q_{n,k,i}^{(a)}) = \sum_{m=0}^n \sum_{j=0}^m \sum_{h=0}^1 c_{m,j,h} w_R^{(a-\tilde{a},0)} Q_{m,j,h}^{(a-\tilde{a})},$$

where the coefficients are

$$c_{m,j,h} = \left\| Q_{m,j,h}^{(a-\tilde{a})} \right\|_{Q^{(a-\tilde{a})}}^{-2} \left\langle \Delta_S(w_R^{(a,0)} Q_{n,k,i}^{(a)}), Q_{m,j,h}^{(a-\tilde{a})} \right\rangle_{Q^{(0)}}.$$



Using equation (4.14), and integrating by parts thrice, we then have that

$$\begin{aligned}
& \left\langle \Delta_S(w_R^{(a,0)} Q_{n,k,i}^{(a)}, Q_{m,j,h}^{(a-\tilde{a})}) \right\rangle_{Q^{(0)}} \\
&= \left( \int_0^{2\pi} Y_{k,i}(\theta) Y_{j,h}(\theta) d\theta \right) \\
&\quad \cdot \left( \int_\alpha^1 R_{m-j}^{(a-\tilde{a},2j)} w_R^{(a-2,k+j)} \left\{ R_{n-k}^{(a,2k)} [-k(k+1)w_R^{(2,0)} - 2a(k+1)z w_R^{(1,0)} + a(a-1)\rho^2] \right. \right. \\
&\quad \quad \quad + R_{n-k}^{(a,2k)'} [-2(k+1)z w_R^{(2,0)} + 2a w_R^{(1,2)}] \\
&\quad \quad \quad \left. \left. + R_{n-k}^{(a,2k)''} w_R^{(2,2)} \right\} dz \right) \\
&= \pi \delta_{k,j} \delta_{i,h} \int_\alpha^1 \left\{ R_{m-k}^{(a-\tilde{a},2k)} R_{n-k}^{(a,2k)} w_R^{(a-2,2k)} [-k(k+1)w_R^{(2,0)} - 2a(k+1)z w_R^{(1,0)} + a(a-1)\rho^2] \right. \\
&\quad \quad \quad + a R_{n-k}^{(a,2k)'} R_{m-k}^{(a-\tilde{a},2k)} w_R^{(a-1,2k+2)} \\
&\quad \quad \quad \left. + R_{m-k}^{(a-\tilde{a},2k)} \frac{d}{dz} [R_{n-k}^{(a,2k)'} w_R^{(a,2k+2)}] \right\} dz \\
&= \pi \delta_{k,j} \delta_{i,h} \int_\alpha^1 \left\{ R_{m-k}^{(a-\tilde{a},2k)} R_{n-k}^{(a,2k)} w_R^{(a-2,2k)} [-k(k+1)w_R^{(2,0)} - 2a(k+1)z w_R^{(1,0)} + a(a-1)\rho^2] \right. \\
&\quad \quad \quad + a R_{n-k}^{(a,2k)'} R_{m-k}^{(a-\tilde{a},2k)} w_R^{(a-1,2k+2)} \\
&\quad \quad \quad \left. + R_{n-k}^{(a,2k)} w_R^{(a-1,2k)} [R_{m-k}^{(a-\tilde{a},2k)''} w_R^{(1,2)} + a R_{m-k}^{(a-\tilde{a},2k)'} \rho^2 - 2(k+1)z R_{m-k}^{(a-\tilde{a},2k)} w_R^{(1,0)}] \right\} dz \\
&= \pi \delta_{k,j} \delta_{i,h} \int_\alpha^1 \left\{ R_{m-k}^{(a-\tilde{a},2k)} R_{n-k}^{(a,2k)} w_R^{(a-2,2k)} [-k(k+1)w_R^{(2,0)} - 2a(k+1)z w_R^{(1,0)} + a(a-1)\rho^2] \right. \\
&\quad \quad \quad + R_{n-k}^{(a,2k)} w_R^{(a-1,2k+2)} [R_{m-k}^{(a-\tilde{a},2k)''} \rho^2 - 2(k+1)z R_{m-k}^{(a-\tilde{a},2k)'}] \\
&\quad \quad \quad \left. + a [R_{n-k}^{(a,2k)} R_{m-k}^{(a-\tilde{a},2k)'} + R_{m-k}^{(a-\tilde{a},2k)} R_{n-k}^{(a,2k)'}] w_R^{(a-1,2k+2)} \right\} dz \\
&= \pi \delta_{k,j} \delta_{i,h} \int_\alpha^1 \left\{ R_{m-k}^{(a-\tilde{a},2k)} R_{n-k}^{(a,2k)} w_R^{(a-2,2k)} [-k(k+1)w_R^{(2,0)} - 2a(k+1)z w_R^{(1,0)} + a(a-1)\rho^2] \right. \\
&\quad \quad \quad + R_{n-k}^{(a,2k)} w_R^{(a-1,2k+2)} [R_{m-k}^{(a-\tilde{a},2k)''} \rho^2 - 2(k+1)z R_{m-k}^{(a-\tilde{a},2k)'}] \\
&\quad \quad \quad \left. - a R_{n-k}^{(a,2k)} R_{m-k}^{(a-\tilde{a},2k)} w_R^{(a-2,2k)} [(a-1)\rho^2 - 2(k+1)z w_R^{(1,0)}] \right\} dz \\
&= \pi \delta_{k,j} \delta_{i,h} \int_\alpha^1 R_{n-k}^{(a,2k)} w_R^{(a,2k)} r_{m-k} dz
\end{aligned}$$

where  $r_{m-k}$  is a degree  $m-k$  polynomial in  $z$ , and so the above is zero for  $n-k > m-k \iff m < n$ .

Finally, fix  $a = 1$ . For the operator  $\Delta_W^{(1)}$  for the Laplacian on the weighted space, the entries of the operator are given by the coefficients in the expansion  $\Delta_S(w_R^{(1,0)} Q_{n,k,i}^{(1)}) = \sum_{m=0}^{n+2} \sum_{j=0}^m \sum_{h=0}^1 c_{m,j,h} Q_{m,j,h}^{(1)}$ , where the coefficients are given by

$$c_{m,j,h} = \left\| Q_{m,j,h}^{(1)} \right\|_{Q^{(1)}}^{-2} \left\langle \Delta_S(w_R^{(1,0)} Q_{n,k,i}^{(1)}), Q_{m,j,h}^{(1)} \right\rangle_{Q^{(1)}}.$$

Using equation (4.14) with  $a = 1$ , and Lemma 9, we then have that

$$\begin{aligned} & \left\langle \Delta_S(w_R^{(1,0)} Q_{n,k,i}^{(1)}), Q_{m,j,h}^{(1)} \right\rangle_{Q^{(1)}} \\ &= \left( \int_0^{2\pi} Y_{k,i}(\theta) Y_{j,h}(\theta) d\theta \right) \\ & \quad \cdot \left( \int_\alpha^1 R_{m-j}^{(1,2j)} \left\{ R_{n-k}^{(1,2k)} [-k^2 w_R^{(1,k)} - w_R^{(1,k)} - 2(k+1)z w_R^{(0,k)}] \right. \right. \\ & \quad \quad \quad + R_{n-k}^{(1,2k)'} [2w_R^{(0,k+2)} - 2(k+1)z w_R^{(1,k)}] \\ & \quad \quad \quad \left. \left. + R_{n-k}^{(1,2k)''} w_R^{(1,k+2)} \right\} w_R^{(1,j)} dz \right) \\ &= \pi \delta_{k,j} \delta_{i,h} \int_\alpha^1 R_{m-k}^{(1,2k)} \left\{ R_{n-k}^{(1,2k)} [-k(k+1)w_R^{(1,0)} - 2(k+1)z + c_{n,k}] \right. \\ & \quad \quad \quad \left. + c_{n-1,k} R_{n-k-1}^{(1,2k)} + c_{n+1,k} R_{n-k+1}^{(1,2k)} \right\} w_R^{(1,2k)} dz \\ &= -\pi \delta_{k,j} \delta_{i,h} (\delta_{m,n-1} + \delta_{m,n} + \delta_{m,n+1}) \int_\alpha^1 \left\{ R_{n-k}^{(1,2k)} R_{m-k}^{(1,2k)} (k(k+1)w_R^{(1,0)} + 2(k+1)z) \right. \\ & \quad \quad \quad \left. + R_{n-k}^{(1,2k)'} R_{m-k}^{(1,2k)'} w_R^{(2,2(k+1))} \right\} dz \end{aligned}$$

where the  $c_{n-1,k}$ ,  $c_{n,k}$ ,  $c_{n+1,k}$  are those derived in Lemma 9. □

By applying these differential operators, we are (in some cases) incrementing or decrementing the parameter value  $a$ . It is therefore necessary to also be able to raise or lower the parameter by way of an independent operator. There exist conversion operators that do exactly this, transforming the OPs from one (weighted or non-weighted) parameter space to another.

**Definition 16.** Define the operator matrices  $T^{(a) \rightarrow (a+\tilde{a})}$ ,  $T_W^{(a) \rightarrow (a-\tilde{a})}$  for conversion between non-weighted spaces and weighted spaces respectively according to

$$\begin{aligned} \mathbb{Q}_N^{(a)}(x, y, z) &= \left( T^{(a) \rightarrow (a+\tilde{a})} \right)^\top \mathbb{Q}_N^{(a+\tilde{a})}(x, y, z) \\ \mathbb{W}_N^{(a)}(x, y, z) &= \left( T_W^{(a) \rightarrow (a-\tilde{a})} \right)^\top \mathbb{W}_N^{(a-\tilde{a})}(x, y, z) \end{aligned}$$

**Lemma 10.** The operator matrices in Definition 16 are sparse, with banded-block-banded structure. More specifically:

- $T^{(a) \rightarrow (a+\tilde{a})}$  is block-diagonal with sub-block bandwidths  $(0, 2\tilde{a})$
- $T_W^{(a) \rightarrow (a-\tilde{a})}$  is block-diagonal with sub-block bandwidths  $(2\tilde{a}, 0)$

*Proof.* We proceed with the case for the non-weighted operators  $T^{(a) \rightarrow (a+\tilde{a})}$ . Since  $\{Q_{m,j,h}^{(a+\tilde{a})}\}$  for  $m = 0, \dots, n$ ,  $j = 0, \dots, m$ ,  $h = 0, 1$  is an orthogonal basis for any degree  $n$  polynomial, we can expand  $Q_{n,k,i}^{(a)} = \sum_{m=0}^n \sum_{j=0}^m t_{m,j} Q_{m,j,h}^{(a+\tilde{a})}$ . The coefficients of the expansion are then the entries of the operator matrix. We will show that the only non-zero coefficients are for  $k = j$ ,  $i = h$  and  $m \geq n - \tilde{a}$ . Note

that

$$t_{m,j} = \left\| Q_{m,j,h}^{(a+\tilde{a})} \right\|_{Q^{(a+\tilde{a})}}^{-2} \left\langle Q_{n,k,i}^{(a)}, Q_{m,j,h}^{(a+\tilde{a})} \right\rangle_{Q^{(a+\tilde{a})}}.$$


where

$$\begin{aligned} \left\langle Q_{n,k,i}^{(a)}, Q_{m,j,h}^{(a+\tilde{a})} \right\rangle_{Q^{(a+\tilde{a})}} &= \left( \int_0^{2\pi} Y_{k,i}(\theta) Y_{j,h}(\theta) d\theta \right) \cdot \left( \int_{\alpha}^1 R_{n-k}^{(a,2k)} R_{m-j}^{(a+\tilde{a},2j)} \rho^{k+j} w_R^{(a+\tilde{a},0)} dz \right) \\ &= \pi \delta_{k,j} \delta_{i,h} \int_{\alpha}^1 R_{n-k}^{(a,2k)} R_{m-k}^{(a+\tilde{a},2k)} w_R^{(a+\tilde{a},2k)} dz \end{aligned}$$

which is zero for  $n > m + \tilde{a} \iff m < n - \tilde{a}$ . The sparsity argument for the weighted parameter transformation operator follows similarly.  $\square$

#### 4.4.1 Further partial differential operators

General linear partial differential operators with polynomial variable coefficients can be constructed by composing the sparse representations for partial derivatives, conversion between bases, and Jacobi operators. As a canonical example, we can obtain the matrix operator for the  $\rho^2$ -factored spherical Laplacian  $\rho(z)^2 \Delta_S$ , that will take us from coefficients for expansion in the weighted space  $\mathbb{W}_N^{(1)}(x, y, z) = w_R^{(1,0)}(z) \mathbb{Q}_N^{(1)}(x, y, z)$  to coefficients in the non-weighted space  $\mathbb{Q}_N^{(1)}(x, y, z)$ . Note that this construction will ensure the imposition of the Dirichlet zero boundary conditions on  $\Omega$ , similar to how the Dirichlet zero boundary conditions would be imposed for the operator  $\Delta_W^{(1)}$  in Definition 15. The matrix operator for this  $\rho^2$ -



**Figure 4.1:** “Spy” plots of (differential) operator matrices, showing their sparsity. For (c), the weighted variable coefficient Helmholtz operator is  $\Delta_W^{(1)} + k^2 T^{(0) \rightarrow (1)} V(J_x^{(0)\top}, J_y^{(0)\top}, J_z^{(0)\top}) T_W^{(1) \rightarrow (0)}$  for  $v(x, y, z) = 1 - (3(x - x_0)^2 + 5(y - y_0)^2 + 2(z - z_0)^2)$  where  $(x_0, z_0) := (0.7, 0.2)$ ,  $y_0 := \sqrt{1 - x_0^2 - z_0^2}$  and  $k = 200$ .

factored spherical Laplacian acting on the coefficients vector is then given by

$$D_{\varphi}^{(0)} W_{\varphi}^{(1)} + T^{(0) \rightarrow (1)} T_W^{(1) \rightarrow (0)} (D_{\theta})^2.$$

Importantly, this operator will have banded-block-banded structure, and hence will be sparse, as seen in Figure 4.1.

Another desirable operator is the Biharmonic operator  $\Delta_S^2$ , for which we assume zero Dirichlet and Neumann conditions. That is,

$$u(x, y, z) = 0, \quad \frac{\partial u}{\partial n}(x, y, z) = \nabla_S u(x, y, z) \cdot \hat{\mathbf{n}}(x, y, z) = 0 \quad \text{for } (x, y, z) \in \partial\Omega$$

where  $\partial\Omega$  is the  $z = \alpha$  boundary, and  $\hat{\mathbf{n}}(x, y, z)$  is the outward unit normal vector at the point  $(x, y, z)$  on the boundary, i.e.  $\hat{\mathbf{n}}(x, y, z) = \hat{\mathbf{n}}(\mathbf{x}) := \frac{\mathbf{x}}{\|\mathbf{x}\|} = \mathbf{x}$ . The matrix operator for the Biharmonic operator will take us from coefficients in the space  $\mathbb{W}^{(2)}(x, y, z)$  to coefficients in the space  $\mathbb{Q}_N^{(2)}(x, y, z)$ . To construct this, we can simply multiply together two of the spherical Laplacian operators defined in Definition 15, namely  $\mathcal{L}^{(0) \rightarrow (2)}$  and  $\mathcal{L}_W^{(2) \rightarrow (0)}$ :

$$\mathcal{B}_W^{(2)} := \mathcal{L}^{(0) \rightarrow (2)} \mathcal{L}_W^{(2) \rightarrow (0)}.$$

Since the operator  $\mathcal{L}_W^{(2) \rightarrow (0)}$  acts on coefficients in the  $\mathbb{W}^{(2)}(x, y, z)$  space, we ensure that we satisfy the zero Dirichlet and Neumann boundary conditions – such a function could be written  $u(x, y, z) = w_R^{(2,0)}(z) \tilde{u}(x, y, z)$  and thus its spherical gradient would be zero on the boundary  $z = \alpha$ . This allows us to apply the


$\mathcal{L}^{(0) \rightarrow (2)}$  operator after, safe in the knowledge that boundary conditions have been accounted for. The sparsity and structure of this biharmonic operator are seen in Figure 4.1.

#### 4.4.2 Stability of the Laplacian operator

Denote the condition number of the matrix  $A$ :

$$\kappa(A) := \|A\|_2 \|A^{-1}\|_2,$$

which encodes how accurate the solution to  $A\mathbf{x} = \mathbf{b}$  is under perturbation. Since the Laplacian operator is block-diagonal one can determine the condition number from the condition numbers of each block on the diagonal, and hence deduce the accuracy in solving the Poisson equation, where graphical representations are seen in Figure 4.2 for the  $\Delta_W^{(1)}$  operator matrix. As the degree  $N$  increases, the condition numbers of these blocks becomes grows algebraically fast (at least, the condition number for the first block where  $m = 0$  does, which will be the largest for each  $N$ ). Fortunately however, by applying a trivial diagonal preconditioning matrix similar to that seen for the ultraspherical method [44], we can significantly bound this growth of the condition numbers – this is also seen in Figure Figure 4.2. The preconditioning matrix chosen is  $P^{-1}$ , where  $P$  is the matrix of the diagonal of  $\Delta_W^{(1)}$ .



**Figure 4.2:** Plots of the condition number for the Laplacian operator matrix  $\Delta_W^{(1)}$  and the preconditioned matrix  $P^{-1}\Delta_W^{(1)}$  where  $P$  is the matrix of the diagonal of  $\Delta_W^{(1)}$ . Top: The condition numbers of each diagonal block of  $\Delta_W^{(1)}$  for  $N = 200$ , as well as those for  $P^{-1}\Delta_W^{(1)}$ . Bottom: The maximum condition number of all diagonal blocks of the Laplacian and the preconditioned Laplacian operators, as the degree  $N$  increases.



## 4.5 Computational aspects

In this section we discuss how to expand and evaluate functions in our proposed basis, and take advantage of the sparsity structure in partial differential operators in practical computational applications.

### 4.5.1 Constructing $R_n^{(a,b)}(x)$

It is possible to recursively obtain the recurrence coefficients for the  $\{R_n^{(a,b)}\}$  OPs in (4.5), see [58], by careful application of the Christoffel–Darboux formula [42, 18.2.12].

### 4.5.2 Quadrature rule on the spherical cap

In this section we construct a quadrature rule that is exact for polynomials on the spherical cap  $\Omega$  that can be used to expand functions in the OPs  $Q_{n,k,i}^{(a)}(x, y, z)$  for a given parameter  $a$ .

**Theorem 5.** *Let  $M_1, M_2 \in \mathbb{N}$  and denote the  $M_1$  Gauss quadrature nodes and weights on  $[\alpha, 1]$  with weight  $(t - \alpha)^a$  as  $(t_j, w_j^{(t)})$ . Further, denote the  $M_2$  Gauss quadrature nodes and weights  $[-1, 1]$  with weight  $(1 - x^2)^{-\frac{1}{2}}$  as  $(s_j, w_j^{(s)})$ . Define*

for  $j = 1, \dots, M_1$ ,  $l = 1, \dots, M_2$ :

$$\begin{aligned} x_{l+(j-1)M_2} &:= \rho(t_j) s_l, \\ y_{l+(j-1)M_2} &:= \rho(t_j) \sqrt{1 - s_l^2}, \\ z_{l+(j-1)M_2} &:= t_j, \\ w_{l+(j-1)M_2} &:= w_j^{(t)} w_l^{(s)}. \end{aligned}$$

Let  $f(x, y, z)$  be a function on  $\Omega$ , and  $N \in \mathbb{N}$ . The quadrature rule is then

$$\int_{\Omega} f(x, y, z) w_R^{(a,0)}(z) dA \approx \sum_{j=1}^M w_j [f(x_j, y_j, z_j) + f(-x_j, -y_j, z_j)],$$

where  $M = M_1 M_2$ , and the quadrature rule is exact if  $f(x, y, z)$  is a polynomial of degree  $\leq N$  with  $M_1 \geq \frac{1}{2}(N+1)$ ,  $M_2 \geq N+1$ .

**Remark:** Note that the Gauss quadrature nodes and weights  $(t_j, w_j^{(t)})$  will have to be calculated, however the Gauss quadrature nodes and weights  $(s_j, w_j^{(s)})$  are simply the Chebyshev–Gauss quadrature nodes and weights given explicitly [42, 3.5.23] as  $s_j := \cos\left(\frac{2j-1}{2M_2}\pi\right)$ ,  $w_j^{(s)} := \frac{\pi}{M_2}$ .

*Proof.* Let  $f : \Omega \rightarrow \mathbb{R}$ . Define the functions  $f_e, f_o : \Omega \rightarrow \mathbb{R}$  by

$$\begin{aligned} f_e(x, y, z) &:= \frac{1}{2} \left( f(x, y, z) + f(-x, -y, z) \right), \quad \forall (x, y, z) \in \Omega \\ f_o(x, y, z) &:= \frac{1}{2} \left( f(x, y, z) - f(-x, -y, z) \right), \quad \forall (x, y, z) \in \Omega \end{aligned}$$

so that  $\mathbf{x} \mapsto f_e(\mathbf{x}, z)$  for fixed  $z$  is an even function, and  $\mathbf{x} \mapsto f_o(\mathbf{x}, z)$  for fixed

$z$  is an odd function. Note that if  $f$  is a polynomial, then  $f_e(\rho(t)x, \rho(t)y, t)$  is a polynomial in  $t \in [\alpha, 1]$  for fixed  $(x, y) \in \mathbb{R}^2$ .

Firstly, we note that

$$\begin{aligned} \int_0^{2\pi} g(\cos \theta, \sin \theta) \, d\theta &= \int_{-1}^1 \left[ g(x, \sqrt{1-x^2}) + g(x, -\sqrt{1-x^2}) \right] \frac{dx}{\sqrt{1-x^2}} \\ &= \int_{-1}^1 \left[ g(x, \sqrt{1-x^2}) + g(-x, -\sqrt{1-x^2}) \right] \frac{dx}{\sqrt{1-x^2}} \end{aligned}$$

for any function  $g$ , using a change of variables  $x \rightarrow -x$  for the second term in the integral. Then, integrating the even function  $f_e$  we have

$$\begin{aligned} &\int_{\Omega} f_e(x, y, z) w_R^{(a,0)}(z) \, dA \\ &= \int_{\alpha}^1 w_R^{(a,0)}(z) \left( \int_0^{2\pi} f_e(\rho(z) \cos \theta, \rho(z) \sin \theta, z) \, d\theta \right) dz \\ &= \int_{\alpha}^1 w_R^{(a,0)}(z) \left( \int_{-1}^1 \left[ f_e(\rho(z)x, \rho(z)\sqrt{1-x^2}, z) \right. \right. \\ &\quad \left. \left. + f_e(-\rho(z)x, -\rho(z)\sqrt{1-x^2}, z) \right] \frac{dx}{\sqrt{1-x^2}} \right) dz \\ &= 2 \int_{\alpha}^1 w_R^{(a,0)}(z) \left( \int_{-1}^1 f_e(\rho(z)x, \rho(z)\sqrt{1-x^2}, z) \frac{dx}{\sqrt{1-x^2}} \right) dz \\ &\approx \int_{\alpha}^1 w_R^{(a,0)}(z) \left( \sum_{l=1}^{M_2} w_l^{(s)} f_e(\rho(z)s_l, \rho(z)\sqrt{1-s_l^2}, z) \right) dz \quad (\star) \\ &\approx \sum_{j=1}^{M_1} w_j^{(t)} \sum_{l=1}^{M_2} w_l^{(s)} f_e(\rho(t_j)s_l, \rho(t_j)\sqrt{1-s_l^2}, t_j) \quad (\star\star) \\ &= \sum_{k=1}^{M_1 M_2} w_j f_e(x_j, y_j, z_j). \end{aligned}$$

Suppose  $f$  is a polynomial in  $x, y, z$  of degree  $N$ , and hence that  $f_e$  is a degree

$\leq N$  polynomial. It follows that  $s \mapsto f_e(\rho(z)s, \rho(z)\sqrt{1-s^2}, z)$  for fixed  $z$  is then a polynomial of degree  $\leq N$ . We therefore achieve equality at  $(\star)$  if  $2M_2 - 1 \geq N$  and we achieve equality at  $(\star\star)$  if also  $2M_1 - 1 \geq N$ .

Integrating the odd function  $f_o$  results in

$$\begin{aligned}
 & \int_{\Omega} f_o(x, y, z) w_R^{(a,0)}(z) \, dA \\
 &= \int_{\alpha}^1 w_R^{(a,0)}(z) \left( \int_0^{2\pi} f_o(\rho(z) \cos \theta, \rho(z) \sin \theta, z) \, d\theta \right) dz \\
 &= \int_{\alpha}^1 w_R^{(a,0)}(z) \left( \int_{-1}^1 \left[ f_o(\rho(z)x, \rho(z)\sqrt{1-x^2}, z) \right. \right. \\
 &\quad \left. \left. + f_o(-\rho(z)x, -\rho(z)\sqrt{1-x^2}, z) \right] \frac{dx}{\sqrt{1-x^2}} \right) dz \\
 &= 0.
 \end{aligned}$$

since  $f_o(x, y, z) = -f_o(-x, -y, z)$ . Hence, for a polynomial  $f$  in  $x, y, z$  of degree  $N$ ,

$$\begin{aligned}
 \int_{\Omega} f(x, y, z) w_R^{(a,0)}(z) \, dA &= \int_{\Omega} \left( f_e(x, y, z) + f_o(x, y, z) \right) w_R^{(a,0)}(z) \, dA \\
 &= \int_{\Omega} f_e(x, y, z) w_R^{(a,0)}(z) \, dA \\
 &= \sum_{j=1}^M w_j f_e(x_j, y_j, z_j),
 \end{aligned}$$

where  $M = M_1 M_2$  and  $2M_1 - 1 \geq N, 2M_2 - 1 \geq N$ . □

### 4.5.3 Obtaining the coefficients for expansion of a function on the spherical cap

Fix  $a \in \mathbb{R}$ . Then for any function  $f : \Omega \rightarrow \mathbb{R}$  we can express  $f$  by

$$f(x, y, z) \approx \sum_{k=0}^N \mathbb{Q}_{N,k}^{(a)}(x, y, z)^\top \mathbf{f}_k = \mathbb{Q}_N^{(a)}(x, y, z)^\top \mathbf{f}$$

for  $N$  sufficiently large, where  $\mathbb{Q}_{N,k}^{(a)}, \mathbb{Q}_N^{(a)}$  is defined in equations (4.10–4.12) and where

$$\mathbf{f}_k := \begin{pmatrix} f_{k,k,0} \\ f_{k,k,1} \\ \vdots \\ f_{N,k,0} \\ f_{N,k,1} \end{pmatrix} \in \mathbb{R}^{2(N-k+1)} \quad \text{for } n = 1, 2, \dots, N, \quad \mathbf{f}_0 := \begin{pmatrix} f_{0,0,0} \\ \vdots \\ f_{N,0,0} \end{pmatrix} \in \mathbb{R}^{N+1},$$

$$\mathbf{f} := \begin{pmatrix} \mathbf{f}_0 \\ \vdots \\ \mathbf{f}_N \end{pmatrix} \in \mathbb{R}^{2(N+1)^2}, \quad f_{n,k,i} := \left\langle f, \mathbb{Q}_{n,k,i}^{(a)} \right\rangle_{Q^{(a)}} \left\| \mathbb{Q}_{n,k,i}^{(a)} \right\|_{Q^{(a)}}^{-2}.$$

Recall from equation (4.4) that  $\left\| \mathbb{Q}_{n,k,i}^{(a)} \right\|_{Q^{(a)}}^2 = \omega_R^{(a,2k)} \pi$ . Using the quadrature rule detailed in Section 4.5.2 for the inner product, we can calculate the coefficients

$f_{n,k,i}$  for each  $n = 0, \dots, N$ ,  $k = 0, \dots, n$ ,  $i = 0, 1$ :

$$\begin{aligned} f_{n,k,i} &= \frac{1}{2 \omega_R^{(a,2k)} \pi} \sum_{j=1}^M w_j [f(x_j, y_j, z_j) Q_{n,k,i}^{(a)}(x_j, y_j, z_j) + f(-x_j, -y_j, z_j) Q_{n,k,i}^{(a)}(-x_j, -y_j, z_j)] \\ &= \frac{1}{M_2 \omega_R^{(a,2k)}} \sum_{j=1}^M [f(x_j, y_j, z_j) Q_{n,k,i}^{(a)}(x_j, y_j, z_j) + f(-x_j, -y_j, z_j) Q_{n,k,i}^{(a)}(-x_j, -y_j, z_j)] \end{aligned}$$

where the quadrature nodes and weights are those from Theorem 5, and  $M = M_1 M_2$  with  $2M_1 - 1 \geq N$ ,  $M_2 - 1 \geq N$  (i.e. we can choose  $M_2 := N + 1$  and  $M_1 := \lceil \frac{N+1}{2} \rceil$ ).

**Remark:** While it may be possible to leverage fast transforms, as in [54, 55], to speed up the calculation of the coefficients, this remains an open question.

#### 4.5.4 Function evaluation

For a function  $f$ , with coefficients vector  $\mathbf{f}$  for expansion in the  $\{Q_{n,k,i}\}$  basis as determined via the method in Section 4.5.3 up to order  $N$ , we can use the Clenshaw algorithm to evaluate the function at a point  $(x, y, z) \in \Omega$  as follows. Let  $A_n, B_n, D_n^\top, C_n$  be the Clenshaw matrices from Definition 14, and define the

rearranged coefficients vector  $\tilde{\mathbf{f}}$  via

$$\mathbf{f}_n := \begin{pmatrix} f_{n,0,0} \\ f_{n,1,0} \\ f_{n,1,1} \\ \vdots \\ f_{n,n,0} \\ f_{n,n,1} \end{pmatrix} \in \mathbb{R}^{2(N+1)} \quad \text{for } n = 1, 2, \dots, N, \quad \mathbf{f}_0 = f_{0,0,0} \in \mathbb{R},$$

$$\mathbf{f} := \begin{pmatrix} \mathbf{f}_0 \\ \vdots \\ \mathbf{f}_N \end{pmatrix} \in \mathbb{R}^{(N+1)^2}.$$

The trivariate Clenshaw algorithm works similar to the bivariate Clenshaw algorithm introduced in [45] for expansions in the triangle:

1) Set  $\boldsymbol{\xi}_{N+2} = \mathbf{0}$ ,  $\boldsymbol{\xi}_{N+1} = \mathbf{0}$ .

2) For  $n = N : -1 : 0$

$$\text{set } \boldsymbol{\xi}_n^T = \mathbf{f}_n^T - \boldsymbol{\xi}_{n+1}^T D_n^T (B_n - G_n(x, y, z)) - \boldsymbol{\xi}_{n+2}^T D_{n+1}^T C_{n+1}$$

3) Output:  $f(x, y, z) \approx \boldsymbol{\xi}_0^T \tilde{\mathbf{Q}}_0^{(a)} = \xi_0 Q_0^{(a)}$

### 4.5.5 Calculating non-zero entries of the operator matrices

The proofs of Theorem 4 and Lemma 10 provide a way to calculate the non-zero entries of the operator matrices given in Definition 15 and Definition 16. We can simply use quadrature to calculate the 1D inner products, which has a complexity of  $\mathcal{O}(N^2)$ . This proves much cheaper computationally than using the 3D quadrature rule to calculate the surface inner products, which has a complexity of  $\mathcal{O}(N^3)$ .

### 4.5.6 Obtaining operator matrices for variable coefficients

The Clenshaw algorithm outlined in Section 4.5.4 can also be used with Jacobi matrices  $J_x^{(a)}, J_y^{(a)}, J_z^{(a)}$  replacing the point  $(x, y, z)$ . Let  $v : \Omega \rightarrow \mathbb{R}$  be the function that we wish to obtain an operator matrix  $V$  for  $v$ , so that

$$v(x, y, z) f(x, y, z) = v(x, y, z) \mathbf{f} \tilde{\mathbb{Q}}^{(a)}(x, y, z) = (V \mathbf{f})^\top \tilde{\mathbb{Q}}^{(a)}(x, y, z),$$

i.e.  $V \mathbf{f}$  is the coefficients vector for the function  $v(x, y, z) f(x, y, z)$ .

To this end, let  $\tilde{\mathbf{v}}$  be the coefficients for expansion up to order  $N$  in the  $\{Q_{n,k,i}\}$  basis of  $v$  (rearranged as in Section 4.5.4 so that  $v(x, y, z) = \tilde{\mathbf{v}}^\top \tilde{\mathbb{Q}}^{(a)}(x, y, z)$ ). Denote  $X := (J_x^{(a)})^\top$ ,  $Y := (J_y^{(a)})^\top$ ,  $Z := (J_z^{(a)})^\top$ . The operator  $V$  is then the



result of the following:

1) Set  $\boldsymbol{\xi}_{N+2} = \mathbf{0}$ ,  $\boldsymbol{\xi}_{N+2} = \mathbf{0}$ .

2) For  $n = N : -1 : 0$


$$\text{set } \boldsymbol{\xi}_n^T = \tilde{\mathbf{v}}_n^T - \boldsymbol{\xi}_{n+1}^T D_n^T (B_n - G_n(X, Y, Z)) - \boldsymbol{\xi}_{n+2}^T D_{n+1}^T C_{n+1}$$

3) Output:  $V = v(X, Y, Z) \approx \boldsymbol{\xi}_0 \tilde{Q}_0^{(a)} = \xi_0 Q_0^{(a)}$

where at each iteration,  $\boldsymbol{\xi}_n$  is a vector of matrices.

## 4.6 Examples on spherical caps with zero Dirichlet conditions

We now demonstrate how the sparse linear systems constructed as above can be used to efficiently solve PDEs with zero Dirichlet conditions on the spherical cap defined by  $\Omega$ . We consider Poisson, inhomogeneous variable coefficient Helmholtz equation and the Biharmonic equation, as well as a time dependent heat equation, demonstrating the versatility of the approach.



**Figure 4.3:** Top: The computed solution to  $\Delta u = f$  with zero boundary conditions with  $f(x, y, z) = -2e^x yz(2 + x) + w_R^{(1,0)}(z)e^x(y^3 + z^2y - 4xy - 2y)$ . Bottom: The norms of each block of the computed solution of the Poisson equation with right hand side function  $f(x, y, z) = \|\mathbf{x} - (\epsilon + 1/\sqrt{3})(1, 1, 1)^\top\|$  for different  $\epsilon$  values. This indicates spectral convergence.

### 4.6.1 Poisson

The Poisson equation is the classic problem of finding  $u(x, y, z)$  given a function  $f(x, y, z)$  such that:

$$\begin{cases} \Delta_S u(x, y, z) = f(x, y, z) & \text{in } \Omega \\ u(x, y, z) = 0 & \text{on } \partial\Omega \end{cases} \quad (4.15)$$

noting the imposition of zero Dirichlet boundary conditions on  $u$ .

We can tackle the problem as follows. Choose an  $N \in \mathbb{N}$  large enough for the problem, and denote the coefficient vector for expansion of  $u$  in the  $\mathbb{W}_N^{(1)}$  OP basis up to degree  $N$  by  $\mathbf{u}$ , and the coefficient vector for expansion of  $f$  in the  $\mathbb{Q}_N^{(1)}$  OP basis up to degree  $N$  by  $\mathbf{f}$ . Since  $f$  is known, we can obtain  $\mathbf{f}$  using the quadrature rule in Section 4.5.3. In matrix-vector notation, our system hence becomes:

$$\Delta_W^{(1)} \mathbf{u} = \mathbf{f}$$

which can be solved to find  $\mathbf{u}$ . In Figure 4.3 we see the solution to the Poisson equation with zero boundary conditions given in equation (4.15) in the disk-slice  $\Omega$ . In Figure 4.3 we also show the norms of each block of calculated coefficients of the approximation for four right-hand sides of the Poisson equation with  $N = 200$ , that is,  $(N + 1)^2 = 40,401$  unknowns. The right hand sides we choose here are

given by

$$f(x, y, z) = \left\| \left( x - (\epsilon + 1/\sqrt{3}), y - (\epsilon + 1/\sqrt{3}), z - (\epsilon + 1/\sqrt{3}) \right)^\top \right\|$$

for differing choices of  $\epsilon$  – this parameter serves to alter the distance from which we would have a singularity. In the plot, a “block” is simply the group of coefficients corresponding to OPs of the same degree, and so the plot shows how the norms of these blocks decay as the degree of the expansion increases. Thus, the rate of decay in the coefficients is a proxy for the rate of convergence of the computed solution: as typical of spectral methods, we expect the numerical scheme to converge at the same rate as the coefficients decay. We see that we achieve spectral convergence for these examples.


### 4.6.2 Inhomogeneous variable-coefficient Helmholtz

Find  $u(x, y)$  given functions  $v, f : \Omega \rightarrow \mathbb{R}$  such that:


$$\begin{cases} \Delta_S u(x, y, z) + k^2 v(x, y, z) u(x, y, z) = f(x, y, z) & \text{in } \Omega \\ u(x, y, z) = 0 & \text{on } \partial\Omega \end{cases} \quad (4.16)$$

where  $k \in \mathbb{R}$ , noting the imposition of zero Dirichlet boundary conditions on  $u$ .

We can tackle the problem as follows. Denote the coefficient vector for expansion of  $u$  in the  $\mathbb{W}_N^{(1)}$  OP basis up to degree  $N$  by  $\mathbf{u}$ , and the coefficient vector for



**Figure 4.4:** Top: The computed solution to  $\Delta u + k^2 v u = f$  with zero boundary conditions with  $f(x, y, z) = y e^x (z - \alpha)$ ,  $v(x, y, z) = 1 - (3(x - x_0)^2 + 5(y - y_0)^2 + 2(z - z_0)^2)$  where  $(x_0, z_0) := (0.7, 0.2)$ ,  $y_0 := \sqrt{1 - x_0^2 - z_0^2}$  and  $k = 100$ . Bottom: The norms of each block of the computed solution of the Helmholtz equation with the right hand side function  $f(x, y, z) = 1$  and the same function  $v(x, y, z)$ , for various  $k$  values. This indicates spectral convergence.



**Figure 4.5:** Time in seconds to build and solve the system  $[\Delta_S + v(x, y, z)] u(x, y, z) = f(x, y, z)$ , for a rotationally invariant  $v(x, y, z) = v(z)$ . This demonstrates that the approach is roughly of order  $\mathcal{O}(N^2)$ , where  $N$  is the degree to which we approximate the solution (the number of unknowns is then  $(N + 1)^2$ ). Here, we used  $f = -2e^x yz(2 + x) + (z - \alpha)e^x(y^3 + z^2y - 4xy - 2y)$  and  $v(x, y, z) = v(z) = \cos(z)$ .

expansion of  $f$  in the  $\mathbb{Q}_N^{(1)}$  OP basis up to degree  $N$  by  $\mathbf{f}$ . Since  $f$  is known, we can obtain the coefficients  $\mathbf{f}$  using the quadrature rule in Section 4.5.3.

Define  $X := (J_x^{(0)})^\top$ ,  $Y := (J_y^{(0)})^\top$ ,  $Z := (J_z^{(0)})^\top$ . We can obtain the matrix operator for the variable-coefficient function  $v(x, y, z)$  by using the Clenshaw algorithm with matrix inputs as the Jacobi matrices  $X, Y, Z$ , yielding an operator matrix of the same dimension as the input Jacobi matrices a la the procedure introduced in [45]. We can denote the resulting operator acting on coefficients in the  $\mathbb{Q}_N^{(0)}$  space by  $v(X, Y, Z)$ . In matrix-vector notation, our system hence becomes:

$$(\Delta_W^{(1)} + k^2 T^{(0) \rightarrow (1)} V T_W^{(1) \rightarrow (0)}) \mathbf{u} = \mathbf{f}$$

which can be solved to find  $\mathbf{u}$ . We can see the sparsity and structure of this matrix system in Figure 4.1 with  $v(x, y, z) = zxy^2$  as an example. In Figure 4.4 we see

the solution to the inhomogeneous variable-coefficient Helmholtz equation with zero boundary conditions given in equation (4.16) in the spherical cap  $\Omega$ , with  $f(x, y, z) = ye^x w_R^{(1,0)}(z)$ ,  $v(x, y, z) = 1 - (3(x - x_0)^2 + 5(y - y_0)^2 + 2(z - z_0)^2)$  where  $(x_0, z_0) := (0.7, 0.2)$ ,  $y_0 := \sqrt{1 - x_0^2 - z_0^2}$  and  $k = 100$ . In Figure 4.4 we also show the norms of each block of calculated coefficients for the approximation of the solution to the inhomogeneous variable-coefficient Helmholtz equation with various  $k$  values. Here, we use  $N = 200$ , that is,  $(N + 1)^2 = 40,401$  unknowns. Once again, the rate of decay in the coefficients is a proxy for the rate of convergence of the computed solution, and we see that we achieve spectral convergence.

In Figure 4.5 we plot the time taken<sup>3</sup> to construct the operator for  $\Delta_S + v(x, y, z)$ , with a rotationally invariant  $v(x, y, z) = v(z) = \cos z$ , and solve a zero boundary condition Helmholtz problem. The plot demonstrates that as we increase the degree of approximation  $N$ , we achieve a complexity of an optimal  $\mathcal{O}(N^2)$ .

What about other boundary conditions? One simple extension is the case where the value on the boundary takes that of a function depending only on  $x$  and  $y$ , i.e.  $c = c(x, y)$ . In this case, the problem

$$\begin{cases} \Delta_S u(x, y, z) + k^2 v(x, y, z) u(x, y, z) = f(x, y, z) & \text{in } \Omega \\ u(x, y, z) = c(x, y) & \text{on } \partial\Omega \end{cases}$$

---

<sup>3</sup>measured using the “@belapsed” macro from the BenchmarkTools.jl package [12] in Julia.

is equivalent to letting  $u(x, y, z) = \tilde{u}(x, y, z) + c(x, y)$  and solving

$$\begin{cases} \Delta_S \tilde{u}(x, y, z) + k^2 v(x, y, z) \tilde{u}(x, y, z) = g(x, y, z) & \text{in } \Omega \\ \tilde{u}(x, y, z) = 0 & \text{on } \partial\Omega \end{cases}$$

for  $\tilde{u}$ , where

$$g(x, y, z) := f(x, y, z) - k^2 v(x, y, z) c(x, y) - \Delta_S c(x, y)$$

for  $(x, y, z) \in \Omega$ . This new problem is then a zero boundary condition Helmholtz problem with right hand side  $g$ . Notice that the spherical Laplacian applied to  $c(x, y)$ , expanded in the  $\mathbb{Q}_N^{(1)}$  basis with coefficients vector  $\mathbf{c} = (c_{n,k,i})$ , is just

$$\begin{aligned} \Delta_S c(x, y) &= \frac{1}{\rho(z)^2} \frac{\partial^2}{\partial \theta^2} c(x, y) \\ &= \frac{1}{\rho(z)^2} \sum_{n=0}^N \sum_{i=0}^1 c_{n,n,i} \frac{\partial^2}{\partial \theta^2} Y_{n,i}(\theta) \\ &= -\frac{1}{\rho(z)^2} \sum_{n=0}^N \sum_{i=0}^1 n^2 c_{n,n,i} Y_{n,i}(\theta) \end{aligned}$$

since the coefficients  $\{c_{n,k,i}\}$  for such a function are zero for  $k < n$  due to the dependence on  $x$  and  $y$  only, which are precisely the Fourier coefficients of  $c(\cos \theta, \sin \theta)$ . Thus, since the function  $c(x, y)$  is known, it is simple to evaluate  $\frac{\partial^2}{\partial \theta^2} c(x, y)$  and hence one can obtain the coefficients for the expansion of  $g(x, y, z)$  in the  $\mathbb{Q}_N^{(1)}$  basis in the usual manner.



### 4.6.3 Biharmonic equation


Our last example is the biharmonic equation: find  $u(x, y, z)$  given a function  $f(x, y, z)$  such that:

$$\begin{cases} \Delta_S^2 u(x, y, z) = f(x, y, z) & \text{in } \Omega \\ u(x, y, z) = 0, \quad \frac{\partial u}{\partial n}(x, y, z) = \nabla_S u(x, y, z) \cdot \hat{\mathbf{n}}(x, y, z) = 0 & \text{on } \partial\Omega \end{cases} \quad (4.17)$$

where  $\Delta_S^2$  is the Biharmonic operator, noting the imposition of zero Dirichlet and Neumann boundary conditions on  $u$ . For clarity, we reiterate that the unit normal vector in this sense is simply  $\hat{\mathbf{n}}(x, y, z) = \hat{\mathbf{n}}(\mathbf{x}) := \frac{\mathbf{x}}{\|\mathbf{x}\|} = \mathbf{x}$  (see Section ??). In Figure 4.6 we see the solution to the Biharmonic equation (4.17) in the spherical cap  $\Omega$ . In Figure 4.6 we also show the norms of each block of calculated coefficients of the approximation for four more complex right-hand sides of the biharmonic equation with  $N = 200$ , that is,  $(N + 1)^2 = 40,401$  unknowns. Once again, the rate of decay in the coefficients is a proxy for the rate of convergence of the computed solution, and we see that we achieve exponential convergence for these more complex functions.

### 4.6.4 Other boundary conditions

Other boundary conditions beyond those already presented here are beyond the scope of this paper. However, the approach established for the case of a triangle domain in 2D shows how Neumann conditions can be considered as a system of equations in partial derivatives involving Dirichlet conditions [45]. This approach



**Figure 4.6:** Left: The computed solution to  $\Delta^2 u = f$  with zero Dirichlet and Neumann boundary conditions with  $f(x, y, z) = (1 + \operatorname{erf}(5(1 - 10((x - 0.5)^2 + y^2))))\rho(z)^2$ . Right: The norms of each block of the computed solution of the biharmonic equation with the right hand side function  $f(x, y, z) = \exp(-\epsilon((x - x_0)^2 + (y - y_0)^2 + (z - z_0)^2))$  where  $(x_0, z_0) := (0.7, 0.2)$ ,  $y_0 := \sqrt{1 - x_0^2 - z_0^2}$ , for various  $\epsilon$  values. This demonstrates algebraic convergence.

may also work here for the spherical cap. Ideally, one would properly tackle the tangent space (which again is beyond the scope of this paper) which would lower the degrees of freedom.

## 4.7 Conclusions

We have shown that trivariate orthogonal polynomials can lead to sparse discretizations of general linear PDEs on spherical cap domains, with Dirichlet boundary conditions on the  $z = \alpha \in (0, 1)$  boundary. We have provided a detailed practical framework for the application of the methods described for quadratic surfaces of revolution [47], by utilising the non-classical 1D OPs on the interval  $[\alpha, 1]$  with the weight  $(z - \alpha)^a (1 - z^2)^{b/2}$  defined for the disk-slice case [58]. Generalisation to spherical bands ( $\alpha \leq z \leq \beta$ ) is straightforward. This work thus forms a building block in developing an *hp*-finite element method to solve PDEs on the sphere by using spherical band and spherical cap shaped elements.

This work also serves as a stepping stone to constructing similar methods to solve partial differential equations on other 3D sub-domains of the sphere such as spherical triangles. It is clear from the construction in this paper that discretizations of spherical gradients and Laplacian's are sparse on other suitable sub-components of the sphere. The resulting sparsity in high-polynomial degree discretizations presents an attractive alternative to methods based on bijective mappings (e.g., [7, 53, 9]). Constructing these sparse spectral methods for surface PDEs on spherical triangles is future work, and has applications in weather

prediction [59], though it is not yet clear how to directly construct, or perhaps *compute*, the necessary orthogonal polynomials.

The next stage is to develop an orthogonal basis for the tangent space of the spherical cap (or band), and obtain sparse differential operators for gradient, divergence etc. On the complete sphere, the vector spherical harmonics that form the orthogonal basis are simply the gradients and perpendicular gradients of the scalar spherical harmonics [2] which has been used effectively for solving PDEs on the sphere [66, 31] – however, we do not have that luxury for the spherical cap or band, and hence the construction of a basis will not be as straightforward.

# Chapter 5

## Summary and future directions

### 5.1 Summary

In this thesis, we have developed sparse spectral methods for solving partial differential equations (PDEs) on disk-slices and trapeziums in 2D, and spherical caps as a surface 3D. The work can also be used as a template for developing similar methods on other such similar domains, in particular other subdomains of the sphere.

We began with an introduction to multidimensional sparse spectral methods by looking at the spherical harmonics on the whole sphere, explaining how they can be written as orthogonal polynomials in  $x, y, z$ , and how Jacobi and differential operators that apply to coefficient vectors will hence be banded-block-banded.

For the disk-slice in 2D, we defined the OPs that allow similar sparse and banded-

block-banded operators required for solving PDEs in the domain, deriving their structure and providing a method to efficiently calculate numerically their entries. The reason they need to be calculated numerically is due to the non-classical univariate OPs that are involved in the 2D OP definitions. Finally, we moved on to use the same arguments for the spherical cap, a 3D surface that is a subdomain of the unit sphere. We once again defined the 3D OPs as orthogonal polynomials in  $x, y, z$ , and showed how the differential operator matrices continue to elicit similar banded-block-banded structure.

## 5.2 Future directions

We hope that this thesis can serve as somewhat of a blueprint for formalising sparse spectral methods, including how one should define the OPs and derive the accompanying Jacobi and differential operators. The motivation behind this work was to develop sparse spectral methods for subdomains of the unit sphere in 3D space. Thus, the natural direction from this point would be to formalise the framework for spherical bands, which would simply involve slightly more complex arithmetic and the sub-block bands would be slightly larger (this due to the  $R$  polynomials we defined now having no zero coefficients in their three-term recurrences, meaning the expressions for multiplication by  $x, y, z$  would gain some extra terms).

Ideally, we would like to pair this with a spectral element method for the sphere, where the elements would be the spherical subdomains of spherical bands and

spherical caps. The goal here would be to test this method against the full sphere spectral method that is in place in the ECMWF model, with the hypothesis being that by reducing the size of the transforms, we can improve the overall efficiency while still maintaining the accuracy that we achieve from a spectral approach.

## .1 Vector-valued functions in the tangent space

We can extend the methodology detailed in Chapter ?? to the tangent space of the spherical cap  $\Omega$ , which we call  $\Omega_T$ . By creating a basis of orthogonal vector polynomials (OVPs) for the spherical cap  $\Omega$ , we can expand vector-valued functions that lie in the tangent space in this basis. Such functions that are useful for the sphere are gradients and perpendicular-gradients of scalar functions.

Any function in the tangent space of  $\Omega$  can be written as  $\nabla f + \hat{\mathbf{r}} \times \nabla g$  for some scalar functions  $f, g$  on  $\Omega$ . Let  $f$  be a scalar function on  $\Omega$  that satisfies zero boundary conditions. Then there exist coefficients  $\mathbf{f}$  such that  $f$  can be expanded in the OP basis  $\mathbb{W}_N^{(1)}$  for large enough  $N$ , i.e.  $f = w_R^{(1,0)} \sum_{n,k,i} f_{n,k,i} Q_{n,k,i}^{(1)}$ . The gradient and perpendicular gradient of  $f$  are then

$$\begin{aligned}\nabla f &= \sum_{n,k,i} f_{n,k,i} \nabla \left( w_R^{(1,0)} Q_{n,k,i}^{(1)} \right) \\ \hat{\mathbf{r}} \times \nabla f &= \sum_{n,k,i} f_{n,k,i} \hat{\mathbf{r}} \times \nabla \left( w_R^{(1,0)} Q_{n,k,i}^{(1)} \right).\end{aligned}$$

For the whole sphere, recall from Section 2.5 that we simply chose the vector spher-

ical harmonics as our tangent space basis, defined as the gradient and perpendicular gradients of the scalar spherical harmonics. We could make that choice because the vector spherical harmonics as defined were naturally orthogonal. Unfortunately, we do not have the same luxury here, with the sets  $\{\nabla(w_R^{(a,0)} Q_{n,k,i}^{(a)}), \hat{\mathbf{r}} \times \nabla(w_R^{(a,0)} Q_{n,k,i}^{(a)})\}$  and  $\{\nabla Q_{n,k,i}^{(a)}, \hat{\mathbf{r}} \times \nabla Q_{n,k,i}^{(a)}\}$  not being orthogonal for any parameter  $a$ . The game for future work is to find an orthogonal basis that spans the tangent space of the spherical cap  $\Omega$ , such that we can sparsely expand the likes of  $\nabla Q_{n,k,i}^{(a)}$ .

Appendix numbering not working?



# Bibliography

- [1] Paul Appel and J Kampe De Feriet. Fonctions hypergéométriques et hypersphériques. *Polynomes D’Hermite*, 140, 1926.
- [2] Rubén G Barrera, GA Estevez, and J Giraldo. Vector spherical harmonics and their application to magnetostatics. *European Journal of Physics*, 6(4):287, 1985.
- [3] Peter Bauer et. al. The ecmwf scalability programme: Progress and plans. Technical memorandum 857, ECMWF, <https://www.ecmwf.int/node/19380>, 02 2020.
- [4] S. L. Belousov. *Tables of Normalized Associated Legendre Polynomials*, volume 18. Pergamon Press, 1962.
- [5] Sven Beuchler and Joachim Schoeberl. New shape functions for triangular p-FEM using integrated Jacobi polynomials. *Numerische Mathematik*, 103(3):339–366, 2006.
- [6] Florian Beyer, Boris Daszuta, Jörg Frauendiener, and Ben Whale. Numerical evolutions of fields on the 2-sphere using a spectral method based on spin-

- weighted spherical harmonics. *Classical and Quantum Gravity*, 31(7):075019, 2014.
- [7] Boris Bonev, Jan S Hesthaven, Francis X Giraldo, and Michal A Kopera. Discontinuous Galerkin scheme for the spherical shallow water equations with applications to tsunami modeling and prediction. *Journal of Computational Physics*, 362:425–448, 2018.
- [8] John P Boyd. *Chebyshev and Fourier spectral methods*. Courier Corporation, 2001.
- [9] John P Boyd. A Chebyshev/rational Chebyshev spectral method for the Helmholtz equation in a sector on the surface of a sphere: defeating corner singularities. *Journal of Computational Physics*, 206(1):302–310, 2005.
- [10] John P Boyd and Fu Yu. Comparing seven spectral methods for interpolation and for solving the poisson equation in a disk: Zernike polynomials, logan–shepp ridge polynomials, chebyshev–fourier series, cylindrical robert functions, bessel–fourier expansions, square-to-disk conformal mapping and radial basis functions. *Journal of Computational Physics*, 230(4):1408–1438, 2011.
- [11] B Carrascal, GA Estevez, Peilian Lee, and V Lorenzo. Vector spherical harmonics and their application to classical electrodynamics. *European Journal of Physics*, 12(4):184, 1991.
- [12] Jiahao Chen and Jarrett Revels. Robust benchmarking in noisy environments. *arXiv preprint arXiv:1608.04295*, Aug 2016.

- [13] Hyeong-Bin Cheong. A dynamical core with double fourier series: Comparison with the spherical harmonics method. *Monthly weather review*, 134(4):1299–1315, 2006.
- [14] James W Cooley and John W Tukey. An algorithm for the machine calculation of complex fourier series. *Mathematics of computation*, 19(90):297–301, 1965.
- [15] M Diamantakis, P Bauer, W Deconinck, P Dueben, C Kuenhlein, S Malardel, A Mueller, P Smolarkiewicz, F Vana, and N Wedi. The ecmwf model dynamical core and visions of its future. *NUMERICAL WEATHER*, page 19, 2018.
- [16] Eid H Doha and Ali H Bhrawy. Efficient spectral-galerkin algorithms for direct solution for second-order differential equations using jacobi polynomials. *Numerical Algorithms*, 42(2):137–164, 2006.
- [17] Charles F Dunkl and Yuan Xu. *Orthogonal Polynomials of Several Variables*. Number 155. Cambridge University Press, 2014.
- [18] K Franklin Evans. The spherical harmonics discrete ordinate method for three-dimensional atmospheric radiative transfer. *Journal of Atmospheric Sciences*, 55(3):429–446, 1998.
- [19] Martin J Fengler and Willi Freeden. A nonlinear galerkin scheme involving vector and tensor spherical harmonics for solving the incompressible navier–stokes equation on the sphere. *SIAM Journal on Scientific Computing*, 27(3):967–994, 2005.

- [20] Toshio Fukushima. Numerical computation of spherical harmonics of arbitrary degree and order by extending exponent of floating point numbers. *Journal of Geodesy*, 86(4):271–285, 2012.
- [21] Quoc T Le Gia, Ming Li, and Yu Guang Wang. Favest: Fast vector spherical harmonic transforms. *arXiv preprint arXiv:1908.00041*, 2019.
- [22] Timon S Gutleb and Sheehan Olver. A sparse spectral method for volterra integral equations using orthogonal polynomials on the triangle. *SIAM Journal on Numerical Analysis*, 58(3):1993–2018, 2020.
- [23] Masters Guy and Richards-Dinger Keith. On the efficient calculation of ordinary and generalized spherical harmonics. *Geophysical Journal International*, 135(1):307–309, 1998.
- [24] EL Hill. The theory of vector spherical harmonics. *American Journal of Physics*, 22(4):211–214, 1954.
- [25] Rainer Hollerbach, Caroline Nore, Philippe Marti, Stijn Vantieghem, Francky Luddens, and J Léorat. Parity-breaking flows in precessing spherical containers. *Physical Review E*, 87(5):053020, 2013.
- [26] Mayer Humi. Factorisation of separable partial differential equations. *Journal of Physics A: Mathematical and General*, 20(14):4577, 1987.
- [27] Arieh Iserles. *A First Course in the Numerical Analysis of Differential Equations*, 2nd edition. Number 44. Cambridge university press, 2009.
- [28] Mourad EH Ismail and Ruiming Zhang. A review of multivariate orthogonal

- polynomials. *Journal of the Egyptian Mathematical Society*, 25(2):91–110, 2017.
- [29] Keith Julien and Mike Watson. Efficient multi-dimensional solution of PDEs using chebyshev spectral methods. *Journal of Computational Physics*, 228(5):1480–1503, 2009.
- [30] Tom Koornwinder. Two-variable analogues of the classical orthogonal polynomials. In *Theory and application of special functions*, pages 435–495. Academic Press, 1975.
- [31] Daniel Lecoanet, Geoffrey M Vasil, Keaton J Burns, Benjamin P Brown, and Jeffrey S Oishi. Tensor calculus in spherical coordinates using jacobi polynomials. Part-II: Implementation and examples. *Journal of Computational Physics: X*, 3:100012, 2019.
- [32] Huiyuan Li and Jie Shen. Optimal error estimates in Jacobi-Weighted Sobolev spaces for polynomial approximations on the triangle. *Mathematics of Computation*, 79(271):1621–1646, 2010.
- [33] Colin B Macdonald and Steven J Ruuth. Level set equations on surfaces via the closest point method. *Journal of Scientific Computing*, 35(2):219–240, 2008.
- [34] Colin B Macdonald and Steven J Ruuth. The implicit closest point method for the numerical solution of partial differential equations on surfaces. *SIAM Journal on Scientific Computing*, 31(6):4330–4350, 2010.
- [35] Alphonse P Magnus. Painlevé-type differential equations for the recurrence

- coefficients of semi-classical orthogonal polynomials. *Journal of Computational and Applied Mathematics*, 57(1-2):215–237, 1995.
- [36] Sylvie Malardel, Nils Wedi, Willem Deconinck, Michail Diamantakis, Christian Kühnlein, George Mozdzynski, Mats Hamrud, and Piotr Smolarkiewicz. A new grid for the ifs. *ECMWF Newsletter*, 146:23–28, 2016.
- [37] John C Mason and David C Handscomb. *Chebyshev Polynomials*. CRC press, 2002.
- [38] Martin J Mohlenkamp. A fast transform for spherical harmonics. *Journal of Fourier analysis and applications*, 5(2-3):159–184, 1999.
- [39] Jonathan T Moon, Bruce Walter, and Steve Marschner. Efficient multiple scattering in hair using spherical harmonics. *ACM SIGGRAPH Proceedings*, pages 1–7, 2008.
- [40] HE Moses. The use of vector spherical harmonics in global meteorology and aeronomy. *Journal of the Atmospheric Sciences*, 31(6):1490–1499, 1974.
- [41] Tables of Clebsch-Gordan Coefficients of SU3. Mcnamee, p and chilton, frank. *Reviews of Modern Physics*, 36(4):1005, 1964.
- [42] Frank WJ Olver, Daniel W Lozier, Ronald F Boisvert, and Charles W Clark. *NIST Handbook of Mathematical Functions*. Cambridge University Press, 2010.
- [43] Sheehan Olver. ApproxFun.jl v0.12, Julia package. <https://github.com/JuliaApproximation/ApproxFun.jl>, 2019.

- [44] Sheehan Olver and Alex Townsend. A fast and well-conditioned spectral method. *SIAM Review*, 55(3):462–489, 2013.
- [45] Sheehan Olver, Alex Townsend, and Geoff Vasil. A sparse spectral method on triangles. *SIAM J. Sci. Comput.*, 41(6):A3728–A3756, 2019.
- [46] Sheehan Olver, Alex Townsend, and Geoffrey M Vasil. Recurrence relations for a family of orthogonal polynomials on a triangle. In *Spectral and High Order Methods for Partial Differential Equations ICOSAHOM 2018*, pages 79–92. Springer, Cham, 2020.
- [47] Sheehan Olver and Yuan Xu. Orthogonal polynomials in and on a quadratic surface of revolution. *Mathematics of Computation*, 89:2847–2865, 2020.
- [48] William H Press, Saul A Teukolsky, William T Vetterling, and Brian P Flannery. *Numerical Recipes 3rd Edition: The Art of Scientific Computing*. Cambridge university press, 2007.
- [49] Martin Reinecke and Dag Sverre Seljebotn. Libsharp—spherical harmonic transforms revisited. *Astronomy & Astrophysics*, 554:A112, 2013.
- [50] Robert Rubinstein, Susan Kurien, and Claude Cambon. Scalar and tensor spherical harmonics expansion of the velocity correlation in homogeneous anisotropic turbulence. *Journal of Turbulence*, 16(11):1058–1075, 2015.
- [51] Jun John Sakurai and Eugene D Commins. *Modern Quantum Mechanics, revised edition*. American Association of Physics Teachers, 1995.

- [52] Jie Shen, Tao Tang, and Li-Lian Wang. *Spectral Methods: Algorithms, Analysis and Applications*, volume 41. Springer Science & Business Media, 2011.
- [53] J Shipton, TH Gibson, and CJ Cotter. Higher-order compatible finite element schemes for the nonlinear rotating shallow water equations on the sphere. *Journal of Computational Physics*, 375:1121–1137, 2018.
- [54] Richard Mikaël Slevinsky. On the use of hahn’s asymptotic formula and stabilized recurrence for a fast, simple and stable Chebyshev–Jacobi transform. *IMA Journal of Numerical Analysis*, 38(1):102–124, 2018.
- [55] Richard Mikaël Slevinsky. Fast and backward stable transforms between spherical harmonic expansions and bivariate fourier series. *Applied and Computational Harmonic Analysis*, 47(3):585–606, 2019.
- [56] Richard Mikaël Slevinsky, Hadrien Montanelli, and Qiang Du. A spectral method for nonlocal diffusion operators on the sphere. *Journal of Computational Physics*, 372:893–911, 2018.
- [57] Peter-Pike Sloan. Efficient spherical harmonic evaluation. *Journal of Computer Graphics Techniques*, 2(2):84–90, 2013.
- [58] Ben Snowball and Sheehan Olver. Sparse spectral and p-finite element methods for partial differential equations on disk slices and trapeziums. *Studies in Applied Mathematics*, 145:3–35, 2020.
- [59] Andrew Staniforth and John Thuburn. Horizontal grids for global weather and climate prediction models: a review. *Quarterly Journal of the Royal Meteorological Society*, 138(662):1–26, 2012.



- [60] Reiji Suda and Masayasu Takami. A fast spherical harmonics transform algorithm. *Mathematics of computation*, 71(238):703–715, 2002.
- [61] Paul N Swarztrauber. The vector harmonic transform method for solving partial differential equations in spherical geometry. *Monthly Weather Review*, 121(12):3415–3437, 1993.
- [62] Paul N Swarztrauber. Spectral transform methods for solving the shallow-water equations on the sphere. *Monthly Weather Review*, 124(4):730–744, 1996.
- [63] Lloyd N Trefethen. *Spectral Methods in MATLAB*. SIAM, 2000.
- [64] Dmitriĭ Aleksandrovich Varshalovich, Anatoli Nikolaevitch Moskalev, and Valerii Kel'manovich Khersonskii. *Quantum Theory of Angular Momentum*. World Scientific, 1988.
- [65] Geoffrey M Vasil, Keaton J Burns, Daniel Lecoanet, Sheehan Olver, Benjamin P Brown, and Jeffrey S Oishi. Tensor calculus in polar coordinates using Jacobi polynomials. *Journal of Computational Physics*, 325:53–73, 2016.
- [66] Geoffrey M Vasil, Daniel Lecoanet, Keaton J Burns, Jeffrey S Oishi, and Benjamin P Brown. Tensor calculus in spherical coordinates using Jacobi polynomials. Part-I: Mathematical analysis and derivations. *Journal of Computational Physics: X*, 3:100013, 2019.
- [67] Nils P Wedi, Mats Hamrud, and George Mozdzynski. A fast spherical harmonics transform for global NWP and climate models. *Monthly Weather Review*, 141(10):3450–3461, 2013.

- 
- [68] David L Williamson. The evolution of dynamical cores for global atmospheric models. *Journal of the Meteorological Society of Japan. Ser. II*, 85:241–269, 2007.
- [69] David L Williamson, John B Drake, James J Hack, Rüdiger Jakob, and Paul N Swarztrauber. A standard test set for numerical approximations to the shallow water equations in spherical geometry. *Journal of Computational Physics*, 102(1):211–224, 1992.
- [70] Frits Zernike. Diffraction theory of the knife-edge test and its improved form, the phase-contrast method. *Monthly Notices of the Royal Astronomical Society*, 94:377–384, 1934.