LECTURE 6: THE KALMAN FILTER

STAT 545: INTRODUCTION TO COMPUTATIONAL STATISTICS


Vinayak Rao

Department of Statistics, Purdue University

September 5, 2018

Marginalization:

$$\begin{bmatrix} X \\ Y \end{bmatrix} \sim \mathcal{N} \left(\begin{bmatrix} \mu_X \\ \mu_Y \end{bmatrix}, \begin{bmatrix} \Sigma_{XX} & \Sigma_{XY} \\ \Sigma_{YX} & \Sigma_{YY} \end{bmatrix} \right), \quad X \sim \textbf{?}$$



$$\mu = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \quad \Sigma = \begin{bmatrix} 10 & 4 \\ 4 & 2 \end{bmatrix}$$

Marginalization:


$$\begin{bmatrix} X \\ Y \end{bmatrix} \sim \mathcal{N} \left(\begin{bmatrix} \mu_X \\ \mu_Y \end{bmatrix}, \begin{bmatrix} \Sigma_{XX} & \Sigma_{XY} \\ \Sigma_{YX} & \Sigma_{YY} \end{bmatrix} \right), \quad X \sim ?$$

$$\mu = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \quad \Sigma = \begin{bmatrix} 10 & 4 \\ 4 & 2 \end{bmatrix}$$

$$X \sim \mathcal{N} \left(\mu_X, \Sigma_{XX} \right)$$

Conditioning:


$$\begin{bmatrix} X \\ Y \end{bmatrix} \sim \mathcal{N} \left(\begin{bmatrix} \mu_X \\ \mu_Y \end{bmatrix}, \begin{bmatrix} \Sigma_{XX} & \Sigma_{XY} \\ \Sigma_{YX} & \Sigma_{YY} \end{bmatrix} \right), \quad Y | (X = a) \sim ?$$



$$\mu = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \quad \Sigma = \begin{bmatrix} 10 & 4 \\ 4 & 2 \end{bmatrix}$$


Conditioning:


$$\begin{bmatrix} X \\ Y \end{bmatrix} \sim \mathcal{N} \left(\begin{bmatrix} \mu_X \\ \mu_Y \end{bmatrix}, \begin{bmatrix} \Sigma_{XX} & \Sigma_{XY} \\ \Sigma_{YX} & \Sigma_{YY} \end{bmatrix} \right), \quad Y | (X = a) \sim ?$$



$$\mu = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \quad \Sigma = \begin{bmatrix} 10 & 4 \\ 4 & 2 \end{bmatrix}$$


$$Y|X \sim \mathcal{N}\left(\mu_Y + \Sigma_{XY}\Sigma_{XX}^{-1}(a - \mu_X), \Sigma_{YY} - \Sigma_{XY}\Sigma_{XX}^{-1}\Sigma_{YX}\right)$$





We have a Gaussian 'prior' on X_1 .

We observe a noisy measurement $Y_1|X_1 \sim \mathcal{N}(AX_1, \Sigma_E)$.




We have a Gaussian 'prior' on X_1 .

We observe a noisy measurement $Y_1|X_1 \sim \mathcal{N}(AX_1, \Sigma_E)$.

$$\begin{bmatrix} X \\ \epsilon \end{bmatrix} \to \begin{bmatrix} X \\ Y \end{bmatrix}$$

X and Y jointly Gaussian: what is its mean and covariance?




We have a Gaussian 'prior' on X_1 .

We observe a noisy measurement $Y_1|X_1 \sim \mathcal{N}(AX_1, \Sigma_E)$.

$$\begin{bmatrix} X \\ \epsilon \end{bmatrix} \to \begin{bmatrix} X \\ Y \end{bmatrix}$$

X and Y jointly Gaussian: what is its mean and covariance?

Y is marginally Gaussian: what is its mean and covariance?



We have a Gaussian 'prior' on X_1 .

We observe a noisy measurement $Y_1|X_1 \sim \mathcal{N}(AX_1, \Sigma_E)$.

$$\begin{bmatrix} X \\ \epsilon \end{bmatrix} \to \begin{bmatrix} X \\ Y \end{bmatrix}$$


X and Y jointly Gaussian: what is its mean and covariance?

Y is marginally Gaussian: what is its mean and covariance?

X|Y is Gaussian: what is its mean and covariance?

PRODUCT OF GAUSSIAN DENSITIES:


Product of Gaussian densities is Gaussian a Gaussian density (upto a multiplication constant)



Intuition: sum of two quadratic functions is a quadratic

PRODUCT OF GAUSSIAN DENSITIES:

Product of Gaussian densities is Gaussian a Gaussian density (upto a multiplication constant)



Intuition: sum of two quadratic functions is a quadratic

Aside: need only specify prob. distrib. to a constant

- p(x) and $C \cdot p(x)$ represents the same, if C is independent of x
- · Probabilities must integrate to 1

$$P_1(X_1) \qquad P_i(X_{i+1}|X_i)$$

$$X_1 \longrightarrow X_2 \longrightarrow X_3 \longrightarrow X_4$$

A sequence of random variables such that

$$P(X_{i+1}|X_i, X_{i-1}, \dots, X_1) = P(X_{i+1}|X_i)$$

We'll stick to homogeneous chains:

$$P_1(X_1) \qquad P(X_{i+1}|X_i)$$

$$X_1 \rightarrow X_2 \rightarrow X_3 \rightarrow X_4$$

We'll stick to homogeneous chains:

$$P_1(X_1) \qquad P(X_{i+1}|X_i)$$

$$X_1 \rightarrow X_2 \rightarrow X_3 \rightarrow X_4$$

In fact, with $X_i \in \Re^D$, we will consider:

$$X_1 \sim \mathcal{N}(\mu_0, \Sigma_0)$$

$$X_{i+1} = AX_i + \epsilon_i, \quad \epsilon_i \sim \mathcal{N}(0, \Sigma_E)$$

We'll stick to homogeneous chains:

$$P_1(X_1) \qquad P(X_{i+1}|X_i)$$

$$X_1 \rightarrow X_2 \rightarrow X_3 \rightarrow X_4$$


In fact, with $X_i \in \Re^D$, we will consider:

$$X_1 \sim \mathcal{N}(\mu_0, \Sigma_0)$$
 $X_{i+1} = AX_i + \epsilon_i, \quad \epsilon_i \sim \mathcal{N}(0, \Sigma_E)$

If our chain has T steps, a TD-dimensional Gaussian! In the figure, T=4. In practice: thousands to millions.

A HIDDEN MARKOV MODEL


We don't observe the chain directly:



$$Y_i = BX_i + \zeta_i, \quad \zeta \sim \mathcal{N}(0, \Sigma_z), \quad Y_i \in \Re^d$$

A HIDDEN MARKOV MODEL

We don't observe the chain directly:




$$Y_i = BX_i + \zeta_i, \quad \zeta \sim \mathcal{N}(0, \Sigma_z), \quad Y_i \in \Re^d$$

We want to answer questions like: What is $p(X_i|Y_1,\dots,Y_T)$? $\{X_i,Y_i\}$ is a (D+d)T-dimensional Gaussian.

A HIDDEN MARKOV MODEL

We don't observe the chain directly:



$$Y_i = BX_i + \zeta_i, \quad \zeta \sim \mathcal{N}(0, \Sigma_z), \quad Y_i \in \Re^d$$

We want to answer questions like: What is $p(X_i|Y_1,\dots,Y_T)$? $\{X_i,Y_i\}$ is a (D+d)T-dimensional Gaussian. We 'just' have to look at conditionals?

THE KALMAN FILTER

[board]