

Prognostic implications of HER2 changes after neoadjuvant chemotherapy in patients with HER2-zero and HER2-low breast cancer

Sora Kang, So Heun Lee, Hee Jin Lee, Hye Hyun Jeong, Jae Ho Jeong, Jeong Eun Kim, Jin-Hee Ahn, Kyung Hae Jung, Gyungyub Gong, Hak Hee Kim, Saebyeol Lee, Jongwon Lee, Sung-Bae Kim



PII: S0959-8049(23)00308-8

DOI: <https://doi.org/10.1016/j.ejca.2023.112956>

Reference: EJC112956

To appear in: *European Journal of Cancer*

Please cite this article as: Sora Kang, So Heun Lee, Hee Jin Lee, Hye Hyun Jeong, Jae Ho Jeong, Jeong Eun Kim, Jin-Hee Ahn, Kyung Hae Jung, Gyungyub Gong, Hak Hee Kim, Saebyeol Lee, Jongwon Lee and Sung-Bae Kim, Prognostic implications of HER2 changes after neoadjuvant chemotherapy in patients with HER2-zero and HER2-low breast cancer, *European Journal of Cancer*, (2023) doi:<https://doi.org/10.1016/j.ejca.2023.112956>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2023 Published by Elsevier.

Prognostic implications of HER2 changes after neoadjuvant chemotherapy in patients with HER2-zero and HER2-low breast cancer.

Sora Kang^{a,b}, So Heun Lee^a, Hee Jin Lee^c, Hye Hyun Jeong^a, Jae Ho Jeong^a, Jeong Eun Kim^a, Jin-Hee Ahn^a, Kyung Hae Jung^a, Gyungyub Gong^c, Hak Hee Kim^d, Saebyeol Lee^e, Jongwon Lee^e, Sung-Bae Kim^a

^aDepartment of Oncology, University of Ulsan College of Medicine, Asan Medical Center, 88, Olympic 43-gil, Songpa-gu, Seoul, 05505, Republic of Korea

^bDivision of Hemato-oncology, Department of Internal Medicine, Chungnam National University Hospital, 282 Munhwa-ro, Jung-gu, Daejeon, 35015, Republic of Korea

^cDepartment of Pathology, University of Ulsan College of Medicine, Asan Medical Center, 88, Olympic 43-gil, Songpa-gu, Seoul, 05505, Republic of Korea

^dDepartment of Radiology, Convergence Medicine, University of Ulsan College of Medicine, Asan Medical Center, 88, Olympic 43-gil, Songpa-gu, Seoul, 05505, Republic of Korea

^eDivision of Breast Surgery, Department of Surgery, University of Ulsan College of Medicine, Asan Medical Center, 88, Olympic 43-gil, Songpa-gu, Seoul, 05505, Republic of Korea

E-mail addresses of the authors

Sora Kang: sorakang0515@gmail.com

So Heun Lee: soso0225@naver.com

Hee Jin Lee: backlila@gmail.com

Hye Hyun Jeong: 0315jhh@gmail.com

Jae Ho Jeong: jaeho.jeong@amc.seoul.kr

Jeong Eun Kim: jeongeunkim@amc.seoul.kr

Jin-Hee Ahn: drjiny@amc.seoul.kr

Kyung Hae Jung: khjung@amc.seoul.kr

Gyungyub Gong: gygong@amc.seoul.kr

Hak Hee Kim: radhkim@daum.net

Saeyeol Lee: newstar153@hanmail.net

Jongwon Lee: leejuw@amc.seoul.kr

Correspondence to: Sung-Bae Kim, M.D. Ph.D.,

Department of Oncology, Asan Medical Center, University of Ulsan College of Medicine

88, Olympic-ro 43-gil, Songpa-gu, Seoul 05505, South Korea

E-mail: sbkim3@amc.seoul.kr

Funding

This study did not receive any dedicated funding from public, commercial, or non-profit entities.

Abstract

Background: The transition of HER2 status after neoadjuvant chemotherapy (NAC) in HER2-low breast cancer has not been thoroughly evaluated. Here, we evaluated the HER2 transition among HER2-zero and HER2-low breast cancer cases post-NAC and its impact on clinical outcome.

Methods: We included 1,288 patients with HER2-low or zero breast cancer who underwent NAC and surgery between 2014 and 2018 and had paired pre- and post-therapeutic HER2 status results.

Results: Among patients who were HER2-zero pre-NAC ($n = 650$), 68% and 29% were HER2-zero and HER2-low, respectively, post-NAC. Among patients who were HER2-low pre-NAC ($n = 638$), 32% of patients showed HER2 changes (low to zero), and 59% of patients had a constant HER2-low status post-NAC. Patients with constant HER2-low or transitions from HER2-low to zero had a higher proportion of hormone receptor positivity (84% and 79%) than those with changes from HER2-zero to low (77%) or with constant HER2-zero (56%), respectively. Multivariable logistic regression analysis revealed that patients with oestrogen receptor-positivity had a higher probability of gaining HER2-low expression than those with oestrogen receptor-negativity (odds ratio 2.48). No significant differences were observed in terms of overall survival or disease-free survival between patients with and without HER2-

changes according to their hormone receptor status, except in the post-therapeutic HER2-low, hormone receptor- negativity subset.

Conclusion: Temporal heterogeneity of HER2-low expression is observed in substantial numbers of post-NAC breast cancer patients. Clinical outcomes show no significant associations, except in the post-therapeutic HER2-low, hormone receptor negativity subset. The prognostic implications of HER2 transition in HER2-low breast cancer require further investigation.

Keywords:

HER2-low breast cancer, neoadjuvant chemotherapy, HER2-zero breast cancer, temporal heterogeneity, prognosis

List of abbreviations

HER2: Human epidermal growth factor receptor 2

NAC: Neoadjuvant chemotherapy

HR+: Hormone receptor-positivity

HR-: Hormone receptor-negativity

1 Introduction

Breast cancer is a heterogeneous disease with distinct biology and treatment outcomes, and the hormone receptor and human epidermal growth factor receptor 2 (HER2) statuses of tumours are essential characteristics guiding treatment strategies[1]. Newly developed anti-HER2 antibody-drug conjugates (ADC; trastuzumab deruxtecan and trastuzumab duocarmazine) have shown promising efficacy in patients with HER2-low breast cancer (tumours with immunohistochemistry[IHC] 1+ or IHC 2+ and in situ hybridization [ISH] negativity), and these observations have prompted ongoing research into the biology and clinical implications of HER2-low breast cancer[2,3]. However, it remains debatable whether HER2-low breast cancer is a separate entity, and there is conflicting evidence about the prognostic role of HER2-low expression in breast cancer[4-9].

Neoadjuvant chemotherapy (NAC) before surgery is a standard treatment option for locally advanced breast cancer to induce tumour shrinkage and increase the possibility of breast-conserving surgery[10]. In fact, HER2 status changes (positive to negative, negative to positive) of residual tumours after NAC have been frequently reported, and the association between HER2 changes and prognostic implications has been evaluated[11-14]. However, HER2 status changes, including the concept of HER2-low and its possible transitions (HER2-zero to low or positive, or vice versa), have not been well delineated. Anti-HER2 ADCs have shown promising efficacy for previously treated metastatic HER2-low breast cancer[2]; therefore, re-testing of HER2 status in residual disease may assist in the identification of potential candidates for anti-HER2 targeting agents. Additionally, investigations into HER2 status changes between pre-therapeutic tissue and post-therapeutic residual tumours may help clarify the treatment-resistant mechanisms. Therefore, we aimed to evaluate the changes in HER2 status after NAC and the implications of HER2 changes for clinical outcomes in patients with HER2-negative (HER2-zero and HER2-low) breast cancer as a follow-up on


our initial report on pathologic complete response, long-term outcomes, and recurrence patterns between these two groups post-NAC[7].

2 Methods

2.1 Patients

This retrospective study was performed at Asan Medical Centre, Seoul, Republic of Korea. Figure 1 shows the CONSORT diagram of this study. We identified 1,572 patients who were treated with NAC and underwent curative resection between 2014–2018; the details of these patients have been published previously[7]. Among the 1,572 patients, we excluded patients who showed a pathological complete response ($n = 195$) and those with bilateral breast cancer categorized post-operatively ($n = 64$), multifocal or multicentric breast cancer ($n = 4$), and unavailable HER2 status ($n = 21$). Finally, 1,288 patients with available HER2 status in paired pre-and post-operative samples were included in this analysis. We collected the patients' data from their medical records, including their baseline characteristics, tumour characteristics, and survival outcomes.

Figure 1. CONSORT diagram of the included patients who were treated.



The Allred scoring system was used to assess the oestrogen receptor (ER) and progesterone receptor (PgR) statuses of the tumours. The cut-off value for defining ER or PgR positivity was an Allred score of 3 or above. We defined a tumour as hormone receptor-positive when it was either ER or PgR-positive, or both.

HER2 status was determined following the 2018 American Society of Clinical Oncology/College of American Pathologists (ASCO/CAP) guidelines[15]. HER2-low tumour was defined as a tumour with HER2 IHC scores 1+ or 2+ accompanied by ISH negativity,

and a HER2-zero tumour was defined as a tumour with HER2 IHC score 0. The ER, PgR, and HER2 statuses were assessed using pre-treatment core biopsy tissue and post-therapeutic samples after resection.

2.2 Outcomes

The study outcomes were disease-free survival (DFS) and overall survival (OS) according to HER2 changes post-NAC. DFS was calculated from the date of surgery to the date of recurrence or occurrence of a secondary malignancy or death from any cause (whichever occurred first). OS was calculated from the date of surgery to death from any cause.

2.3 Statistical analysis

We used descriptive statistics to show the patients' baseline and tumour characteristics. Chi-square and Fisher's exact tests were used to analyse the categorical variables, as appropriate. Additionally, a Cohen's Kappa analysis was used to assess the level of concordance of HER2 expression between the primary tumour and residual disease. The agreement between the two lesions was defined as follows: poor, ≤ 0.2 ; mild, 0.2–0.4; moderate, 0.4–0.6; substantial, 0.6–0.8; and almost perfect, > 0.8 [16]. We also used Kaplan-Meier methods and log-rank tests to estimate the DFS and OS and compare the subgroups. Bivariable logistic regression analysis was used to evaluate the association between the HER2 transition and other variables, while univariate and multivariate analyses using a Cox

proportional hazard model were used to estimate the prognostic implications of the HER2 status and other variables. For the multivariate analysis, age, histologic grade, Ki-67 expression before NAC, ER status before NAC, clinical T stage, clinical N stage, and HER2 transition were included as prognostic factors. The ypT stage, ypN stage, and ER status after NAC were not included in the multivariate analysis because of the interactions between other variables (clinical T stage, clinical N stage, and ER status before NAC). All statistical analyses were performed using statistical software R, version 4.2.2 (R Core Development Team, Vienna, Austria), and statistical significance was set at $p < 0.05$.

2.4 Ethical approval and consent to participate

This study was reviewed and approved by the Institutional Review Board of Asan Medical Centre (approval number 2022-0541). The requirement for informed consent was waived owing to this study's retrospective nature.

3 Results


3.1 Changes in HER2 status after NAC

Among the 650 patients who were HER2-zero before NAC, 68% ($n = 445$) and 29% ($n = 189$) of patients showed HER2-zero and HER2-low after NAC, respectively. Among the 638 patients who were HER2-low before NAC, 32% ($n = 201$) of patients showed HER2 changes (low to zero) and 59% ($n = 374$) of patients showed constant HER2-low status after NAC. Additionally, 16 (2.5%) and 63 (9.9%) patients developed HER2 positivity after NAC

in the HER2-zero and HER2-low groups, respectively (Figure 2A). The discordance rate was 36.5%, and Cohen's kappa value was 0.27.

HER2-low status is strongly associated with hormone receptor status; therefore, we analysed the HER2 transition after NAC separately by hormone receptor status. In the hormone receptor-positive (HR+) groups, 36% (n = 145) of HER2-zero before NAC and 30% (n = 158) of HER2-low before NAC showed a HER2 transition (zero to low, or vice versa) after NAC (Figure 2B). The discordance rate was 33.8%, and Cohen's kappa value was 0.31. In contrast, in the hormone receptor-negative (HR-) group, 40% (n = 43) of patients with initial HER2-low tumours before NAC developed HER2-zero tumours after NAC, whereas 18% (n = 44) of patients showed a HER2 transition (zero to low) in the initial HER2-zero group (Figure 2C). The discordance rate was 25.6%, and Cohen's kappa value was 0.40.

Figure 2. Changes in HER2 status from the primary tumour to residual disease after neoadjuvant chemotherapy. (A) Overall cohort, (B) Patients with hormone receptor - positivity, (C) Patients with hormone receptor-negativity. Abbreviation: NAC, neoadjuvant chemotherapy



3.2 Baseline patient characteristics

Table 1 shows the baseline characteristics of the patients with HER2-zero or HER2-low after NAC according to their pre- and post-therapeutic HER2 statuses. Among the patients with HER2-zero after NAC, those with constant HER2-zero tumours showed a higher proportion of nuclear and histologic grades, less HR+ (pre-therapeutic samples, 56% vs. 79%; post-therapeutic samples, 51% vs. 73%), and higher Ki-67 expression compared to those with HER2 transition (low to zero). Among the patients with HER2-low after NAC, those with constant HER2-low tumours showed more HR+ (pre-therapeutic samples, 84% vs. 77%; post-therapeutic samples, 83% vs. 77%) and less Ki-67 expression than those with HER2 changes after NAC (zero to low). However, no differences were observed in terms of nuclear and histologic grades among these patients. Furthermore, no differences were found in the distribution of residual cancer burden categories between patients with and without a HER2-change after NAC (HER2-zero post-NAC group, $p = 0.6$; HER2-low post-NAC group, $p = 0.4$).

Supplementary Table 1 presents the baseline characteristics of the patients with HER2-positive tumours after NAC according to their HER2 status before NAC. No significant differences were observed among these patients according to their HER2 status before NAC.

Table 1. Baseline characteristics of patients according to their pre-therapeutic and post-therapeutic HER2 status

Characteristic	HER2-zero after NAC (n = 646)			HER2-low after NAC (n = 563)		
	Constantly HER2-zero, N = 445 (%)	HER2-low to HER2-zero, N = 201 (%)	p-value	HER2-zero to HER2-low, N = 189 (%)	Constantly HER2-low, N = 374 (%)	p-value
Age, median (range)	46 (24–74)	46 (24–77)	0.2	46 (27–77)	47 (24–78)	0.7
Histology			0.4			0.8
Invasive ductal carcinoma	409 (92%)	185 (92%)		174 (92%)	350 (94%)	
Invasive lobular carcinoma	13 (2.9%)	9 (4.5%)		7 (3.7%)	11 (2.9%)	
Others	23 (5.2%)	7 (3.5%)		8 (4.2%)	13 (3.5%)	
Nuclear grade before NAC			0.011			0.2
1	5 (1.1%)	3 (1.5%)		3 (1.6%)	3 (0.8%)	
2	301 (68%)	157 (78%)		141 (75%)	305 (82%)	
3	139 (31%)	41 (20%)		43 (23%)	64 (17%)	
Missing	0	0		2	2	
Histologic grade before NAC			0.010			0.3
1	7 (1.6%)	3 (1.5%)		3 (1.6%)	4 (1.1%)	
2	298 (67%)	157 (78%)		141 (75%)	304 (81%)	
3	140 (31%)	41 (20%)		43 (23%)	64 (17%)	
Missing	0	0		2	2	
Clinical Stage before NAC			0.7			0.7
Stage IA	3 (0.7%)	2 (1.0%)		1 (0.5%)	1 (0.3%)	

Characteristic	HER2-zero after NAC (n = 646)			HER2-low after NAC (n = 563)		
	Constantly HER2- zero, N = 445 (%)	HER2- low to HER2- zero, N = 201 (%)	p- value	HER2- zero to HER2- low, N = 189 (%)	Constantly HER2-low, N = 374 (%)	p- value
Stage IIA	130 (29%)	54 (27%)		57 (30%)	110 (29%)	
Stage IIB	140 (31%)	75 (37%)		69 (37%)	123 (33%)	
Stage IIIA	71 (16%)	31 (15%)		24 (13%)	58 (16%)	
Stage IIIB	14 (3.1%)	8 (4.0%)		8 (4.2%)	15 (4.0%)	
Stage IIIC	86 (19%)	31 (15%)		29 (15%)	67 (18%)	
Unknown	1 (0.2%)	0 (0%)		1 (0.5%)	0 (0%)	
ER positivity before NAC	247 (56%)	158 (79%)	<0.001	145 (77%)	316 (84%)	0.024
Ki-67 status before NAC			<0.001			0.025
≤15.0%	63 (14%)	43 (21%)		38 (20%)	94 (25%)	
15.1–35.0	91 (20%)	70 (35%)		54 (29%)	133 (36%)	
>35.0	291 (65%)	88 (44%)		97 (51%)	147 (39%)	
Tumour-infiltrating lymphocytes before NAC			0.14			0.14
≤10.0%	314 (71%)	156 (78%)		137 (72%)	300 (80%)	
10.0-59.0	76 (17%)	28 (14%)		33 (17%)	47 (13%)	
>59.0%	12 (2.7%)	1 (0.5%)		2 (1.1%)	6 (1.6%)	
unknown	43 (9.7%)	16 (8.0%)		17 (9.0%)	21 (5.6%)	

Characteristic	HER2-zero after NAC (n = 646)		p-value	HER2-low after NAC (n = 563)		p-value
	Constantly HER2-zero, N = 445 (%)	HER2-low to HER2-zero, N = 201 (%)		HER2-zero to HER2-low, N = 189 (%)	Constantly HER2-low, N = 374 (%)	
Pathologic T stage			0.2			0.4
T0 or Tis	20 (4.5%)	4 (2.0%)		2 (1.1%)	10 (2.7%)	
T1 or T1mi	199 (45%)	99 (49%)		74 (39%)	149 (40%)	
T2	169 (38%)	67 (33%)		72 (38%)	154 (41%)	
T3	47 (11%)	28 (14%)		37 (20%)	55 (15%)	
T4	10 (2.2%)	3 (1.5%)		4 (2.1%)	6 (1.6%)	
Pathologic N stage			0.11			0.2
N0	175 (39%)	66 (33%)		64 (34%)	121 (32%)	
N1 or N1mi	165 (37%)	92 (46%)		75 (40%)	175 (47%)	
N2 or N3	105 (24%)	43 (21%)		50 (26%)	78 (21%)	
Neoadjuvant chemotherapy regimen			0.9			0.3
AC based	415 (93%)	188 (94%)		178 (94%)	359 (96%)	
AC	106 (24%)	55 (27%)		58 (31%)	109 (29%)	
AC → docetaxel, carboplatin 4	7 (1.6%)	2 (1.0%)		2 (1.1%)	2 (0.5%)	
AC → weekly paclitaxel 12	10 (2.2%)	1 (0.5%)		3 (1.6%)	3 (0.8%)	
AC → docetaxel 4	285 (64%)	130 (65%)		113 (60%)	241 (64%)	

Characteristic	HER2-zero after NAC (n = 646)			HER2-low after NAC (n = 563)		
	Constantly HER2- zero, N = 445 (%)	HER2- low to HER2- zero, N = 201 (%)	p- value	HER2- zero to HER2- low, N = 189 (%)	Constantly HER2-low, N = 374 (%)	p- value
Non-AC based						
AD 4	5 (1.1%)	0 (0%)		2 (1.1%)	4 (1.1%)	
FAC 6	2 (0.4%)	0 (0%)		0 (0%)	0 (0%)	
FEC → docetaxel 3	30 (6.7%)	13 (6.5%)		11 (5.8%)	15 (4.0%)	
ER status after NAC			<0.001			0.076
Negative	218 (49%)	54 (27%)		43 (23%)	62 (17%)	
Positive	226 (51%)	147 (73%)		146 (77%)	312 (83%)	
Unknown	1	0		0	0	
ER Allred score after NAC	N = 226	N = 147	0.12	N = 146	N = 312	0.5
Low (3-4)	10 (4.4%)	7 (4.8%)		8 (5.5%)	11 (3.5%)	
Moderate (5-6)	22 (9.7%)	6 (4.1%)		5 (3.4%)	8 (2.6%)	
High (7-8)	194 (86%)	134 (91%)		133 (91%)	293 (94%)	
PgR status after NAC			0.002			0.4
Negative	310 (70%)	115 (57%)		87 (46%)	188 (50.2%)	
Positive	133 (30%)	86 (43%)		102 (54%)	186 (49.7%)	
Unknown	2	0		0	0	
PgR Allred score after NAC	N = 133	N = 86	<0.001	N = 102	N = 186	0.7
Low (3-4)	47 (35%)	23 (27%)		34 (33%)	69 (37%)	
Moderate (5-6)	53 (40%)	28 (33%)		42 (41%)	68 (37%)	

Characteristic	HER2-zero after NAC (n = 646)		p-value	HER2-low after NAC (n = 563)		p-value
	Constantly HER2-zero, N = 445 (%)	HER2-low to HER2-zero, N = 201 (%)		HER2-zero to HER2-low, N = 189 (%)	Constantly HER2-low, N = 374 (%)	
High (7-8)	33 (25%)	35 (41%)		26 (25%)	49 (26%)	
Residual cancer burden			0.6			0.4
Class I	39 (8.8%)	15 (7.5%)		9 (4.8%)	22 (5.9%)	
Class II	260 (59%)	129 (64%)		111 (59%)	236 (63%)	
Class III	128 (29%)	50 (25%)		65 (34%)	105 (28%)	
Unknown	18	7		4	11	

Abbreviations: NAC, neoadjuvant chemotherapy; ER, oestrogen receptor; AC, Doxorubicin plus cyclophosphamide; AD, Doxorubicin plus docetaxel; FAC, 5-fluorouracil, Doxorubicin, cyclophosphamide; FEC, 5-fluorouracil, epirubicin, cyclophosphamide; PgR, progesterone receptor status

3.3 Association between HER2 changes and other variables

We conducted bivariable and multivariable logistic regression analyses to evaluate the association between a HER2 transition and other variables. We included the variables that showed significant differences between patients who did and did not show HER2 status changes. Among the patients with HER2-zero before NAC, patients with HR+ showed a higher probability of HER2 transition (zero to low) than those with HR- (odds ratio [OR], 2.48; 95% confidence interval [CI], 1.62–3.87) upon multivariable analysis. However, histologic grade and Ki-67 expression were not associated with a HER2 transition (Table 2A). Among the patients with HER2-low before NAC, none of the variables showed a significant association with HER2 changes (low to zero), including ER status (OR, 0.7; 95% CI, 0.42–1.15; $p=0.2$, Table 2B).

Table 2A. Bivariable logistic regression analysis of HER2 status after NAC (HER2-zero vs. HER2-low) among the patients with HER2-zero before NAC by hormone receptor status and other variables.

Characteristic	Bivariable			Multivariable		
	OR	95% CI	p-value	OR	95% CI	p-value
Histologic grade G3 (vs. G1 or G2)	0.67	0.45–0.97	0.036	1.13	0.71–1.78	0.6
ER positivity before NAC (vs. negative)	2.72	1.88–3.99	<0.001	2.48	1.62–3.87	<0.001
Ki-67 expression						
15.1–35.0 (vs. ≤15%)	1.05	0.63–1.76	0.86	1.08	0.64–1.84	0.8
>35.0 (≤15%)	0.59	0.38–0.94	0.025	0.8	0.48–1.34	0.4

Abbreviations: NAC, neoadjuvant chemotherapy; ER, oestrogen receptor; OR, odds ratio; CI, confidence interval

Table 2B. Bivariable logistic regression analysis of HER2 status after NAC (HER2-low vs. HER2-zero) among the patients with HER2-low before NAC by hormone receptor status and other variables.

Characteristic	Bivariable			Multivariable		
	OR	95% CI	p-value	OR	95% CI	p-value
Histologic grade G3 (vs. G1 or G2)	1.25	0.82–1.91	0.30	0.97	0.58–1.61	>0.9
ER positivity before NAC (vs. negative)	0.67	0.43–1.03	0.066	0.7	0.42–1.15	0.2
Ki-67 expression						
15.1–35.0 (vs. ≤15%)	1.16	0.74–1.83	0.53	1.14	0.72–1.82	0.6
>35.0 (≤15%)	1.33	0.86–2.07	0.21	1.19	0.73–1.95	0.5

Abbreviations: NAC, neoadjuvant chemotherapy; ER, oestrogen receptor; OR, odds ratio; CI, confidence interval

3.4 Prognostic implications of HER2 changes for clinical outcomes


Supplementary Figure 1 shows the OS and DFS according to the post-therapeutic HER2 status. The 5-year OS rates were 82.4% (95% CI, 79.4–85.5), 90.2% (95% CI, 87.6–92.8), and 92.9% (95% CI, 87.0–99.1) in the HER2-zero, HER2-low, and HER2-positive

groups, respectively. Furthermore, the 5-year DFS rates were 71.2% (95% CI, 67.4–75.2), 71.7% (95% CI, 67.7–75.9), and 77.4% (95% CI, 68.5–87.5) in the HER2-zero, HER2-low, and HER2-positive groups, respectively.

To investigate the prognostic implications of HER2 changes, we compared OS and DFS according to pre- and post-therapeutic HER2 statuses, as well as hormone receptor status (Figures 3, 4). Among the patients with post-therapeutic HER2-zero tumours, those who showed a HER2 transition (low to zero) had better OS and DFS compared to those with constant HER2-zero tumours (OS, $p = 0.0009$, Figure 3A; DFS, $p = 0.00028$, Figure 3B). However, when analysing hormone receptor status separately, no differences were observed in terms of OS and DFS according to HER2 changes among the HR+ subsets (OS, $p = 0.064$, Figure 3C; DFS, $p = 0.053$, Figure 3D) and HR- subsets (OS, $p = 0.29$, Figure 3E; DFS, $p = 0.12$, Figure 3F).

Figure 3. Clinical outcomes of patients with HER2-zero after NAC according to HER2 status before NAC. (A) OS of all patients, (B) DFS of all patients, (C) OS of patients with hormone receptor positivity, (D) DFS of patients with hormone receptor positivity, (E) OS of patients with hormone receptor negativity, (F) DFS of

patients with hormone receptor negativity. Abbreviations: NAC, neoadjuvant chemotherapy; OS, overall survival; DFS, disease-free survival




Furthermore, among the patients with HER2-low in the residual tumours, no significant differences were observed in terms of OS and DFS according to HER2 transition

in the overall cohort (OS, $p = 0.067$, Figure 4A; DFS, $p = 0.51$, Figure 4B) and HR+ subsets (OS, $p = 0.4$, Figure 4C; DFS, $p = 0.17$, Figure 4D). However, among patients with HR-tumours, those with a HER2 transition (zero to low) showed significantly worse OS ($p = 0.0054$) and DFS ($p = 0.0041$) than those with constant HER2-low statuses (Figure 4E, 4F).

Figure 4. Clinical outcomes of patients with HER2-low after NAC according to HER2 status before NAC. (A) OS of all patients, (B) DFS of all patients, (C) OS of patients with hormone receptor positivity, (D) DFS of patients with hormone receptor positivity, (E) OS of patients with hormone receptor negativity, (F) DFS of

patients with hormone receptor negativity. Abbreviations: NAC, neoadjuvant chemotherapy; OS, overall survival; DFS, disease-free survival



3.5 Factors associated with clinical outcomes

Tables 3A and 3B show the results of the univariate and multivariate analyses in terms of OS and DFS among patients with HER2-zero and HER2-low after NAC, respectively. Among the patients with HER2-zero in the residual tumours, a HER2 transition (low to zero) was significantly associated with OS (hazard ratio [HR] 0.61; 95% CI, 0.38–0.99; $p = 0.047$) and DFS (HR 0.64; 95% CI, 0.44–0.93; $p = 0.02$, Table 3A). However, among the patients with post-therapeutic HER2-low, a HER2 transition (zero to low) was not associated with OS (HR 1.16; 95% CI, 0.70–1.91; $p = 0.6$) or DFS (HR 1.02; 95% CI, 0.72–1.44; $p > 0.9$) (Table 3B).

Table 3A. Univariable and multivariable analyses of patients with HER2-zero after NAC

Characteristic	Overall survival						Disease-free survival					
	Univariable			Multivariable			Univariable			Multivariable		
	H R	95 % CI	p- value	H R	95 % CI	p- value	H R	95 % CI	p- value	H R	95 % CI	p- value
Age < 40 (vs. ≥ 40)	1.42	0.97–2.09	0.075	1.47	0.99–2.17	0.057	1.33	0.97–1.83	0.076	1.38	1.00–1.92	0.052
Clinical T stage – T3, T4 (vs. T1, T2)	1.75	1.21–2.55	0.003	1.57	1.07–2.29	0.021	1.33	0.97–1.83	0.078	1.25	0.91–1.73	0.2
Clinical N stage – N+ (vs. N0)	2.21	1.35–3.62	0.002	2.55	1.54–4.22	<0.001	1.77	1.22–2.55	0.002	2.07	1.42–3.02	<0.001
Histologic grade G3 (vs. G1, G2)	1.76	1.21–2.55	0.003	1.02	0.68–1.52	>0.9	1.66	1.22–2.25	0.001	1.15	0.82–1.61	0.4

Characteristic	Overall survival						Disease-free survival					
	Univariable			Multivariable			Univariable			Multivariable		
	H R	95 % CI	p- value	H R	95 % CI	p- value	H R	95 % CI	p- value	H R	95 % CI	p- value
Ki-67 expression (%)												
15.1–35.0 (vs. ≤15.0)	5.99	1.38–25.9	0.017	5.88	1.35–25.5	0.018	2.16	1.09–4.27	0.026	1.91	0.96–3.81	0.065
>35.0 (vs. ≤5.0)	15.2	3.76–61.8	<0.001	9.76	2.35–40.5	0.002	3.91	2.12–7.24	<0.001	2.43	1.26–4.66	0.008
ER negative before NAC (vs. positive)	3.07	2.11–4.47	<0.001	2.12	1.40–3.22	<0.001	2.51	1.86–3.37	<0.001	2.04	1.46–2.85	<0.001
Pathologic T stage												
T1mi, T1, T2 (vs. T0, Tis)	1.02	0.38–2.78	0.97				1.03	0.45–2.33	0.95			
T3, T4 (vs. T0, Tis)	1.75	0.61–5.07	0.30				1.69	0.71–4.04	0.23			
Pathologic N stage												
N1, N1mi (vs. N0)	1.24	0.78–1.97	0.37				0.96	0.66–1.40	0.85			
N2, N3 (vs. N0)	2.63	1.66–4.17	<0.001				2.30	1.61–3.29	<0.001			
ER negative after NAC (vs. positive)	5.26	3.42–8.10	<0.001				3.27	2.40–4.46	<0.001			
HER2 transition from low to zero (vs. constantly zero)	0.46	0.29–0.74	0.001	0.61	0.38–0.99	0.047	0.52	0.36–0.74	<0.001	0.64	0.44–0.93	0.02

Abbreviations: NAC, neoadjuvant chemotherapy; ER, oestrogen receptor; HR, hazard ratio; CI, confidence interval

Table 3B. Univariable and multivariable analyses of patients with HER2-low after NAC

		Overall survival						Disease-free survival					
		Univariable			Multivariable			Univariable			Multivariable		
Characteristic		HR	95% CI	p-value	HR	95% CI	p-value	HR	95% CI	p-value	HR	95% CI	p-value
Age <40 (vs. ≥40)		1.6	0.98	0.057	1.1	0.68	0.6	1.6	1.20	0.003	1.4	1.06	0.023
		3	–		4	–		8	–		9	–	
			2.69			1.92			2.37			2.10	
Clinical stage – T3, T4 (vs. T1, T2)	T	2.1	1.34	0.002	2.0	1.27	0.004	1.6	1.17	0.004	1.5	1.12	0.010
		7	–		9	–		4	–		6	–	
			3.50			3.43			2.28			2.19	
Clinical stage – N+ (vs. N0)	N	4.2	1.95	<0.00	5.6	2.52	<0.00	2.7	1.79	<0.00	3.0	1.96	<0.00
		7	–	1	3	–	1	8	–	1	7	–	1
			9.35			12.5			4.30			4.80	
Histologic grade G3 (vs. G1, G2)		3.9	2.44	<0.00	1.9	1.04	0.036	2.1	1.52	<0.00	1.6	1.03	0.037
		3	–	1	5	–		6	–	1	2	–	
			6.34			3.64			3.06			2.56	
Ki67 expression (%)													
15.1-35.0 (vs. ≤15.0)		1.1	0.46	0.72	0.8	0.34	0.8	1.1	0.70	0.59	0.9	0.59	0.9
		9	–		8	–		5	–		6	–	
			3.08			2.30			1.87			1.58	
>35.0 (vs. ≤15.0)		4.4	2.01	<0.00	1.9	0.83	0.12	2.0	1.33	0.001	1.2	0.79	0.3
		4	–	1	9	–		7	–		9	–	
			9.79			4.73			3.21			2.10	

		Overall survival						Disease-free survival					
		Univariable			Multivariable			Univariable			Multivariable		
Characteristic		HR	95% CI	p-value	HR	95% CI	p-value	HR	95% CI	p-value	HR	95% CI	p-value
ER negative before NAC (vs. positive)		4.4	2.77	<0.00	2.8	1.58	<0.00	2.1	1.50	<0.00	1.7	1.09	0.019
		6	–	1	9	–	1	5	–	1	1	–	
			7.20			5.31			3.09			2.68	
Pathologic T stage													
	T1mi, T1,	0.5	0.12	0.36				0.8	0.27	0.79			
	T2 (vs. T0, Tis)	2	–					6	–				
			2.13						2.71				
	T3, T4 (vs. T0, Tis)	1.4	0.33	0.64				2.2	0.69	0.18			
		2	–					1	–				
			6.01						7.12				
Pathologic N stage													
	N1, N1mi (vs. N0)	1.6	0.81	0.17				1.8	1.16	0.008			
		5	–					1	–				
			3.36						2.82				
	N2, N3 (vs. N0)	5.1	2.59	<0.00				4.0	2.55	<0.00			
		3	–	1				1	–	1			
			10.2						6.32				
ER negative after NAC (vs. positive)		4.5	2.84	<0.00				2.3	1.63	<0.00			
		8	–	1				2	–	1			
			7.38						3.29				
HER2 transition from zero to		1.5	0.97	0.069	1.1	0.70	0.6	1.1	0.80	0.51	1.0	0.72	>0.9
		6	–		6	–		2	–		2	–	
			2.53			1.91			1.57			1.44	

Overall survival							Disease-free survival					
		Univariable			Multivariable		Univariable			Multivariable		
Characteristi c		HR	95% CI	p- value	HR	95% CI	HR	95% CI	p- value	HR	95% CI	p- value
low	(vs.											
constantly low)												

Abbreviations: NAC, neoadjuvant chemotherapy; ER, oestrogen receptor; HR, hazard ratio; CI, confidence interval

4 Discussion

In this study, we evaluated the changes in HER2-low expression among patients who underwent NAC and curative surgical resection. The discordance rate between the primary tumour and residual disease was 36.5%, indicating the temporal heterogeneity of HER2-low expression. Additionally, we found a higher proportion of HR+ among patients with constant HER2-low (84%) or a transition from HER2-low to zero (79%) than in those with changes from HER2-zero to low (77%) or with constant HER2-zero (56%), respectively, suggesting that HER2-low expression has a strong relationship with HR+. These results are consistent with previous reports that showed high instability of HER2-low expression in residual disease after NAC[17,18] and an association between HER2-low and HR+[19].

Previously, two studies evaluated the evolution of HER2-low expression among patients with relapsed or metastatic breast cancer and reported a significant discordance rate between primary and recurred or metastatic tumours (38% and 66%, respectively)[20,21]. Notably, these studies observed that HER2-expression was more unstable among the HR+

subgroups compared to the HR- subgroups[22]. For example, in Miglietta et al.'s study[20], in the HR+ subgroup, 47.3% and 53.8% were HER2-low in the primary and matched relapsed tumours, respectively. In contrast, in the HR- subgroup, 35.4% and 36.2% were HER2-low. These results are similar to those of Tarantino et al.[21], which showed a greater degree of instability in HER2-low expression among patients with HR+ (57% vs. 64% in primary and metastatic tumours, respectively) than in those with HR- (36% vs. 38%, respectively). Our results are similar to their findings, as follows: in the HR+ subgroup, 56.7% (531/936) were HER2-low before NAC, and 49.2% (461/936) were HER2-low after NAC, whereas 30.3% (107/352) were HER2-low before NAC and 28.9% (102/352) were HER2-low after NAC in the HR- subgroup. Additionally, as shown in the multivariable logistic regression analysis, patients with ER-positivity had a higher probability of gaining HER2-low expression than those with ER-negativity (OR, 2.48; $p < 0.001$). Therefore, we hypothesize that HER2-low expression appears to be highly unstable both in residual disease and advanced breast cancer, and HR+ may affect changes in HER2-low after anti-cancer treatment.

The mechanisms underlying the dynamic changes in HER2 expression after NAC remain unclear; however, one possible explanation is that therapeutic selective pressure due to chemotherapy results in the up- or down-regulation of HER2 expression as resistance mechanisms[21,23]. Furthermore, inter-observer variations of HER2-low expression may be related to the discordance rate of HER2 status. In fact, distinguishing HER2-low, especially IHC0 vs. IHC1+, is highly variable among pathologists[24] because current ASCO/CAP 2018 guidelines for HER2 measurements have not been developed for distinguishing lower levels of HER2 expression[15]. The results of the Destiny-Breast 06 trial (NCT04494425), which includes a mandatory central assessment of HER2 status, is expected to provide a

better understanding of this limitation[25]. Additionally, developing reliable quantitative HER2 testing methods are required for accurately identifying lower levels of HER2 expression in the future.

Currently, only a few studies have reported the prognostic implications of a HER2-change (including HER2-low) following NAC[19,26]. According to Miglietta et al., there was no significant association between changes in HER2 status and DFS when compared to patients who showed concordant HER2 status (constant HER2-low vs. loss of HER2-low [low to zero], $p = 0.23$; constant HER2-zero vs. gaining HER2-low [zero to low], $p = 0.77$)[19]. In contrast, at the 2022 San-Antonio Breast Cancer Symposium, the German Breast Group (GBG) presented new findings suggesting that a shift from HER2-zero to HER2-low was significantly associated with reduced invasive DFS (iDFS) relative to the constant HER2-low group (constant HER2-low vs. gaining HER2-low [zero to low], HR 1.43; 95% CI, 1.01–2.01; $p = 0.04$) in patients with HR+ tumours after NAC[26].

These inconsistent results might be attributable to the differences in the included patients' characteristics. For example, Miglietta et al. showed the results of the entire population without separately analysing the clinical outcomes based on hormone receptor status. In contrast, the GBG study focused exclusively on patients with HR+ tumours. Additionally, Miglietta et al. compared within the patients who showed initial HER2-zero or HER2-low, whereas the GBG study compared among the patients with post-therapeutic HER2-low. Considering the differences in the included patients' characteristics and the inconsistent results between the two aforementioned studies[19,26], it is challenging to directly compare these findings and reach definitive conclusions regarding the prognostic role of dynamic changes of HER2-low expression after NAC.

In our study, we observed that the transition from HER2-low to HER2-zero was associated with better OS and DFS in the overall cohort compared to those with constant HER2-zero tumours (Figure 3A–3B). These results are also supported by the multivariate analyses. However, when we analysed the data separately according to the hormone receptor status, no significant differences were observed in terms of OS and DFS, as depicted in Figure 3C–3F. Notably, in the HR+ subset, the shift from HER2-zero to HER2-low was not significantly associated with a decreased DFS or OS compared to constant HER2-low (Figure 4C–4D), which contrasts with the findings of the GBG study. An intriguing finding from our study is that tumours gaining HER2-low expression after NAC exhibited significantly worse OS and DFS compared to the constant HER2-low group in the HR- subset (Figure 4E–4F).

To the best of our knowledge, this is the first study that analysed the clinical outcomes separately according to hormone receptor status and the transition of HER2-low expression after NAC. Based on our findings, we assume that gaining HER2-low from HER2-zero tumour may not be a significant prognostic factor in HR+ tumours, but it might be related to DFS and OS in HR- tumours. While our study provides valuable insights, it is important to note that definitive conclusions cannot be drawn at this time due to the limited evidence available. Further investigation is warranted to understand the prognostic role of the transition of HER2-low after NAC. We believe that our study's findings will contribute to designing future studies in this field.

Lastly, HER2-low is now considered a predictive biomarker of the efficacy of T-DXd in metastatic breast cancer[2], and the application of T-DXd is now moving from an advanced setting to an earlier setting, such as neoadjuvant chemotherapy[27,28]. As of today, no anti-HER2-targeting antibodies have been approved for adjuvant therapy of HER2-low tumours,

since trastuzumab did not show clinically significant benefits among these patients[29]. Owing to the poorer outcomes of patients who failed to achieve pathologically complete responses after NAC, there are still unmet needs for better adjuvant chemotherapy for HER2-low breast cancer, and a future clinical trial is required to investigate the efficacy of anti-HER2 ADCs as adjuvant chemotherapy for HER2-low breast cancer. However, as shown in our study, HER2-low expression is highly discordant between the primary tumour and residual disease. Therefore, it is largely unknown which patients would benefit most from anti-HER2 ADC adjuvant chemotherapy: those expressing HER2-low continuously or those gaining or losing HER2-low expression after NAC, or both. In future clinical trials, it may be necessary to test the HER2 status of both the primary tumour and residual disease after NAC to answer this question, and this may help with identifying optimal candidates for adjuvant chemotherapy with anti-HER2 ADC.

Our study had several limitations. First, this was a single-centre retrospective study, which is inherently susceptible to selection bias. However, the relatively large number of patients included in this analysis and the similar treatment strategy used for most patients may overcome this limitation. Additionally, while there was no central review of HER2 expression for the present study, experienced pathologists reviewed the HER2 status within the same institution based on current ASCO/CAP guidelines; therefore, our findings are considered reliable. Second, we did not evaluate the genomic profiles and molecular statuses in our cohort, which may help with understanding the biology of HER2-low tumours. Third, we only included Korean patients. Considering the ethnic disparities in the characteristics and clinical outcomes of breast cancer, our results may be insufficient to generalize to all patients.

In conclusion, temporal heterogeneity of HER2-low expression is observed in substantial numbers of post-NAC breast cancer patients. The clinical outcomes showed no significant associations with changes in HER2 status, except in the post-therapeutic HER2-low, HR- subset. Therefore, the prognostic implications of a HER2 transition in HER2-low breast cancer require further investigation.

Authors' contributions

Sung-Bae Kim contributed to the study conception and design. Acquisition of data was performed by So Heun Lee, Hee Jin Lee, Hye Hyun Jeong, Jae Ho Jeong, Jeong Eun Kim, Jin-Hee Ahn, Kyung Hae Jung, Gyungyub Gong, Hak Hee Kim, Saebyeol Lee, Jongwon Lee, and Sora Kang. Data analysis and interpretation were performed by Sora Kang and Sung-Bae Kim. The first draft of the manuscript was written by Sora Kang, and all authors commented on subsequent versions of the manuscript. All authors read and approved the final manuscript.

Data statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

References

- [1] Harbeck N, Penault-Llorca F, Cortes J, et al. Breast cancer. *Nature Reviews Disease Primers* 2019;5(1):66. <https://doi.org/10.1038/s41572-019-0111-2>.
- [2] Modi S, Jacot W, Yamashita T, et al. Trastuzumab Deruxtecan in Previously Treated HER2-Low Advanced Breast Cancer. *New England Journal of Medicine* 2022;387(1):9-20. <https://doi.org/10.1056/NEJMoa2203690>.
- [3] Tarantino P, Hamilton E, Tolaney SM, et al. HER2-Low Breast Cancer: Pathological and Clinical Landscape. *Journal of Clinical Oncology* 2020;38(17):1951-62. <https://doi.org/10.1200/JCO.19.02488>.
- [4] Schettini F, Chic N, Brasó-Maristany F, et al. Clinical, pathological, and PAM50 gene expression features of HER2-low breast cancer. *npj Breast Cancer* 2021;7(1):1. <https://doi.org/10.1038/s41523-020-00208-2>.
- [5] Agostinetto E, Rediti M, Fimereli D, et al. HER2-Low Breast Cancer: Molecular Characteristics and Prognosis. *Cancers* 2021;13(11). <https://doi.org/10.3390/cancers13112824>.
- [6] Tarantino P, Jin Q, Tayob N, et al. Prognostic and Biologic Significance of ERBB2-Low Expression in Early-Stage Breast Cancer. *JAMA Oncology* 2022;8(8):1177-83. <https://doi.org/10.1001/jamaoncol.2022.2286>.
- [7] Kang S, Lee SH, Lee HJ, et al. Pathological complete response, long-term outcomes, and recurrence patterns in HER2-low versus HER2-zero breast cancer after neoadjuvant chemotherapy. *European Journal of Cancer* 2022;176:30-40. <https://doi.org/10.1016/j.ejca.2022.08.031>.
- [8] Denkert C, Seither F, Schneeweiss A, et al. Clinical and molecular characteristics of HER2-low-positive breast cancer: pooled analysis of individual patient data from four prospective, neoadjuvant clinical trials. *The Lancet Oncology* 2021;22(8):1151-61. [https://doi.org/10.1016/S1470-2045\(21\)00301-6](https://doi.org/10.1016/S1470-2045(21)00301-6).
- [9] Mutai R, Barkan T, Moore A, et al. Prognostic impact of HER2-low expression in hormone receptor positive early breast cancer. *The Breast* 2021;60:62-9. <https://doi.org/https://doi.org/10.1016/j.breast.2021.08.016>.
- [10] Untch M, Konecny GE, Paepke S, et al. Current and future role of neoadjuvant therapy for breast cancer. *The Breast* 2014;23(5):526-37. <https://doi.org/https://doi.org/10.1016/j.breast.2014.06.004>.

- [11] Parinyanitikul N, Lei X, Chavez-MacGregor M, et al. Receptor Status Change From Primary to Residual Breast Cancer After Neoadjuvant Chemotherapy and Analysis of Survival Outcomes. *Clinical Breast Cancer* 2015;15(2):153-60. <https://doi.org/10.1016/j.clbc.2014.09.006>.
- [12] Yoshida A, Hayashi N, Suzuki K, et al. Change in HER2 status after neoadjuvant chemotherapy and the prognostic impact in patients with primary breast cancer. *Journal of Surgical Oncology* 2017;116(8):1021-8. <https://doi.org/https://doi.org/10.1002/jso.24762>.
- [13] Ven Svd, Smit VTHBM, Dekker TJA, et al. Discordances in ER, PR and HER2 receptors after neoadjuvant chemotherapy in breast cancer. *Cancer Treatment Reviews* 2011;37(6):422-30. <https://doi.org/10.1016/j.ctrv.2010.11.006>
- [14] Niikura N, Tomotaki A, Miyata H, et al. Changes in tumor expression of HER2 and hormone receptors status after neoadjuvant chemotherapy in 21 755 patients from the Japanese breast cancer registry. *Annals of Oncology* 2016;27(3):480-7. <https://doi.org/https://doi.org/10.1093/annonc/mdv611>.
- [15] Wolff AC, Hammond MEH, Allison KH, et al. Human Epidermal Growth Factor Receptor 2 Testing in Breast Cancer: American Society of Clinical Oncology/College of American Pathologists Clinical Practice Guideline Focused Update. *J Clin Oncol* 2018;36(20):2105-22. <https://doi.org/10.1200/jco.2018.77.8738>.
- [16] Cohen J. A Coefficient of Agreement for Nominal Scales. *Educational and Psychological Measurement* 1960;20(1):37-46. <https://doi.org/10.1177/001316446002000104>.
- [17] Bo J, Yu B, Bi R, et al. Conversion of ER and HER2 status after neoadjuvant therapy in Chinese breast cancer patients. *Clinical Breast Cancer* 2023. <https://doi.org/10.1016/j.clbc.2023.03.002>.
- [18] Ma Y, Zhu M, Zhang J, et al. Prognostic Value of the Evolution of HER2-Low Expression after Neoadjuvant Chemotherapy. *Cancer Research and Treatment* 2023. <https://doi.org/10.4143/crt.2022.1633>.
- [19] Miglietta F, Griguolo G, Bottosso M, et al. HER2-low-positive breast cancer: evolution from primary tumor to residual disease after neoadjuvant treatment. *npj Breast Cancer* 2022;8(1):66. <https://doi.org/10.1038/s41523-022-00434-w>.
- [20] Miglietta F, Griguolo G, Bottosso M, et al. Evolution of HER2-low expression from primary to recurrent breast cancer. *npj Breast Cancer* 2021;7(1):137. <https://doi.org/10.1038/s41523-021-00343-4>.

- [21] Tarantino P, Gandini S, Nicolò E, et al. Evolution of low HER2 expression between early and advanced-stage breast cancer. *European Journal of Cancer* 2022;163:35-43.
<https://doi.org/10.1016/j.ejca.2021.12.022>.
- [22] Molinelli C, Jacobs F, Marchiò C, et al. HER2-Low Breast Cancer: Where Are We? *Breast Care* 2022;17(6):533-45. <https://doi.org/10.1159/000527391>.
- [23] Kan S, Koido S, Okamoto M, et al. Gemcitabine treatment enhances HER2 expression in low HER2-expressing breast cancer cells and enhances the antitumor effects of trastuzumab emtansine. *Oncology Reports* 2015;34(1):504-10. <https://doi.org/10.3892/or.2015.3974>.
- [24] Fernandez AI, Liu M, Bellizzi A, et al. Examination of Low ERBB2 Protein Expression in Breast Cancer Tissue. *JAMA Oncology* 2022;8(4):1-4. <https://doi.org/10.1001/jamaoncol.2021.7239>.
- [25] Study of Trastuzumab Deruxtecan (T-DXd) vs Investigator's Choice Chemotherapy in HER2-low, Hormone Receptor Positive, Metastatic Breast Cancer (DB-06).
<https://clinicaltrials.gov/ct2/show/NCT04494425>; [accessed June 14 2023].
- [26] Denkert C, Martín M, Untch M, et al. Abstract HER2-06: HER2-06 Outcome analysis of HER2-zero or HER2-low hormone receptor-positive (HR+) breast cancer patients - characterization of the molecular phenotype in combination with molecular subtyping. *Cancer Research* 2023;83(5_Supplement):HER2-06-HER2-. <https://doi.org/10.1158/1538-7445.Sabcs22-her2-06>.
- [27] Hurvitz SA, Peddi PF, Tetef ML, et al. TRIO-US B-12 TALENT: Phase II neoadjuvant trial evaluating trastuzumab deruxtecan with or without anastrozole for HER2-low, HR+ early stage breast cancer. *Journal of Clinical Oncology* 2021;39(15_suppl):TPS603-TPS.
https://doi.org/10.1200/JCO.2021.39.15_suppl.TPS603.
- [28] TRUDI: T-DXd+Durvalumab in HER2+/Low IBC (TRUDI). *ClinicalTrials.gov* identifier NCT05795101. Available here. <https://clinicaltrials.gov/ct2/show/NCT05795101>; [accessed 23, May 2023].
- [29] Fehrenbacher L, Cecchini RS, Geyer CE, Jr., et al. NSABP B-47/NRG Oncology Phase III Randomized Trial Comparing Adjuvant Chemotherapy With or Without Trastuzumab in High-Risk Invasive Breast Cancer Negative for HER2 by FISH and With IHC 1+ or 2. *Journal of Clinical Oncology* 2020;38(5):444-53. <https://doi.org/10.1200/jco.19.01455>.

Figure legends

Supplementary material

Supplementary Table 1. Baseline characteristics of patients with HER2-positive tumours after neoadjuvant chemotherapy according to HER2-status before NAC.

Supplementary Figure 1. (A) Overall survival and (B) Disease-free survival according to post-therapeutic HER2 status

Author contributions

Sora Kang: data curation, formal analysis, writing, review and editing of the manuscript

So Heun Lee: data curation, writing, review and editing of the manuscript

Hee Jin Lee: review and editing of the manuscript

Hychyun Jeong: data curation, review and editing of the manuscript

Jae Ho Jeong: data curation, review and editing of the manuscript

Jeong Eun Kim: data curation, review and editing of the manuscript

Jin-Hee Ahn: data curation, review and editing of the manuscript

Kyung Hae Jung: data curation, review and editing of the manuscript

Gyungyub Gong: review and editing of the manuscript

Hak Hee Kim: review and editing of the manuscript

Saebyeol Lee: review and editing of the manuscript

Jongwon Lee: review and editing of the manuscript

Sung-Bae Kim: conceptualization, methodology, project administration, supervision, writing, review and editing of the manuscript

Declaration of Competing Interest

☐ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☒ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Kyung Hae Jung has advisory roles at Astra-Zeneca, BIXINK, MSD, Novartis, Pfizer, Roche, Takeda and Everest Medicine. Sung-Bae Kim is a consultant on the advisory boards of Novartis, AstraZeneca, Lilly, Dae Hwa Pharmaceutical Co. Ltd, ISU Abxis, and Daiichi-Sankyo, and has received research funding from Novartis, Sanofi-Aventis, and DongKook Pharm Co., and owns stock in Genopeaks and NeogeneTC. None of the other authors has any conflicts of interest to declare in relation to this study.

Highlights

- HER2-low expression is discordant between the primary tumour and residual disease
- HER2-low expression is unstable both in residual disease and advanced breast cancer
- HER2-low expression has a strong relationship with hormone-receptor positivity
- Changes in HER2 expression do not reflect a significant prognostic role