

Low Locoregional Recurrence Rate Among Node-Negative Breast Cancer Patients With Tumors 5 cm or Larger Treated by Mastectomy, With or Without Adjuvant Systemic Therapy and Without Radiotherapy: Results From Five National Surgical Adjuvant Breast and Bowel Project Randomized Clinical Trials

Alphonse G. Taghian, Jong-Hyeon Jeong, Eleftherios P. Mamounas, David S. Parda, Melvin Deutsch, Joseph P. Costantino, and Norman Wolmark

From the National Surgical Adjuvant Breast and Bowel Project Operations and Biostatistical Centers; Department of Biostatistics, Graduate School of Public Health, University of Pittsburgh; Department of Radiation Oncology, University of Pittsburgh Medical Center; Departments of Radiation Oncology and Human Oncology, Allegheny General Hospital, Pittsburgh, PA; Department of Radiation Oncology, Massachusetts General Hospital and Boston Medical Center, Boston, MA; Aultman Cancer Center, Canton; and Northeastern Ohio Universities College of Medicine, Kootstown, OH.

Submitted April 5, 2006; accepted June 16, 2006.

Supported by Public Health Service Grants No. U10CA-12027, U10CA-37377, U10CA-69974, and U10CA-69651 from the National Cancer Institute, Department of Health and Human Services.

Authors' disclosures of potential conflicts of interest and author contributions are found at the end of this article.

Address reprint requests to Alphonse G. Taghian, MD, PhD, Department of Radiation Oncology, Massachusetts General Hospital; 55 Fruit St, Boston, MA 02114; e-mail: ataghian@partners.org.

© 2006 by American Society of Clinical Oncology

0732-183X/06/2424-3927/\$20.00

DOI: 10.1200/JCO.2006.06.9054

ABSTRACT

Purpose

Lymph node (LN) –negative breast cancer tumors ≥ 5 cm occur infrequently, and their optimal management is not well defined. In this study, we assess patterns of locoregional failure (LRF) in LN-negative patients who underwent mastectomy, either with or without adjuvant chemotherapy or hormonal therapy and without postmastectomy radiation therapy (PMRT).

Patients and Methods

Of 8,878 breast cancer patients enrolled onto National Surgical Adjuvant Breast and Bowel Project B-13, B-14, B-19, B-20, and B-23 node-negative trials, 313 had tumors that were 5 cm or larger (median, 5.5 cm; range, 5.0 to 15.5 cm) at pathology and were treated by mastectomy. Median follow-up time was 15.1 years. Therapy included adjuvant chemotherapy in 34.2% of patients; tamoxifen in 21.1%; chemotherapy plus tamoxifen in 19.2%; and no systemic therapy in 25.5%.

Results

Twenty-eight patients experienced LRF. The overall 10-year cumulative incidences of isolated LRF, LRF with and without distant failure (DF), and DF alone as first event were 7.1%, 10.0%, and 23.6%, respectively. Cumulative incidences for isolated LRF as first event for patients with tumors of 5 cm or more than 5 cm were 7.0% and 7.2%, respectively ($P = .9$). For patients who underwent no systemic treatment, chemotherapy alone, tamoxifen alone, or chemotherapy plus tamoxifen, the incidences were 12.6%, 5.6%, 4.6%, and 5.3%, respectively ($P = .2$). The majority of failures occurred on the chest wall (24 of 28 patients). Multivariate analysis did not identify significant prognostic factors for LRF.

Conclusion

In patients with LN-negative tumors ≥ 5 cm who are treated by mastectomy with or without adjuvant systemic therapy and no PMRT, LRF as first event remains low. PMRT should not be routinely used for these patients.

J Clin Oncol 24:3927-3932. © 2006 by American Society of Clinical Oncology

INTRODUCTION

Patients with node-negative breast cancer and primary tumors 5 cm or larger represent less than 1% of new breast cancer patients,¹ and optimal management for these patients is not well defined. Although standard recommendations for these patients include systemic treatment in addition to surgery, the addition of radiation therapy for improved local control or survival is debatable. The postmastectomy radiation therapy (PMRT) guidelines of the

American Society of Clinical Oncology² provide no recommendations for T3N0 tumors because of a lack of information and conflicting data. Yet, the majority of practicing radiation oncologists recommend PMRT for these tumors.³

In this study, we examine the rates and patterns of locoregional failure (LRF) in 313 breast cancer patients whose tumors were ≥ 5 cm at pathology and who, per protocol, did not receive PMRT as part of five randomized clinical trials conducted by the National Surgical Adjuvant Breast and

Bowel Project (NSABP). These patients underwent axillary lymph node (LN) dissection and were treated with modified radical mastectomy (MRM), with or without adjuvant chemotherapy or adjuvant hormonal therapy.

PATIENTS AND METHODS

Study Population

Patients included in this report were treated in five NSABP node-negative trials (B-13, B-14, B-19, B-20, and B-23) evaluating adjuvant chemotherapy and/or hormonal therapy but not allowing for adjuvant PMRT. To be included in this analysis, a patient had to be eligible with follow-up, had to have undergone MRM, and had to have had a pathologic tumor ≥ 5 cm in diameter. A brief summary of each of the five trials is provided in Table 1. Treatment details, outcome, and other aspects of each trial have been reported in previous publications (B-13,⁴ B-14,^{5,6} B-19,⁷ B-20,⁸ and B-23⁹). All trials were approved by local human investigations committees or institutional review boards in accordance with assurances filed with and approved by the Department of Health and Human Services, when appropriate. Written informed consent was required for participation in the trials.

Statistical Methods

In NSABP studies, local failure after mastectomy is defined as any recurrence of tumor in the ipsilateral chest wall or in the mastectomy scar. Regional failure is defined as any recurrence of tumor in the ipsilateral supraclavicular, infraclavicular, axillary, or internal mammary nodes. Recurrence in any other site is considered distant failure (DF). For our study, isolated LRF was defined as any first local or regional failure without evidence of simultaneous DF. Simultaneous DF is defined as any subsequent DF that occurred within 4 months after the diagnosis of the isolated LRF. The occurrence of contralateral events and other second primary cancers was ignored in the determination of (isolated) locoregional recurrences. LRF with or without DF (LRF \pm DF) was defined as any first LRF with or without any subsequent DF.

Time to isolated LRF was defined as time from definitive surgery to the first diagnosis of an isolated LRF. Time to DF was defined as time from surgery to the first DF. Local or regional failures occurring without a subsequent DF but with additional follow-up of less than 4 months were counted as isolated events.

The nonparametric method¹⁰ was used to estimate 10-year cumulative incidence for isolated LRF, LRF \pm DF, and DF alone. For the analysis of cumulative incidence for each end point, competing events were defined as (1) any first isolated recurrence, (2) any first recurrence followed by a DF within 4 months, (3) any first DF, and (4) any death without evidence of previous recurrence. Gray's K-sample statistic was used to test whether any statistical significance of differences existed in cumulative incidence among groups stratified by protocols (univariate analysis).¹¹ The Cox proportional hazards model¹² was used to test the association between

cause-specific hazard functions and selected patient and tumor characteristics and to estimate the magnitude of such association, stratified by protocols. All *P* values were two tailed, and *P* $\leq .05$ was considered to be significant. Analyses were based on all follow-up information received at the NSABP Biostatistical Center as of June 30, 2005.

RESULTS

A total of 8,878 node-negative patients enrolled onto these five NSABP trials were eligible with follow-up. Of these, 313 patients met our predetermined criteria and are included in this analysis (Table 1). Patients in this study were treated between 1981 and 1998. Table 2 lists patient and tumor characteristics. Two hundred sixteen and 169 patients were at risk for any first recurrence at 5 and 10 years, respectively. Of the 313 patients considered for this analysis, 28 (8.9%) presented with isolated LRF as the first event. The overall 10-year cumulative incidences of isolated LRF, LRF \pm DF, and DF alone as first event were 7.1%, 10.0%, and 23.6%, respectively. The median times to develop isolated LRF and DF were 2.3 and 1.8 years, respectively. The majority of LRFs (62.2%) occurred within the first 4 years, and 18.9%, 10.8%, and 8.1% of LRFs occurred at 4 to 8, 8 to 12, and more than 12 years of follow-up, respectively.

The rates of isolated LRF, LRF \pm DF, and DF according to different patient and tumor characteristics are listed in Table 3. Results from univariate analyses indicated that age, menopausal status, estrogen receptor status, pathologic tumor size, number of LNs removed, and the type of treatment were not significant predictors for either LRF or DF. Patients who had one to five LNs removed had a 10-year isolated LRF cumulative incidence estimate of 16.7% compared with 7.3% for patients who had ≥ 10 LNs removed. However, this difference was not significant (*P* = .21). Also, patients who were randomly assigned to no adjuvant systemic treatment arms had a 12.6% isolated LRF rate compared with a 5.3% rate in patients who received systemic treatment with chemotherapy, tamoxifen, or both (*P* = .2). Using multivariate analysis, we were unable to identify any independent prognostic factors for LRF \pm DF. The majority of recurrences (24 of 28 patients) occurred in the chest wall and around the mastectomy scar (85.7%); two patients experienced recurrences in the supraclavicular fossa, one patient experienced a recurrence in the axilla, and one patient experienced a recurrence in the parasternal area.

Table 1. Summary of NSABP Treatment Protocols and No. of Patients Included in This Report

Trial	No. of Patients With Tumors ≥ 5 cm	Total No. of Patients	Period of Accrual	Patient ER Status (fmol/mg)	Trial Treatment Groups
B-13	51	731	1981-1988	< 10	No further therapy v M \rightarrow F + LV
B-14	103	2,817	1982-1994	> 10	Placebo v Tam (5 or 10 years)
B-19	54	1,074	1988-1990	< 10	M \rightarrow F + LV v CMF
B-20	45	2,301	1988-1993	> 10	M \rightarrow F + Tam v CMF + Tam v Tam
B-23	60	1,955	1991-1998	< 10	Six CMF \pm Tam v 4 AC \pm Tam
Total	313	8,878*	1981-1998	—	—

Abbreviations: NSABP, National Surgical Adjuvant Breast and Bowel Project; ER, estrogen receptor; M, methotrexate; F, fluorouracil; LV, leucovorin; AC, doxorubicin 60 mg/m² intravenously and cyclophosphamide 600 mg/m² intravenously; CMF, cyclophosphamide 100 mg/m² orally, methotrexate 40 mg/m² intravenously, and fluorouracil 600 mg/m² intravenously; Tam, tamoxifen.

*No. of patients eligible with follow-up and analyzed.

Table 2. Patient, Tumor, and Treatment Characteristics of the Study Population

Characteristics	No. of Patients	%
Age, years		
< 50	162	51.8
50-59	89	28.4
\geq 60	62	19.8
Median	49	
Range	29-74	
Menopausal status		
Premenopausal	164	52.4
Postmenopausal	145	46.3
Unknown	4	1.3
Tumor ER status		
Positive	147	47.0
Negative	166	53.0
Pathologic tumor size, cm		
5	144	46.0
5.1-7.0	130	41.5
7.1-10.0	34	10.9
> 10.0	5	1.6
Median	5.5	
Range	5.0-15.5	
No. of LN removed		
1-5	12	3.8
6-9	39	12.5
\geq 10	262	83.7
Median	16	
Range	1.0-39.0	
Treatment regimen		
No systemic treatment	80	25.5
Chemotherapy only	107	34.2
Tam only	66	21.1
Chemotherapy + Tam	60	19.2

Abbreviations: ER, estrogen receptor; LN, lymph nodes; Tam, tamoxifen.

DISCUSSION

Because of the rarity of reports and information about the efficacy of radiotherapy for the treatment of patients with T3N0 breast cancer, Recht et al² avoided giving recommendations for this group of patients in the American Society of Clinical Oncology guidelines for PMRT. Nevertheless, many clinicians choose to treat T3N0 patients with radiotherapy. In a recent survey³ of radiation oncologists, 1,137 respondents from North America and Europe (88.3% and 84.8%, respectively) would recommend PMRT for patients with T3N0 breast cancer despite the lack of clinical evidence regarding its benefit; furthermore, of these oncologists, 47.7% and 20.2%, respectively, would recommend supraclavicular radiation.

The study we carried out was designed to contribute to the base of clinical decision-making knowledge about the use of radiation therapy in node-negative patients with large tumors. We found 169 patients with tumors (T3N0) larger than 5 cm among 8,878 patients enrolled onto five node-negative trials (Table 1), representing less than 2% of node-negative patients. We decided to include patients with 5-cm tumors (large T2N0) to increase the number of patients to 313, which represented 3.5% of LN-negative patients enrolled onto these trials (Table 1). The isolated LRF and LRF \pm DF rates were similar whether the tumor was 5 cm or larger than 5 cm (Table 3). Although the total

number of patients in this series is relatively small, to our knowledge, because of the infrequency of this clinical scenario, this is the largest series to date on patients with tumors of 5 cm or larger with negative LNs treated by mastectomy without PMRT.

In Finland, Helinto et al¹ identified 38 patients with T3N0 tumors treated by MRM from among 4,190 new breast cancer patients (representing < 1% of the patients). Mignano et al¹³ reported the LRF rate in his series of 101 patients with T3N0 breast cancer treated from 1974 to 1994 with MRM who received no radiation and of whom, only 9% received adjuvant systemic therapy. Isolated LRF as first event was reported in 12 patients (12%); six of these patients experienced chest wall recurrence, and six experienced regional LN failure. The authors concluded that the risk of LRF in T3N0 is low and does not warrant the routine use of PMRT. In a combined analysis from the Massachusetts General Hospital, M.D. Anderson Cancer Center, and Yale University, Floyd et al,¹⁴ with a median follow-up time of 85 months, found that the 5-year actuarial cumulative rates of LRF in 70 node-negative patients with tumors \geq 5 cm was 7.6%; four of the five failures occurred in the chest wall, and one occurred in the axilla. Tumor size and menopausal status were not found to be significant prognostic factors of LRF, but in a multivariate analysis, lymphatic vessel invasion (LVI) was found to be an independent prognostic factor for LRF, disease-free survival, and overall survival (OS). Although the low LRF rate in that series is in agreement with what we found, the study by Floyd et al¹⁴ was a retrospective study, and the omission of PMRT for the patients in their series was determined by patient and physician biases that may have affected the selection of the cohort analyzed.

All the patients in our series received no PMRT, as required by the different protocols. One might argue that, between 1981 and 1998, physicians who had patients at particularly high risk for LRF when PMRT was not an option might have been disinclined to enroll them in these NSABP studies. This is an unlikely scenario because in the same period (1984 to 1994) 5,758 patients with MRM and positive LNs (ie, considered higher risk patients) were accrued to node-positive NSABP trials that also did not allow PMRT.¹⁵ Almost 50% of the latter group had \geq four positive LNs, and 11% had larger than 5 cm tumors with positive LNs. Therefore, it is unlikely that only node-negative, low-risk patients for LRF were selected to enroll onto the node-negative NSABP trials included in this analysis.

Several randomized studies have suggested that PMRT reduces the risk of LRF by two thirds and potentially increases OS.¹⁶⁻¹⁸ These trials have targeted mainly patients with positive LNs. However, the randomized clinical trial reported by Overgaard et al,^{16,17} in which patients were randomly assigned to cyclophosphamide, methotrexate, and fluorouracil chemotherapy versus cyclophosphamide, methotrexate, and fluorouracil plus PMRT, included 135 premenopausal¹⁶ and 132 postmenopausal¹⁷ patients with T3N0 tumors. The results in that group showed that PMRT decreased the LRF rate from 17% to 3% in premenopausal patients and from 23% to 6% in postmenopausal patients.^{16,17} In addition, PMRT significantly increased the OS rate from 70% to 82% in premenopausal patients¹⁶ but did not significantly increase OS in postmenopausal patients. However, in those studies, axillary LN dissection was considered suboptimal because an average of only seven LNs were removed^{16,17} compared with 15 to 17 LNs removed in other studies.^{15,19-21} Thus, it is possible that some of the T3N0 patients in the Overgaard et al^{16,17} series may have been found to be node positive if additional axillary LNs had been removed

Table 3. Ten-Year Cumulative Incidence Rates and Raw Estimates at Year 10 of ILRF, LRF \pm DF, and DF

Factor	No. of Patients	ILRF			LRF \pm DF			DF		
		Incidence		No. of Events	Incidence		No. of Events	Incidence		No. of Events
		%	SE		%	SE		%	SE	
Age, years*										
< 49	162	8.1	2.2	13	10.0	2.4	16	25.5	3.5	41
50-59	89	5.7	2.5	5	11.4	3.4	10	22.8	4.5	20
≥ 60	62	6.5	3.2	4	8.1	3.5	5	19.5	5.1	12
P		.71			.28			.98		
Menopausal status*										
Pre-/perimenopausal	164	7.4	2.1	12	9.3	2.3	15	22.8	3.3	37
Postmenopausal	145	6.2	2.0	9	10.4	2.6	15	23.7	3.6	34
P		.29			.58			.82		
ER status*										
Positive	147	7.7	2.2	11	9.7	2.5	14	23.2	3.5	34
Negative	166	6.6	1.9	11	10.3	2.4	17	23.7	3.3	39
P		.57			.60			.26		
Pathologic tumor size, cm*										
5.0	144	7.0	2.1	10	9.8	2.5	14	26.7	3.7	38
5.1-7.0	130	8.6	2.5	11	10.1	2.7	13	20.2	3.6	26
> 7.0	39	2.6	2.6	1	10.3	4.9	4	23.1	6.9	9
P		.34			.90			.42		
No. of axillary LN dissected*										
1-5	12	16.7	11.4	2	16.7	11.4	2	41.7	15.1	5
6-9	39	2.6	2.6	1	5.1	3.6	2	18.3	6.4	7
≥ 10	262	7.3	1.6	19	10.4	1.9	27	23.5	2.6	61
P		.21			.28			.28		
Treatment regimen										
No systemic treatment	80	12.6	3.7	10	16.3	4.2	13	27.7	5.1	22
Chemotherapy only	107	5.6	2.3	6	7.5	2.6	8	21.5	4.0	23
Tam only	66	4.6	2.6	3	7.7	3.3	5	25.9	5.5	17
Chemotherapy + Tam	60	5.3	3.0	3	8.6	3.8	5	18.6	5.1	11
P		.21			.32			.60		

NOTE. P values compare overall cumulative incidence during the entire follow-up period. The events were counted only up to year 10 for a fair comparison with the corresponding 10-year cumulative incidence estimates.

Abbreviations: ILRF, isolated local and/or regional failure as first events with no distant failure within the first 4 months of diagnosis; LRF \pm DF, local and/or regional failure with or without distant failure; DF, distant failure; ER, estrogen receptor; LN, lymph nodes; Tam, tamoxifen.

*Unknowns, if any, were omitted from each analysis.

at surgery. In our series, patients who had had one to five axillary LNs removed had a 16.7% isolated LRF rate, in contrast to 7.3% in patients who had had 10 or more LNs removed. Although the difference was not statistically significant because of the small number of patients who had one to five LNs removed (12 patients), the trend suggests that adequate LN dissection may be able to identify T3N0 patients with a lower risk for LRF.

As discussed earlier, the rate of LRF in T3N0 patients randomly assigned to the nonradiation arm in the Overgaard et al^{16,17} trials was higher (17% to 23%) than the rate observed in our series (7.1%) or in other similar series (Floyd et al,¹⁴ 7.6%; and Mignano et al,¹³ 12%). This discrepancy between the randomized trials^{16,17,22} and other series^{15,19-21} in the LRF rate in patients who did not receive PMRT was also found in patients with one to three and \geq four positive axillary LNs.¹⁵ Thus, the applicability of the findings from the randomized trials to current practice about the routine use of PMRT in patients with tumors larger than 5 cm and negative axillary nodes (by an adequate axillary dissection) who are treated with systemic therapy has been questioned.

As mentioned earlier, the univariate and multivariate analyses in our study did not show any significant prognostic factors for

LRF or LRF \pm DF. Although an adequate number of removed axillary LNs and the use of systemic treatment were found to be associated with better locoregional control, these associations were not statistically significant. This could be because the number of patients and, more importantly, the number of events were relatively small in our study. It is also possible that some potentially prognostic factors were not studied because of the lack of information in our database. These factors include LVI,^{14,21,23,24} tumor grade,^{21,23} human epidermal growth factor receptor 2 expression,^{25,26} close margins,²⁴ p53 expression,²⁷ and presence of extensive intraductal component.²⁸ Floyd et al¹⁴ found that LVI was an independent prognostic factor for LRF. Nevertheless, tumor size, as long as the LNs were negative, was not found to be an independent prognostic factor for LRF in either of our series or the study by Floyd et al.¹⁴ In a recent retrospective analysis of 877 patients with T1-3 node-negative tumors treated at Massachusetts General Hospital with mastectomy without adjuvant PMRT, Jaggi et al²⁹ suggested that patients with multiple risk factors, including close margins, T2 or larger tumors, premenopausal status, and LVI, are at higher risk for LRF. However, only 25 patients (0.55%) had tumors larger than 5 cm.

Large breast tumors that present without the involvement of regional LNs may be a distinct clinical and biologic entity. It is possible that the inability of these tumors to spread to regional nodes despite their ability to grow to large size may indicate a more indolent biologic nature. This clinical presentation is clearly in contrast to the one in which small tumors have already spread to regional LNs and/or distant sites at the time of clinical presentation. The study of biologic differences and gene fingerprinting³⁰ between these extremes of tumor presentation may help in the understanding of the nature of the invasive and metastatic potential of breast tumors and in the identification of patients who would benefit from PMRT.

The most common PMRT approach used by radiation oncologists and in the cooperative group studies includes the area of the chest wall and regional LNs.^{16,17,22} The risk of radiation pneumonitis increases with the number of radiation fields used as well as with the use of chemotherapy.^{31,32} Furthermore, the risk for arm lymphedema increases when radiotherapy includes the regional lymph node.^{33,34} All these facts raise the question of the need for regional radiation should radiation be used in this group of patients. Because the chest wall was by far the most common site of failure in this study (86%) as well as in the study by Floyd et al,¹⁴ it may be reasonable to consider treating the chest wall only, without radiating the regional LNs, in subsets of patients thought to be at higher risk, thereby minimizing the adverse effects of PMRT.

In our series, the 10-year cumulative incidence for LRF \pm DF in patients who did not receive systemic treatment was 16.3% (13 of 80 patients) compared with 7.5% (eight of 107 patients), 7.7% (five of 66 patients), and 8.6% (five of 60 patients) in patients who received

systemic adjuvant chemotherapy, adjuvant tamoxifen, or both, respectively (Table 3). However, the number of events was small, and these differences were not statistically significant. Most patients with ≥ 5 -cm N0 tumors would now receive systemic treatment.⁸ Therefore, with systemic therapy, the overall LRF \pm DF rate without PMRT in our study would be in the range of 6% to 7.7%, and the isolated LRF as first event rate would be in the range of 4.6% to 5.6% (Table 3). This rate is low enough that the benefit from routine PMRT might not outweigh its potential adverse effects. Furthermore, it is possible that a majority of this population might be treated with neoadjuvant chemotherapy because there would be a better chance for them under that circumstance to undergo conservative surgery.³⁵ To identify these low-risk patients, it might be reasonable for them to undergo up-front sentinel-node surgery; if the node is negative, the patient might not require PMRT or regional radiation if conservative approaches are used. A recent meta-analysis³⁶ suggested that an absolute reduction in LRF rate of approximately 20% is needed to improve survival by approximately 5%. Thus, it is unlikely that PMRT in this population will yield a significant improvement in survival.

In conclusion, the locoregional recurrence rate in this small population of node-negative patients with large tumors is lower than what radiation oncologists have perceived.³ We found no statistically significant risk factors for locoregional recurrence in this study. These data should be interpreted with caution because of the small number of patients and events and because LVI, grade, and margins were not among the factors that were included in our database, but the data suggest that pathologic tumor size alone, in the absence of positive LNs, should not be a basis for PMRT.

REFERENCES

- Helinto M, Blomqvist C, Heikkilä P, et al: Post-mastectomy radiotherapy in pT3N0M0 breast cancer: Is it needed? *Radiother Oncol* 52:213-217, 1999
- Recht A, Edge SB, Solin LJ, et al: Postmastectomy radiotherapy: Clinical practice guidelines of the American Society of Clinical Oncology. *J Clin Oncol* 19:1539-1569, 2001
- Ceille E, Jagsi R, Goldberg S, et al: Radiotherapy for invasive breast cancer in North America and Europe: Results of a survey. *Int J Radiat Oncol Biol Phys* 61:365-373, 2005
- Fisher B, Redmond C, Dimitrov NV, et al: A randomized clinical trial evaluating sequential methotrexate and fluorouracil in the treatment of patients with node-negative breast cancer who have estrogen-receptor-negative tumors. *N Engl J Med* 320:473-478, 1989
- Fisher B, Dignam J, Bryant J, et al: Five versus more than five years of tamoxifen therapy for breast cancer patients with negative lymph nodes and estrogen receptor-positive tumors. *J Natl Cancer Inst* 88:1529-1542, 1996
- Fisher B, Costantino J, Redmond C, et al: A randomized clinical trial evaluating tamoxifen in the treatment of patients with node-negative breast cancer who have estrogen-receptor-positive tumors. *N Engl J Med* 320:479-484, 1989
- Fisher B, Dignam J, Mamounas E, et al: Sequential methotrexate and fluorouracil for the treatment of node-negative breast cancer patients with estrogen receptor-negative tumors: Eight-year results from National Surgical Adjuvant Breast and Bowel Project (NSABP) B-13 and first report of findings from NSABP B-19 comparing methotrexate and fluorouracil with conventional cyclophosphamide, methotrexate, and fluorouracil. *J Clin Oncol* 14:1982-1992, 1996
- Fisher B, Dignam J, Wolmark N, et al: Tamoxifen and chemotherapy for lymph node-negative, estrogen receptor-positive breast cancer. *J Natl Cancer Inst* 89:1673-1682, 1997
- Fisher B, Anderson S, Tan-Chiu E, et al: Tamoxifen and chemotherapy for axillary node negative, estrogen receptor-negative breast cancer: Findings from the National Surgical Breast and Bowel Project B-23. *J Clin Oncol* 19:931-942, 2001
- Kalbfleisch J, Prentice RL: *The Statistical Analysis of Failure Time Data*. New York, NY, John Wiley and Sons, 1980
- Gray R: A class of K-sample tests for comparing the cumulative incidence of a competing risk. *Ann Stat* 16:1141-1154, 1988
- Cox D: Regression models and life-tables. *J R Stat Soc B* 34:187-220, 1972
- Mignano J, Gage I, Piantatosi S, et al: Local recurrence after mastectomy in patients with T3N0 breast carcinoma treated without postoperative irradiation. *Breast Cancer Res Treat* 41:255, 1996 (abstr 230)
- Floyd S, Buchholz TA, Haffty B, et al: Low local recurrence rate without post-mastectomy radiation in node negative breast cancer patients with tumors 5 cm and larger. *Int J Radiat Oncol Biol Phys* (in press)
- Taghian A, Jeong J-H, Mamounas E, et al: Patterns of locoregional failure in patients with operable breast cancer treated by mastectomy and adjuvant chemotherapy with or without tamoxifen and without radiotherapy: Results from five National Surgical Adjuvant Breast and Bowel Project randomized clinical trials. *J Clin Oncol* 22:4247-4254, 2004
- Overgaard M, Hansen PS, Overgaard J, et al: Postoperative radiotherapy in high-risk premenopausal women with breast cancer who receive adjuvant chemotherapy: Danish Breast Cancer Cooperative Group 82b Trial. *N Engl J Med* 337:949-955, 1997
- Overgaard M, Jensen MB, Overgaard J, et al: Postoperative radiotherapy in high-risk postmenopausal breast cancer patients given adjuvant tamoxifen: Danish Breast Cancer Cooperative Group DBCG 82c randomized trial. *Lancet* 353:1641-1648, 1999
- Ragaz J, Jackson SM, Le N, et al: Adjuvant radiotherapy and chemotherapy in node-positive premenopausal women with breast cancer. *N Engl J Med* 337:956-962, 1997
- Recht A, Gray R, Davidson N, et al: Locoregional failure 10 years after mastectomy and adjuvant chemotherapy with or without tamoxifen without irradiation: Experience of the Eastern Cooperative Oncology Group. *J Clin Oncol* 17:1689-1700, 1999
- Katz A, Strom E, Buchholz T, et al: Locoregional recurrence patterns after mastectomy and doxorubicin-based chemotherapy: Implications for postoperative irradiation. *J Clin Oncol* 18:2817-2827, 2000
- Wallgren A, Bonetti M, Gelber RD, et al: Risk factors for locoregional recurrence among breast cancer patients: Results from International Breast Cancer Study Group trials I through VII. *J Clin Oncol* 21:1205-1213, 2003
- Ragaz J, Olivetto IA, Spinelli JJ, et al: Locoregional radiation therapy in patients with high-risk

breast cancer receiving adjuvant chemotherapy: 20-year results of the British Columbia randomized trial. *J Natl Cancer Inst* 97:116-126, 2005

23. Pisansky T, Ingle JN, Schaid DJ, et al: Patterns of tumor relapse following mastectomy and adjuvant systemic therapy in patients with axillary lymph node-positive breast cancer. *Cancer* 72:1247-1260, 1993

24. Katz A, Strom EA, Buchholz TA, et al: The influence of pathologic tumor characteristics on locoregional recurrence rates following mastectomy. *Int J Radiat Oncol Biol Phys* 50:735-742, 2001

25. Paik S, Bryant J, Park C, et al: ErbB-2 and response to doxorubicin in patients with axillary lymph node-positive, hormone receptor-negative breast cancer. *J Natl Cancer Inst* 90:1361-1370, 1998

26. Paik S, Bryant J, Tan-Chiu E, et al: HER2 and choice of adjuvant chemotherapy for invasive breast cancer: National Surgical Adjuvant Breast and Bowel Project Protocol B-15. *J Natl Cancer Inst* 92:1991-1998, 2000

27. Zellars R, Clark, GM, Allred, DC, et al: Prognostic value of p53 for local failure in mastectomy treated breast cancer patients. *Proc Am Soc Clin Oncol* 17:104a, 1998 (abstr 401)

28. Voogd A, Nielsen M, Peterse JL, et al: Differences in risk factors for local and distant recurrence after breast-conserving therapy or mastectomy for stage I and II breast cancer: Pooled results of two large European randomized trials. *J Clin Oncol* 19:1688-1697, 2001

29. Jagsi R, Raad RA, Goldberg S, et al: Locoregional recurrence rates and prognostic factors for failure in node-negative patients treated with mastectomy: Implications for postmastectomy radiation. *Int J Radiat Oncol Biol Phys* 62:1035-1039, 2005

30. Paik S, Shak S, Tang G, et al: A multigene assay to predict recurrence of tamoxifen-treated, node-negative breast cancer. *N Engl J Med* 351:2817-2826, 2004

31. Lingos T, Recht A, Vicini F, et al: Radiation pneumonitis in breast cancer patients treated with conservative surgery and radiation therapy. *Int J Radiat Oncol Biol Phys* 21:355-360, 1991

32. Taghian A, Assaad S, Niemierko A, et al: Risk of pneumonitis in breast cancer patients treated with radiation therapy and combination chemotherapy with paclitaxel. *J Natl Cancer Inst* 93:1806-1811, 2001

33. Coen JJ, Taghian AG, Kachnic LA, et al: Risk of lymphedema after regional nodal irradiation with breast conservation therapy. *Int J Radiat Oncol Biol Phys* 55:1209-1215, 2003

34. Højris I, Anderson J, Overgaard M, et al: Late treatment related morbidity in breast cancer patients randomized to postmastectomy radiotherapy and systemic treatment versus systemic treatment alone. *Eur J Cancer* 35:S206, 1999 (suppl 4, abstr 803)

35. Fisher B, Bryant J, Wolmark N, et al: Effect of preoperative chemotherapy on the outcome of women with operable breast cancer. *J Clin Oncol* 16:2672-2685, 1998

36. Early Breast Cancer Trialists' Collaborative Group: Effects of radiotherapy and of differences in the extent of surgery for early breast cancer on local recurrence and 15-year survival: An overview of the randomised trials. *Lancet* 366:2087-2106, 2005

Acknowledgment

We thank Barbara C. Good, PhD, and Wendy L. Rea for editorial assistance.

Authors' Disclosures of Potential Conflicts of Interest

Although all authors completed the disclosure declaration, the following author or immediate family members indicated a financial interest. No conflict exists for drugs or devices used in a study if they are not being evaluated as part of the investigation. For a detailed description of the disclosure categories, or for more information about ASCO's conflict of interest policy, please refer to the Author Disclosure Declaration and the Disclosures of Potential Conflicts of Interest section in Information for Contributors.

Authors	Employment	Leadership	Consultant	Stock	Honoraria	Research Funds	Testimony	Other
Eleftherios P. Mamounas			AstraZeneca (A)		AstraZeneca (A)			
Dollar Amount Codes (A) < \$10,000 (B) \$10,000-99,999 (C) ≥ \$100,000 (N/R) Not Required								

Author Contributions

Conception and design: Alphonse G. Taghian, Eleftherios P. Mamounas, Melvin Deutsch, Joseph P. Costantino, Norman Wolmark

Administrative support: Joseph P. Costantino, Norman Wolmark

Collection and assembly of data: Joseph P. Costantino

Data analysis and interpretation: Alphonse G. Taghian, Jong-Hyeon Jeong, Eleftherios P. Mamounas, David S. Parda, Joseph P. Costantino

Manuscript writing: Alphonse G. Taghian, Eleftherios P. Mamounas, David S. Parda, Melvin Deutsch, Joseph P. Costantino

Final approval of manuscript: Alphonse G. Taghian, Jong-Hyeon Jeong, Eleftherios P. Mamounas, David S. Parda, Melvin Deutsch, Joseph P. Costantino, Norman Wolmark

JOURNAL OF CLINICAL ONCOLOGY

Official Journal of the American Society of Clinical Oncology

Vol 24, No 24

C O N T E N T S

August 20, 2006

Editorials

Thoracic Radiation Therapy in Limited Stage Small-Cell Lung Cancer: Timing Is Everything... Isn't It?

Michael C. Perry (see article on page 3823) 3815

Neoadjuvant Chemotherapy for Disseminated Colorectal Cancer: Changing the Paradigm

Lawrence Leichman (see article on page 3939) 3817

■ Are Nomograms Better Than Currently Available Stage Groupings in Bladder Cancer?

Cora N. Sternberg (see article on page 3967) 3819

Some Thoughts on the Reporting of Adverse Events in Phase II Cancer Clinical Trials

Stewart J. Anderson (see article on page 3933) 3821

Original Reports

LUNG CANCER

① Early Compared With Late Radiotherapy in Combined Modality Treatment for Limited Disease Small-Cell Lung Cancer: A London Lung Cancer Group Multicenter Randomized Clinical Trial and Meta-Analysis

Stephen G. Spiro, Lindsay E. James, Robin M. Rudd, Colin W. Trask, Jeffrey S. Tobias, Michael Snee, David Gilligan, Philip A. Murray, Mary Carmen Ruiz de Elvira, Katy M. O'Donnell, Nicole H. Gower, Peter G. Harper, and Allan K. Hackshaw (see editorial on page 3815) 3823

Symptom Improvement in Lung Cancer Patients Treated With Erlotinib: Quality of Life Analysis of the National Cancer Institute of Canada Clinical Trials Group Study BR.21

Andrea Bezjak, Dongsheng Tu, Lesley Seymour, Gary Clark, Aleksandra Trajkovic, Mauro Zukin, Joseph Ayoub, Sergio Lago, Ronaldo de Albuquerque Ribeiro, Alexandra Gerogianni, Arnold Cyjon, Jonathan Noble, Francis Laberge, Raymond Tsz-Tong Chan, David Fenton, Joachim von Pawel, Martin Reck, and Frances A. Shepherd 3831

(continued on following page)

Journal of Clinical Oncology (ISSN 0732-183X) is published 36 times a year, three times monthly, by American Society of Clinical Oncology, 1900 Duke St, Suite 200, Alexandria, VA 22314. Periodicals postage is paid at Alexandria, VA, and at additional mailing offices. Publication Mail Agreement Number 863289.

Editorial correspondence should be addressed to Daniel G. Haller, MD, *Journal of Clinical Oncology*, 330 John Carlyle St, Suite 300, Alexandria, VA 22314. Telephone: (703) 797-1900; Fax: (703) 684-8720. E-mail: jco@asco.org. Internet: www.jco.org.

POSTMASTER: ASCO members send change of address to American Society of Clinical Oncology, 1900 Duke St, Suite 200, Alexandria, VA 22314. Nonmembers send change of address to *Journal of Clinical Oncology* Customer Service, 330 John Carlyle St, Suite 300, Alexandria, VA 22314.

2006 annual subscription rates, effective September 1, 2005: United States and possessions: individual, \$435; single issue, \$35. International: individual, \$605; single issue, \$45. Institutions: Tier 1: \$615 US, \$870 Int'l; Tier 2: \$715 US, \$970 Int'l; Tier 3: \$1,035 US, \$1,290 Int'l; Tier 4: \$1,140 US, \$1,395 Int'l; Tier 5: contact JCO for a quote. See <http://www.jco.org/subscriptions/tieredpricing.shtml> for descriptions of each tier. Student and resident: United States and possessions: \$215; all other countries, \$300. To receive student/resident rate, orders must be accompanied by name of affiliated institution, date of term, and the signature of program/residency coordinator on institution letterhead. Orders will be billed at individual rate until proof of status is received. Current prices are in effect for back volumes and back issues. Back issues sold in conjunction with a subscription rate are on a prorated basis. Subscriptions are accepted on a 12-month basis. Prices are subject to change without notice. Single issues, both current and back, exist in limited quantities and are offered for sale subject to availability. JCO Legacy Archive (electronic back issues from January 1983 through December 1998) is also available; please inquire.

PEDIATRIC ONCOLOGY

Local Control in Pelvic Ewing Sarcoma: Analysis From INT-0091—A Report From the Children's Oncology Group

Torunn I. Yock, Mark Krailo, Christopher J. Fryer, Sarah S. Donaldson, James S. Miser, Zhengjia Chen, Mark Bernstein, Fran Laurie, Mark. C. Gebhardt, Holcombe E. Grier, and Nancy J. Tarbell 3838

Analysis of Prognostic Factors in Patients With Nonmetastatic Rhabdomyosarcoma Treated on Intergroup Rhabdomyosarcoma Studies III and IV: The Children's Oncology Group

Jane L. Meza, James Anderson, Alberto S. Pappo, and William H. Meyer 3844

Suicidal Ideation and Attempts in Adult Survivors of Childhood Cancer

Christopher J. Recklitis, Rebecca A. Lockwood, Monica A. Rothwell, and Lisa R. Diller 3852

Comparison of Long-Term Neurocognitive Outcomes in Young Children With Acute Lymphoblastic Leukemia Treated With Cranial Radiation or High-Dose or Very High-Dose Intravenous Methotrexate

Brenda J. Spiegler, Kimberly Kennedy, Ronnen Maze, Mark L. Greenberg, Sheila Weitzman, Johann K. Hitzler, and Paul C. Nathan 3858

NEUROONCOLOGY

High-Dose Chemotherapy With Autologous Stem-Cell Transplantation and Hyperfractionated Radiotherapy As First-Line Treatment of Primary CNS Lymphoma

Gerald Illerhaus, Reinhard Marks, Gabriele Ihorst, Roland Guttenberger, Christoph Ostertag, Günther Derigs, Norbert Frickhofen, Friedrich Feuerhake, Benedikt Volk, and Jürgen Finke 3865

Phase III Trial of Carmustine and Cisplatin Compared With Carmustine Alone and Standard Radiation Therapy or Accelerated Radiation Therapy in Patients With Glioblastoma Multiforme: North Central Cancer Treatment Group 93-72-52 and Southwest Oncology Group 9503 Trials

Jan C. Buckner, Karla V. Ballman, John C. Michalak, Gary V. Burton, Terrence L. Cascino, Paula J. Schomberg, Roland B. Hawkins, Bernd W. Scheithauer, Howard M. Sandler, Randolph S. Marks, and Judith R. O'Fallon 3871

HEMATOLOGIC MALIGNANCIES

Multicenter Phase II Trial of Immunotherapy With the Humanized Anti-CD22 Antibody, Epratuzumab, in Combination With Rituximab, in Refractory or Recurrent Non-Hodgkin's Lymphoma

Sandra J. Strauss, Frank Morschhauser, Juergen Rech, Roland Repp, Philippe Solal-Celigny, Pier L. Zinzani, Andreas Engert, Bernard Coiffier, Dieter F. Hoelzer, William A. Wegener, Nick K.W. Teoh, David M. Goldenberg, and T. Andrew Lister 3880

Disclosure of Candidate Genes in Acute Myeloid Leukemia With Complex Karyotypes Using Microarray-Based Molecular Characterization

Frank G. Rücker, Lars Bullinger, Carsten Schwaenen, Daniel B. Lipka, Swen Wessendorf, Stefan Fröhling, Martin Bentz, Simone Miller, Claudia Scholl, Richard F. Schlenk, Bernhard Radlwimmer, Hans A. Kestler, Jonathan R. Pollack, Peter Lichter, Konstanze Döhner, and Hartmut Döhner 3887

Further Analysis of Trials With Azacitidine in Patients With Myelodysplastic Syndrome: Studies 8421, 8921, and 9221 by the Cancer and Leukemia Group B

Lewis R. Silverman, David R. McKenzie, Bercedis L. Peterson, James F. Holland, Jay T. Backstrom, C.L. Beach, and Richard A. Larson 3895

(continued on following page)

① Adverse Prognostic Significance of *KIT* Mutations in Adult Acute Myeloid Leukemia With inv(16) and t(8;21): A Cancer and Leukemia Group B Study

Peter Paschka, Guido Marcucci, Amy S. Ruppert, Krzysztof Mrózek, Hankui Chen, Rick A. Kittles, Tamara Vukosavljevic, Danilo Perrotti, James W. Vardiman, Andrew J. Carroll, Jonathan E. Kolitz, Richard A. Larson, and Clara D. Bloomfield 3904

BREAST CANCER

① Lack of Benefit of Maintenance Paclitaxel in First-Line Chemotherapy in Metastatic Breast Cancer

Alessandra Gennari, Dino Amadori, Mario De Lena, Oriana Nanni, Paolo Bruzzi, Vito Lorusso, Luigi Manzione, and Pier Franco Conte 3912

Randomized Trial of Single Compared With Tandem High-Dose Chemotherapy Followed by Autologous Stem-Cell Transplantation in Patients With Chemotherapy-Sensitive Metastatic Breast Cancer

Nicolaus Kröger, Markus Frick, Oleg Gluz, Svjetlana Mohrmann, Bernd Metzner, Christian Jackisch, Yon Ko, Hans-Walter Lindemann, Carl Richard Meier, Hans Peter Lohrmann, Ute Ruffert, Matthias Hänel, Heinrich Bodenstern, Andreas Neubauer, Gerhard Ehninger, Hans-Heinrich Wolf, Kathrin Kolbe, Karin Burock, Axel R. Zander, and Ulrike Nitz 3919

Low Locoregional Recurrence Rate Among Node-Negative Breast Cancer Patients With Tumors 5 cm or Larger Treated by Mastectomy, With or Without Adjuvant Systemic Therapy and Without Radiotherapy: Results From Five National Surgical Adjuvant Breast and Bowel Project Randomized Clinical Trials

Alphonse G. Taghian, Jong-Hyeon Jeong, Eleftherios P. Mamounas, David S. Parda, Melvin Deutsch, Joseph P. Costantino, and Norman Wolmark 3927

CLINICAL TRIALS

Adverse Event Reporting in Publications Compared With Sponsor Database for Cancer Clinical Trials

Orit Scharf and A. Dimitrios Colevas ([see editorial on page 3821](#)) 3933

GASTROINTESTINAL CANCER

Complete Response of Colorectal Liver Metastases After Chemotherapy: Does It Mean Cure?

Stéphane Benoist, Antoine Brouquet, Christophe Penna, Catherine Julié, Mostafa El Hajjam, Sophie Chagnon, Emmanuel Mitry, Philippe Rougier, and Bernard Nordlinger ([see editorial on page 3817](#)) 3939

① Randomized Phase III Trial of Gemcitabine Plus Cisplatin Compared With Gemcitabine Alone in Advanced Pancreatic Cancer

Volker Heinemann, Detlef Quetzsch, Frank Gieseler, Michael Gonnermann, Herbert Schönekeas, Andreas Rost, Horst Neuhaus, Caroline Haag, Michael Clemens, Bernard Heinrich, Ursula Vehling-Kaiser, Martin Fuchs, Doris Fleckenstein, Wolfgang Gesierich, Dirk Uthgenannt, Hermann Einsele, Axel Holstege, Axel Hinke, Andreas Schalhorn, and Ralf Wilkowski 3946

Phase II Trial of Preoperative Chemoradiation in Patients With Localized Gastric Adenocarcinoma (RTOG 9904): Quality of Combined Modality Therapy and Pathologic Response

Jaffer A. Ajani, Kathryn Winter, Gordon S. Okawara, John H. Donohue, Peter W.T. Pisters, Christopher H. Crane, John F. Greskovich, P. Rani Anne, Jeffrey D. Bradley, Christopher Willett, and Tyvin A. Rich 3953

(continued on following page)

BONE MARROW TRANSPLANTATION

① Results of Genoidentical Hemopoietic Stem Cell Transplantation With Reduced Intensity Conditioning for Acute Myelocytic Leukemia: Higher Doses of Stem Cells Infused Benefit Patients Receiving Transplants in Second Remission and Beyond—The Acute Leukemia Working Party of the European Cooperative Group for Blood and Marrow Transplantation

Norbert-Claude Gorin, Myriam Labopin, Jean-Michel Boiron, Niklas Theorin, Tim Littlewood, Shimon Slavin, Hildegard Greinix, Jean Yves Cahn, E. Paolo Alessandrino, Alessandro Rambaldi, Arnon Nagler, Emmanuelle Polge, and Vanderson Rocha 3959

GENITOURINARY CANCER

■ Postoperative Nomogram Predicting Risk of Recurrence After Radical Cystectomy for Bladder Cancer

International Bladder Cancer Nomogram Consortium ([see editorial on page 3819](#)) 3967

Defining Biochemical Recurrence of Prostate Cancer After Radical Prostatectomy: A Proposal for a Standardized Definition

Andrew J. Stephenson, Michael W. Kattan, James A. Eastham, Zohar A. Dotan, Fernando J. Bianco Jr, Hans Lilja, and Peter T. Scardino ([see article on page 3984](#)) 3973

Metabolic Syndrome in Men With Prostate Cancer Undergoing Long-Term Androgen-Deprivation Therapy

Milena Braga-Basaria, Adrian S. Dobs, Denis C. Muller, Michael A. Carducci, Majnu John, Josephine Egan, and Shehzad Basaria 3979

Absolute Prostate-Specific Antigen Value After Androgen Deprivation Is a Strong Independent Predictor of Survival in New Metastatic Prostate Cancer: Data From Southwest Oncology Group Trial 9346 (INT-0162)

Maha Hussain, Catherine M. Tangen, Celestia Higano, Paul F. Schelhammer, James Faulkner, E. David Crawford, George Wilding, Atif Akdas, Eric J. Small, Bryan Donnelly, Gary MacVicar, and Derek Raghavan ([see article on page 3973](#)) 3984

PREVENTION

① Influence of Hormone Replacement Therapy on Tamoxifen-Induced Vasomotor Symptoms

Ivana Sestak, Roseann Kealy, Robert Edwards, John Forbes, and Jack Cuzick 3991

SARCOMAS

① Impact of High-Dose Busulfan Plus Melfalan As Consolidation in Metastatic Ewing Tumors: A Study by the Société Française des Cancers de l'Enfant

Odile Oberlin, Annie Rey, Anne Sophie Desfachelles, Thierry Philip, Dominique Plantaz, Claudine Schmitt, Emmanuel Plouvier, Odile Lejars, Hervé Rubie, Philippe Terrier, and Jean Michon 3997

SUPPORTIVE CARE AND QUALITY OF LIFE

Prognosis of Critically Ill Patients With Cancer and Acute Renal Dysfunction

Márcio Soares, Jorge I.F. Salluh, Marília S. Carvalho, Michael Darmon, José R. Rocco, and Nelson Spector 4003

Review Article

Systematic Review on the Efficacy of Cytoreductive Surgery Combined With Perioperative Intraperitoneal Chemotherapy for Peritoneal Carcinomatosis From Colorectal Carcinoma

Tristan D. Yan, Deborah Black, Renaldo Savady, and Paul H. Sugarbaker 4011

(continued on following page)

Special Article

① Shaping Your Career to Maximize Personal Satisfaction in the Practice of Oncology

Tait Shanafelt, Harold Chung, Heather White, and Laurie Jean Lyckholm 4020

Diagnosis in Oncology

Isolated Splenic Sinus Histiocytosis (Rosai-Dorfman disease) in Association With Myelodysplastic Syndrome

Stacey K. Knox, Paul J. Kurtin, and David P. Steensma 4027

Isolated Blast Crisis in CNS in a Patient With Chronic Myelogenous Leukemia Maintaining Major Cytogenetic Response After Imatinib

Hyun Jung Kim, Chul Won Jung, Kihyun Kim, Jin Seok Ahn, Won Seog Kim, Kwan Park, Young Hye Ko, Won Ki Kang, and Keunchil Park 4028

Azygo-Tracheal Fistula in a Complete Implantable Central Venous System

Carlo Cosimo Quattrocchi, Matteo Sammarra, Massimiliano Carassiti, Bruno Vincenzi, Giuseppe Tonini, Rosario Francesco Grasso, and Bruno Beomonte Zobel 4029

Alveolar Rhabdomyosarcoma Mimicking a Lymphoma at Presentation

Steve C. Tsai, Lisa D. Reale, Neal Flomenberg, Roland Schwarting, and Robert E. Enck 4031

History of Oncology

Arnoldus Goudsmit, MD, PhD: Chemotherapist, Visionary, Founder of the American Society of Clinical Oncology, 1909-2005

Gretchen M. Krueger, Lori L. Alexander, Deborah A. Whippen, and Charles M. Balch 4033

Correspondence

Influence of Whole-Brain Radiotherapy on Remission of Brain Metastases

Carsten Nieder 4037

In Reply

Istvan Petak 4037

Quality of Life in a Randomized Control Trial? Comments Concerning MA 17

Pascale This, Anne de la Rochefordière, Anne Bredart, Bernard Asselain, Rollon Poinot, Sylvie Dolbeault, and Brigitte Sigal-Zafrani 4038

In Reply

Timothy J. Whelan, Paul E. Goss, James N. Ingle, Donsheng Tu, Lois Shepherd, and Joseph L. Pater 4038

On the Origin and Nature of Elevated Levels of Circulating Endothelial Cells After Treatment With a Vascular Disrupting Agent

Yuval Shaked, Francesco Bertolini, Urban Emmenegger, Christina R. Lee, and Robert S. Kerbel 4040

In Reply

Laurens V. Beerepoot, Els O. Witteveen, and Emile E. Voest 4040

(continued on following page)

Primary Cutaneous B-Cell Lymphoma

Lynn D. Wilson and Benjamin D. Smith 4041

In Reply

Pier Luigi Zinzani, Pietro Quaglino, Nicola Pimpinelli, Emilio Berti, Gianandrea Baliva, Serena Rupoli, Maurizio Martelli, Mauro Alaibac, Giovanni Borroni, Sergio Chimenti, Renato Alterini, Lapo Alinari, Maria Teresa Fierro, Nazario Cappello, Alessandro Pileri, Davide Soligo, Marco Paulli, Stefano Pileri, Marco Santucci, and Maria Grazia Bernengo 4041

Specification on the Definition of Adult-Type Soft Tissue Sarcoma

Andrea Ferrari and Michela Casanova 4042

In Reply

Sheri L. Spunt and Alberto S. Pappo 4043

Measuring the Impact of Chemotherapy on Fertility in Women With Breast Cancer

Kutluk Oktay, Ozgur Oktem, Andrea Reh, and Linda Vahdat 4044

Erratum 4047

Also in This Issue

[Announcements](#)

[Information for Contributors](#)

[Current Abstracts](#)

[Calendar of Oncology Events](#)



Online supplementary information available at www.jco.org



Article was published online ahead of print at www.jco.org

www.jco.org

www.asco.org