aineva.it

Monday 12.05.2025

Published 11 05 2025, 17:00



1	2	3	4	5
low	moderate	considerable	high	very high

Monday 12.05.2025

Published 11 05 2025, 17:00

Danger Level 2 - Moderate



2500m

Tendency: Decreasing avalanche danger
on Tuesday 13 05 2025

Wet snow

2500m

Snowpack stability: poor
Frequency: some
Avalanche size: medium

Wind slab

2900m

Snowpack stability: poor
Frequency: some
Avalanche size: medium

Sunshine and high temperatures will give rise as the day progresses to increasing moistening of the snowpack.

In some localities 5 cm of snow has fallen above approximately 2600 m. As a consequence of warming during the day and the solar radiation, the likelihood of moist and wet avalanches being released will increase gradually in particular on steep slopes above approximately 2500 m. The new snow can be released by a single winter sport participant. This applies in particular in gullies and bowls on steep slopes at high altitudes and in high Alpine regions.
Backcountry touring calls for experience in the assessment of avalanche danger and careful route selection.

Snowpack

Sunshine and high temperatures will give rise as the day progresses to increasing moistening of the snowpack in all aspects. These conditions will cause a gradual weakening of the snowpack. Below approximately 1900 m hardly any snow is lying.

Monday 12.05.2025

Published 11 05 2025, 17:00

Danger Level 1 - Low



Tendency: Constant avalanche danger
on Tuesday 13 05 2025

Wet snow

Snowpack stability: poor
Frequency: few
Avalanche size: small

Sunshine and high temperatures will give rise as the day progresses to increasing moistening of the snowpack.

As a consequence of warming during the day and the solar radiation, the likelihood of moist and wet avalanches being released will increase gradually in particular on steep slopes above approximately 2100 m.
Backcountry touring calls for caution and restraint.

Snowpack

Sunshine and high temperatures will give rise as the day progresses to increasing moistening of the snowpack in all aspects. These conditions will cause a gradual weakening of the snowpack. Below approximately 2000 m hardly any snow is lying.