# Wednesday 11.03.2026

Published 10 03 2026, 17:00




| 1 | 2 | 3 | 4 | 5 |
|---|---|---|---|---|
| low | moderate | considerable | high | very high |

## Danger Level 2 - Moderate

**Tendency: Constant avalanche danger**
on Thursday 12 03 2026



Persistent weak layer

**2000m**

Wet snow

**2200m**

### Weakly bonded old snow requires caution. Slight increase in danger of moist avalanches in the course of the day.

Winter sport participants can release avalanches in isolated cases. These can reach large size. The avalanche prone locations for dry avalanches are to be found in particular on very steep northwest, north and northeast facing slopes above approximately 2000 m and on very steep west and east facing slopes above approximately 2200 m. Caution is to be exercised in particular at transitions from a shallow to a deep snowpack, when entering gullies and bowls for example.

As the day progresses the likelihood of moist slab avalanches being released will increase on very steep west and east facing slopes below approximately 2200 m. Caution is to be exercised in particular in areas where the snow cover is rather shallow.

### Snowpack

**Danger patterns** dp.7: snow-poor zones in snow-rich surrounding | dp.10: springtime scenario

Distinct weak layers exist in the bottom section of the snowpack on west, north and east facing slopes, in particular above approximately 2000 m.

Towards its surface, the snowpack consists of faceted crystals, especially on shady slopes at elevated altitudes.

Outgoing longwave radiation during the night will be severely restricted. As a consequence of high relative humidity the snowpack can not consolidate, especially below approximately 2200 m.

In steep terrain there is a danger of falling on the hard snow surface. This applies at high altitude.

### Tendency

> Weakly bonded old snow requires caution. Gradual increase in danger of wet avalanches in the course of the day.

## Danger Level 1 - Low




1

**Tendency: Constant avalanche danger**
on Thursday 12 03 2026

### Low avalanche danger will prevail.

As a consequence of warming during the day only isolated wet avalanches are possible.

Only isolated dry avalanches are possible. They can be released, in particular by large loads, but they will be small in most cases. Individual avalanche prone locations are to be found on steep shady slopes at high altitude.

### Snowpack

**Danger patterns** dp.10: springtime scenario

Shady slopes: Avalanche prone weak layers exist in the old snowpack in particular on northwest, north and northeast facing slopes.

Outgoing longwave radiation during the night will be severely restricted. As a consequence of high relative humidity the snowpack can not consolidate. Sunny slopes: The weather conditions gave rise to moistening of the snowpack.

### Tendency

Low avalanche danger will prevail.