Wednesday 18.03.2026

Published 17 03 2026, 17:00



1	2	3	4	5
low	moderate	considerable	high	very high

Danger Level 3 - Considerable

Treeline

Tendency: Increasing avalanche danger
on Thursday 19 03 2026



Wind slab

Treeline

New snow

Treeline

The fresh snow and the wind slabs must be evaluated with care and prudence.

The fresh snow and the extensive wind slabs must be evaluated with care and prudence. Avalanches can in isolated cases be released in the old snowpack and reach quite a large size on very steep northwest, north and northeast facing slopes.
As a consequence of warming during the day and solar radiation small and, in isolated cases, medium-sized moist loose snow avalanches are possible.

Snowpack

Danger patterns dp.6: cold, loose snow and wind | dp.1: deep persistent weak layer

Very steep, little used shady slopes: Avalanche prone weak layers exist in the old snowpack in particular on northwest, north and northeast facing slopes. This applies at high altitude.

Outgoing longwave radiation during the night will be good in some case. Sunshine and high temperatures will give rise as the day progresses to increasing moistening of the snowpack.

Tendency

Fresh wind slabs are in some cases still prone to triggering in particular on steep shady slopes. The avalanche danger will decrease gradually.

Danger Level 3 - Considerable

Treeline

Tendency: Increasing avalanche danger
on Thursday 19 03 2026



Wind slab

Treeline

Persistent
weak layer

2000m

The fresh snow and the wind slabs can be released easily.

The fresh snow and the sometimes deep wind slabs can be released easily in particular on northwest to north to northeast facing aspects above the tree line. Avalanches can in isolated cases release deeper layers of the snowpack and reach large size.
Weakly bonded old snow requires caution. These avalanche prone locations are to be found in particular in northwest to north to northeast facing aspects. Caution is to be exercised in particular on steep shady slopes, as well as at transitions from a shallow to a deep snowpack, when entering gullies and bowls for example.
As a consequence of warming during the day and solar radiation small and medium-sized natural avalanches are possible.

Snowpack

Danger patterns dp.6: cold, loose snow and wind | dp.1: deep persistent weak layer

Over a wide area 30 to 50 cm of snow, and even more in some localities, fell in the last two days above approximately 1700 m. Distinct weak layers exist in the bottom section of the snowpack in particular on steep shady slopes. This applies especially above approximately 2000 m.
Below approximately 1800 m: The old snowpack remains moist.

Tendency

Fresh wind slabs are in some cases still prone to triggering in particular on steep shady slopes. The avalanche danger will decrease gradually.