Thursday 05.03.2026

Published 04 03 2026, 17:00

AM



PM



1	2	3	4	5
low	moderate	considerable	high	very high

Danger Level 2 - Moderate

AM:



Tendency: Constant avalanche danger
on Friday 06 03 2026

Persistent weak layer

2400m

PM:



2400m

Tendency: Constant avalanche danger
on Friday 06 03 2026



Wet snow

2400m
2000m

Persistent weak layer

2400m

The backcountry touring conditions are generally favourable.

The surface of the snowpack will freeze to form a strong crust only at high altitudes, in the event of an overcast night in particular.
Slight increase in danger as a consequence of warming during the day. In particular at intermediate altitudes individual mostly small moist and wet avalanches are possible as the day progresses, in particular in case of releases originating from extremely steep starting zones that still retain some snow.
Weak layers exist deeper in the snowpack. Avalanches can in very isolated cases be released in deep layers and reach medium size in isolated cases. These avalanche prone locations are to be found especially at transitions from a shallow to a deep snowpack, when entering gullies and bowls for example and in extremely steep terrain.

Snowpack

Danger patterns dp.1: deep persistent weak layer | dp.10: springtime scenario

The weather conditions fostered a strengthening of the snowpack in particular on sunny slopes. Towards its surface, the snowpack is largely stable and its surface has a melt-freeze crust that is strong in many cases. In steep terrain there is a danger of falling here.
Especially steep shady slopes high altitudes and the high Alpine regions: Weak layers exist in the old snowpack. Towards its surface, the snowpack consists of faceted crystals; its surface consists of faceted crystals.
Adjacent to ridgelines and in pass areas as well as at high altitude snow depths vary greatly, depending on

the infuence of the wind.

Tendency

The backcountry and freeriding conditions are quite favourable. In steep terrain there is a danger of falling on the hard snow surface. As a consequence of warming during the day there will be no slight decrease in the danger of moist and wet avalanches in the late morning.