

**2019 Math 340:101. Quiz 5** Friday, 2019 November 8 IN CLASS

Last name .....

First name .....

**Student number** ..... .....

**Grade**

1. (3 points) Given the following dictionary of a primal problem, find the corresponding **dual dictionary**.

$$\begin{array}{rcl}
 x_2 & = & 1 \quad -x_1 \quad -x_3 \\
 x_4 & = & 2 \quad -2x_1 \quad +x_3 \\
 x_5 & = & 2 \quad +x_1 \quad +x_3 \\
 \hline
 z & = & 3 \quad -x_1 \quad +x_3
 \end{array}$$

**Solution:**

- First put the objective function at the first row:

$$\begin{array}{rcl}
 z & = & 3 \quad -x_1 \quad +x_3 \\
 \hline
 x_2 & = & 1 \quad -x_1 \quad -x_3 \\
 x_4 & = & 2 \quad -2x_1 \quad +x_3 \\
 x_5 & = & 2 \quad +x_1 \quad +x_3
 \end{array}$$

- Recall the correspondence between the primal and dual variables. As there are 2 original variables and 3 slack variables, we have

$$\begin{aligned}
 x_j &\longleftrightarrow y_{3+j} \text{ for } j = 1, 2, \\
 x_{2+i} &\longleftrightarrow y_i \text{ for } i = 1, 2, 3.
 \end{aligned}$$

- Then, use the “– transpose” property of the dual dictionary

$$\begin{array}{rcl}
 -w & = & -3 \quad -y_5 \quad -2y_2 \quad -2y_3 \\
 \hline
 y_4 & = & 1 \quad +y_5 \quad +2y_2 \quad -y_3 \\
 y_1 & = & -1 \quad +y_5 \quad -y_2 \quad -y_3
 \end{array}$$

2. (2 points) (1 mark per two correct answers) We use the notation in the class. You are given  $A, \vec{b}, \vec{c}, \vec{x}_B, \vec{x}_N, B$  and  $N$ . Let  $\vec{x}^*$  be the basic solution of the corresponding primal dictionary, and let  $\vec{y}^*$  be the corresponding dual basic solution (i.e. the basic solution of the dual dictionary). Check whether the following statements are true or false.
- True / False** If  $\vec{x}^*, \vec{y}^*$  are primal/dual feasible, respectively, then they are primal/dual optimal, respectively.
  - True / False** It is possible in some examples, that  $\vec{x}^*$  is not primal feasible and  $\vec{y}^*$  is not dual feasible.
  - True / False** If  $\vec{x}^*$  is the unique primal optimal solution, then  $\vec{y}^*$  is the unique dual optimal solution.
  - True / False** If we change both  $\vec{b}$  and  $\vec{c}$  to different vectors  $\vec{b}'$  and  $\vec{c}'$ , respectively, while  $A, \vec{x}_B, \vec{x}_N, B$  and  $N$  remain the same, then the resulting primal and dual basic solutions  $\vec{x}^{**}, \vec{y}^{**}$  will have to be different from  $\vec{x}^*, \vec{y}^*$ , respectively.

**Solution.**

- (a) True.
- (b) True.
- (c) False.
- (d) False.

3. We use the notation as in the class. You are given  $A, \vec{b}, \vec{c}, \vec{x}_B, B$  and  $B^{-1}$ .

$$[A|I] = \begin{array}{c|ccccc} & x_1 & x_2 & x_3 & x_4 & x_5 \\ \hline & 2 & 1 & 1 & 0 & 0 \\ & 1 & 2 & 0 & 1 & 0 \\ & 0 & 1 & 0 & 0 & 1 \end{array}, \quad \vec{b} = \begin{bmatrix} 4 \\ 3 \\ 1 \end{bmatrix}, \quad \vec{x}_B = \begin{bmatrix} x_3 \\ x_1 \\ x_2 \end{bmatrix}$$

$$\vec{c}^T = \begin{array}{ccccc} & x_1 & x_2 & x_3 & x_4 & x_5 \\ \hline & 3 & 2 & 0 & 0 & 0 \end{array}, \quad B = \begin{array}{ccc} & x_3 & x_1 & x_2 \\ \hline & 1 & 2 & 1 \\ & 0 & 1 & 2 \\ & 0 & 0 & 1 \end{array}, \quad B^{-1} = \begin{bmatrix} 1 & -2 & 3 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix}$$

Let  $\vec{x}^*$  be the basic solution of the corresponding primal dictionary, and let  $\vec{y}^*$  be the corresponding dual basic solution (i.e. the basic solution of the dual dictionary).

(i) (3 points) Find  $x_1^*, x_2^*, x_3^*, x_4^*, x_5^*$  and  $y_1^*, y_2^*, y_3^*, y_4^*, y_5^*$ .

**Solution:** As  $\vec{x}^*$  is the basic solution

$$\vec{x}_B^* = \begin{bmatrix} x_3^* \\ x_1^* \\ x_2^* \end{bmatrix} = B^{-1}\vec{b} = \begin{bmatrix} 1 & -2 & 3 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 4 \\ 3 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

and

$$\vec{x}_N^* = \begin{bmatrix} x_4^* \\ x_5^* \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

As  $\vec{y}^*$  is the dual basic solution,  $(\vec{y}^*)^T B = \vec{c}_B^T$ , so

$$(\vec{y}^*)^T = \vec{c}_B^T B^{-1} = \begin{array}{ccc} & x_3 & x_1 & x_2 \\ \hline & 0 & 3 & 2 \end{array} \begin{bmatrix} 1 & -2 & 3 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix} = \begin{array}{ccc} y_1 & y_2 & y_3 \\ \hline 0 & 3 & -4 \end{array}$$

So,  $(y_1^*, y_2^*, y_3^*) = (0, 3, -4)$ . Note that  $\vec{y}_B = (y_1, y_4, y_5)$  using the primal/dual variable correspondence. Note that because  $\vec{y}^*$  is dual basic,  $\vec{y}_B^* = (y_1^*, y_4^*, y_5^*) = \vec{0}$ . Therefore,  $(y_1^*, y_2^*, y_3^*, y_4^*, y_5^*) = (0, 3, -4, 0, 0)$ .

(ii) (2 points) Is  $\vec{x}^*$  optimal? To earn credit, you have to justify your answer.

**Solution.**

Note that  $\vec{x}_B^* > \vec{0}$  so  $\vec{x}^*$  is non-degenerate. if it was optimal then the dual basic solution  $\vec{y}^*$  must be dual feasible. However,  $(y_1^*, y_2^*, y_3^*, y_4^*, y_5^*) \not\geq \vec{0}$ , so not dual feasible.

Blank page