

## SPEECH PROCESSING

### Linear Predictive Coding (LPC)

Patrick A. Naylor  
Spring Term 2018/19


Imperial College London

## PART 1

- This lecture studies one of the most important concepts underpinning many applications of speech processing, namely LPC
  - Concept of Linear Prediction
  - Derivation of Linear Prediction Equations
  - Autocorrelation method of LPC
  - Interpretation of LPC filter as a spectral whitener

Imperial College London

## Concept of Linear Prediction



- $u(n)$  volume flow at the glottis
- $u_l(n)$  volume flow at the lips
- $s(n)$  pressure at the microphone
- $V(z) = \frac{Gz^{-p/2}}{1 - \sum_{j=1}^p a_j z^{-j}} = \frac{Gz^{-p/2}}{A(z)}$  vocal tract transfer function
- $R(z) = 1 - z^{-1}$  lip radiation model
- The aim of Linear Prediction Analysis (LPC) is to estimate  $V(z)$  from the speech signal  $s(n)$ .

Imperial College London

3

## Notes

- We will neglect the pure delay term  $z^{-1/p}$  in the numerator of  $V(z)$ .
- 50% of the world puts a + sign in the denominator of  $V(z)$  (this is almost essential when using MATLAB).

$$V(z) = \frac{Gz^{-p/2}}{1 - \sum_{j=1}^p a_j z^{-j}}$$

Imperial College London

4

## Preview ... in straightforward terms

- Predict sample  $s(n)$  from samples  $s(n-1), s(n-2), \dots, s(n-p)$
- Consider prediction of 4 samples from their previous 2

$$s(2) = a_1s(1) + a_2s(0)$$

$$s(3) = a_1s(2) + a_2s(1)$$


$$s(4) = a_1s(3) + a_2s(2)$$

$$s(5) = a_1s(4) + a_2s(3)$$


- This is an overdetermined system of simultaneous equations
  - If we try to predict only 2 samples, the exact solution for the coefficients can be found
  - Otherwise we consider a least squares solution
- Call the prediction  $\hat{s}(n)$

- Important points to consider in determining the least squares solution
  - The frame  $\{F\}$  of samples over which to solve
  - Method of solution
    - Formulate the linear algebra problem in the form  $\mathbf{X}\mathbf{a}=\mathbf{b}$
    - Solve by matrix inversion
- These issues are the main points to discuss in this talk
- What should  $p$  be to predict successfully:
  - A sinusoid?
  - Voiced speech?
  - Unvoiced speech?
  - The stock market?
- LPC captures the harmonic content of a signal.
  - Anything not harmonic is unpredictable and gives a prediction error.

## Linearity



- We can reverse the order of  $V(z)$  and  $R(z)$  since both are linear and  $V(z)$  doesn't change substantially during the impulse response of  $R(z)$  or vice-versa ( $R(z)$  has an impulse duration of typically 2 samples):



Imperial College London

7

## Prediction Error

$$s(n) = Gu'(n) + \sum_{j=1}^p a_j s(n-j)$$

Typically use  $i$  or  $j$  for counter

- If the vocal tract resonances have high gain, the second term will dominate:

$$s(n) \approx \sum_{j=1}^p a_j s(n-j)$$

- The right hand side of this expression is a *prediction* of  $s(n)$  as a *linear sum* of past speech samples. Define the *prediction error* at sample  $n$  as

$$e(n) = s(n) - \sum_{j=1}^p a_j s(n-j) = s(n) - a_1 s(n-1) - a_2 s(n-2) - \dots - a_p s(n-p)$$

- In terms of z-transforms

$$E(z) = S(z)A(z)$$

Imperial College London

8

## Error Minimization

- Given a frame of speech  $\{F\}$ , we would like to find the values  $a_i$  that minimize

$$Q_E = \sum_{n \in \{F\}} e^2(n) \quad [1]$$

- To do so, we differentiate w.r.t each  $a_i$

$$\frac{\partial Q_E}{\partial a_i} = \sum_{n \in \{F\}} \frac{\partial(e^2(n))}{\partial a_i} = \sum_{n \in \{F\}} 2e(n) \frac{\partial e(n)}{\partial a_i} = - \sum_{n \in \{F\}} 2e(n)s(n-i)$$

Imperial College London

9

- The optimum values of  $a_i$  must satisfy  $p$  equations:

$$\begin{aligned} & \sum_{n \in \{F\}} e(n)s(n-i) = 0 \quad \text{for } i = 1, \dots, p \\ \Rightarrow & \sum_{n \in \{F\}} \left( s(n)s(n-i) - \sum_{j=1}^p a_j s(n-j)s(n-i) \right) = 0 \quad \text{for } i = 1, \dots, p \\ \Rightarrow & \sum_{j=1}^p a_j \sum_{n \in \{F\}} s(n-j)s(n-i) = \sum_{n \in \{F\}} s(n)s(n-i) \\ \Rightarrow & \sum_{j=1}^p \phi_{ij} a_j = \phi_{i0} \quad \text{where } \phi_{ij} = \sum_{n \in \{F\}} s(n-i)s(n-j) \end{aligned}$$

- which can be written in matrix form

$$\Phi \mathbf{a} = \mathbf{c} \Rightarrow \mathbf{a} = \Phi^{-1}\mathbf{c} \quad \text{providing } \Phi^{-1} \text{ exists}$$


– the matrix  $\Phi$  is symmetric and positive semi-definite.

Imperial College London

10

## Frame-based Processing

- Consider frame-based processing of a speech signal
  - Extract a set of frames of the speech signal employing a tapered window of duration 20 – 30 ms typically overlapping by 50%



Imperial College London

13

## Autocorrelation LPC

- Take  $\{F\}$  in equation [1] to be of infinite extent

$$\phi_{ij} = \sum_{n=-\infty}^{+\infty} s(n-i)s(n-j)$$

- Because of the symmetry and the infinite sum, we have

$$\phi_{ij} = \phi_{|i-j|,0} = R_{|i-j|}$$

- where the sequence  $R_k$  is the autocorrelation of the windowed speech

- The matrix  $\Phi$  is now Toeplitz (constant diagonals) and the equations

$$\Phi \mathbf{a} = \mathbf{c}$$


are called the *Yule-Walker* equations.

- Inverting a symmetric, positive definite, Toeplitz  $p \times p$  matrix takes  $O(p^2)$  operations instead of the normal  $O(p^3)$ . Inversion procedure is known as the Levinson or Levinson-Durbin algorithm.

Imperial College London

14

## Autocorrelation LPC example: /a/ from “father”




$e(n)$  is the result of inverse filtering  $s(n)$  with the prediction filter

$$E(z) = S(z)A(z)$$

Imperial College London

15

## Resulting Spectra and Poles



Imperial College London

16

## Spectral Flatness

- Autocorrelation LPC finds the filter of the form
$$A(z) = 1 - a_1 z^{-1} - \dots - a_p z^{-p}$$
that minimizes the energy of the prediction error.
- We will show that we can also interpret this in terms of flattening the spectrum of the error signal
- Define the normalized power spectrum of the prediction error signal  $e(n)$

$$P_E(\omega) = \frac{|E(e^{j\omega})|^2}{Q_E} \quad Q_E = \sum e^2(n) = \frac{1}{2\pi} \int_{\omega=0}^{2\pi} |E(e^{j\omega})|^2 d\omega$$

- where  $E(z)$  is the  $Z$ -transform of the signal and  $Q_E$  is the signal energy. The average value of  $P_E$  is equal to 1.


Imperial College London

17

- We define the *spectral roughness* of the signal as:

$$R_E = \frac{1}{2\pi} \int_{\omega=0}^{2\pi} P_E(\omega) - 1 - \log(P_E(\omega)) d\omega$$

- $R_E$  is similar to the variance of  $P_E$  since
  - the integrand is similar to  $\frac{1}{2}(P_E - 1)^2$  where  $\text{mean}(P_E) = 1$ .



Imperial College London

18

- We can find an alternative expression for  $R_E$

$$\begin{aligned}
 R_E &= \frac{1}{2\pi} \int_{\omega=0}^{2\pi} P_E(\omega) - 1 - \log(P_E(\omega)) d\omega \\
 &= \frac{1}{2\pi} \int_{\omega=0}^{2\pi} -\log(P_E(\omega)) d\omega \quad \text{since } \int P_E(\omega) d\omega = 1 \\
 &= \log(Q_E) - \frac{1}{2\pi} \int_{\omega=0}^{2\pi} \log\left(\left|E(e^{j\omega})\right|^2\right) d\omega \quad \text{since } P_E(\omega) = \frac{\left|E(e^{j\omega})\right|^2}{Q_E}
 \end{aligned}$$

- Thus the spectral roughness of a signal equals the difference between its log energy and the average of its log energy spectrum.

- We know that  $E(z) = S(z) \times A(z)$ , hence

$$\log\left(\left|E(e^{j\omega})\right|^2\right) = \log\left(\left|S(e^{j\omega})\right|^2\right) + \log\left(\left|A(e^{j\omega})\right|^2\right)$$

- Substituting this in the expression for  $R_E$  gives

$$\begin{aligned}
 R_E &= \log(Q_E) - \frac{1}{2\pi} \int_{\omega=0}^{2\pi} \log\left(\left|E(e^{j\omega})\right|^2\right) d\omega \\
 &= \log(Q_E) - \frac{1}{2\pi} \int_{\omega=0}^{2\pi} \log\left(\left|S(e^{j\omega})\right|^2\right) d\omega - \frac{1}{2\pi} \int_{\omega=0}^{2\pi} \log\left(\left|A(e^{j\omega})\right|^2\right) d\omega
 \end{aligned}$$



- We saw in the section on filter properties that the term involving  $A$  is zero since  $a_0=1$  and all roots of  $A$  lie inside the unit circle. Hence

$$R_E = \log(Q_E) - \frac{1}{2\pi} \int_{\omega=0}^{2\pi} \log\left(\left|S(e^{j\omega})\right|^2\right) d\omega$$

- The term involving  $S$  is independent of  $A$ . It follows that if  $A$  is chosen to minimize  $Q_E$ , it will also minimize  $R_E$ , the spectral roughness of  $e(n)$ . The filter  $A(z)$  is a whitening filter because it makes the spectrum flatter.

## Example



- These two graphs show a windowed speech signal, /a/, and the error signal after filtering by  $A(z)$



Imperial College London

21

- These graphs show the log energy spectrum of each signal
  - The two horizontal lines on each graph are the mean value (same for both graphs) and the log of the total energy.
  - The spectral roughness is the difference between the two



Imperial College London

22

## PART 2

- In this lecture, we look at further elements under the general heading of Linear Prediction
  - Covariance method of LPC
  - Preemphasis
  - Closed Phase Covariance LPC
  - Alternative LPC parameter sets:
  - Pole positions
  - Reflection Coefficients
  - Log Area Ratios

Imperial College London

23

## Variants of LPC

We consider two variants of LPC analysis which differ only in their choice of speech frame,  $\{F\}$

- Autocorrelation LPC Analysis
  - Requires a windowed signal
    - tradeoff between spectral resolution and time resolution
  - Requires >20 ms of data
  - Has a fast algorithm because  $\Phi$  is Toeplitz symmetric
  - Guarantees a stable filter  $V(z)$
- Covariance LPC Analysis (Prony's method)
  - No windowing required
  - Gives infinite spectral resolution
  - Requires >2 ms of data
  - Slower algorithm because  $\Phi$  is not Toeplitz symmetric
  - Sometimes gives an unstable filter  $V(z)$

Imperial College London

24

## Covariance LPC

- Already seen that  $\sum_{j=1}^p \phi_{ij} a_j = \phi_{i0}$  where  $\phi_{ij} = \sum_{n \in \{F\}} s(n-i)s(n-j)$
- Now we choose  $\{F\}$  to be a finite segment of speech:  
 $\{F\} = s(n) \text{ for } 0 \leq n \leq (N-1)$   
then we have:  
$$\phi_{ij} = \sum_{n=0}^{N-1} s(n-i)s(n-j)$$
- The matrix  $\Phi$  is still symmetric but is no longer Toeplitz
  - Since the matrix is not Toeplitz, the computation involved in inverting  $\Phi$  is  $\propto p^3$  rather than  $p^2$  and so takes longer
- Covariance LPC generally gives better results than Autocorrelation LPC but is more sensitive to the precise position of the frame in relation to the vocal fold closures.

Imperial College London

25

## Recursive Computation

- The entire matrix  $\Phi$  can be calculated recursively from its first row or column.


$$\begin{aligned}\phi_{ij} &= \sum_{n=-1}^{N-2} s(n-i+1)s(n-j+1) \\ &= s(-i)s(-j) - s(N-i)s(N-j) + \sum_{n=0}^{N-1} s(n-i+1)s(n-j+1) \\ &= s(-i)s(-j) - s(N-i)s(N-j) + \phi_{i-1,j-1}\end{aligned}$$

Imperial College London

26

## Unstable Poles

- Covariance LPC does not necessarily give a stable filter  $V(z)$ 
  - (though it usually does).
- We can force stability by replacing an unstable pole at  $z = p$  by a stable pole at  $z = 1/p^*$



See analysis of reflection in the unit circle from previous lecture material

Imperial College London

27

- As we have seen in the section on filter properties, reflecting a pole in the unit circle leaves the magnitude response unchanged except for multiplying by a constant (equal to the magnitude of the pole).
  - spectral flattening property of LPC is unaltered by this pole reflection.
- Discovering which poles lie outside the unit circle is quite expensive in terms of computation
  - this is a further computational disadvantage of covariance LPC.

Imperial College London


28

## Pre-emphasis

- The matrix  $\Phi$  is always non-singular, but not necessarily by very much.
- A measure of how close a matrix is to being singular is given by its condition number
  - for a symmetric +ve definite matrix, this is the ratio of its largest to its smallest eigenvalue.
- For large  $p$ , the condition number of  $\Phi$  tends to the ratio  $S_{\max}(\omega)/S_{\min}(\omega)$ .
- We can thus improve the numerical properties of the LPC analysis procedure by flattening the speech spectrum before calculating the matrix  $\Phi$ .
- For voiced speech, the input to  $V(z)$  is  $u'(n)$  whose magnitude spectrum reduces at high frequencies with a gradient of about -6 dB/octave
  - This can be compensated with a 1st-order high-pass filter with a zero near  $z=1$

$$P(z) = 1 - \alpha z^{-1}$$

- $P(z)$  is approximately a differentiator
- The normalised corner frequency of  $P(z)$  is approximately  $(1-\alpha)/2\pi$
- This is typically placed in the range 0 to 150 Hz.
- From a spectral flatness point of view, the optimum value of  $\alpha$  is  $\phi_{10}/\phi_{00}$  (obtained from autocorrelation LPC with  $p = 1$ ).



Variable	Description	Spectral Envelope
$u(n)$	Glottal volume velocity.	$\sim -12$ dB/oct
$u'(n)$	Derivative of glottal volume velocity / voice source signal.	$\sim -6$ dB/oct
$u''(n)$	2 <sup>nd</sup> derivative of glottal volume velocity / derivative of voice source signal / whitened voice source signal.	Flat
$s'(n)$	Preemphasised speech signal / whitened speech signal.	$\sim V(e^{j\omega})$
$s(n)$	Speech signal.	$\sim V(e^{j\omega})$ with -6dB/oct tilt


## Closed-phase Covariance LPC

- We have already seen that  $s(n) = Gu'(n) + \sum_{j=1}^p a_j s(n-j)$
- We have neglected the term  $Gu'(n)$  because we don't know what it is and it is assumed to be much smaller than the second term
- If we knew when the vocal folds were closed, we could restrict  $\{F\}$  to those particular intervals
  - In those intervals the assumption  $Gu'(n)=0$  is substantially valid
- We can estimate the times of vocal fold closure in two ways
  - Search for spikes (impulsive events) in the prediction residual signal  $e(n)$
  - Using a Laryngograph (or Electroglottograph or EGG): this instrument measures the radio-frequency conductance across the larynx.
    - Conductance  $\propto$  Vocal fold contact area.
    - Accurate but inconvenient.

Imperial College London

31


- In Closed-Phase LPC, we choose our analysis interval  $\{F\}$  to consist of one or more closed phase intervals
  - (not necessarily contiguous)
- No preemphasis is necessary because the excitation now has a flat spectrum



Imperial College London

32

## Closed Phase Analysis of /i/ from 'bee'



## Alternative Parameter Sets

- The vocal tract filter is defined by  $p+1$  parameters:

$$V(z) = \frac{G}{1 - \sum_{k=1}^p a_k z^{-k}}$$

- The LPC (or AR) coefficients  $a_k$  have some bad properties:
  - The frequency response of  $V(z)$  is very sensitive to small changes in  $a_k$  (such as quantizing errors in coding)
  - There is no easy way to verify that the filter is stable
  - Interpolating between the parameters that correspond to two different filters will not vary the frequency response smoothly from one to the other: stability is not even guaranteed.
- There are several alternative parameter sets that are equivalent to the  $a_k$ 
  - most require  $G$  to be specified as well

## Pole Positions

- We can factorize the denominator of  $V(z)$  to give its poles:

$$1 - \sum_{k=1}^p a_k z^{-k} = \prod_{k=1}^p (1 - x_k z^{-1})$$

- The polynomial roots  $x_k$  are either real or occur in complex conjugate pairs.  $|x_k|$  must be  $< 1$  for stability
- Factorizing polynomials is computationally expensive
- The frequency response is (very) sensitive to pole position errors near  $|z|=1$ .

## Reflection Coefficients

- Any all-pole filter is equivalent to a tube with  $p$  sections: this is characterized by  $p$  reflection coefficients (assuming  $r_0=1$ )
- We can convert between the reflection coefficients and the polynomial coefficients by using the formulae given earlier in the course<sup>[1]</sup>
- Properties:
  - An all-pole filter is stable iff the corresponding reflection coefficients all lie between -1 and +1
  - Interpolating between two reflection coefficient sets will give a smoothly changing frequency response
  - High coefficient sensitivity near  $\pm 1$
- The negative reflection coefficients are sometimes called the *PARCOR* coefficients (PARCOR = partial correlation)

[1] The Levinson-Durbin Recursion

## Log Area Ratios

- Log area ratios are derived from the lossless tube model

$$g_i = \log\left(\frac{A_{i+1}}{A_i}\right) = \log\left(\frac{1+r_i}{1-r_i}\right) \Leftrightarrow r_i = \frac{e^{g_i} - 1}{e^{g_i} + 1} = \tanh(g_i)$$

- Stability is guaranteed for any values of  $g_i$ .

Imperial College London

37

## PART 3

- In this lecture, we at more alternative sets of LPC coefficients and their applications
  - Cepstral Coefficients
    - Relation to pole positions
    - Relation to LPC filter coefficients
  - Line Spectrum Frequencies
    - Relation to pole positions and to formant frequencies
  - Summary of LPC parameter sets

Imperial College London

38

## Cepstral Coefficients

- Most *speech recognisers* describe the spectrum of speech sounds using *cepstral coefficients*
  - good at discriminating between different phonemes
  - substantially independent of each other
  - have approximately Gaussian distributions for a particular phoneme
- Cepstrum is defined as inverse Fourier transform of log spectrum
  - (periodic spectrum -> discrete cepstrum)

$$c_n = \frac{1}{2\pi} \int_{\omega=-\pi}^{+\pi} \log(V(e^{j\omega})) e^{j\omega n} d\omega$$

- Can be computed either from roots of the prediction filter polynomial or from the coefficients of the prediction filter polynomial.

Imperial College London

39

## Computation from Roots $x_k$

- Define the cepstral coefficients  $c_n$  in terms of

$$C(z) = \sum_{n=-\infty}^{+\infty} c_n z^{-n} \Rightarrow c_n = \frac{1}{2\pi} \int_{\omega=-\pi}^{+\pi} C(e^{j\omega}) e^{j\omega n} d\omega$$

- This is the standard inverse z-transform derived by taking the inverse Fourier transform of both sides of the first equation.
- By equating the Fourier transforms of the two expressions for  $c_n$ , we get

$$\begin{aligned} C(z) &= \log(V(z)) \\ &= \log\left(\frac{G}{A(z)}\right) = \log(G) - \log(A(z)) \end{aligned}$$

$$\text{where } A(z) = 1 - \sum_{k=1}^P a_k z^{-k} = \prod_{k=1}^P (1 - x_k z^{-1})$$

Imperial College London

40

- Next, using the Taylor series  $\log(1-y) = -\sum_{n=1}^{\infty} \frac{y^n}{n}$  for  $|y| < 1$

$$C(z) = \log(G) - \log(A(z))$$

$$\begin{aligned} &= \log(G) - \sum_{k=1}^p \log(1-x_k z^{-1}) \\ &= \log(G) + \sum_{k=1}^p \sum_{n=1}^{\infty} \frac{x_k^n}{n} z^{-n} \end{aligned}$$

- By collecting all the terms in  $z^{-n}$ , we obtain  $c_n$  in terms of  $x_k$ :

$$c_n = \begin{cases} 0 & \text{for } n < 0 \\ \log(G) & \text{for } n = 0 \\ \sum_{k=1}^p \frac{x_k^n}{n} & \text{for } n > 0 \end{cases}$$

- Because  $|x_k| < 1$  the  $c_n$  decrease exponentially with  $n$ .

Imperial College London

41

## Computation from Coefficients $a_k$

- Differentiating  $C(z) = \log(G) - \log(A(z))$  with respect to  $z$ :

$$\begin{aligned} C'(z) &= \frac{-A'(z)}{A(z)} \Rightarrow A(z)C'(z) = -A'(z) \\ &\Rightarrow A(z)zC'(z) = -zA'(z) \end{aligned}$$

- Gives

$$\begin{aligned} &\left(1 - \sum_{k=1}^p a_k z^{-k}\right) \left(z \sum_{m=0}^{\infty} -mc_m z^{-(m+1)}\right) = -z \sum_{n=1}^p n a_n z^{-(n+1)} \\ &\Rightarrow \left(1 - \sum_{k=1}^p a_k z^{-k}\right) \left(\sum_{m=1}^{\infty} mc_m z^{-m}\right) = \sum_{n=1}^p n a_n z^{-n} \\ &\Rightarrow \sum_{n=1}^{\infty} n c_n z^{-n} - \sum_{k=1}^p \sum_{m=1}^{\infty} mc_m a_k z^{-(m+k)} = \sum_{n=1}^p n a_n z^{-n} \end{aligned}$$

- Replacing  $m$  by  $n-k$  (to make the  $z$  exponent uniform) gives:

$$\Rightarrow \sum_{n=1}^{\infty} n c_n z^{-n} = \sum_{n=1}^p n a_n z^{-n} + \sum_{k=1}^p \sum_{n=k+1}^{\infty} (n-k) c_{(n-k)} a_k z^{-n}$$

Imperial College London

42

- Now take the coefficient of  $z^{-n}$  in the above equation noting that  
 $n \geq k+1 \Rightarrow k \leq n-1$

$$nc_n = na_n + \sum_{k=1}^{\min(p,n-1)} (n-k)c_{(n-k)}a_k$$

$$\Rightarrow c_n = a_n + \frac{1}{n} \sum_{k=1}^{\min(p,n-1)} (n-k)c_{(n-k)}a_k$$

- Thus we have a recurrence relation to calculate the  $c_n$  from the  $a_k$  coefficients

$$c_n = a_n + \frac{1}{n} \sum_{k=1}^{\min(p,n-1)} (n-k)c_{(n-k)}a_k$$

- From which

$$c_1 = a_1$$

$$c_2 = a_2 + \frac{1}{2}c_1a_1$$


$$c_3 = a_3 + \frac{1}{3}(2c_2a_1 + c_1a_2)$$

$$c_4 = a_4 + \frac{1}{4}(3c_3a_1 + 2c_2a_2 + c_1a_3)$$

$$c_5 = \dots$$

- These coefficients are called the *complex cepstrum* coefficients
  - even though they are real
- The *cepstrum* coefficients use  $\log|V|$  instead of  $\log(V)$ 
  - half as big, except for  $c_0$
- Note the cute names:

- spectrum  $\rightarrow$  cepstrum ; frequency  $\rightarrow$  quefrency ; filter  $\rightarrow$  lifter ; etc



## Line Spectrum Frequencies (LSF)

- Consider  $A(z) = G \times V^{-1}(z) = 1 - \sum_{j=1}^p a_j z^{-j} = 1 - a_1 z^{-1} - a_2 z^{-2} - \dots - a_p z^{-p}$
- We can form symmetric and antisymmetric polynomials:
$$P(z) = A(z) + z^{-(p+1)} A^*(z^{*-1})$$

$$= 1 - (a_1 + a_p) z^{-1} - (a_2 + a_{p-1}) z^{-2} - \dots - (a_p + a_1) z^{-p} + z^{-(p+1)}$$


$$Q(z) = A(z) - z^{-(p+1)} A^*(z^{*-1})$$

$$= 1 - (a_1 - a_p) z^{-1} - (a_2 - a_{p-1}) z^{-2} - \dots - (a_p - a_1) z^{-p} - z^{-(p+1)}$$
- $V(z)$  is **stable** if and only if the roots of  $P(z)$  and  $Q(z)$  all lie on the unit circle and they are interleaved.


Imperial College London

45

- Example Poles:



LSFs:



- If the roots of  $P(z)$  are at  $\exp(2\pi j f_i)$  for  $i=1, 3, \dots$  and the roots of  $Q(z)$  are at  $\exp(2\pi j f_i)$  for  $i=0, 2, \dots$  with  $f_{i+1} > f_i \geq 0$ 
  - then the LSF frequencies are defined as  $f_1, f_2, \dots, f_p$ .
- Note that it is always true that  $f_0 = +1$  and  $f_{p+1} = -1$

E.g.

$$A(z) = 1 - 0.7z^{-1} + 0.5z^{-2} \quad P(z) = 1 - 0.2z^{-1} - 0.2z^{-2} + z^{-3}$$

$$z^{-3} A^*(z^{*-1}) = 0.5z^{-1} - 0.7z^{-2} + z^{-3} \quad Q(z) = 1 - 1.2z^{-1} + 1.2z^{-2} - z^{-3}$$

Imperial College London

46

## Proof that roots of $P(z)$ and $Q(z)$ lie on the unit circle

- Given  $P(z) = 0 \Leftrightarrow A(z) = -z^{-(p+1)}A^*(z^{*-1}) \Leftrightarrow H(z) = -1$
- $Q(z) = 0 \Leftrightarrow A(z) = +z^{-(p+1)}A^*(z^{*-1}) \Leftrightarrow H(z) = +1$
- where  $H(z) = \frac{A(z)}{z^{-(p+1)}A^*(z^{*-1})} = z \prod_{i=1}^p \frac{(1-x_i z^{-1})}{z^{-1}(1-x_i^* z)} = z \prod_{i=1}^p \frac{(z-x_i)}{(1-x_i^* z)}$
- here the  $x_i$  are the roots of  $A(z) = V^{-1}(z)$ .
- Providing all the  $x_i$  lie inside the unit circle, the absolute values of the terms making up  $H(z)$  are either all  $> 1$  or else all  $< 1$ .

Imperial College London

47

- Taking || of a typical term:

$$\begin{aligned}
 & \left| \frac{(z-x_i)}{(1-x_i^* z)} \right| > 1 \Leftrightarrow |1-x_i^* z| < |z-x_i| \\
 & \Leftrightarrow (1-x_i^* z)(1-x_i^* z)^* < (z-x_i)(z-x_i)^* \\
 & \Leftrightarrow (1-x_i^* z)(1-x_i^* z)^* < (z-x_i)(z^* - x_i^*) \\
 & \Leftrightarrow 1 - x_i^* z - x_i^* z^* + x_i x_i^* z z^* < z z^* - x_i^* z - x_i^* z^* + x_i x_i^* \\
 & \Leftrightarrow 1 - x_i x_i^* - z z^* + x_i x_i^* z z^* < 0 \\
 & \Leftrightarrow (1 - |x_i|^2)(1 - |z|^2) < 0 \Leftrightarrow |z| > 1 \quad \text{since each } |x_i| < 1
 \end{aligned}$$

- Thus each term is greater or less than 1 according to whether  $|z| > 1$  or  $|z| < 1$
- Hence  $|H(z)| = 1$  if and only if  $|z| = 1$  and so the roots of  $P(z)$  and  $Q(z)$  must lie on the unit circle.


Imperial College London

48

## Proof that the roots of $P(z)$ and $Q(z)$ are interleaved

- We want to find the values of  $z = e^{j\omega}$  that make  $H(z) = \pm 1$  or equivalently that make  $\arg(H(z)) = \pm \pi$ .

- If  $z = e^{j\omega}$  then




$$\begin{aligned}\arg(H(e^{j\omega})) &= \arg\left(e^{j(1-p)\omega} \prod_{i=1}^p \frac{(e^{j\omega} - x_i)}{(e^{-j\omega} - x_i^*)}\right) \\ &= (1-p)\omega + \sum_{i=1}^p (\arg(e^{j\omega} - x_i) - \arg(e^{-j\omega} - x_i^*)) \\ &= (1-p)\omega + 2 \sum_{i=1}^p \arg(e^{j\omega} - x_i)\end{aligned}$$

- As  $\omega$  goes from 0 to  $2\pi$ ,  $\arg(z-a)$  changes monotonically by  $+2\pi$  if  $|a|<1$
- Therefore as  $\omega$  goes from 0 to  $2\pi$ ,  $\arg(H(e^{j\omega}))$  increases by

$$(1-p) \times 2\pi + 2p \times 2\pi = (1+p) \times 2\pi$$

- Since  $H(e^{j\omega})$  goes round the unit circle  $(1+p)$  times, it must pass through each of the points  $+1$  and  $-1$  alternately  $(1+p)$  times



- $\arg(H(z))$  varies most rapidly when  $z$  is near one of the  $x_i$  so the LSF frequencies will cluster near the formants

## Summary of LPC parameter sets

- Filter Coefficients:  $a_i$ 
  - Stability check difficult; Sensitive to errors; Cannot interpolate
- Pole Positions:  $x_i$ 
  - + Stability check easy; Can interpolate but unordered.
  - Hard to calculate; Sensitive to errors near  $|x_i|=1$
- Reflection Coefficients:  $r_i$ 
  - + Stability check easy; Can interpolate
  - Sensitive to errors near  $\pm 1$
- Log Area Ratios:  $g_i$ 
  - + Stability guaranteed; Can interpolate
- Cepstral Coefficients :  $c_i$ 
  - + Good for speech recognition
  - Stability check difficult
- Line Spectrum Frequencies:  $f_i$ 
  - + Stability check easy; Can interpolate; Vary smoothly in time; Strongly correlated  $\Rightarrow$  better coding; Related to spectral peaks (formants).
  - Awkward to calculate