



从中子星天文观测到致密物质物态

报告人：文德华

华南理工大学物理与光电学院

2024年11月

目 录

1. 中子星研究概述
2. 关联性分析与普适性关系
3. 机器学习算法
4. 含暗物质中子星
5. 结论

目 录

1. 中子星研究概述

2. 关联性分析与普适性关系




3. 机器学习算法

4. 含暗物质中子星

5. 结论


中子星研究概述

- Neutron stars are a remarkable marriage of Einstein's theory of general relativity with nuclear physics.




中子星研究概述

- The interiors of neutron star harbor extreme matter that cannot be probed in the laboratory. At such high densities and pressures, their cores may consist predominantly of exotic matter such as free quarks or hyperons.



中子星研究概述 observations

- Neutron star observation related installation: Gravitational wave observations from the Laser Interferometer Gravitational-wave Observatory (LIGO), Neutron star observations from Five-hundred-meter Aperture Spherical radio Telescope (FAST), and X-ray observations from the Neutron Star Interior Composition Explorer (NICER), are beginning to run.





➤ Atmospheres (only a few centimeters closest to the surface) can have atoms.

➤ At densities greater than $\sim 10^7 \text{g/cm}^3$, the Fermi energy of electrons becomes high enough that the matter becomes progressively richer in neutrons.

➤ At densities greater than $4 \times 10^{11} \text{g/cm}^3$ it becomes possible for neutrons to “drip out” of the nucleus, which means that matter is a mix of free neutrons, free electrons, and nuclei. At about “nuclear saturation density” $2.7 \times 10^{14} \text{g/cm}^3$, there are no longer any isolated nuclei, and we now have the neutron soup.

➤ Pushing to yet higher densities, it may be energetically favorable to form other baryons with at least one strange quark, such as hyperons, until eventually close to the center of the star, quarks may become deconfined and one may encounter a degenerate quark-gluon plasma.





Mass–radius relations, where the gray area denotes the range of maximum mass accurately observed so far. The light red area denotes the marginalized posterior (M_g, R) of the merged binary neutron star released by LIGO and VIRGO collaboration in GW170817. The light blue and green areas denote the marginalized posterior (M_g, R_e) released by NICER for PSR J0030 + 0451 and PSR J0740 + 6620, respectively.

目 录

1. 中子星研究概述


2. 关联性分析与普适性关系

3. 机器学习算法

4. 含暗物质中子星

5. 结论

Kent Yagi* and Nicolás Yunes, Science 341, 365 (2013)



- These universal relations can be used to learn about neutron star deformability through observations of the moment of inertia (and vice versa), distinguish neutron stars from quark stars, and test general relativity.

One of the few purely theoretical studies of neutron stars to be published in SCIENCE.

Why do we study the **universal relations**?

- There is considerable uncertainty in understanding the equation of state for the super dense nuclear matter in neutron stars.
- The universal relation reflects the nature that some relations between the global properties of NS are not dependent on the EOS.
- This relation can be used to constrain the properties that cannot be directly observed (such as the gravitational binding energy) or cannot be accurately observed (such as the radius).
- The universal relation can be further used to constrain the EOS (such as the symmetry energy).

参量之间的关联性

The linear correlation, which reflects the strength of linear correlation between two quantities, can be expressed as


$$r(X, Y) = \frac{N \sum_{i=1}^N X_i Y_i - \sum_{i=1}^N X_i \sum_{i=1}^N Y_i}{\sqrt{N \sum_{i=1}^N X_i^2 - \left(\sum_{i=1}^N X_i \right)^2} \sqrt{N \sum_{i=1}^N Y_i^2 - \left(\sum_{i=1}^N Y_i \right)^2}}$$

Briefly:

$$r(X, Y) = \frac{\text{Cov}(X, Y)}{\sqrt{D(X)} \sqrt{D(Y)}}$$

The closer the absolute value of the correlation coefficient $|r|$ is to 1, the greater the correlation strength between the two quantities will be.



中子星性质参量之间的关联性



Linear correlation properties of the neutron star properties. It is shown that the quantities possessing desired linear correlation properties are the compactness β , moment of inertia I , gravitational redshift z and gravitational binding energy E_g .

中子星性质参量之间的关联性

Predicting the new universal (I/M^3 - β) relations based on the linear correlation analysis



The universal relation I/M^3 - β can be expressed as

$$\log I/M^3 = -1.261 \log \beta + 0.277$$

中子星性质与核物质参量的关联分析

- 饱和点处的核物质参量, L : 对称能斜率

$$e(\rho, \delta) = e(\rho) + S(\rho)\delta^2,$$
$$e(\rho) = e_0(\rho_0) + \frac{K_0}{2}x^2 + \frac{J_0}{6}x^3 + O(x^4), \quad x = \frac{\rho - \rho_0}{3\rho_0},$$
$$S(\rho) = E_{\text{sym}}(\rho_0) + \underbrace{L}_{\text{circled}}x + \frac{K_{\text{sym}}}{2}x^2 + O(x^3).$$

- 目标: 对中子星各种性质与核物质参量的关联特征进行系统性分析, 找出较好的关联, 并联合地面实验给出的核物质参量数据来对中子星性质进行约束, 并同天文观测数据进行比较分析其中的物理机制。
- 方法: Pearson 线性关联系数

对称能斜率 L 对 $1.4M_{\odot}$ 中子星性质的约束






TABLE I: The correlations between NS properties and $M_0 + aL$ (a represents the coefficient of maximal correlation). r represents the linear correlation coefficient, and $L_1 = 58.7 \pm 28.1\text{MeV}$, $L_2 = 106 \pm 37\text{MeV}$, $L_3 = 54 \pm 8\text{MeV}$, respectively.

	$M_0 + aL$				
	a	r	L_1	L_2	L_3
$R_{1.4}$	73.5	0.958	11.63-13.18	12.56-14.54	12.00-12.58
$\Lambda_{1.4}$	55.5	0.972	182.1-782.9	527.3-1285.1	321.7-561.3

由地面实验提取的 $L = 54 \pm 8\text{MeV}$ (Reinhard2021PRL)给出的中子星 $R_{1.4}$, $\Lambda_{1.4}$ 的约束范围能够与GW170817($R_{1.4} < 13.5\text{km}$, $70 \leq \Lambda_{1.4} \leq 580$),NICER的天文观测数据较好的符合

对称能斜率 L 对低质量中子星引力结合能 $|E_g|$ 的约束



$$\begin{aligned} |E_{g,0.4}| &= -3.653 \times 10^{-5} L + 0.019 \\ &= 0.017 M_{\odot}, \\ |E_{g,0.6}| &= -7.864 \times 10^{-5} L + 0.041 \\ &= 0.036 - 0.037 M_{\odot}, \\ |E_{g,0.8}| &= -1.311 \times 10^{-4} L + 0.072 \\ &= 0.064 - 0.066 M_{\odot}, \\ |E_{g,1.0}| &= -1.968 \times 10^{-4} L + 0.112 \\ &= 0.100 - 0.103 M_{\odot}. \end{aligned}$$

低质量中子星的引力结合能 $|E_g|$ 在理论上可以被 L 的约束范围较好的约束，其能够与超新星爆发中的总释放能量联系起来以用来预测低质量中子星的形成。


目 录

1. 中子星研究概述
2. 关联性分析与普适性关系
3. 机器学习算法
4. 含暗物质中子星
5. 结论

神经网络算法下中子星质量半径对EoS的约束

EOS: $P(\rho)$ $\xrightleftharpoons[\text{逆TOV映射}]{\text{TOV映射}}$ TOV方程: $\frac{dP}{dr} = -\frac{G\rho m(r)}{r^2} \left(1 + \frac{P}{\rho c^2}\right) \left(1 + \frac{4\pi P r^3}{m(r)c^2}\right) \left(1 - \frac{2Gm(r)}{rc^2}\right)^{-1}$

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho$$



如果将EOS到M-R的映射称为TOV映射，则M-R到EOS的映射可视为**逆TOV映射**

1.神经网络算法下中子星质量半径对EoS的约束

Parameter range (MeV)		
parameter	range	interval
L	30 ~ 143	+1
K _s	-400 ~ 100	+1
J _s	-200 ~ 800	+100
J ₀	-300~400	+100
K ₀	240	
E _{sym} (ρ ₀)	31.7	

总样本量约: 5,026,032

$$E_0(\rho) = E_0(\rho_0) + \frac{K_0}{2} \left(\frac{\rho - \rho_0}{3\rho_0} \right)^2 + J_0 \left(\frac{\rho - \rho_0}{3\rho_0} \right)^3$$


$$E_{sym}(\rho) = E_{sym}(\rho_0) + L \left(\frac{\rho - \rho_0}{3\rho_0} \right) + \frac{K_{sym}}{2} \left(\frac{\rho - \rho_0}{3\rho_0} \right)^2 + \frac{J_{sym}}{6} \left(\frac{\rho - \rho_0}{3\rho_0} \right)^3$$

筛选条件


Maximum mass	2.14M _⊙
tidal deformation	70~580(1.4M _⊙)
Mass radius corresponding	1.29~1.59M _⊙ /1.2~13.3km

筛选后剩余样本量884,294

神经网络算法下中子星质量半径对EoS的约束



40对 (M,R) 点作为输入, NN能够给出比较准确的结果, 当输入点降低至35点时相对误差显著变大。




只变化 $L-K_{sym}$ 时, 生成样本量约8,000。输入点数降低至**4**



目 录

1. 中子星研究概述
2. 关联性分析与普适性关系
3. 机器学习算法
4. 含暗物质中子星
5. 结论

含暗物质中子星





由于中子星极强的引力，它可能会吸收其附近的暗物质，或者与纯暗物质星合并形成含暗物质中子星。




暗物质对中子星潮汐形变以及半径的影响

自相互作用玻色暗物质

理想费米气暗物质



1. 潮汐形变 Λ 与普通物质半径 R_N 随 F_X 的变化;
2. 不同暗物质模型下含暗物质中子星的差别。



R_N 和 Λ 的负相关关系


当暗物质质量分数较大时，会出现普通物质半径 R_N 和潮汐形变 Λ 的负相关关系（虚线）

本质原因：增大暗物质质量分数时 R_N 减小，但是 Λ 随 $R = R_D$ 的增大而增大（III和IV）。


含暗物质中子星的宏观性质 —— R_N 和 Λ 负相关关系的普遍性

Hongyi Sun and DHW*, Phys. Rev. D 109, 123037 (2024)

自相互作用玻色暗物质



理想费米气暗物质



R_N 和 Λ 的负相关关系在改变暗物质模型及其物态参数的情况下是普遍存在的，而且可以利用该负相关关系来限制暗物质物态的参数。

目 录

1. 中子星研究概述
2. 关联性分析与普适性关系
3. 机器学习算法
4. 含暗物质中子星
5. 结论

结 论

- 采用线性相关性的分析方法可以更高效、更精确地预测中子星性质参量的普遍性关系。
- 利用构建的普适性关系，可以对无法观测或难以观测的中子星性质参量进行有效约束。
引力结合能是一种构建中子星性质参量普适性关系的理想参量。
- 利用神经网络算法，根据一定数量、一定精度的中子星质量-半径观测值可以给出较为精确的物态方程。
- 含暗物质中子星潮汐形变与普通物质半径之间的负相关关系可以作为中子星内是否含有暗物质的观测证据。

参考文献:

1. Zehan Wu, Dehua Wen*, From masses and radii of neutron stars to EOS of nuclear matter through neural network, **Chinese Physics C** 48, (2024) 024101
2. Hongyi Sun and Dehua Wen*, Implications of supermassive neutron stars for the form of the equation of state of hybrid stars, **Phys. Rev. C** 108, 025801 (2023)
3. Shen Yang, Dehua Wen*, Systematic correlation analysis between the nuclear matter parameters and neutron star properties within relativistic mean-field theory, **Phys. Rev. D** 107, 063009 (2023)
4. Jue Wang, Shen Yang, Dehua Wen*, Constraining properties of massive neutron star through R-mode, **Eur. Phys. J. A**, 58, 132 (2022)
5. Shen Yang, Dehua Wen*, Jue Wang and Jing Zhang, Exploring the universal relations with the correlation analysis of neutron star properties, **Phys. Rev. D** 105, 063023 (2022)
6. Yuxi Li, Jue Wang, Zehan Wu and Dehua Wen*, Inferring the gravitational binding energy and moment of inertia of PSR J0030 + 0451 and PSR J0740 + 6620 from new universal relations, **Class. Quantum Grav.** 39, 035014 (14pp) (2022)
7. Jing Zhang, Dehua Wen* and Yuxi Li, Constraint on nuclear symmetry energy imposed by f-mode oscillation of neutron stars, **Commun. Theor. Phys.** 73, 115302 (1-9) (2021)
8. Y. X. Li, H. Y. Chen, D. H. Wen* and J. Zhang, Constraining the nuclear symmetry energy and properties of the neutron star from GW170817 by Bayesian analysis, **Eur. Phys. J. A** 57, 31(2021)
9. W. J. Sun, D. H. Wen* and J. Wang, New quasi-universal relations for static and rapid rotating neutron stars, **Phys. Rev. D** 102, 023039 (2020)
10. R. R. Jiang, D. H. Wen* and H. Y. Chen, Universal behavior of compact star based upon gravitational binding energy, **Phys. Rev. D** 100, 123010 (2019).
11. B. A. Li, P. G. Krastev, D. H. Wen and N. B. Zhang, Towards understanding astrophysical effects of nuclear symmetry energy, **Eur. Phys. J. A** 55, 117 (2019)
12. D. H. Wen, B. A. Li, * H. Y. Chen and N. B. Zhang, GW170817 implications on the frequency and damping time of f -mode oscillations of neutron stars, **Phys. Rev. C** 99, 045806 (2019)
13. H. Y. Chen, D. H. Wen* and N. Zhang, Suitable resolution of EOS tables for neutron star investigations, **Chin. Phys. C** 43, 054108(2019)
14. N. Zhang, D. H. Wen* and H. Y. Chen, Imprint of the speed of sound in nuclear matter on global properties of neutron stars, **Phys. Rev. C** 99, 035803 (2019)



感谢聆听!