A Brief Crash Course in the Einstein Field Equations

Saavan Shah

October 27, 2026

Abstract

In the early twentieth century, modern physics was mainly led by two developing theories: Quantum Physics and Relativistic Physics. Albert Einstein was credited with the creation of the latter field with his papers on special and general relativity. General relativity unifies special relativity and the laws of universal gravitation by examining gravity as an intrinsically geometric property of spacetime. Particularly, it examines how energy, mass, momentum, and stress of various systems affect the curvature of spacetime. We understand this relationship through Einstein's Field Equations:

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R + \Lambda g_{\mu\nu} = 8\pi G T_{\mu\nu}$$

In this paper, we will derive this equation from fundamental physical principles and results found in special relativity. Along the way, we will examine the consequences of the equation and the effects of general relativity as we consider the universe.

1 Where We Start: First Principles

1.1 Special Relativity and Flat Space Time

Einstein's journey toward a theory of general relativity began with his work in special relativity, which unified space and time into a single geometric object known as spacetime. It was founded on the postulate that the speed of light in a vacuum is the same for all inertial observers (in a vacuum), and what was observed was that light became the cosmic speed limit. In special relativity, an event is labeled by four coordinates $x^\mu = (t, x, y, z)$ and the invariant interval between local events is given by the Minkowski metric:

$$ds^2 = \eta_{\mu\nu}dx^\mu dx^\nu = -dt^2 + dx^2 + dy^2 + dz^2$$

The key feature of this interval is that it remains invariant under Lorentz transformations (e.g., rotations or velocity boosts), which relate inertial reference frames moving at constant velocities relative to one another. The invariance of this line element reflects our postulates of the speed of light and the geometric structure of flat spacetime.

For a free massive particle, proper time is defined through $d\tau^2 = -ds^2$. Thus, we can find the action of a free particle by using the principle of stationary action and extremizing the action, yielding the equation of motion:

$$S = -m\int d\tau \implies \frac{d^2x^\mu}{d\tau^2} = 0$$

This equation of motion describes straight-line motion through flat spacetime. Thus, in special relativity, freely moving particles must follow straight worldlines. However, special relativity applies only to inertial frames, and it cannot, as is, describe accelerating observers or gravitational effects. It can, however, be naturally lifted to do so. In fact, the foundation of all of general relativity was that gravity itself might emerge from considering noninertial coordinate systems and curved spacetime geometry.

1.2 Spacetime Becomes Curved

One of the most important considerations guiding Einstein toward general relativity was the universality of gravitational acceleration. In Newtonian mechanics, inertial mass and gravitational mass are conceptually distinct quantities:

$$F = m\vec{a} \iff F = \frac{GMm}{r^2}$$

While m represents the same measurable value, they are treated in two distinct circumstances. Experimentally, all objects ($M_{grav} >>> m_{fall}$) fall with the same acceleration regardless of mass or composition. Einstein recognized that this equality was a significant result. Suppose one observer sees a free particle moving inertially (y), and another observer uses accelerated coordinates $y^\rho = y^\rho(x)$. The particle in respective coordinates becomes:

$$\frac{d^2y^\rho}{d\tau^2} = 0 \quad \longmapsto \quad \frac{d^2x^\lambda}{d\tau^2} + (\frac{\partial x^\lambda}{\partial y^\rho}\frac{\partial^2 y^\rho}{\partial x^\mu \partial x^\nu})\frac{dx^\mu}{d\tau}\frac{dx^\nu}{d\tau} = \frac{d^2x^\lambda}{d\tau^2} + \Gamma^\lambda_{\mu\nu}\frac{dx^\mu}{d\tau}\frac{dx^\nu}{d\tau} = 0$$

These terms resemble gravitational accelerations, even though they arise purely from differential geometry and coordinate choice. Einstein, thus, concluded that gravity is not a fundamental force in the same vein as electromagnetism. This equation should also look familiar, because it is precisely the geodesic formula. Instead of objects freely falling, they simply follow the s̈hortest" paths (geodesics) through curved spacetime. It is here that we can begin postulating on the conclusion of general relativity: Gravity emerges from the curvature of spacetime.

1.3 The Equivalence Principle

Einstein's equivalence principle simply states that locally, a gravitational field is indistinguishable from some general acceleration. This principle can be illustrated through thought experiments involving rocket science. It shouldn't be hard to imagine, after all, it isn't actually rocket science.

Consider an observer inside a rocket accelerating upward in empty space (no external gravitational field). A beam of light fired horizontally appears to bend downward (from the perspective of an external viewer) because the floor accelerates upward while the light travels. Now compare this with an observer standing in a gravitational field. A horizontally traveling light beam also bends downward. Einstein reasoned that the two situations are experimentally indistinguishable and, as such, concluded that gravity and acceleration are locally equivalent.

The equivalence principle also predicts gravitational redshift. Suppose light travels upward through a gravitational field. Since clocks deeper in a gravitational well run slower, the frequency of the emitted light differs from the received frequency. The fractional frequency shift is approximately:

$$\frac{\omega_{reciever} - \omega_{emmiter}}{\omega_{emmiter}} = \frac{\Phi_{reciever} - \Phi_{emmiter}}{c^2}$$

Where Φ describes the Newtonian gravitational potential. Gravity, therefore, not only affects motion, but the flow of time as well (since spacetime is a unified manifold). This gravitational redshift means clocks in stronger gravitational fields tick more slowly than those in weaker gravitational fields. This has been experimentally verified many times over through precision measurement involving atomic clocks and gravitational redshifts.

1.4 The Universe and Cosmology in Curved Spacetime

Einstein generalized flat spacetime completely by replacing the Minkowski metric $\eta_{\mu\nu}$ with the general spacetime metric $g_{\mu\nu}(x)$:

$$ds^2 = \eta_{\mu\nu}dx^\mu dx^\nu \quad \longmapsto \quad ds^2 = g_{\mu\nu}(x)dx^\mu dx^\nu$$

The metric determines all geometric information about spacetime, including distances, times, causal structure, and trajectories (this will be evident in our final equation). Unlike special relativity, where

spacetime is globally flat, our general relativity metric is a function of local coordinates, thereby encoding gravitational effects geometrically.

1.4.1 What Happens to Distances?

In a general curved spacetime, defining spatial distance requires a hint more nuance than for flat Euclidean (or even Minkowskian) geometry because the metric itself determines how distance is defined on a point-by-point basis. An operational definition of spatial distance comes from the idea of radar communication, where an observer sends light pulses to a distant object and measures the round-trip travel time (since, as we will see, light follows null trajectories). Thus, the proper spatial distance between local points is purely the proper spatial line differential between nearby points at a fixed coordinate time component:

$$dl^2 = (g_{ij} - \frac{g_{0i}g_{0j}}{g_{00}})dx^i dx^j$$

And if we wanted to determine larger distances, say across galaxies, we must integrate this line differential with respect to a proper time differential along a given contour.

1.4.2 Cons to Having a Long Distance Relationship in an Expanding Universe

If we assume, as we always tend to do, that cosmological space is homogeneous and isotropic on large scales, we obtain a metric of the following form: $ds^2 = -dt^2 + a(t)^2(d\vec{x})^2$, where $a(t)$ is a cosmological scale factor. In this spacetime, spatial distances evolve dynamically with time. Therefore, the corresponding proper distance differential becomes $dl^2 = a(t)^2(d\vec{x})^2$. This equation shows that the physical separation between comoving objects increases proportionally to the scale factor.

We have determined through experimental observations and theoretical fact-checks that in our recent dark-energy dominated history (the past 4-5 billion years), our cosmological factor is given by:

$$a(t) = e^{Ht} \implies ds^2 = -dt^2 + e^{2Ht}(d\vec{x})^2$$

An important consequence of exponential expansion is the existence of a de Sitter horizon. Because spacetime expands so rapidly, sufficiently distant regions recede faster than light can ever traverse the intervening space. Consequently, observers possess a cosmological event horizon beyond which no information can ever reach them, and this radius is halved if you want information to be returned as well. This is only possible with scale factors whose inverse have convergent integrals ($a(t) \sim t^p,\ p > 1$). This is good and bad news. Maybe the alien you met on that one universal dating app literally can't text you back.

To the smart alecks wryly raising their hands, thinking we broke special relativity ("I thought nothing can move faster than light?"), put your hands down. We didn't break special relativity since "faster than light" actually references the geometry of local light cones. While that is true in local light cones, in sufficiently rapidly expanding universes, spacetime expands faster than global light cones can grow.

Speaking of light cones, certain physicists like to preserve the structure of light cones at clean 45° angles. Thus, we can switch from standard time to conformal time η (not to be confused with the Minkowskian metric), which resembles flat Minkowski space up to an overall scale factor:

$$d\eta = \frac{dt}{a(t)} \implies ds^2 = a(\eta)^2(-d\eta^2 + d\vec{x}^2)$$

1.4.3 Redshift From an Expanding Universe

As spacetime expands, light traveling through the universe experiences cosmological redshift. The wavelength of light stretches together with spacetime itself, producing the relation

$$1 + z = \frac{a(t_{\text{obs}})}{a(t_{\text{emit}})}$$

Unlike ordinary Doppler redshift, cosmological redshift arises from the expansion of spacetime rather than motion through space.

1.4.4 Curvature of Space?

You might think that the section header is a typo, but it is a serious consideration in modern astrophysics. Georges LeMaitre posed the tedious, but valid, question: why can we naively assume space must be flat? Why can't space itself have intrinsic curvature? While we are comfortable assuming that space is flat, by readings from the Planck telescope, we aren't completely confident. The geometry of the universe could possess positive, zero, or negative (but constant by isotropy of space) spatial curvature. By incorporating spatial curvature, we quickly arrive at the Friedmann-Robertson-Walker metric:

$$ds^2 = -dt^2 + a(t)^2\left(\frac{dr^2}{1 - kr^2/L^2} + r^2 d\Omega^2\right),\ k = -1 \text{ (open)},\ 0 \text{ (flat)}, +1 \text{ (closed)}$$

These cases describe the hyperbolic, flat, and 3-spherical geometries. And we get the following distance functions:

$$k = -1 \implies d = a(t) L \sinh^{-1}\left(\frac{R}{L}\right);\ k = 0 \implies d = a(t)R;\ k = 1 \implies d = a(t) L \sin^{-1}\left(\frac{R}{L}\right)$$

1.5 Motion in Curved Spacetime

1.5.1 Massive Particles and Recovering the Newtonian Limit

In special relativity, free massive particles move along straight lines through flat Minkowski spacetime. Mathematically, their trajectories satisfy:

$$\frac{d^2 x^\mu}{d\tau^2} = 0,$$

Once spacetime becomes curved, the notion of a "straight line" must be generalized. In general relativity, free particles move along geodesics, which are the straightest possible paths in curved spacetime. Using the universal principle of stationary action, the trajectory of a massive particle is determined by extremizing the proper time, $S = -m \int d\tau$,

If we substitute the curved spacetime interval ($ds^2 = g_{\mu\nu} dx^\mu dx^\nu$) into our action and then apply the principles of variational calculus, we get that the massive particle must satisfy the geodesic equation:

$$\frac{d^2 x^\lambda}{d\tau^2} + \Gamma^\lambda_{\mu\nu} \frac{dx^\mu}{d\tau} \frac{dx^\nu}{d\tau} = 0$$

Note that the Christoffel symbols encode how coordinates vary across curved spacetime, and while they, themselves, are not tensors, they determine how vectors are parallel transported and how freely falling particles move. This equation demonstrates one of the central ideas of general relativity: gravity is not a conventional force. Instead, freely falling particles experience no force locally and simply follow geodesics determined by spacetime geometry.

To ensure we haven't lost the already established plot of physics, we should have a sanity check and recover motion in Newtonian gravity. To do so, we consider the weak-field approximation in which the metric differs only slightly from flat spacetime:

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}, \qquad |h_{\mu\nu}| \ll 1.$$

For weak gravitational fields and slowly moving particles, the dominant contribution comes from the g_{00} component of the metric ($g_{00} \approx -(1 + 2\Phi)$, where Φ is the Newtonian gravitational potential. Therefore, the geodesic equation gives us back cleanly that

$$\frac{d^2\vec{x}}{dt^2} = -\nabla\Phi$$

This is precisely Newton's law of gravitation, suggesting Newtonian gravity emerges naturally as the low-energy limit of general relativity.

1.5.2 Massless Particles and Relativistic Gravitational Redshifts

Light behaves fundamentally differently from massive particles because photons possess no rest mass. Consequently, photons do not experience proper time along their trajectories. Instead of timelike paths satisfying $ds^2 < 0$, light follows null geodesics satisfying

$$ds^2 = 0$$

Because proper time vanishes for light rays, photon trajectories cannot be parameterized using τ. Instead, one introduces an affine parameter ζ, which labels points along the light ray. Rather than boring you with another action integral, I will simply state that the null geodesic equation is

$$\frac{d^2x^\rho}{d\zeta^2} + \Gamma^\rho_{\mu\nu}\frac{dx^\mu}{d\zeta}\frac{dx^\nu}{d\zeta} = 0$$

Thus, light also follows geodesics determined by the spacetime metric. However, you should note the key difference is that these trajectories lie on the light cone defined by $ds^2 = 0$. An immediate consequence is gravitational lensing. Since spacetime curvature alters null geodesics, light rays bend when passing near massive objects. This effect was famously confirmed during the 1919 Eddington Experiment, which measured the deflection of starlight by the Sun's gravitational field, and it served as one of the earliest confirmations of the results of general relativity.

Now, for the third time, let us talk about Redshifts. We are no strangers to the fact that gravity affects the frequency of light (if you need more convincing, consider a photon climbing out of a gravitational potential well). Because clocks deeper in a gravitational field run more slowly, the emitted and received frequencies differ. If we redo our classic thought experiment of two observers. Emmett, who sends light signals, and Reverie, who receives them. Accounting for changes in observed time in curved space-time, we see:

$$\Delta\tau_R = \sqrt{-g_{00}(\vec{x}_R)}\Delta t_E = \Delta\tau_E(\sqrt{g_{00}(\vec{x}_R)}/\sqrt{g_{00}(\vec{x_E})} \implies \omega_R = \omega_E\sqrt{\frac{g_{00}(\vec{x}_E)}{g_{00}(\vec{x}_R)}}$$

Note, if we want to, we can recover the weak-field limit from section 1.3, by simply stating that $g_{00}(\vec{x}) = 1 - 2\Phi(\vec{x})$.

2 A Hop, Skip, and a Jump to Einstein's Field Equations

From our previous sections, we have already discovered a plethora of phenomena emerging from treating our kinematics as geodesics in curved spacetime. This is great, but we are heavily reliant on the promised existence of some fabled metric. We have the intuition that gravity is a manifestation of the curvature of spacetime, but we have to show precisely how. So, let us finally ask the question: How do we find $g_{\mu\nu}$?

2.1 What Do We Want?

As physicists, we have a privilege that our mathematician colleagues don't: guess and check. We don't need rigor or to show a formal chain of logic; all we need is a guess and a prayer. We can make a suite of assumptions, some less sound than others, but all that matters is that we get a working model. Therefore, let us abuse this privilege.

In ordinary field theory, actions usually look something like:

$$S = \int d^4x[T - V]$$

For scalar fields ϕ, the kinetic term is quadratic in derivatives: $(\partial_\mu\phi)(\partial^\mu\phi)$. For electromagnetic systems, the action is quadratic in field strength: $S_{EM} = -\frac{1}{4}\int d^4x F_{\mu\nu}F^{\mu\nu}$. Therefore, making an informed guess, we should expect gravitational field equations to arise from an action quadratic in field derivatives. Our field in this case? $g_{\mu\nu}$!

However, we run into a problem: $\partial_\lambda g_{\mu\nu}$ is not coordinate invariant. We can always choose locally inertial coordinates at a point p such that:

$$g_{\mu\nu}(p) = \eta_{\mu\nu}, \quad \partial_\lambda g_{\mu\nu} = 0$$

Thus, first derivatives can always be transformed away locally, and they do not contain true, generally covariant gravitational information. What we want is an action to be coordinate invariant.

$$S_{grav} = \int d^4x\sqrt{-g}(\text{ Scalar in Terms of } g_{\mu\nu}\ )$$

Note, the reason we must add the extra coefficient is that our differential itself isn't coordinate invariant. At this point, the reader may already be screaming at their screen: the action must depend on curvature. After all, this connects almost one-to-one to Riemann's search for a covariant description of geometry. However, if you are more skeptical, we will discuss why scalar curvature must be the only simple, nontrivial scalar we should consider.

2.2 Curvature!

The Christoffel symbols are given by:

$$\Gamma^\rho_{\mu\nu} = \frac{1}{2}g^{\rho\sigma}\left(\partial_\mu g_{\nu\sigma} + \partial_\nu g_{\mu\sigma} - \partial_\sigma g_{\mu\nu}\right)$$

They contain first derivatives of the metric, but they are not tensors. This means that they can always be made to vanish at a point through an appropriate coordinate choice. The true tensorial measure of curvature is obtained by examining how covariant derivatives fail to commute. That is:

$$[D_\mu, D_\nu]V^\rho = R^\rho{}_{\sigma\mu\nu}V^\sigma$$

Where we define the Riemann curvature tensor:

$$R^\rho{}_{\sigma\mu\nu} = \partial_\mu\Gamma^\rho_{\nu\sigma} - \partial_\nu\Gamma^\rho_{\mu\sigma} + \Gamma^\rho_{\mu\lambda}\Gamma^\lambda_{\nu\sigma} - \Gamma^\rho_{\nu\lambda}\Gamma^\lambda_{\mu\sigma}.$$

If we examine the structure of R (to see if it satisfies our action rule of thumb)

$$\Gamma \sim \partial g, \qquad R \sim \partial\Gamma + \Gamma\Gamma \sim \partial^2 g + (\partial g)^2$$

This has exactly the form expected for a field theory: action that is quadratic in derivatives. Further, Curvature is tensorial and cannot be transformed away, so it contains the true gravitational degrees of freedom. It must be the answer to our search.

2.2.1 Contracting Curvature

The Riemann tensor has four indices, but matter is described by the stress-energy tensor $T_{\mu\nu}$, a rank-2 tensor. Thus, we contract indices of the Riemann tensor:

Contracting gives the Ricci tensor:

$$R_{\mu\nu} = R^{\rho}{}_{\mu\rho\nu}.$$

Contracting once more gives the scalar curvature:

$$R = g^{\mu\nu} R_{\mu\nu}.$$

This scalar curvature is the simplest invariant scalar available for the gravitational action.

2.3 Lights! Camera! Action?

We therefore propose the Einstein–Hilbert action:

$$S_{\text{EH}} = \frac{1}{16\pi G} \int d^4x \sqrt{-g}\, R$$

The factor $1/(16\pi G)$ is chosen so that the weak-field limit reproduces Newtonian gravity:

$$\nabla^2 \Phi = 4\pi G \rho$$

This, however, is only one piece of the picture. The full action is

$$S = S_{\text{EH}} + S_m,$$

We include a term to consider the action of matter because spacetime is not empty. It contains particles, radiation, fluids, electromagnetic fields, etc. Therefore, their physics cannot be ignored, so we add the S_m term to the total action. The next section covers more about this form of action. To solve for the dynamics of this field, we must do what we always do: vary the action, but now we vary it with respect to the metric:

$$g_{\mu\nu} \mapsto g_{\mu\nu} + \delta g_{\mu\nu}, \quad \delta S = 0$$

2.4 The Stress Energy Tensor

The stress-energy tensor is defined through the variation of the matter action:

$$\delta S_{\text{matter}} = -\frac{1}{2} \int d^4x \sqrt{-g}\, T_{\mu\nu} \delta g^{\mu\nu} \implies T_{\mu\nu} = -\frac{2}{\sqrt{-g}} \frac{\delta S_{\text{matter}}}{\delta g^{\mu\nu}}$$

The stress-energy tensor measures how matter responds to changes in spacetime geometry. Its components have direct physical interpretations:

- T_{00} is the energy density
- T_{0i} is momentum density or energy flux
- T_{ij} contains pressure and stress

Thus, gravity is sourced not merely by mass, but by all forms of energy, momentum, pressure, and stress. This is the white lie that most people say when they discuss general relativity, mass is not the only craftsman of curved spacetime, but it truly takes a whole tensor.

2.5 Varying Einstein-Hilbert Action

Ending our brief tangent into understanding the Stress-Energy Tensor, let us now finally vary the action and see what shakes out of the equation. To vary the action, we use the identity:

$$R = g^{\mu\nu}R_{\mu\nu} \implies \delta(\sqrt{-g}R) = R\delta\sqrt{-g} + \sqrt{-g}R_{\mu\nu}\delta g^{\mu\nu} + \sqrt{-g}g^{\mu\nu}\delta R_{\mu\nu}$$

Using the Palantini identity and the following identity:

$$\delta\sqrt{-g} = -\frac{1}{2}\sqrt{-g}\, g_{\mu\nu}\delta g^{\mu\nu}$$

We can observe that the boundary terms vanish, leaving

$$\delta S_{EH} = \frac{1}{16\pi G}\int d^4x\sqrt{-g}\left(R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R\right)\delta g^{\mu\nu}$$

This motivates the definition of the Einstein tensor:

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R$$

Thus,

$$\delta S_{EH} = \frac{1}{16\pi G}\int d^4x\sqrt{-g}\, G_{\mu\nu}\delta g^{\mu\nu}$$

The Einstein tensor satisfies the contracted Bianchi identity:

$$D_\mu G^{\mu\nu} = 0$$

The Bianchi identity is also consistent with local energy conservation, which bodes especially well for our next step:

$$D_\mu T^{\mu\nu} = 0$$

2.6 Einstein's Field Equations (Almost)

The total variation becomes

$$\delta S = \delta S_{EH} + \delta S_{matter}$$

Substituting both variations gives

$$\delta S = \int d^4x\sqrt{-g}\left[\frac{1}{16\pi G}G_{\mu\nu} - \frac{1}{2}T_{\mu\nu}\right]\delta g^{\mu\nu}$$

Since $\delta g^{\mu\nu}$ is arbitrary, what remains in our brackets must be constantly zero:

$$\frac{1}{16\pi G}G_{\mu\nu} - \frac{1}{2}T_{\mu\nu} = 0$$

Therefore, we are left with (almost) the full Einstein Field Equations

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = 8\pi G T_{\mu\nu}$$

Before we continue lets take a breath and see what we actually derived. The left-hand side describes spacetime curvature, being a function of our curvature and spacetime metric. The right-hand side describes matter and energy with the Stress-Energy Tensor. So, our (or rather Einstein's) intuition rings true.

2.7 The Cosmological Constant

If you can't tell by the not-so-subtle hints in the section prior, we are missing something. Usually, every class has that one smart aleck, who, at this point, would be at the edge of their seat, waving their hand in the air and feeling the urge to correct the lecturer. Since I know you, reader, are too kind to correct me, I will have to be both the lecturer and the annoying student.

"Hey! Constant vacuum energy can't be ignored because spacetime volume depends on the metric:"

$$d^4x\sqrt{-g}$$

As much as I hate to give my kudos to that guy, they are right. We can't simply forget the ambient vacuum energy of a spacetime. Thus, vacuum energy contributes to gravity and to our field equations. Therefore, the most general covariant gravitational action becomes:

$$S = \frac{1}{16\pi G}\int d^4x\sqrt{-g}(R - 2\Lambda)$$

(The two-factor gets added to make the final form more eye-catching). The cosmological constant acts like an intrinsic vacuum energy density of spacetime. Varying the cosmological constant term produces an additional contribution:

$$+\Lambda g_{\mu\nu}$$

Thus, we find the completed, full field equations:

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R + \Lambda g_{\mu\nu} = 8\pi G T_{\mu\nu}$$

$\therefore$ ■ Q.E.D

3 Where Do We Go Now?

The Einstein field equations transform spacetime from a passive backdrop in physics into a dynamical, physical object, reframing gravity as a study of geometry rather than as a study of forces acting in fixed space and time. While it, like most famous physical equations, looks beautifully simple, it is relatively weak, not in terms of significance but rather in solving potential. We can really only solve for spacetime metrics of highly symmetrized systems. Nonetheless, those cases, the Schwarzschild, Kerr, and Reissner–Nordström metrics, are interesting to examine because they examine the physics behind black holes, event horizons, and relativistic astrophysical objects.

Applying the equations to homogeneous and isotropic universes yields the Friedmann equations, forming the foundation of modern cosmology and explaining cosmic expansion, the Big Bang, the cosmic microwave background, inflation, and dark-energy-driven acceleration. Moreover, small perturbations of the metric ($g'_{\mu\nu} = g_{\mu\nu} + \epsilon h_{\mu\nu}$) pave the way into the study of gravitational-wave and signature observations of supermassive objects.

My favorite consideration, and hopefully someday future research, is studying how interfacing curved spacetime and quantum fields leads to questions of quantum gravity, dark energy, and grand unification of general relativity and quantum field theory. Hopefully, this was as elucidating a journey for you as it was for me, and I hope you enjoyed reading. Thank you.

References

[1] Zee, A. Einstein Gravity in a Nutshell. Princeton University Press, 2013.

[2] "Einstein Field Equations." Wikipedia, Wikimedia Foundation, 25 Apr. 2026, en.wikipedia.org/wiki/Einstein$_{f}$ield$_{e}$quations.

[2] "General Relativity." Wikipedia, Wikimedia Foundation, 14 May 2026, en.wikipedia.org/wiki/General$_{r}$elativity.

[2] Carroll, Bradley W., and Dale A. Ostlie. An Introduction to Modern Astrophysics. Cambridge University Press, 2024.

[3] Kenyon, I. R. Introduction to General Relativity and Cosmology. IOP Publishing, 2023.

[4] YouTube, YouTube, www.youtube.com/watch?v=XRr1kaXKBsUt=366s. Accessed 15 May 2026.

[5] YouTube, YouTube, www.youtube.com/watch?v=xodtfM1r9FA. Accessed 15 May 2026.

[6] Cao, Xiaodong, Cornell University Math 4540 Notes