

Mathematics I (BSM 101)

Manju Subedi

Gandaki University
Bachelor in Information Technology(BIT)
BSM 101

manjusubedi.061@gmail.com

January 4, 2023

Extreme Value Function:


The Max-Min Theorem for Continuous Functions:

If f is continuous at every point of a closed interval I , then f assumes both an absolute maximum value M and an absolute minimum value m somewhere in I . That is, there are numbers x_1 and x_2 in I with $f(x_1) = m$, $f(x_2) = M$, and $m \leq f(x) \leq M$ for every other x in I


Extreme Value Function:

The Max-Min Theorem for Continuous Functions:


If f is continuous at every point of a closed interval I , then f assumes both an absolute maximum value M and an absolute minimum value m somewhere in I . That is, there are numbers x_1 and x_2 in I with $f(x_1) = m$, $f(x_2) = M$, and $m \leq f(x) \leq M$ for every other x in I




Maximum and minimum
at interior points



Maximum and minimum
at endpoints



Maximum at interior point,
minimum at endpoint



Minimum at interior point,
maximum at endpoint

Absolute Extreme Values

Let f be a function with domain D . Then f has an **absolute maximum** value on D at a point c if

$$f(x) \leq f(c) \quad \text{for all } x \text{ in } D$$

Absolute Extreme Values

Let f be a function with domain D . Then f has an **absolute maximum** value on D at a point c if

$$f(x) \leq f(c) \quad \text{for all } x \text{ in } D$$

and an **absolute minimum** value on D at c if

$$f(x) \geq f(c) \quad \text{for all } x \text{ in } D.$$

Absolute maximum and minimum values are called absolute extrema. Absolute extrema are also called global extrema.

Local Extreme Values:

A function f has a **local maximum** value at an interior point c of its domain if $f(x) \leq f(c)$ for all x in some open interval containing c .


Local Extreme Values:

A function f has a **local maximum** value at an interior point c of its domain if $f(x) \leq f(c)$ for all x in some open interval containing c .

A function f has a **local minimum** value at an interior point c of its domain if $f(x) \geq f(c)$ for all x in some open interval containing c .

Local Extreme Values:

A function f has a **local maximum** value at an interior point c of its domain if $f(x) \leq f(c)$ for all x in some open interval containing c .
A function f has a **local minimum** value at an interior point c of its domain if $f(x) \geq f(c)$ for all x in some open interval containing c .



The First Derivative Theorem for Local Extreme Values:

If f has a local maximum or minimum value at an interior point c of its domain, and if f' is defined at c , then

$$f'(c) = 0.$$

Finding Extrema

The First Derivative Theorem for Local Extreme Values:

If f has a local maximum or minimum value at an interior point c of its domain, and if f' is defined at c , then

$$f'(c) = 0.$$

Critical Point: An interior point of the domain of a function f where f' is zero or undefined is a critical point of f .

Finding Extrema

The First Derivative Theorem for Local Extreme Values:

If f has a local maximum or minimum value at an interior point c of its domain, and if f' is defined at c , then

$$f'(c) = 0.$$

Critical Point: An interior point of the domain of a function f where f' is zero or undefined is a critical point of f .

Note: Maximum and minimum of the function are occurs only at critical points.

Finding Extrema

The First Derivative Theorem for Local Extreme Values:

If f has a local maximum or minimum value at an interior point c of its domain, and if f' is defined at c , then

$$f'(c) = 0.$$

Critical Point: An interior point of the domain of a function f where f' is zero or undefined is a critical point of f .

Note: Maximum and minimum of the function are occurs only at critical points.

Finding Absolute Extrema: In any Continuous Function f on a Closed Interval

1. Find all critical points by setting $f'(x) = 0$
2. Evaluate f at all critical points and endpoints.
3. Take the largest and smallest of these values.

Finding Extrema

The First Derivative Theorem for Local Extreme Values:

If f has a local maximum or minimum value at an interior point c of its domain, and if f' is defined at c , then

$$f'(c) = 0.$$

Critical Point: An interior point of the domain of a function f where f' is zero or undefined is a critical point of f .

Note: Maximum and minimum of the function are occurs only at critical points.

Finding Absolute Extrema: In any Continuous Function f on a Closed Interval

1. Find all critical points by setting $f'(x) = 0$
2. Evaluate f at all critical points and endpoints.
3. Take the largest and smallest of these values.

Example 1: Find the absolute maximum and minimum values of $f(x) = x^2$ on $[-2, 1]$.

Increasing and Decreasing Functions

Definition: Let f be a function defined on an interval I and let x_1 and x_2 be any two points in I .

1. f increases on I if $x_1 < x_2 \Rightarrow f(x_1) < f(x_2)$.
2. f decreases on I if $x_1 < x_2 \Rightarrow f(x_1) > f(x_2)$.

Increasing and Decreasing Functions


Definition: Let f be a function defined on an interval I and let x_1 and x_2 be any two points in I .

1. f increases on I if $x_1 < x_2 \Rightarrow f(x_1) < f(x_2)$.
2. f decreases on I if $x_1 < x_2 \Rightarrow f(x_1) > f(x_2)$.

The First Derivative Test for Increasing and Decreasing: Suppose that f is continuous on $[a, b]$ and differentiable on (a, b) .

If $f'(x) > 0$ at each point of (a, b) , then f increases on $[a, b]$.

If $f'(x) < 0$ at each point of (a, b) , then f decreases on $[a, b]$.



Intervals of Increasing or Decreasing

Let f be a differentiable function on an interval I .

To find intervals on which f is increasing and decreasing:

- 1 Find the critical values of f . That is, find all c in I where $f'(c) = 0$ or f' is not defined.

Intervals of Increasing or Decreasing

Let f be a differentiable function on an interval I .

To find intervals on which f is increasing and decreasing:

- 1 Find the critical values of f . That is, find all c in I where $f'(c) = 0$ or f' is not defined.
- 2 Use the critical values to divide I into subintervals.

Intervals of Increasing or Decreasing

Let f be a differentiable function on an interval I .

To find intervals on which f is increasing and decreasing:

- 1 Find the critical values of f . That is, find all c in I where $f'(c) = 0$ or f' is not defined.
- 2 Use the critical values to divide I into subintervals.
- 3 Pick any point p in each subinterval, and find the sign of $f'(p)$.

Intervals of Increasing or Decreasing

Let f be a differentiable function on an interval I .

To find intervals on which f is increasing and decreasing:

- ① Find the critical values of f . That is, find all c in I where $f'(c) = 0$ or f' is not defined.
- ② Use the critical values to divide I into subintervals.
- ③ Pick any point p in each subinterval, and find the sign of $f'(p)$.
 - a. If $f'(p) > 0$, then f is increasing on that subinterval.

Intervals of Increasing or Decreasing

Let f be a differentiable function on an interval I .

To find intervals on which f is increasing and decreasing:

- ① Find the critical values of f . That is, find all c in I where $f'(c) = 0$ or f' is not defined.
- ② Use the critical values to divide I into subintervals.
- ③ Pick any point p in each subinterval, and find the sign of $f'(p)$.
 - a. If $f'(p) > 0$, then f is increasing on that subinterval.
 - b. If $f'(p) < 0$, then f is decreasing on that subinterval.

Intervals of Increasing or Decreasing

Let f be a differentiable function on an interval I .

To find intervals on which f is increasing and decreasing:

- ① Find the critical values of f . That is, find all c in I where $f'(c) = 0$ or f' is not defined.
- ② Use the critical values to divide I into subintervals.
- ③ Pick any point p in each subinterval, and find the sign of $f'(p)$.
 - a. If $f'(p) > 0$, then f is increasing on that subinterval.
 - b. If $f'(p) < 0$, then f is decreasing on that subinterval.

Example:2 Let $f(x) = x^3 + x^2 - x + 1$. Find intervals on which f is increasing or decreasing.

Finding Local Extrema(Maximum/Minimum)

First Derivative Test: Let f be a function defined on an open interval I . Let f be continuous at a critical point c in I . Then

- 1 If $f'(x)$ changes sign from positive to negative as x increases through c , i.e., if $f'(x) > 0$ at every point sufficiently close to and to the left of c , and $f'(x) < 0$ at every point sufficiently close to and to the right of c , then c is a point of local maxima.

Finding Local Extrema(Maximum/Minimum)

First Derivative Test: Let f be a function defined on an open interval I . Let f be continuous at a critical point c in I . Then

- 1 If $f'(x)$ changes sign from positive to negative as x increases through c , i.e., if $f'(x) > 0$ at every point sufficiently close to and to the left of c , and $f'(x) < 0$ at every point sufficiently close to and to the right of c , then c is a point of local maxima.
- 2 If $f'(x)$ changes sign from negative to positive as x increases through c , i.e., if $f'(x) < 0$ at every point sufficiently close to and to the left of c , and $f'(x) > 0$ at every point sufficiently close to and to the right of c , then c is a point of local minima.

Finding Local Extrema(Maximum/Minimum)

First Derivative Test: Let f be a function defined on an open interval I . Let f be continuous at a critical point c in I . Then

- 1 If $f'(x)$ changes sign from positive to negative as x increases through c , i.e., if $f'(x) > 0$ at every point sufficiently close to and to the left of c , and $f'(x) < 0$ at every point sufficiently close to and to the right of c , then c is a point of local maxima.
- 2 If $f'(x)$ changes sign from negative to positive as x increases through c , i.e., if $f'(x) < 0$ at every point sufficiently close to and to the left of c , and $f'(x) > 0$ at every point sufficiently close to and to the right of c , then c is a point of local minima.
- 3 If $f'(x)$ does not change sign as x increases through c , then c is neither a point of local maxima nor a point of local minima. Infact, such a point is called point of inflection.

Finding Local Extrema(Maximum/Minimum)



First Derivative Test: Let f be a function defined on an open interval I . Let f be continuous at a critical point c in I . Then

- 1 If $f'(x)$ changes sign from positive to negative as x increases through c , i.e., if $f'(x) > 0$ at every point sufficiently close to and to the left of c , and $f'(x) < 0$ at every point sufficiently close to and to the right of c , then c is a point of local maxima.
- 2 If $f'(x)$ changes sign from negative to positive as x increases through c , i.e., if $f'(x) < 0$ at every point sufficiently close to and to the left of c , and $f'(x) > 0$ at every point sufficiently close to and to the right of c , then c is a point of local minima.
- 3 If $f'(x)$ does not change sign as x increases through c , then c is neither a point of local maxima nor a point of local minima. Infact, such a point is called point of inflection.

Example 3: Find all points of local maxima and local minima of the function f given by

$$f(x) = x^3 - 3x + 3$$

Finding Local Extrema(Maximum/Minimum)



Finding Local Extrema(Maximum/Minimum)

Second Derivative Test: Let f be a function defined on an interval I and $c \in I$. Let f be twice differentiable at c . Then

- If $f'(c) = 0$ and $f''(c) < 0$, then f has a local maximum at $x = c$.
- If $f'(c) = 0$ and $f''(c) > 0$, then f has a local minimum at $x = c$.
- The test fails if $f'(c) = 0$ and $f''(c) = 0$.

Finding Local Extrema(Maximum/Minimum)

Second Derivative Test: Let f be a function defined on an interval I and $c \in I$. Let f be twice differentiable at c . Then

- If $f'(c) = 0$ and $f''(c) < 0$, then f has a local maximum at $x = c$.
- If $f'(c) = 0$ and $f''(c) > 0$, then f has a local minimum at $x = c$.
- The test fails if $f'(c) = 0$ and $f''(c) = 0$.

Example 4: Find local maximum and local minimum values of the function f given by

$$f(x) = 3x^4 + 4x^3 - 12x^2 + 12$$

Example 5: Find all the points of local maxima and local minima of the function f given by

$$f(x) = 2x^3 - 6x^2 + 6x + 5$$

Thank you

