

Ch2. Gaussian Processes (Part 1)

Bayesian Optimization Seminar

Sungwoo Park

Uncertainty Quantification Lab
Seoul National University

January 22, 2026

1. Review of Chapter 1: Introduction
2. Definition and Basic Properties
3. Inference with Exact and Noisy Observations
4. Joint Gaussian Processes
5. Summary

1. Review of Chapter 1: Introduction
2. Definition and Basic Properties
3. Inference with Exact and Noisy Observations
4. Joint Gaussian Processes
5. Summary

The Optimization Problem

Goal

Find the global optimum of an objective function $f : \mathcal{X} \rightarrow \mathbb{R}$:

$$x^* = \arg \max_{x \in \mathcal{X}} f(x); \quad f^* = \max_{x \in \mathcal{X}} f(x)$$

Setting:

- Objective function f is **expensive** to evaluate (time, cost, resources)
- We can only access f through **sequential observations**
- Observations may be **noisy**: $y = f(x) + \varepsilon$

Challenge: How do we decide where to observe next, given limited budget?

Observation Model

Observations are realized by a stochastic mechanism:

$$p(y \mid x, \phi), \quad \text{where } \phi = f(x)$$

Common model — Additive Gaussian noise:

$$y = \phi + \varepsilon, \quad \varepsilon \sim \mathcal{N}(0, \sigma_n^2)$$

$$\Rightarrow p(y \mid x, \phi, \sigma_n) = \mathcal{N}(y; \phi, \sigma_n^2)$$

Assumption: Multiple observations are **conditionally independent** given the objective function values:

$$p(\mathbf{y} \mid \mathbf{x}, \phi) = \prod_i p(y_i \mid x_i, \phi_i)$$

The Bayesian Approach: Key Idea

Core Principle

Treat the unknown objective function f as a **random variable** and use **Bayesian inference** to reason about it.

Bayesian Inference Refresher:

1. Start with a **prior** $p(\phi \mid x)$ encoding initial beliefs
2. Observe data $\mathcal{D} = (x, y)$ via the **likelihood** $p(y \mid x, \phi)$
3. Update to the **posterior** via Bayes' rule:

$$p(\phi \mid \mathcal{D}) = \frac{p(y \mid x, \phi) p(\phi \mid x)}{p(y \mid x)}$$

The posterior captures what we now believe about ϕ after seeing the data.

Inference of the Objective Function

To reason about the *entire* objective function $f : \mathcal{X} \rightarrow \mathbb{R}$, we need a **stochastic process** — a probability distribution over functions.

Specifying a Stochastic Process

We specify the distribution of function values $\phi = f(\mathbf{x})$ for any finite set of locations $\mathbf{x} \subset \mathcal{X}$:

$$p(\phi \mid \mathbf{x})$$

Gaussian Processes: The family where all such finite-dimensional distributions are **multivariate Gaussian** — mathematically convenient and widely used in Bayesian optimization.

Posterior Predictive Distribution

After observing data \mathcal{D} , we can predict the outcome of a new observation at x :

Posterior Predictive Distribution

$$p(y' | x, \mathcal{D}) = \int p(y' | x, \phi) p(\phi | x, \mathcal{D}) d\phi$$

Interpretation:

- Integrates over all possible values of $\phi = f(x)$
- Weights by their plausibility under the posterior
- Naturally accounts for **uncertainty** in the objective function

This distribution is *instrumental* for making informed decisions about where to observe next.

Contents

1. Review of Chapter 1: Introduction
- 2. Definition and Basic Properties**
3. Inference with Exact and Noisy Observations
4. Joint Gaussian Processes
5. Summary

What is a Gaussian Process?

A **Gaussian process (GP)** extends the multivariate normal distribution to model functions on infinite domains.

Key Idea

We model an objective function $f : \mathcal{X} \rightarrow \mathbb{R}$ as an infinite collection of random variables, one for each point in the domain. The **Kolmogorov extension theorem** allows us to specify this distribution through finite-dimensional marginals.

GPs inherit convenient mathematical properties of the multivariate normal distribution while remaining computationally tractable.

Recall: Kolmogorov's Extension Theorem

Question

What is the *consistency* property in Kolmogorov's Extension Theorem?

Kolmogorov's Extension Theorem [Durrett, 2019]

Suppose we are given probability measures μ_n on $(\mathbb{R}^n, \mathcal{B}(\mathbb{R}^n))$ that are *consistent*, that is,

$$\mu_{n+1}((a_1, b_1] \times \cdots \times (a_{n+1}, b_{n+1}] \times \mathbf{R}) = \mu_n((a_1, b_1] \times \cdots \times (a_n, b_n])$$

Then there is a unique probability measure P on $(\mathbb{R}^{\mathbf{N}}, \mathcal{B}(\mathbb{R}^{\mathbf{N}}))$ with

$$P(\omega : \omega_i \in (a_i, b_i], 1 \leq i \leq n) = \mu_n((a_1, b_1] \times \cdots \times (a_n, b_n])$$

where $\mathbf{N} = \{1, 2, \dots\}$ and $\mathcal{B}(\mathbb{R}^{\mathbf{N}}) = \{(\omega_1, \omega_2, \dots) : \omega_i \in \mathcal{B}(\mathbb{R})\}$

In [Durrett, 2019], theorem is not complete to use in our senario because P defines on countable index set \mathbf{N} .

Q&A: Countable vs. Uncountable Domains

Question

What are the characteristics of modeling on countable vs uncountable domains?

Definition: Sample path [Le Gall, 2016]

Let $(X_t)_{t \in \mathcal{X}}$ be a random process with values in E . The *sample paths* of X are the mappings $\mathcal{X} \ni t \mapsto X_t(\omega)$ obtained when fixing $\omega \in \Omega$. The sample paths of X is thus form a collection of mappings from \mathcal{X} into E indexed by $\omega \in \Omega$.

Countable infinite domain (e.g., $\mathcal{X} = \mathbb{Z}$):

- GP reduces to specifying consistent MVN distributions on all finite subsets
- Sample paths are sequences $\{f(x_i)\}_{i=1}^{\infty}$
- No notion of “continuity” in the classical sense

Q&A: Countable vs. Uncountable Domains

Question

What are the characteristics of modeling on countable vs uncountable domains?

Definition: Sample path [Le Gall, 2016]

Let $(X_t)_{t \in \mathcal{X}}$ be a random process with values in E . The *sample paths* of X are the mappings $\mathcal{X} \ni t \mapsto X_t(\omega)$ obtained when fixing $\omega \in \Omega$. The sample paths of X is thus form a collection of mappings from \mathcal{X} into E indexed by $\omega \in \Omega$.

Uncountable infinite domain (e.g., $\mathcal{X} = \mathbb{R}^d$):

- Sample paths are actual functions
- Continuity, differentiability become meaningful properties
- Requires careful treatment (measurability issues, sample path properties)

Kolmogorov Extension Theorem: The Foundation

Question

What is the *consistency* property in Kolmogorov's Extension Theorem?

Kolmogorov's Extension Theorem : Stochastic process ver. [Oksendal,]

For all $t_1, \dots, t_k \in T$, $k \in \mathbb{N}$ let ν_{t_1, \dots, t_k} be probability measures on \mathbb{R}^{nk} s.t.

$$\nu_{t_{\sigma(1)}, \dots, t_{\sigma(k)}}(F_1 \times \dots \times F_k) = \nu_{t_1, \dots, t_k}(F_{\sigma^{-1}(1)} \times \dots \times F_{\sigma^{-1}(k)}) \quad (\text{K1})$$

for all permutations σ on $\{1, 2, \dots, k\}$ and

$$\nu_{t_1, \dots, t_k}(F_1 \times \dots \times F_k) = \nu_{t_1, \dots, t_k, t_{k+1}, \dots, t_{k+m}}(F_1 \times \dots \times F_k \times \mathbb{R}^n \times \dots \times \mathbb{R}^n) \quad (\text{K2})$$

for all $m \in \mathbb{N}$.

Kolmogorov Extension Theorem: The Foundation

Kolmogorov's Extension Theorem : Stochastic process ver. (cont.) [Oksendal,]

Then there exists a probability space (Ω, \mathcal{F}, P) and a stochastic process $\{X_t\}$ on Ω , $X_t : \Omega \rightarrow \mathbb{R}^n$, s.t.

$$\nu_{t_1, \dots, t_k}(F_1 \times \dots \times F_k) = P[X_{t_1} \in F_1, \dots, X_{t_k} \in F_k],$$

for all $t_i \in T$, $k \in \mathbb{N}$ and all Borel sets F_i .

Consistency Conditions:

1. **Permutation invariance** (K1): The joint distribution is unchanged by reordering indices
2. **Marginalization consistency** (K2): If $\mathbf{x} \subset \mathbf{x}'$, then marginalizing $p(\phi' \mid \mathbf{x}')$ over $\phi' \setminus \phi$ yields $p(\phi \mid \mathbf{x})$

For GPs, these are automatically satisfied because the MVN satisfies them (the marginal of a MVN is MVN with the corresponding submatrix of the covariance).

GP Specification: Mean and Covariance Functions

A GP on f is specified by:

$$p(f) = \mathcal{GP}(f; \mu, K)$$

- **Mean function** $\mu : \mathcal{X} \rightarrow \mathbb{R}$: determines the expected function value

$$\mu(x) = \mathbb{E}[\phi \mid x], \quad \text{where } \phi = f(x)$$

- **Covariance function (kernel)** $K : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$: encodes the correlation structure

$$K(x, x') = \text{cov}[\phi, \phi' \mid x, x'], \quad \text{where } \phi' = f(x')$$

The covariance function must be **symmetric** and **positive semidefinite**.

Q&A: Extension from Countable Dense Subsets

Question

If f is a.s. continuous and $D \subset \mathcal{X}$ is countable dense, can $p(f \mid D)$ determine $p(f)$?

Proof.

Let f and g be two continuous functions on \mathcal{X} . Suppose they agree on a dense subset $D \subseteq \mathcal{X}$. For any $x \in \mathcal{X}$, there exists a sequence $\{q_n\} \subset D$ such that $q_n \rightarrow x$. By continuity,

$$f(x) = f\left(\lim_{n \rightarrow \infty} q_n\right) = \lim_{n \rightarrow \infty} f(q_n) = \lim_{n \rightarrow \infty} g(q_n) = g\left(\lim_{n \rightarrow \infty} q_n\right) = g(x)$$

Thus, $f(x) = g(x)$ for all $x \in \mathcal{X}$. Since sample paths are a.s. continuous, the process is uniquely determined by its values on D . □

Finite-Dimensional Marginals

For any finite set of points $\mathbf{x} \subset \mathcal{X}$, the corresponding function values $\phi = f(\mathbf{x})$ follow a multivariate normal distribution:

$$p(\phi \mid \mathbf{x}) = \mathcal{N}(\phi; \boldsymbol{\mu}, \boldsymbol{\Sigma})$$

where

$$\boldsymbol{\mu} = \mathbb{E}[\phi \mid \mathbf{x}] = \mu(\mathbf{x}); \quad \boldsymbol{\Sigma} = \text{cov}[\phi \mid \mathbf{x}] = K(\mathbf{x}, \mathbf{x})$$

Gram Matrix

$K(\mathbf{x}, \mathbf{x})$ is the matrix formed by evaluating K for each pair of points:

$$\Sigma_{ij} = K(x_i, x_j)$$

Example: Squared Exponential Covariance




Figure 1: Example of Gaussian Process [Garnett, 2023]

Consider $\mathcal{X} = [0, 30]$ with:

- Mean function: $\mu \equiv 0$ (constant central tendency)
- Covariance function (squared exponential):

$$K(x, x') = \exp\left(-\frac{1}{2}|x - x'|^2\right)$$

Example: Squared Exponential Covariance




Figure 1: Example of Gaussian Process [Garnett, 2023]

Properties:

- $\text{var}[\phi \mid x] = K(x, x) = 1$ at every point
- Correlation decreases with distance: nearby values are highly correlated, distant values are nearly independent
- This encodes a statistical notion of **continuity**

Sampling from a Gaussian Process (Appendix A.2)

To sample from a GP with mean μ and covariance K :

1. Choose a finite grid of points $\mathbf{x} = (x_1, \dots, x_n)$
2. Compute $\boldsymbol{\mu} = \mu(\mathbf{x})$ and $\boldsymbol{\Sigma} = K(\mathbf{x}, \mathbf{x})$
3. Factor: $\boldsymbol{\Sigma} = \mathbf{L}\mathbf{L}^\top$ (Cholesky decomposition)
4. Sample $\mathbf{z} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$
5. Compute $\boldsymbol{\phi} = \boldsymbol{\mu} + \mathbf{L}\mathbf{z}$

The resulting sample $\boldsymbol{\phi}$ represents function values at the chosen grid points, respecting the correlation structure encoded by K .

1. Review of Chapter 1: Introduction
2. Definition and Basic Properties
- 3. Inference with Exact and Noisy Observations**
4. Joint Gaussian Processes
5. Summary

General Framework: Jointly Gaussian Observations

We can condition a GP $p(f) = \mathcal{GP}(f; \mu, K)$ on any vector \mathbf{y} sharing a joint Gaussian distribution with f :

$$p(f, \mathbf{y}) = \mathcal{GP} \left(\begin{bmatrix} f \\ \mathbf{y} \end{bmatrix}; \begin{bmatrix} \mu \\ \mathbf{m} \end{bmatrix}, \begin{bmatrix} K & \boldsymbol{\kappa}^\top \\ \boldsymbol{\kappa} & \mathbf{C} \end{bmatrix} \right)$$

where:

- $p(\mathbf{y}) = \mathcal{N}(\mathbf{y}; \mathbf{m}, \mathbf{C})$: marginal distribution of observations
- $\boldsymbol{\kappa}(x) = \text{cov}[\mathbf{y}, \phi \mid x]$: cross-covariance function

Q&A: What Observations \mathbf{y} Are Jointly Gaussian with f ?

Question

Besides function values, what else can \mathbf{y} be?

Any of the following are jointly Gaussian with a GP:

- Function values: $\mathbf{y} = f(\mathbf{x})$ (the basic case)
- Affine transformations: $\mathbf{y} = \mathbf{A}f(\mathbf{x}) + \mathbf{b}$
- Limits of affine transformations:
 - Partial derivatives: $\frac{\partial f}{\partial x_i}(x)$
 - Integrals/expectations: $\int f(x)p(x)dx$
- Any of the above + independent Gaussian noise

Q&A: What Observations \mathbf{y} Are Jointly Gaussian with f ?

Question

Besides function values, what else can \mathbf{y} be?

Other examples: Based on linear (affine) operator, these are joint Gaussian distribution.

- Line/Path Integral Observations [Särkkä, 2011]

$$\mathbf{y} = \int_{\gamma} f(x) ds + \varepsilon, \quad \kappa(x') = \int_{\gamma} K(x, x') ds$$

Used at Tomographic reconstruction (CT/MRI), Measure expected path of moving sensor

- Convolved/Smoothed Observations by smoothing kernel G [Alvarez et al., 2011]

$$\mathbf{y} = (f * G)(x_0) = \int f(x') G(x_0 - x') dx', \quad \kappa(x) = \int G(x_0 - x') K(x', x) dx'$$

Used at Sensor spatial averaging, image blur (PSF), multi-output GP via convolution

Posterior Gaussian Process

Conditioning on observations $\mathcal{D} = \mathbf{y}$ yields a GP posterior:

$$p(f \mid \mathcal{D}) = \mathcal{GP}(f; \mu_{\mathcal{D}}, K_{\mathcal{D}})$$

Posterior Mean and Covariance

$$\mu_{\mathcal{D}}(x) = \mu(x) + \kappa(x)^{\top} \mathbf{C}^{-1}(\mathbf{y} - \mathbf{m})$$

$$K_{\mathcal{D}}(x, x') = K(x, x') - \kappa(x)^{\top} \mathbf{C}^{-1} \kappa(x')$$

Inference procedure:

1. Compute marginal distribution of \mathbf{y}
2. Derive cross-covariance function κ
3. Apply the posterior formulas

Handling Additive Gaussian Noise

Suppose we observe $\mathbf{z} = \mathbf{y} + \varepsilon$ where $\varepsilon \sim \mathcal{N}(\mathbf{0}, \mathbf{N})$ is independent noise.

Then:

$$p(\mathbf{z} \mid \mathbf{N}) = \mathcal{N}(\mathbf{z}; \mathbf{m}, \mathbf{C} + \mathbf{N}); \quad \text{cov}[\mathbf{z}, \phi \mid x] = \kappa(x)$$

Key Result

Simply replace \mathbf{C} with $\mathbf{C} + \mathbf{N}$ in the posterior formulas!

As $\mathbf{N} \rightarrow \mathbf{0}$, the posterior converges to that from direct observation of \mathbf{y} .

Inference with Exact Function Evaluations

Suppose we observe f at locations \mathbf{x} , revealing $\phi = f(\mathbf{x})$.

The posterior is $p(f \mid \mathcal{D}) = \mathcal{GP}(f; \mu_{\mathcal{D}}, K_{\mathcal{D}})$ with:

$$\begin{aligned}\mu_{\mathcal{D}}(x) &= \mu(x) + K(x, \mathbf{x})\Sigma^{-1}(\phi - \mu) \\ K_{\mathcal{D}}(x, x') &= K(x, x') - K(x, \mathbf{x})\Sigma^{-1}K(\mathbf{x}, x')\end{aligned}$$

where $\Sigma = K(\mathbf{x}, \mathbf{x})$ and $\mu = \mu(\mathbf{x})$.

Key properties:

- Posterior mean **interpolates** through observed points
- Posterior variance **vanishes** at observed locations
- Uncertainty remains unchanged far from observations

Inference with Noisy Function Evaluations

Suppose observations are corrupted: $\mathbf{y} = \phi + \varepsilon$ with $\varepsilon \sim \mathcal{N}(\mathbf{0}, \mathbf{N})$.

Common noise models:

- **Homoskedastic:** $\mathbf{N} = \sigma_n^2 \mathbf{I}$ (constant noise)
- **Heteroskedastic:** $\mathbf{N} = \text{diag}(\sigma_n^2(\mathbf{x}))$ (location-dependent)

The posterior formulas become:

$$\begin{aligned}\mu_{\mathcal{D}}(x) &= \mu(x) + K(x, \mathbf{x})(\Sigma + \mathbf{N})^{-1}(\mathbf{y} - \mu) \\ K_{\mathcal{D}}(x, x') &= K(x, x') - K(x, \mathbf{x})(\Sigma + \mathbf{N})^{-1}K(\mathbf{x}, x')\end{aligned}$$

The posterior mean no longer interpolates exactly; extreme values may be “explained away” as noise.

Interpretation of Posterior Moments

Consider a single observation y with distribution $\mathcal{N}(y; m, s^2)$ and define:

- z-score: $z = \frac{y-m}{s}$
- Correlation: $\rho = \text{corr}[y, \phi \mid x] = \frac{\kappa(x)}{\sigma s}$

Posterior Moments (Scalar Case)

Posterior mean of ϕ : $\mu + \sigma \rho z$

Posterior std of ϕ : $\sigma \sqrt{1 - \rho^2}$

Intuition:

- Mean shifts proportionally to z-score and correlation strength
- Variance reduction depends only on correlation $|\rho|$

Q&A: Differential Entropy as Global Uncertainty

Question

Why is differential entropy a measure of global uncertainty?

Definition: Differential Entropy

For a random variable ω with density $p(\omega)$, *differential entropy* is defined as:

$$H[\omega] = - \int p(\omega) \log p(\omega) d\omega$$

Interpretation:

- Measures the “spread” or “uncertainty” in the distribution
- Higher entropy \Rightarrow more uncertain (flatter distribution)
- For Gaussian: $H[\mathcal{N}(\mu, \sigma^2)] = \frac{1}{2} \log(2\pi e \sigma^2)$

Q&A: Differential Entropy as Global Uncertainty

Question

Why is differential entropy a measure of global uncertainty?

Definition: Differential Entropy

For a random variable ω with density $p(\omega)$, *differential entropy* is defined as:

$$H[\omega] = - \int p(\omega) \log p(\omega) d\omega$$

For GPs: The joint differential entropy of function values at \mathbf{x} is:

$$H[\phi \mid \mathbf{x}] = \frac{1}{2} \log |2\pi e K(\mathbf{x}, \mathbf{x})|$$

Observations *reduce* this entropy: $|K_{\mathcal{D}}(\mathbf{x}, \mathbf{x})| \leq |K(\mathbf{x}, \mathbf{x})|$.

Posterior Predictive Distribution

For the latent function value $\phi = f(x)$:

$$p(\phi \mid x, \mathcal{D}) = \mathcal{N}(\phi; \mu_{\mathcal{D}}(x), K_{\mathcal{D}}(x, x))$$

For a **noisy observation** y at location x (with noise variance σ_n^2):

$$p(y \mid x, \mathcal{D}, \sigma_n) = \mathcal{N}(y; \mu_{\mathcal{D}}(x), K_{\mathcal{D}}(x, x) + \sigma_n^2)$$

The predictive credible intervals for noisy measurements are inflated compared to the latent function, reflecting observation uncertainty.

Contents

1. Review of Chapter 1: Introduction
2. Definition and Basic Properties
3. Inference with Exact and Noisy Observations
- 4. Joint Gaussian Processes**
5. Summary

Topics Covered in Remaining Sections

The remainder of Chapter 2 covers more specialized topics:

- **§2.4 Joint Gaussian Processes:** Modeling multiple correlated functions
- **§2.5 Continuity:** Conditions for continuous sample paths
- **§2.6 Differentiability:** Conditions for differentiable sample paths; derivative observations
- **§2.7 Existence/Uniqueness of Global Maxima:** Theoretical guarantees
- **§2.8 Non-Gaussian Observations:** Approximate inference methods

Motivation: Modeling Multiple Functions

In some settings, we need to jointly reason about **multiple related functions**:

- An objective function and its gradient
- An expensive objective and cheaper surrogates (multifidelity)
- Multiple objectives (multiobjective optimization)

Key Idea

“Paste together” multiple functions into a single function on a larger domain, then construct a standard GP on this combined function.

Q&A: Why Disjoint Union for Joint GP Domains?

Question

Is disjoint union the standard way to define joint GP domains?

Yes, mathematically this is the cleanest approach:

$$\bigsqcup f : \mathcal{X} = \bigsqcup_i \mathcal{X}_i \rightarrow \mathbb{R}, \quad \text{where } (\bigsqcup f)|_{\mathcal{X}_i} \equiv f_i$$

Why disjoint union?

- Ensures each point (x, i) uniquely identifies *which* function
- Even if $\mathcal{X}_1 = \mathcal{X}_2 = [0, 1]$, we distinguish $f_1(0.5)$ from $f_2(0.5)$
- The kernel can then define both within-function and cross-function covariance

Alternative: If functions share a domain and we want $f(x) \approx g(x)$, we can use a *separable* kernel structure without explicit disjoint union.

Definition of Joint Gaussian Process

Consider functions $\{f_i : \mathcal{X}_i \rightarrow \mathbb{R}\}$. Define the **disjoint union**:

$$\bigsqcup f : \mathcal{X} \rightarrow \mathbb{R}, \quad \mathcal{X} = \bigsqcup \mathcal{X}_i$$

such that $\bigsqcup f|_{\mathcal{X}_i} \equiv f_i$.

A **joint Gaussian process** is a GP on $\bigsqcup f$:

$$p(\bigsqcup f) = \mathcal{GP}(\bigsqcup f; \mu, K)$$

The mean and covariance functions on \mathcal{X} encode both:

- Marginal behavior of each function
- Cross-correlations between functions

Decomposed Notation

For two functions $f : \mathcal{F} \rightarrow \mathbb{R}$ and $g : \mathcal{G} \rightarrow \mathbb{R}$:

$$p(f, g) = \mathcal{GP} \left(\begin{bmatrix} f \\ g \end{bmatrix} ; \begin{bmatrix} \mu_f \\ \mu_g \end{bmatrix}, \begin{bmatrix} K_f & K_{fg} \\ K_{gf} & K_g \end{bmatrix} \right)$$

Components:

- μ_f, K_f and μ_g, K_g : marginal GP parameters
- $K_{fg}(x, x') = \text{cov}[\phi, \gamma \mid x, x']$: cross-covariance
- $K_{gf} = K_{fg}^\top$

Marginal property: Each function has a marginal GP distribution:

$$p(f) = \mathcal{GP}(f; \mu_f, K_f); \quad p(g) = \mathcal{GP}(g; \mu_g, K_g)$$

Q&A: Designing Cross-Covariance Functions

Question

How do we choose the cross-covariance function K_{fg} ?

Common approaches:

1. **Scaled base kernel:** $K_{fg}(x, x') = \rho \cdot K(x, x')$ where $|\rho| < 1$
 - Simple, interpretable: ρ is the correlation at any point
2. **Linear Model of Coregionalization (LMC):**

$$K_{fg}(x, x') = \sum_{q=1}^Q a_q^{(f)} a_q^{(g)} K_q(x, x')$$

3. **Convolution processes:** Define via convolution with a smoothing kernel

Constraint: The full covariance matrix must remain positive semidefinite!

Example: Correlated Functions




Figure 2: Example of Joint Gaussian Process [Garnett, 2023]

Consider $f, g : [0, 30] \rightarrow \mathbb{R}$ with:

- Same marginal: $\mu \equiv 0$, squared exponential covariance K
- Cross-covariance: $K_{fg}(x, x') = 0.9 \cdot K(x, x')$

Example: Correlated Functions




Figure 2: Example of Joint Gaussian Process [Garnett, 2023]

For any point x , the correlation between $\phi = f(x)$ and $\gamma = g(x)$ is:

$$\text{corr}[\phi, \gamma \mid x] = 0.9$$

Consequence: Samples from the joint distribution show strong coupling, the functions “move together.”

Q&A: Domain Size and Function Influence in Joint GPs

Question

Can domain size differences affect function influence in joint GPs?

Yes, this is a real concern in practice!

- Function with more observations may dominate inference
- Different scales of domains may require different length scales
- Numerical conditioning can suffer from imbalanced data

Mitigation strategies:

1. **Output scaling:** Normalize each function to similar variance
2. **Separate length scales:** Use ARD-style kernels for each function
3. **Weighted likelihoods:** Down-weight abundant data sources
4. **Hierarchical priors:** Place priors on correlation parameters

Inference for Joint GPs

The joint GP construction allows us to condition on observations of **any** of the functions using the standard inference procedure.

Examples

Given observations of f on the left side of the domain and observations of g on the right side:

- Observations of f inform our belief about g (and vice versa)
- Information propagates through the cross-covariance structure
- Strong correlation \Rightarrow strong information transfer

This is particularly useful for **multifidelity optimization**: cheap surrogate evaluations inform our belief about the expensive objective.

Extension to Vector-Valued Functions

A GP on a vector-valued function $\mathbf{f} : \mathcal{X} \rightarrow \mathbb{R}^d$ is defined by a joint GP on its coordinate functions $\{f_i\} : \mathcal{X} \rightarrow \mathbb{R}$.

Notation: $\mathcal{GP}(\mathbf{f}; \mu, K)$ where:

- $\mu : \mathcal{X} \rightarrow \mathbb{R}^d$ (vector-valued mean)
- $K : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}^{d \times d}$ (matrix-valued covariance)

Applications:

- Joint distribution of f and ∇f (gradient)
- Multiobjective optimization with correlated objectives
- Modeling spatial vector fields

Contents

1. Review of Chapter 1: Introduction
2. Definition and Basic Properties
3. Inference with Exact and Noisy Observations
4. Joint Gaussian Processes
5. Summary

Summary of Key Ideas

1. **GP Definition:** Specified by mean μ and covariance K functions; finite marginals are multivariate Gaussian
2. **Exact Inference:** Conditioning on jointly Gaussian observations yields a GP posterior with closed-form mean and covariance
3. **Noisy Inference:** Replace \mathbf{C} with $\mathbf{C} + \mathbf{N}$ to handle additive Gaussian noise
4. **Posterior Interpretation:** Mean update \propto (correlation \times z-score); variance reduction depends on correlation strength
5. **Joint GPs:** Model multiple correlated functions; enable information sharing across related tasks

References



Alvarez, M. A., Rosasco, L., and Lawrence, N. D. (2011).
Kernels for vector-valued functions: A review.



Durrett, R. (2019).
Probability: Theory and examples.
Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press.



Garnett, R. (2023).
Bayesian optimization.
Cambridge University Press.



Le Gall, J.-F. (2016).
Brownian motion, martingales, and stochastic calculus.
Graduate texts in mathematics. Springer International Publishing, 2016 edition.



Oksendal, B.
Stochastic differential equations: An introduction with applications.
Universitext. Springer.



Särkkä, S. (2011).
Linear operators and stochastic partial differential equations in Gaussian process regression, pages

Thank You