	-	ty of a leap ye	ear selected at	random contain 53	
Su	nday is:				
	(a) 53/ 366	(b) 1/7	(c) 2/7	(d) 53/365	
	-			A marble is drawn at	
raı	ndom. The pro	-	_		
	(a) 3/5	(b) 2/5	(c) $0/5$	(d) 1/5	
3.	The probabilit	y that it will r	ain tomorrow i	s 0.85. What is the	
pro	obability that it				
	(a) 0.25	(b) 0.145	(c) 3/20	(d) none of these	
4.				ected from the number	ers
(1,	2, 3,,15) is a multiple	e of 4?		
	(a) 1/5	(b) 4/5	(c) 2/15	(d) 1/3	
5 .	What are the	total outcome	es when we thi	row three coins?	
	(a) 4	(b) 5	(c) 8	(d) 7	
6.	The probabili	ty that a prim	e number sele	cted at random from	the
nu	mbers (1,2,3, .	35) is:			
	(a) 12/35	(b) 11/3	(c) 13/3	35 (d) none of these	е
7 .	The sum of th	e probability	of an event an	d non event is:	
	(a) 2	(b) 1 (c	c) 0 (d) n	one of these.	
8.	The following	probabilities	are given; cho	oose the correct answ	/er
	that which is	not possible.	_		
	(a) 0.15	(b) 2/7	(c) 7/5	(d) none of these.	
9.	, ,	* *		than the probability o	
	tting at least tv	wo heads, is:	-		
	(a) 1/4	(b) 3/8	(c) ½	(d) 1/8	
10	. A letter is ch	nosen at rand	om from the le	etters of the word	Vowe
•	ASSASSINATION	ON . The pro	obability that t	he letter chosen has:	VOVC
	(a) 6/13	(b) 7/13	(c) 1	(d) none of the	se.
		, , ,	` ,	` '	
11	. A dice is thro	wn. Find the I	probability of g	getting an even numb	er.
(A)) 2/3	(B) 1	(C) 5/6	(D) 1/2	
` .		` ,	` ,		
12	. Two coins are	e thrown at th	ie same time. I	Find the probability of	f
ge	tting both head	ds.			
(A)) 3/4 <mark>(B) 1/4</mark>	(C) 1/2	(D) 0		
13	. Two dice are	thrown simul	taneously. The	e probability of gettin	g a

1

sum of 9 is:

(A) 1/10	(B) 3/10	(C) 1/9	(D) 4/9				
14. 100 cards are numbered from 1 to 100. Find the probability of getting a prime number.							
(A) 3/4		(C) 1/4	(D) 29	9/100			
_	contains 5 red ba a blue ball is do n a bag is: (B) 10	uble that of a r		-			
16. A box of 600 bulbs contains 12 defective bulbs. One bulb is taken out at random from this box. Then the probability that it is non-defective bulb is: (A) 143/150 (B) 147/150 (C) 1/25 (D) 1/50							
17. Cards r mixed thor the probab	narked with nun oughly. One card ility that the nun (B) 1/10	nbers 2 to 101 d is drawn fron nber on card is	are placed in n this box ran a perfect squ	a box and domly, then			
18. What is the probability of getting 53 Mondays in a leap year? (A) 1/7 (B) 53/366 (C) 2/7 (D) 7/366							
19. A card is drawn from a well shuffled deck of 52 cards. Find the probability of getting a king of red suit. (A) 1/26 (B) 3/26 (C) 7/52 (D) 1/13							
20. A game of chance consists of spinning an arrow which is equally likely to come to rest pointing to one of the number 1,2,312 ,then the probability that it will point to an odd number is: (A) 1/6 (B) 1/12 (C) 7/12 (D) 5/12 Answer: 0.5							
21. A game consists of tossing a one rupee coin 3 times and noting its outcome each time. Aryan wins if all the tosses give the same result i.e. three heads or three tails and loses otherwise. Then the probability that Aryan will lose the game. (A) $3/4$ (B) $1/2$ (C) 1 (D) $1/4$							

same birthday is	s the same birth	Probability that bo day is: (C) 1/365				
23. A number x 2. Then the prob (A) 1/5 (B) 2	oability that x ² <	2 is?	ımbers -2, -1, 0 , 1,			
24. A jar contains 24 marbles. Some are red and others are white. If a marble is drawn at random from the jar, the probability that it is red is 2/3, then the number of white marbles in the jar is: (A) 10 (B) 6 (C) 8 (D) 7						
25. A number is selected at random from first 50 natural numbers. Then the probability that it is a multiple of 3 and 4 is: (A) 7/50 (B) 4/25 (C) 1/25 (D) 2/25						
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?						
a) $\frac{1}{7}$	b) $\frac{5}{42}$	c) $\frac{1}{21}$	(d) $\frac{4}{21}$			
	_	_	es are 50, 70, 82,			
93, and 20. The a) 25.79			d) 25.69			
28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.						
a) 13, 15	b) 13, 18	c) 18, 15	d) 13, 16			
29. A coin is tossed up 4 times. The probability that tails turn up in 3 cases is						
		$\frac{(c)^{1}}{4}$	d) $^{1}/_{6}$			
30. X is a variate a) 8 b)		d 3. The valu<mark>e</mark> of E) 27 <mark>d)</mark>	E(X²) is			
31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let Z= 5X-2Y. The variance of Z is?						

probability?	he following value $(b) \sum x P(b)$	•	ne is not possi	ble in				
c) $P(x) = 0.5$	d) P(x) =	= - 0.5						
33.If E(x) =	2 and E(z) = 4, t l b) 6	h en E(z - x) c) 0		ufficient data				
34.The cov	ariance of two in	dependent	random variabl	le is				
a) 1	b) 0	c) – 1	d) Un	defined				
35.If Σ P(x) a) 0) = k² - 8 then, th b) 1	e value of k		sufficient data				
` '	36.If P(x) = 0.5 and x = 4, then E(x) = ? a) 1 b) 0.5 c) 4 d) 2							
37.In a discrete probability distribution, the sum of all probabilities is always?								
a) 0	b) Infinite	c) 1	d) Und	efined				
-	obability of hittir	ng the targe	t is 0.4, find m	ean and				
variance. a) 0.4, 0.24	b) 0.6, 0.2	24	c) 0.4, 0.16	d) 0.6, 0.16				
39.If the probability that a bomb dropped from a place will strike the target is 60% and if 10 bombs are dropped, find mean and variance? a) 0.6, 0.24 b) 6, 2.4 c) 0.4, 0.16 d) 4, 1.6 40. Find the mean of tossing 8 coins. a) 2 b) 4 c) 8 d) 1 41. What is the mean and variance for standard normal distribution?								

c) 5

d) 7

a) 3

b) 4

,		variance is o	,						
a) E(X)			iable X is given by c) E(X2) - (E(X))2			d) (E(X))2			
		ndom variab E(X2)	_	•		d) (E(X))2			
	44.Mean of a constant 'a' is a) 0								
	45.Variance of a constant 'a' is . a) 0								
46.Find	46.Find the mean and variance of X?								
	x 0	1	2	3	4				
) 1/9	2/9	3/9	2/9	1/9				
f(x)	179								

	Х	0	1	2	3	
	f(x)	1/6	2/6	2/6	1/6	
a) ().5		b) 1.5		c) 2.5	d) 3.5

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by

c) np2q

d) npq2

- 49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**
- a) P(X = x) = nCx px qx
- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx
- 50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**
- a) \sqrt{np}

- b) \sqrt{pq} c) (np)2 d) \sqrt{npq}