a) See Python notebook

b) i) for
$$N : \text{even} \longrightarrow l_{(\text{levels})} = \frac{N}{2} + 1$$
for $N : \text{odd} \longrightarrow l_{(\text{levels})} = \frac{LN}{2J} + 1$

(ii) the lowest energy will be -N, from then, every time we thip a spin the energy either:

So
$$E_n = -N + 4 \cdot n$$
 $n = 0, 1, ..., \ell - 1$

degeneracy:
$$D = 2 \times {N \choose 2n} = 2 \frac{N!}{2n!(N-2n)!}$$

eg: N=4,
$$n=1$$
 should have $D=12$; $D=2\cdot \binom{4}{2}=2\cdot \frac{4!}{2!2!}=2\cdot 3!=12$

(i)
$$E_n = -N + 2n$$
 where $n = 0, 1, ..., \ell - 1$

$$\mathcal{D}_{j} = \binom{\nu}{N} = \frac{\nu_{i} (N - \nu_{j})}{N_{i}}$$

Question 2

See Python notebook

Question 3

$$E_{\alpha} = A + \frac{B}{N} \left(\Delta_{\alpha} - \frac{C}{12} \right) + O\left(\frac{1}{N^{2}}\right)$$

$$E_{o}^{PBC} \simeq A - \frac{B}{N} C$$

$$\begin{cases} E_{\alpha} - E_{o}^{PBC} = \frac{B}{N} \Delta_{\alpha} \\ E_{\tau}^{PBC} - E_{o}^{PBC} = \frac{B}{N} 2 \end{cases}$$

$$\longrightarrow \Delta_{\alpha} \approx 2 \frac{E_{\alpha} - E_{o}^{PBC}}{E_{\tau}^{PBC} - E_{o}^{PBC}}$$

b) See Python notebook

Question 4:

See Python notebook