

Supplementary Table 2: Stage-specific Spearman correlations between immune checkpoint concentrations in PBMCs stimulated with sEVs and their corresponding concentrations in isolated sEVs.

Stage	Checkpoint	rho	p-value	Interpretation
HC	PD-1	-0,288	0,637	Weak
	PD-L2	-0,774	0,225	Strong
	Galectin-9	-1	NA	Very strong
	Free Active	-0,774	0,225	Strong
	4-1BB	0,894	0,105	Very strong
	PD-L1	0,866	0,057	Very strong
CG	4-1BB	-0,447	0,552	Moderate
	B7.2	-0,894	0,105	Very strong
	PD-L2	-0,288	0,637	Weak
	sCD25	1	NA	Very strong
	PD-1	0,774	0,225	Strong
	Galectin-9	0,866	0,333	Very strong
GC I	PD-1	-0,288	0,637	Weak
	PD-L2	-0,774	0,225	Strong
	CTLA-4	-0,577	0,308	Moderate
	B7.2	0,894	0,105	Very strong
	LAG_3	0,866	0,057	Very strong
	Galectin-9	0,866	0,333	Very strong
	Free Active	0,866	0,333	Very strong
GC II	PD-1	-0,774	0,225	Strong
	Tim-3	-0,894	0,105	Very strong
	B7.2	0,894	0,105	Very strong
	Galectin-9	0,866	0,333	Very strong
GC III	LAG_3	-0,774	0,225	Strong
	PD-L2	-0,288	0,637	Weak
	B7.2	0,894	0,105	Very strong
	PD-L1	0,866	0,057	Very strong
	CTLA-4	0,866	0,057	Very strong
GC IV	B7.2	-0,894	0,105	Very strong
	sCD25	1	NA	Very strong
	PD-L2	0,774	0,225	Strong
	Tim-3	0,894	0,105	Very strong
	Galectin-9	0,866	0,333	Very strong
	sCD27	0,866	0,333	Very strong