Supplementary Table 1: Spearman correlation analysis among soluble immune checkpoint molecules in PBMCs according to clinical group and time point.

Analyte	Time peaks	stage	Checkpoints	rho
sCD25				
	24	ND	ND	ND
sCD27				
	24	HC	Tim-3	1
			LAG-3	-1
B7.2				
	24	HC	4-1BB, PD-1, CTLA-4	1
			PD-1, CTLA-4	-1
		GC III	4-1BB	1
CTLA-4				
	24	HC	PD-1	1
		CG	Galectin-9	1
			PD-1, PD-L2	-1
		GC I	PD-L1	1
		GC IV	PD-L2	-1
TGF-β1				
	24	HC	PD-L1, PD-L2	-1
PD-L1				
	24	HC	PD-L2	1
			Free Active	-1
		GC I	CTLA-4	1
		GC II	4-1BB, LAG-3	-1
		GC III	PD-L2	1
			LAG-3, Tim-3	-1
		GC IV	LAG-3, PD-1	-1
Tim-3				
	24	HC	sCD27	1
			LAG-3	-1
		GC III	4-1BB, LAG-3	1
			PD-L1, PD-L2	-1
Galectin-9				
	24	CG	CTLA-4	1
			PD-1, PD-L2	-1
LAG-3				
	24	HC	Tim-3, sCD27	-1
		GC I	PD-1	1
		GC II	4-1BB, PD-1	1
			PD-L1	-1
		GC III	Tim-3	1
			PD-L1, PD-L2	-1
		GC IV	PD-1	1
			PD-L1	-1
4-1BB				
	48	GC I	PD-L2	1
		GC II	B7.2, Tim-3	-1
		GC IV	PD-1	1
PD-1				
	48	HC	LAG-3	1
			PD-L2	-1
		CG	PD-L1, PD-L2	1
		GC I	CTLA-4	1
			LAG-3, sCD25, PD-L1, Tim-3	-1
		GC II	CTLA-4	1
			LAG-3, PD-L2	-1
		GC III	PD-L1	-1
		GC IV	4-1BB, sCD25	1
PD-L2				
	72	HC	PD-1	1
			4-1BB	-1
		GC I	sCD25, PD-L1	1
		GC III	sCD25, PD-L1	1
			PD-1	-1
		GC IV	B7.2	1