Supplementary Table 2: Stage-specific Spearman correlations between immune checkpoint concentrations in PBMCs stimulated with sEVs and their corresponding concentrations in isolated sEVs.

Stage	Checkpoint	ρ	Interpretation
HC	PD-1	-0,288	Weak
	PD-L2	-0,774	Strong
	Galectin-9	-1	Very strong
	Free active TGF-β1	-0,774	Strong
	4-1BB	0,894	Very strong
	PD-L1	0,866	Very strong
CG	4-1BB	-0,447	Moderate
	B7.2	-0,894	Very strong
	PD-L2	-0,288	Weak
	sCD25	1	Very strong
	PD-1	0,774	Strong
	Galectin-9	0,866	Very strong
GC I	PD-1	-0,288	Weak
	PD-L2	-0,774	Strong
	CTLA-4	-0,577	Moderate
	B7.2	0,894	Very strong
	LAG-3	0,866	Very strong
	Galectin-9	0,866	Very strong
	Free active TGF-β1	0,866	Very strong
GC II	PD-1	-0,774	Strong
	TIM-3	-0,894	Very strong
	B7.2	0,894	Very strong
	Galectin-9	0,866	Very strong
GC III	LAG-3	-0,774	Strong
	PD-L2	-0,288	Weak
	B7.2	0,894	Very strong
	PD-L1	0,866	Very strong
	CTLA-4	0,866	Very strong
GC IV	B7.2	-0,894	Very strong
	sCD25	1	Very strong
	PD-L2	0,774	Strong
	TIM-3	0,894	Very strong
	Galectin-9	0,866	Very strong
	sCD27	0,866	Very strong

Note. For each immune checkpoint, the time point showing the highest stimulation was used: 24 h for sCD25, sCD27, PD-L1, CTLA-4, B7.2, TIM-3, galectin-9, LAG-3, and free active TGF-β1; 48 h for 4-1BB and PD-1; and 72 h for PD-L2. Correlations are presented descriptively without significance testing because the functional assay included a single biological experimental unit.