



Patients per Gender


Patients' participating at the research divided by Gender




Alzheimer's patients percentage based on gender



Female Alzheimer's patients based on age




Male Alzheimer's patients percentage




Female Alzheimer's patients percentage




Male Alzheimer's patients based on age






Caucasian Patients

Data acquired from 2149 patients




Asian Patients Data acquired from 2149 patients




African American Patients

Data acquired from 2149 patients




Caucasian Patients diagnosed with Alzheimer Data acquired from 2149 patients




Asian Patients diagnosed with Alzheimer

Data acquired from 2149 patients




African American Patients diagnosed with Alzheimer

Data acquired from 2149 patients




Patients' Family History of Alzheimer's

Distribution of Patients Reporting Family History of Alzheimer's




Smoking Prevalence Among Patients



Alzheimer's Patients with Reported Family History


Distribution of Patients Reporting Family History of Alzheimer's




Alzheimer's Diagnosis Rate Among Non-Smokers

No Family History

Family History



Alzheimer's Diagnosis Rate Among Smokers



Frequency of Weekly Alcohol Consumption Diagnosed Patients – BMI Patient Distribution by Alcohol Consumption Level (1 = Low, 8 = High) Patient Distribution by Body Mass Index of the patients (1 = Low, 8 = High) ■ TotalPatients ■ DiagnosedPatients TotalPatientsDiagnosedPatients 300 272 265 268 268 264 264 262 260 260 257 247 Number of Patients **Patients** 200 Number of 109 108 108 101 104 93 94 89 87 83 85 100 .80 0 8 Frequency of Alcohol Consumption Frequency of Alcohol Consumption Diagnosis Rate by Drinking Frequency Group - Group 1 Group - Group 5 Group - Group 3 Group - Group 7 97 108 (32.4%)(31.92%)(36.6%) (38.85%)No Diagn... No Diagn... No Diagn... No Diagn... Diagnosis Diagnosis Diagnosis Diagnosis 194 (67.6%) 177 (68.0...) Diagnosis Rate by Drinking Frequency Group - Group 2 Group - Group 4 Group - Group 6 Group - Group 8 (33.46%) (38.55%) (34.7%)(36.58%) No Diagn... No Diagn... No Diagn... No Diagn... Diagnosis Diagnosis Diagnosis Diagnosis

— 163 (63.4...)


[└] 181 (66.5...)

└─ 175 (65.3%)

— 161 (61....)


Frequency of Weekly Physical Activity

Physical Activity (0–10 Hours/Week): Patient Distribution (1 = Low, 4 = High)




Frequency of Alcohol Consumption


Diagnosis Rate by Physical Activity Group - Group 1




Diagnosis Rate by Physical Activity Group - Group 2



Diagnosis Rate by Physical Activity Group - Group 3




Diagnosis Rate by Physical Activity Group - Group 4



Patients by Diet Quality Score

Diet Quality Score (Ranging from 0-10:): Patient Distribution (1 = Bad, 4 = Good)



Patients by Sleep Quality Score

Sleep Quality Score (Ranging from 4-10): Patient Distribution (1 = Bad, 4 = Good)

