

int != Integer; How choosing the right data type could improve your system’s performance

SoftPapers,¹ Amuda A,^{2*}

¹*Sapafy’s SoftPaper*
²*Contributor from Sapafy*
<amuda@sapafy.com>

ABSTRACT.

A fast and highly responsive system doesn’t only come from scaling the system horizontally or vertically, but the underlying implementations plays a huge role which is mostly underestimated. int is never the same as Integer, and applying the wrong data type comes with a hidden cost which could impact your system’s performance negatively, and the goal of this paper is to show you how choosing the right data type could improve your system’s performance.

INT AND INTEGER HISTORY

If you intend to store or read a number, int or Integer data type will mostly be considered, while both might serve the same results for the end users, they are meant to be used correctly. int is a primitive data type which has a 32-bit size, meaning you can store or maintain a number not more or lesser than -2^{32} to 2^{31}

Integer is a reference data type that also maintains the same bit size as int, but has an object reference in its memory with a reference or pointer to another reference.

Properties ^a %	int	Integer
Size	32bits	32 bits
Subclass of Number in Object	No	Yes
Require memory	<= 32bits	64 bits+
Autoboxing	No	Yes
Requires pointer	No	Yes
Comparison operation	==	equals

^a.

Why Integer requires more memory:

Short answer:

It has more properties that need to be maintained once it’s created.

Long answer:

1. It’s a class
2. All classes in Java are a subclass of Object

^{*}Present address: Centre for Glaciology, Institute of Geography and Earth Sciences, University of Wales, Aberystwyth, UK.

3. All subclass in Java automatically extends the Object methods and properties
4. For an object to exist,
 - i. It must be created
 - ii. For it to be created, it must be stored in a temporary or permanent storage
 - iii. For it to be used, it must be referenced from the storage
 - iv. For it to be referenced, it must maintain properties to determine its uniqueness
 - v. For it to be unique, its properties must be stored in a memory which requires additional memory
HashCode of that object
Header details of the objects

```
\title[Short Title]{The Full Title of Your Paper}  
\author[Short Authors]{Author 1, Author 2 and Author 3}
```

Lists

The IGS class file provides for numbered (enumerate) and unnumbered (itemize) lists. Nested lists are not encouraged. The default numbering system is 1., 2., 3., etc.; please do not change this unless there is a good reason. The IGS design removes bullet points from unnumbered lists.

User-defined macros

If possible, please do not define any new macros.

Table 1. One-column table captions will extend beyond the rules in two-column format. Do not try to adjust! Table captions do not have full points at the end

Period ^a	Surface elevation change	Emergence velocity
1975–85	−0.50	0.43
1986–2002	−1.03	0.32
Difference	−0.53	−0.11

^aPlease do not use more than one ‘&’ between columns, and note that if a table includes table footnotes, it must be inside a `minipage` environment.

Tables

Tables may be typeset in either one- or two-column format. To typeset two-column format, add asterisks (`\begin{table*}... \end{table*}`) as shown in Table 2. We may change the format in-house if necessary. Please avoid the use of colour or shading. Note that if you choose to refer to tables using labels, `\caption` must precede `\label`, as in standard L^AT_EX. Vertical rules are not house-style and will be removed. Note the use of the `minipage` environment in Table 1 which enables table footnotes to be output. If the table is two-column, use `{178mm}` instead of `{86mm}` on line 6. The source code for Tables 1 and 2 is shown immediately below the tables.

FIGURES

Figures may be typeset in either one- or two-column format. One-column format allows up to 86 mm (e.g. Fig. 1); two-column format up to 178 mm (e.g. Fig. 2). Please do not provide original graphics files in which the figure is a great deal larger or smaller than what you envisage will be the final printed size. To typeset two-column format, add asterisks (`\begin{figure*}... \end{figure*}`) as shown in Fig. 2. We may change the format in-house if necessary. Please note that if you choose to refer to figures using labels, `\caption` must precede `\label`, as in standard L^AT_EX.

Please send one file for each figure (in other words do not use subfigures) and use a name that clearly identifies it (e.g. ‘72A712Fig03.eps’).

In addition, figures should be eps, ai (illustrator), ps, tif, psd or pdf. Use strong black lines with a width of at least 0.75pt at final printed size (avoid tinting if possible) and SI units in labels. Lettering should ideally be Optima to match the final typeface; Arial or a similar sans serif font for a second choice. Aim to have the final-size lettering at 9pt, if possible. Figures should not be in boxes. The source code for Figs 1 and 2 is shown immediately below the figures.

Equations

We are including some complex equations as examples. Equations should be checked for width by removing the `[review]` option. Note the use of `cases*` in the following equation:

$$\alpha_{t_2} = \begin{cases} \alpha_{t_1} - a_1[\ln(T+1)]e^{(a_2\sqrt{n})} & n_d > 0 \text{ and } T > 0 \\ \alpha_{t_1} - a_3e^{(a_2\sqrt{n})} & n_d > 0 \text{ and } T < 0 \\ \alpha_{t_1} + a_4P_s & n_d = 0 \end{cases} \quad (1)$$

Equations should be aligned on the equals signs where possible. Equations that extend beyond the one-column measure should be turned over before an operator.

$$I_c = I_0 \left(\frac{\bar{R}_m}{R} \right)^2 \psi^{\frac{p}{p_0 \cos Z}} \times [\cos \beta \cos Z + \sin \beta \sin Z \cos(\psi_{\text{sun}} - \psi_{\text{slope}})] \quad (2)$$

Typesetting upright Greek characters

Normal greek: `\alpha\beta\gamma\delta` $\alpha\beta\gamma\delta$

Upright greek: `\upalpha\upbeta\upgamma\updelta` $\alpha\beta\gamma\delta$

Usual partial: `\partial` ∂

Upright partial: `\uppartial` ∂

Marginal notes

The IGS class file redefines the L^AT_EX command `\marginpar`. If you wish to add a marginal note such as the one alongside this text, you would key `\marginpar{Editor! Help!}`. Marginal notes will be removed before printing.

References

All citations in text should include the author name(s) and the year of publication (e.g. ‘Smith, 2014’; ‘Smith and Jones, 2014’; ‘Smith and others, 2015’) and have an entry in the reference list.

References should:

be short;

be complete and accurate;

be arranged in alphabetical order by first author’s surname;

include too much rather than too little information;

Table 2. Two-column table. Seasonal and annual SAT trends (°C decade^{−1}) in the Arctic

Area	1951–2005					1976–2005				
	Dec–Feb	Mar–May	Jun–Aug	Sep–Nov	Annual	Dec–Feb	Mar–May	Jun–Aug	Sep–Nov	Annual
Atlantic region	0.09	0.29	0.10	0.09	0.15	0.470	0.60	0.45	0.53	0.59
Siberian region	0.12	0.29	0.04	0.17	0.16	0.08	0.69	0.29	0.59	0.48
Pacific region	0.45	0.46	0.25	0.26	0.35	0.712	1.08	0.27	0.66	0.52
Canadian region	0.16	0.12	0.14	0.30	0.18	0.20	0.52	0.48	0.94	0.53
Baffin Bay region	−0.02	0.10	0.00	0.15	0.02	0.33	0.62	0.51	0.80	0.57
Arctic 1	0.16	0.21	0.12	0.20	0.18	0.36	200.65	0.42	0.74	0.54
Arctic 2	0.22	0.29	0.14	0.14	0.19	0.38	0.60	0.40	0.51	0.45
Arctic 3	0.28	0.31	0.14	0.13	0.21	0.42	40.53	0.41	0.42	0.43
NH (land + ocean)	0.13	0.13	0.10	0.10	0.12	0.27	0.24	0.25	0.25	0.25




Fig. 1. One-column figures should be ≤86 mm. Good artwork can make or break a paper. Capitalize the first word of a label and use round not square brackets for units.

- include doi numbers where available (note that older bib databases often included doi’s in the page field – in which case they may appear after a comma and without braces);
- include works accepted but not published as ‘in press’;
- not include personal communications, unpublished data or manuscript in preparation or submitted for publication, data published on the web (these should be included in the text).

Automatic references using BibT_EX

To generate automatic references from a bib database, you must first specify the database (we are using `igsrefs.bib`) and then the IGS bibliography style by placing the following

two commands where you would like the references to appear (normally at the end of your paper, before `\end{document}`):

```
\bibliography{igsrefs}
\bibliographystyle{igs}
```

Then run through the following steps:

1. Run your paper through L^AT_EX.
2. Run BibT_EX on your paper.
3. Open the newly-created bbl file containing the cited references and copy the entire contents to just below the `bibliography/bibliographystyle` commands.
4. Then comment them out:

```
%\bibliography{igsrefs}
%\bibliographystyle{igs}
```

5. Run your paper through L^AT_EX *twice* more.

The IGS do not need your bib or bbl files. Note that BibT_EX will lose the second initial in the entry ‘Box JE’, for example, if it has been typed as ‘{J.E.} Box’ in the bib file. This is because any text in an entry enclosed in `{ }` will be treated as a single unit, and will not be further parsed. Prof. Box’s name will typeset correctly if entered as ‘J. E. Box’ in the bib file.

If you have cited 16 references from the bib database, e.g. (Rignot and Steffen, 2008), (Rignot and others, 2008), (Motyka and others, 2011), (Morlighem and others, 2010), (Morlighem and others, 2011), (Seroussi and others, 2011), (Yan and others, 2013), (Rogozhina and others, 2012), (Hanna and others, 2013), (Goelzer and others, 2013), (Lucas-Picher and others, 2012), (Edwards and others, 2014), (Gladstone

and others, 2010), (Morlighem and others, 2013), (Goldberg and Sergienko, 2011) and (Paterson, 1994), the output will be just those 16 references and they will appear at the end of the article.

Citations using natbib commands Note that the standard natbib style file has been modified to fall into line with IGS style. The modified style file is called igsnatbib.sty (included in this distribution), and works exactly the same as natbib.sty. The default IGS house style is (Yan and others, 2013). The following combinations are also available – refer to the natbib documentation if you require any further explanation:

(Yan and others, 2013) `\citep{Yan13}`
 (see Yan and others, 2013, p. 34) `\citep[see] [p.$\$, $34]{Yan13}`
 (e.g. Yan and others, 2013) `\citep[e.g.] []{Yan13}`
 (Yan and others, 2013, Section 2.3) `\citep[Section~2.3]{Yan13}`
 (Yan and others, 2013; Edwards and others, 2014) `\citep{Yan13, Edwards14}`
 Yan and others (2013); Edwards and others (2014) `\cite{Yan13, Edwards14}`
 Yan and others 2013 `\citealt{Yan13}`
 Yan and others (2013) `\cite{Yan13}`
 Yan and others, 2013 `\citealp{Yan13}`
 Yan and others `\citeauthor{Yan13}`
 (2013) `\citeyearpar{Yan13}`
 2013 `\citeyear{Yan13}`

ACKNOWLEDGEMENTS

We would like to thank Jason Amundson, Ed Bueler, Andrew Clifton, Gwenn Flowers, Ralf Greve and Doug MacAyeal for their constructive reviews of the IGS class file and guide. Thanks are also due to Patrick Daly who once again helped to generate the latest version of igs.bst.

REFERENCES

Edwards TL, Fettweis X, Gagliardini O, Gillet-Chaulet F, Goelzer H, Gregory JM, Hoffman M, Huybrechts P, Payne AJ, Perego M, Price S, Quiquet A and Ritz C (2014) Effects of uncertainty in surface mass balance-elevation feedback on projections of the future sea level contribution of the Greenland ice sheet. *The Cryosphere*, **8**, 195–208 (doi: 10.5194/tc-8-195-2014)

Gladstone RM, Lee V, Vieli A and Payne AJ (2010) Grounding line migration in an adaptive mesh ice sheet model. *J. Geophys. Res.-Earth*, **115**, F04014 (doi: 10.1029/2009JF001615)

Goelzer H, Huybrechts P, Fürst JJ, Nick FM, Andersen ML, Edwards TL, Fettweis X, Payne AJ and Shannon S (2013) Sensitivity of Greenland ice sheet projections to model formulations. *J. Glaciol.*, **59**(216), 733–749 (doi: 10.3189/2013JoG12J182)

Goldberg DN and Sergienko OV (2011) Data assimilation using a hybrid ice flow model. *The Cryosphere*, **5**, 315–327 (doi: 10.5194/tc-5-315-2011)

Hanna E, Navarro FJ, Pattyn F, Domingues CM, Fettweis X, Ivins ER, Nicholls RJ, Ritz C, Smith B, Tulaczyk S, Whitehouse PL and Zwally HJ (2013) Ice-sheet mass balance and climate change. *Nature*, **498**, 51–59 (doi: 10.1038/nature12238)

Lucas-Picher P, Wulff-Nielsen M, Christensen JH, Adalgeirsdóttir G, Mottram RH and Simonsen SB (2012) Very high resolution regional climate model simulations over Greenland: identifying added value. *J. Geophys. Res.*, **117**, D02108 (doi: 10.1029/2011JD016267)

Morlighem M, Rignot E, Seroussi H, Larour E, Dhia HB and Aubry D (2010) Spatial patterns of basal drag inferred using control methods from a full-Stokes and simpler models for Pine Island Glacier, West Antarctica. *Geophys. Res. Lett.*, **37**, L14502 (doi: 10.1029/2010GL043853)

Morlighem M, Rignot E, Seroussi H, Larour E, Dhia HB and Aubry D (2011) A mass conservation approach for mapping glacier ice thickness. *Geophys. Res. Lett.*, **38**, L19503 (doi: 10.1029/2011GL048659)

Morlighem M, Seroussi H, Larour E and Rignot E (2013) Inversion of basal friction in Antarctica using exact and incomplete adjoints of a higher-order model. *J. Geophys. Res.*, **118**, 1746–1753 (doi: 10.1002/jgrf.20125)

Motyka RJ, Truffer M, Fahnestock M, Mortensen J, Rysgaard S and Howat I (2011) Submarine melting of the 1985 Jakobshavn Isbrae floating tongue and the triggering of the current retreat. *J. Geophys. Res.*, **116**, F01007 (doi: 10.1029/2009JF001632)

Paterson WSB (1994) *The physics of glaciers*. Butterworth-Heinemann, Oxford, 3rd edition

Rignot E and Steffen K (2008) Channelized bottom melting and stability of floating ice shelves. *Geophys. Res. Lett.*, **35**, L02503 (doi: 10.1029/2007GL031765)

Rignot E, Box JE, Burgess E and Hanna E (2008) Mass balance of the Greenland ice sheet from 1958 to 2007. *Geophys. Res. Lett.*, **35**, L02502 (doi: 10.1029/2008GL035417)

Rogozhina I, Hagedoorn JM, Martinec Z, Fleming K, Soucek O, Greve R and Thomas M (2012) Effects of uncertainties in the geothermal heat flux distribution on the Greenland ice sheet: an assessment of existing heat flow models. *J. Geophys. Res.*, **117**, F02025 (doi: 10.1029/2011JF002098)




Fig. 2. Two-column figures should be ≤ 178 mm. SSA reconstructed components found by projecting the SSA filters found using the whole 2000 traces in Fig. 4, on trace number 1, ordered by magnitude of variance accounted for in the radar trace.

```

\begin{figure*}%fig2, two column
\centering{\includegraphics{72A712Fig02.eps}}
\caption{Two-column figures should be  $\leq 178$  mm. SSA reconstructed components found by
projecting the SSA filters found using the whole 2000 traces in Fig.~4, on trace number 1,
ordered by magnitude of variance accounted for in the radar trace.}
\label{filters}
\end{figure*}

```

Seroussi H, Morlighem M, Rignot E, Larour E, Aubry D, Dhia HB and Kristensen SS (2011) Ice flux divergence anomalies on 79north glacier, Greenland. *Geophys. Res. Lett.*, **38**, L09501 (doi: 10.1029/2011GL047338)

Yan Q, Zhang Z, Gao Y, Wang H and Johannessen OM (2013) Sensitivity of the modeled present-day Greenland ice sheet to climatic forcing and spin-up methods and its influence on future sea level projections. *J. Geophys. Res. Earth Surf.*, **118**, 2174–2189 (doi: 10.1002/jgrf.20156)

APPENDIX

Start an appendix by typing `\appendix\section{Appendix}`. Appendices appear after the references. Equation numbers automatically start again with (3).

$$2\eta\kappa\frac{\partial\bar{u}}{\partial t} + \rho_t g\bar{u} + D\kappa^4\bar{u} = \bar{\sigma}_{zz}. \quad (3)$$

HANDLING MORE THAN ONE APPENDIX

Use the following code to achieve heading APPENDIX A followed by APPENDIX B and APPENDIX C, with appropriate equation numbers:

```
\appendix
\section{Appendix A}

\setcounter{equation}{0}
\renewcommand\theequation{B\arabic{equation}}
\section{Appendix B}

\setcounter{equation}{0}
\renewcommand\theequation{C\arabic{equation}}
\section{Appendix C}
```