

Some Peculiar Lesions of Traumatic Brain Injuries Encountered in our Practice with Brief Discussion-Imaging Manuscript

*Uduma FU¹, Itanyi UD², Akpan S³, Eduwem DU⁴, Emejulu JKC⁵

ABSTRACT

The advent of neuroimaging like computed tomography and magnetic resonance imaging has facilitated the diagnosis of traumatic brain injuries. Emphasizing certain diagnostic features of some peculiar traumatic brain injuries. The following lesions of traumatic brain injuries were pictorially depicted, namely Tension pneumocephalus, Blow out orbit, Bilateral subacute subdural haematomata, Acute-on-chronic subdural haematoma, Middle cranial fossa acute epidural haematoma, Traumatic basal ganglial haematoma and Acute intra-ventricular haematoma. Mount Fuji sign is typical of tension pneumocephalus while herniation of extra-ocular muscles into the maxillary sinus is diagnostic of blow-out orbit. Rabbit ear appearance is observable in bilateral subacute subdural haematomata.

Keywords: Traumatic brain injury, Computed tomography, Magnetic resonance imaging, Pneumocephalus

INTRODUCTION

Traumatic brain injury (TBI) is a more specific terminology for head injury. TBI is an important public health problem worldwide, thus a potentially devastating clinical condition^{1,4}. In Africa, the commonest etiologic factor is road traffic accident (RTA)^{4,5}. RTA as an independent entity is of grievous concern to many developing nations with its untoward economic impacts^{5,6}. TBI is known to be accompanied by significant morbidity and mortality⁴. This essentially calls for expedited and non-invasive detection of any brain injury and its sequelae. The advent of neuroimaging like computed tomography (CT) and magnetic resonance imaging (MRI) has facilitated the elucidation of variable traumatic intracranial lesions. This informs our opting for few salient cases shown pictorially to highlight lesion peculiarities and localizations that may be inimical to the patients using neuro-images.

IMAGE PRESENTATIONS

Case A




Figure 1: Axial brain CT at the level of lateral ventricles showing frontal intracranial, extra-axial (sub-dural) air density lesion overlying and compressing posteriorly both frontal lobes with air dipping into the interhemispheric fissure and separation of the frontal lobes to a peak like configuration (black arrow) characteristic of Mount Fuji sign of Tension pneumocephalus.

Department of Radiology¹, Faculty of Clinical Sciences, University of Uyo, Uyo, Nigeria.

Department of Radiology², University of Abuja Teaching Hospital, Abuja Nigeria.

Department of Ophthalmology³, Faculty of Clinical Sciences, University of Uyo, Uyo, Nigeria.

Department of Radiology⁴, University of Calabar Teaching Hospital, Calabar, Nigeria.

Neurosurgical Unit, Department of Surgery⁵, Nnamdi Azikiwe University Teaching Hospital, Nnewi, Nigeria.

*Corresponding author: felixuduma@yahoo.com


Case B

Figure 2: Bone window images of cranial CT (coronal, sagittal and axial) showing bony defect (black arrow) of floor of the left orbit (25×20 mm in size) complicated by herniation of left extra-ocular muscle and intra-orbital fat into the left maxillary sinus giving a tear drop sign (Blow-out orbit).

Case C

Figure 3: Axial brain CT at the level of lateral ventricles showing bilateral intracranial extra-axial concavo-convex isodense collection (black arrows) of subdural haematoma in the fronto-parietal calvarium. Note associated compression of both ventricles medially leading to narrow, slit-like elongated ventricles (rabbit ear sign) and slight sub-falcine herniation to the left. (Bilateral subacute subdural haematoma).




Figure 4: Axial T1 weighted MR image in another patient with MRI more eloquently showing bilateral hyperintense crescentic shaped collection (black arrows) extending from frontal to occipital region-(Bilateral subacute subdural haematoma).


Case D

Figure 5: Axial brain CT showing right fronto-parietal extracranial crescentic bleed with fluid-fluid level and dependent hyperdense component (white arrow). Note associated significant subfalcine herniation (black arrow) to contralateral side. (Acute-on-chronic subdural haematoma).


Case E

Figure 6: Unenhanced axial brain CT at the base of the skull showing localized biconvex hyperdense bleed in the anterior part of left middle cranial fossa (black oblique arrow) beneath the left greater wing of sphenoid. This is associated with an aerocoele. Both lesions are compressing on the left temporal cerebral lobe. Note another aerocoele (black curved arrow) in the contralateral right middle cranial fossa. (Anterior temporal acute epidural haematoma and pneumocephalus in middle cranial fossa).


Case F

Figure 7: Unenhanced axial brain CT at the level of lateral ventricle showing hyperdense intra-parenchymal ovoid bleed (black arrow) located in the vicinity of right globus pallidus and putamen. Note compression of right lateral ventricle. (Acute traumatic basal ganglial haematoma).


Case G

Figure 8: Axial brain CT at the level of lateral ventricle showing fluid- fluid level with hyperdense (black arrow) collection in the occipital horns of the lateral ventricles (Bilateral acute intraventricular haematomata).

DISCUSSION

A cursory look at some of the pertinent radiological features of selected neuro-radiological images in traumatic brain injuries may be relevant to underscore the need to avert neurological sequelae. Top on our list is Mount Fuji sign of tension pneumocephalus. Pneumocephalus denotes the availability of air inside the cranial cavity as a consequence of pathological communication with - ambient air⁷⁻⁹. Up to 75% of cases of pneumocephalus are caused by trauma⁷⁻⁸. Other known causes are craniotomies, infections, tumors, brain and spine surgeries as well as surgeries involving paranasal sinuses. In terms of localization, pneumocephalus can be in the epidural, subarachnoid, intra-ventricular, intra-cerebral, or subdural spaces. The most frequent localization is subdural and frontal⁹⁻¹⁷. This is seen in our index patient shown in figure 1.

Tension pneumocephalus is regarded as neurosurgical emergency. It occurs when extra-axial air is in sufficient volume to exert a pressure effect on the brain, with attendant intracranial hypertension and brain herniation.^{7,9,11-16} The consequent clinical presentations manifest as generalized weakness, headache, vomiting and seizures. This therefore warrants immediate intervention in the form of cranial decompression using burr hole or needle aspiration to avert fatality.

As little as 25ml of air is sufficient to cause tension pneumocephalus⁸⁻⁹. CT is the golden standard for diagnosis of pneumocephalus since it can detect as small as 0.55mL of air whereas a conventional skull radiograph requires at least 2ml⁸. Two specific signs have been proposed for the diagnosis of subdural pneumocephalus on brain CT namely (1) Peaking sign (2) Mount Fuji sign⁹⁻¹². Peaking sign is characterized by large collection of air over the anterior and lateral portions of bilateral frontal lobes causing its compression. The air does not cause any separation of the tips of the frontal lobe, thereby leading to an appearance of a single peak in the midline^{9,12,17}. The "Mount Fuji sign," differentiates tension pneumocephalus from non-tension pneumocephalus^{7,10-12}. Characteristically, this sign involves extension of air in between the tips of the frontal lobes suggesting that the pressure of the subdural air exceeds the surface tension of cerebrospinal fluid between the frontal lobes^{8,16}. (see figure 1). This sign was first described by Ishiwata *et al.*^{1,2} and named after the highest mountain in Japan.^{8,12,13,16} The Mount Fuji sign indicates more severe pneumocephalus than the peaking sign and calls for need of emergent decompression⁸⁻¹⁰. Nevertheless, its absence does not essentially rule out subdural tension pneumocephalus⁹.

The second lesion in our presentation is blow-out orbit (blow-up orbit or orbital blow-out fracture). This occurs when there is a fracture of one of the orbital walls with intact orbital rim and any bone fragment must be displaced outside the orbit¹⁸⁻²². In children, it is called trap door fracture as the fracture may spring back into place^{18,21}. Blow out orbit results from direct blow to the central orbit from a fist or ball (an object with a diameter exceeding the bony margins of the orbit)^{18,21}. This could result from assault, motor vehicle accident, barotrauma and violent sports like soccer, boxing, wrestling, basketball and rugby. The fracture site of predilection is inferior orbital wall that is posterior and medial to the infra-orbital groove²¹. Other sites in descending order are medial orbital wall (lamina papyracea), superior and lateral walls¹⁸. Inferior and medial walls are affected simultaneously in 50% of cases²¹. In inferior blow-out fractures, orbital fat alone or in combination with inferior rectus

muscle will prolapse into the maxillary sinus. Consequences of this prolapse will be enophthalmos (due to increased orbital volume), diplopia: (due to extra-ocular muscle entrapment), orbital emphysema and malar region numbness (infraorbital nerve anaesthesia).^{18,21,22} Other presentations are exophthalmos, up-gaze restriction of the eye, periocular ecchymosis and painful eye movement, especially with adduction and infraduction^{18,23}.

CT is the best modality for evaluating blow out orbit. CT features of blow out orbit include: herniation of orbital fat inferiorly into the maxillary sinus giving a "tear drop" sign.^{21,22,24} (see figure 2). Other CT features are presence of intra-orbital bleed resulting in stretching or compression of the optic nerve, globe injury or rupture. Extraocular muscle entrapment is suspected if there is an acute change in angle of the muscle²¹. Surgical indications are herniation of the orbital contents into the maxillary sinus, muscle entrapment, enophthalmos of greater than 2 mm, significant diplopia in the primary position or oculo-cardiac reflex²³.

Subdural hematomas are relatively common, seen in about 10-20% of head injured patients and has a mortality as high as 50-85%²⁵. It is mainly the consequences of inertial forces arising from acceleration and deceleration of the brain within the skull with resultant rupturing of veins via a shearing mechanism²⁵. Subdural hematomas are not restricted by dural tethering at the cranial sutures but do not cross the midline because of the meningeal reflections²⁵⁻²⁷. Subdural haematomas can be bilateral. Such bilateral haematomas may cause medial compression of both ventricles leading to narrow, slit-like elongated ventricles. This corresponds to "squeezed ventricle" or "rabbit's ears" appearance as shown in figure 3²⁵.

Epidural haematoma (EDH) in contrast, is the result of direct impact to the skull with greater than 75% of cases resulting from disruption of the middle meningeal artery²⁶. It is seen in 0.2% - 12% of all head trauma patients with fracture detected in more than 90% of cases²⁶⁻²⁸. It is particularly worrisome when epidural haematoma is located in the middle

cranial fossa as seen in our patient (figures 6). This is because it has a potential for rapid enlargement with subsequent temporal lobe herniation (uncal herniation), brainstem compression and poor outcome²⁶. CT poor prognostic findings in such patients are pterional location of a skull fracture, EDH thickness greater than 1.5 cm, EDH volume greater than 30 ml, location within the lateral aspect of the middle cranial fossa, associated midline shift greater than 5 mm, and presence of a swirl sign within the hematoma (suggesting active bleeding and expansion of the collection)²⁶. However, the anterior location of EDH in our patient (figures 6) rather than laterally to the temporal lobe will predispose to posterior as opposed to medial displacement of the temporal lobe

Traumatic basal ganglion hematoma (TBGH) is defined as intra-cerebral hematoma located in basal ganglion (caudate nucleus, putamen and globus pallidus) (see figure 7) and sometimes in neighboring structures like thalamus and internal capsule²⁸⁻³². This contrasts with spontaneous hemorrhages that are solitary and located in the region of thalamus and internal capsule²⁹. TBGH is thought to be due to shearing of lenticulostriate or anterior choroidal blood vessels caused by rapid acceleration and deceleration forces²⁹. TBGH occurs mainly in young patients and it is a rare entity having been reported in only 3% of closed head injuries²⁹. However, a higher incidence of 10%-12% is seen in autopsy series^{29,30}.

TBGH is an evidence of significant primary brain injury and carries worse morbidity. Poor prognosis is seen when it is bilateral, large (>2cm in diameter) or accompanied by diffuse axonal injury, intra-ventricular hemorrhages, contusions, and extra axial hematomas²⁹⁻³³. It rarely needs surgical evacuation but surgery is considered in increasing size of hematoma and deteriorating neurological condition²⁰. Intra-ventricular hemorrhages may be the complications of parenchymal intracerebral bleed or subarachnoid bleed³⁴. However, spontaneous primary intra-ventricular hemorrhage without a recognizable parenchymal component in unenhanced CT or MRI studies is extremely rare as seen in our patient (figure 8)³⁴. Although primary intra-ventricular hemorrhage is

uncommon, it is a serious clinical condition with a high early mortality³⁴.

CONCLUSION

Mount Fuji sign is typical of tension pneumocephalus while herniation of extra-ocular muscles into the maxillary sinus giving a tear drop sign is diagnostic of blow-out orbit. Rabbit ear appearance is observable in bilateral subacute subdural haematoma. Haematoma in the middle cranial fossa may provoke uncal herniation. While traumatic basal ganglia haematoma and primary acute intra-ventricular haematoma may separately be an evidence of significant primary brain injury. These heterogenous subtypes of traumatic brain injuries were selected to spot light their salient neuro-radiological features to facilitate prompt Radiological diagnosis and subsequent clinical management.

DISCLOSURE: The authors have no conflict of interests to declare in relation to this paper.

CONSENT: The consent of the patients were sought for usage of their neuro-radiological images for this study.

AUTHORS CONTRIBUTIONS: UFU conceived the study. UFU, IUD, EDU, EJKC and AS contributed images and searched the literatures. UFU, IUD contributed to text writing and all authors proof read text.

REFERENCES

1. Lee B, Newberg A. Neuroimaging in Traumatic Brain Imaging. *NeuroRx*. 2005;2:372-383. doi: 10.1602/neurorx.2.2.372
2. Langlois JA, Rutland-Brown W, Wald, W. The epidemiology and impact of traumatic brain injury: A brief overview. *Journal of head trauma rehabilitation*. 2006;21:375-8
3. Obajimi MO, Jumah KB, Brakohuapa WO, Iddrisu W. CT features of head injury in Ghanaian children. *The Nigerian Journal of Surgical Research* 2002,(4)84-88.

4. Emejuru JKC, Uduma FU, Ulasi BI. Computed tomographic scan findings in head injury from a tertiary health facility in South east Nigeria. *J. Spine Neurosurg.* 2015(4)doi: <http://dx.doi.org/10.4172/2325-9701.1000196>
5. Uduma F U, Motah M. An accounting of pathology found on head computed tomography of road traffic accident(Rta) patients in Douala, Cameroon. *Global journal of health science.* 2011;3:171-180 DOI:<http://dx.doi.org/10.5539/gjhs.v3n1p171>
6. Newberg AB, Alavi A Neuroimaging in patients with head injury. *Semin Nucl Med.* 2003;33:136-147.
7. Sanyal P, Bele K, Vishwanath Rai SP. Tension pneumocephalus: Mount Fuji sign. *Arch Med Health Sci* [serial online] 2015;3:352-3. Available from: <http://www.amhsjournal.org/text.asp?2015/3/2/352/171948>
DOI: 10.4103/2321-4848.171948
8. Dabdoub CB, Salas G, N. Silveira Ed, Dabdoub CF. Review of the management of pneumocephalus. *Surg Neurol Int.* 2015;6:155
9. Jackhere SG, Yadav DA, Jain DG, Balasubramaniam. Does the Mount Fuji always signify tension pneumocephalus? An exception and Reappraisal. *Europeanian Journal of Radiology Extra* 2011;78:e5-e7
10. Amit Thapa MS, Deepak Agrawal M Ch Mount Fuji Sign in tension pneumocephalus *Indian Journal of Neurotrauma (IJN16T1)* 2009; 6:161-2
11. Michel SJ. The Mount Fuji sign. *Radiology* 2004;232:449-50
12. Vanhoenacker FM, Herz R, Vandervliet EJ, Parizel PM. The Mount Fuji sign in tension pneumocephalus. *JBR-BTR* 2008; 91:175.
13. Por YC, Barcelo CR, Salyer KE. Extradural tension pneumocephalus after posterior cranial vault remodeling for ventriculoperitoneal shunt-induced craniosynostosis. *J Craniofac Surg.* 2005; 16:504-8.
14. Rao V, Fredrisko O, Gulati S. Post-traumatic epidural tension pneumocephalus: a case report. *Journal of Medical case reports.* 2015;9:151 DOI: 10.1186/s13256-015-0633-5
15. Hong W, You C, Park C, Lee S. Two cases of delayed tension pneumocephalus. *J Korean Neurosurg. Soc.* 2005;37:59-62
16. Mark Hudon, Richard Farb, Jasnon Beiko, Patrick Macdonald. Tension Pneumocephalus- the Mount Fuji sign. *Canadian Journal Neurological Sciences* 2005;32: 538-9
17. Heckman JG, Ganslandt O. The Mount Fuji sign. *N. Engl J. Med.* 2004;350:1881
18. Iwona Rzymska-Grala, Piotr Palczewski, Marcin B³aż, Micha³ Zmorzyński, Marek Go³ebiowski, Hubert Wanyura. A peculiar blow-out fracture of the inferior orbital wall complicated by extensive subcutaneous emphysema: A case report and review of the literature. *Pol J Radiol.* 2012;77:64-68.
19. Suzuki H, Furukawa M, Takahashi E, et al. Barotraumatic blowout fracture of the orbit. *Auris Nasus Larynx.* 2001;28:257-59.
20. Lee HJ, Jilani M, Frohman L, et al. CT of orbital trauma. *Emerg Radiol.* 2004;10:168-72.
21. Gaillard F. Orbital blow-out fracture. radiopaedia.org/articles/orbital-blow-out-fracture.
22. Jones AL, Jones KE. Orbital Roof“Blow-in” Fracture: A Case Report and Review. *J Radiol Case Rep.* 2009;3:25-30. doi: 10.3941/jrcr.v3i12.363
23. Mathew R, Al Shammery YK. Surgical Outcome of Blowout Fractures of Floor of Orbit: A Case Series of 5 Patients. *J Clin Exp Ophthalmol.* 2016, 7:518. doi:10.4172/2155-9570.1000518
24. Kim SH, Ahn KJ, Lee JM, Choi KH, Han SH. The usefulness of orbital lines in detecting blow-out fracture on plain radiography. *BJR* 2014; 73(876) DOI: <http://dx.doi.org/10.1259/bjr.73.876.11205669>
25. Wagner AL, Naul LG. Imaging in Subdural Hematoma.

- Emedicine.medscape.com/article/344482-overview/a2 Updated: Oct 08, 2015
26. Gean AD, Fischbein, NJ, Purcell, DD, Aiken, AH, Manley, G T, Stiver SI. Benign Anterior Temporal Epidural Hematoma: Indolent Lesion with a Characteristic CT Imaging Appearance after Blunt Head Trauma. *Neuroradiology*; 2010,257(1). DOI: <http://dx.doi.org/10.1148/radiol.10092075>
 27. Adhiyaman V, Asghar M, Ganeshram KN, BHowmick. Chronic subdural haematoma in the elderly. *Postgrad Med J* 2002;78:71-75
doi:10.1136/pmj.78.916.71
 28. Yanagawa Y, Sakamoto T, Okada Y. Clinical features of temporal tip epidural hematomas. *J Neurosurg* 2007;107:18-20.
 29. Kumar S, Jha D, Abbey P, Mishra V, Handa A. *Outcome Of Traumatic Basal Ganglia Hemorrhage*. The Internet Journal of Neurosurgery. 2008;(6)1.
 30. Pandey N, Mahaparta A, Singh PK. Bilateral large traumatic hemorrhage of the basal ganglion. *Asian J Neurosurg*. 2014;9:240.
doi: 10.4103/1793-5482.146644
 31. Boto GR, Lobeto RD, Rivas J. Basal ganglion hematoma in severely head injured patients: clinicoradiological analysis of 37 cases. *J Neurosurgery* 2001;94:224-232.
 32. Jang KJ, Jwa CS. Bilateral traumatic hemorrhage of the basal ganglia. *J Korean Neurosurg Soc*. 2007;41:4.272.
 33. Kaushal R, Kataria R. Traumatic bilateral (mirror image) basal ganglia bleed. *Indian J Neurotrauma (IJNT)* 2011;8:109-10.
 34. Arboix A, Eroles G-E, Vicens AA, Oliveres M, Massons J. Spontaneous Primary Intraventricular Hemorrhage: Clinical Features and Early Outcome. *Neurology*. 2012 (2012), Article ID 498303, 7 pages
[Http://dx.doi.org/10.5402/2012/498303](http://dx.doi.org/10.5402/2012/498303)